

AD-A101 978

NAVAL RESEARCH LAB WASHINGTON DC

F/6 4/1

NONLINEAR STABILIZATION OF THE FARLEY-BUNEMAN INSTABILITY BY ST--ETC(U)

JUL 81 M J KESKINEN

UNCLASSIFIED

NRL-MR-4564

NL

1 - 1


20

2000000



END
DATE
FILMED
8 8
DTIC

AD A101



REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NRL Memorandum Report 4564	2. GOVT ACCESSION NO. AD-A101978	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) NONLINEAR STABILIZATION OF THE FARLEY- BUNEMAN INSTABILITY BY STRONG E X B TURBULENCE*	5. TYPE OF REPORT & PERIOD COVERED Interim report on a continuing NRL problem.	
	6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(s) M. J. Keskinen	8. CONTRACT OR GRANT NUMBER(s)	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Research Laboratory Washington, DC 20375	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61153N; RR033-02-44; 47-0883-0-1	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217	12. REPORT DATE July 31, 1981	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. NUMBER OF PAGES 13	
	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.	15a. DECLASSIFICATION DOWNGRADING SCHEDULE	
	17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)	
18. SUPPLEMENTARY NOTES *This work was completed while the author was an invited guest at the Centre de Physique Theorique, Ecole Polytechnique, Palaiseau, France.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Strong turbulence Equatorial electrojet Nonlinear saturation Farely-Buneman instability		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) It is shown that through nonlinear mode coupling processes long wavelength low frequency strong E X B turbulence can stabilize short wavelength high frequency Farley- Buneman modes in a weakly ionized low pressure convecting plasma. Favorable comparisons are made with experimental observations.		

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

NONLINEAR STABILIZATION OF THE FARLEY-BUNEMAN INSTABILITY BY STRONG $E \times B$ TURBULENCE

It is well known that in the absence of a magnetic field the two stream instability can occur in a homogeneous plasma when the electron drift velocity with respect to the ions exceeds the electron thermal velocity.¹ Farley² and Buneman² have shown that, in the presence of a magnetic field, the electron drift velocity with respect to the ions has only to exceed the ion acoustic velocity C_s to generate unstable waves traveling perpendicular to the magnetic field. We will consider the nonlinear evolution of the Farley-Buneman instability in a low β , weakly ionized, convecting plasma which is subjected to a magnetic field $B\hat{x}$, an electric field $E_0\hat{z}$ and a density gradient $(\partial n_0/\partial z)\hat{z}$. Differences in the collision frequencies ($\nu_i/\Omega_i \gg 1$, $\nu_e/\Omega_e \ll 1$) of the ions and electrons with the background neutral gas results in the formation of a cross field current $-J_0\hat{y}$ from the $\underline{v}_d = \underline{E} \times \underline{B}/B^2$ electron drift. For weak currents J_0 long wavelength field aligned fluctuations in density $\delta n \propto \exp[i(ky - \omega t)]$ have been found by Simon³ and Hoh³ to be linearly unstable when $\underline{E}_0 \cdot \nabla n_0 > 0$. In the nonlinear regime this $\underline{E} \times \underline{B}$ gradient drift instability evolves into an isotropic two-dimensional strongly turbulent state in the plane perpendicular to the magnetic field as shown previously.⁴ For stronger currents such that $v_d > C_s$ the Farley-Buneman instability will develop at shorter wavelengths. These long and short wavelength modes can coexist simultaneously (see Fig. 1) with the former usually occurring before the latter. Previous studies of the nonlinear evolution and saturation of these short wavelength Farley-Buneman modes have invoked quasilinear effects,⁵ resonance broadening,⁶ and mode coupling.⁷ These works have neglected the effects of the strong large scale background $\underline{E} \times \underline{B}$ turbulence. In this Letter we show that the long wavelength $\underline{E} \times \underline{B}$ turbulence can stabilize the short wavelength high frequency Farley-Buneman instability. Although the following discussion is

Manuscript submitted May 12, 1981.

applicable to any weakly ionized low β current carrying plasma convecting in regions of $\underline{E} \times \underline{B}$ turbulence it has direct bearing on density irregularities in the equatorial electrojet ionospheric plasma.

The basic equations for the electron (N_e) and ion (N_i) fluids in a low β , weakly ionized, collisional plasma can be written

$$\partial N / \partial t + \nabla \cdot N \underline{v}_e = 0 \quad (1)$$

$$eN(\underline{E} + \underline{v}_e \times \underline{B}) + T \nabla N + N m_e \nabla \cdot \underline{v}_e = 0 \quad (2)$$

$$\partial N / \partial t + \nabla \cdot N \underline{v}_i = 0 \quad (3)$$

$$(\partial / \partial t + \underline{v}_i \cdot \nabla) N m_i \underline{v}_i = eN(\underline{E} + \underline{v}_i \times \underline{B}) - T \nabla N - N m_i \nabla \cdot \underline{v}_i \quad (4)$$

$$\nabla \cdot (\underline{J}_i + \underline{J}_e) = 0 \quad (5)$$

where we have assumed quasineutrality ($N_e \approx N_i \approx N$), isothermality ($T_e \approx T_i$), electrostatic fluctuations ($\underline{E} = -\nabla \phi$), and neglected electron inertia.

Linearizing equations (1)-(5) with $N = n_0 + n$, etc. and assuming fluctuations of the form $n, \phi, \underline{v}_e, \underline{v}_i \propto \exp\{i[k_y y + k_z z - (\omega_{kr} + i\gamma(k)t)]\}$ with $\underline{k} \cdot \underline{B} = 0$ and $k_y L \gg 1$ we find⁸ for the frequency and growth rate in the ion frame

$$\omega_{kr} = \underline{k} \cdot \underline{v}_d / (1 + \psi) \quad (6)$$

$$\gamma(k) = [\psi / (1 + \psi)] \{ (\Omega_e / \nu_e) (v_d / L) \cos^2 \theta + \omega_{kr}^2 / \nu_i - k^2 c_s^2 / \nu_i \} \quad (7)$$

where $\psi = \nu_e \nu_i / \Omega_e \Omega_i$, $L^{-1} = (1/n_0) (\partial n_0 / \partial z)$, $c_s^2 = 2kT/m_i$ and θ is the angle defined by \underline{k} and \underline{v}_d . From the expression for the growth rate $\gamma(k)$ in

eq. (7), we note that at low frequencies (long wavelengths and weak currents) such that $\omega_k/v_i < (\Omega_e/v_e)(1/kL)$ the $\underline{E} \times \underline{B}$ gradient drift term will dominate with all modes with $k < k_c = [(\Omega_e/v_e)(V_d/L)(v_i/c_s^2)]^{1/2} \cos\theta$ unstable. At higher frequencies (short wavelengths and strong currents) Farley-Buneman modes will become unstable if $V_d > c_s$ but with no critical wavelength. In the equatorial electrojet plasma $V_d(t)$ is time dependent varying from $V_d < c_s$ to $V_d > c_s$ over a time interval $\Delta t \gg \gamma(k)^{-1}$. As a result the Farley-Buneman instability will be excited in strong $\underline{E} \times \underline{B}$ turbulence. Since these waves are nondispersive they will interact strongly. However as their amplitude increases, they do not steepen appreciably⁴ but are unstable to perturbations perpendicular to their propagation. It must be noted that this fluid approximation is valid for $\omega \ll v_i$. At higher frequencies kinetic effects, e.g., ion Landau damping, will become important and introduce a high frequency cutoff. For lower frequencies both fluid and kinetic treatments are identical.⁹

By writing $N(\underline{x}, t) = n_0 + \sum_{\underline{k}\omega} n(\underline{k}, \omega) \exp[-i(\underline{k} \cdot \underline{x} - \omega t)]$, $\phi(\underline{x}, t) = \varphi_0 + \sum_{\underline{k}\omega} \varphi(\underline{k}, \omega) \exp[-i(\underline{k} \cdot \underline{x} - \omega t)]$, etc., expanding eqs. (1)-(5) in the small parameter $v_e/v_i \ll \Omega_e/v_e \approx \epsilon \ll 1$, and considering high frequencies ω_k such that $\omega_k/v_i > (\Omega_e/v_e)(1/kL) \approx 10^{-2}$ (for $\lambda \approx 3m$, $L \approx 6 km$, $\Omega_e/v_e = 10^2$) we find to second order in $n_{\underline{k}\omega}$ ($n/n_0 \ll 1$)

$$D(\underline{k}, \omega) n(\underline{k}, \omega) = \int d^2k' d\omega' V(\underline{k}, \underline{k}', \omega, \omega') n(\underline{k}', \omega') n(\underline{k} - \underline{k}', \omega - \omega') \quad (8)$$

where

$$D(\underline{k}, \omega) = \omega - \underline{k} \cdot \underline{V}_d (1 + \psi)^{-1} - i\psi\omega^2 (1 + \psi)^{-1} v_i^{-1} + i\psi k^2 c_s^2 / v_i$$

is the Farley-Buneman dielectric and

$$V(\underline{k}, \underline{k}', \omega, \omega') = -(\hat{x} \times \underline{k}' \cdot \underline{k} / k'^2) [(1 + \psi)^{-1} (v_i/v_e) \underline{k}' \cdot \underline{V}_d + \underline{k}' \cdot \hat{x} \times \underline{V}_d + ik'^2 c_s^2 / v_i]$$

By neglecting the nonlinear term on the right hand side of eq. (8) we recover the linear result from $D(\underline{k}, \omega_{kr} + i\gamma(\underline{k})) = 0$ giving $\omega_{kr} = \underline{k} \cdot \underline{v}_d / (1 + \psi)$ and $\gamma(\underline{k}) = (\psi / (1 + \psi)) (1/v_i) [(\underline{k} \cdot \underline{v}_d)^2 - k^2 c_s^2]$.

We solve eq. (8) for the high frequency short wavelength component of $n(\underline{k}, \omega)$ by considering its mode coupling to the low frequency long wavelength well developed strong $\underline{E} \times \underline{B}$ turbulence. Let a Farley-Buneman wave be denoted by $(\underline{k}_I, \omega_I)$ and a turbulent $\underline{E} \times \underline{B}$ mode by $(\underline{k}_{II}, \omega_{II})$. Physically, when a Farley-Buneman mode $(\underline{k}_I, \omega_I)$ grows to such a level that it can couple with the $\underline{E} \times \underline{B}$ turbulence $(\underline{k}_{II}, \omega_{II})$, a beat wave component $(\underline{k}_I \pm \underline{k}_{II}, \omega_I \pm \omega_{II})$ will appear which in turn can beat with $(\underline{k}_{II}, \omega_{II})$ to affect $(\underline{k}_I, \omega_I)$. The evolution of $n(\underline{k}_I, \omega_I)$ can then be written

$$D^I(\underline{k}_I, \omega_I) n^I(\underline{k}_I, \omega_I) = \int d^2 \underline{k}' d\omega' V(\underline{k}_I, \underline{k}', \omega_I, \omega') n^{II}(\underline{k}', \omega') n^{I-II}(\underline{k}_I - \underline{k}', \omega_I - \omega') \quad (9)$$

The beat wave $(\underline{k}_I - \underline{k}_{II}, \omega_I - \omega_{II})$ evolves according to:

$$D^{I-II}(\underline{k}_I - \underline{k}, \omega_I - \omega) n^{I-II}(\underline{k}_I - \underline{k}, \omega_I - \omega) = \int d^2 \underline{k}' d\omega' V(\underline{k}_I - \underline{k}, \underline{k}', \omega_I - \omega, \omega') n^{II}(-\underline{k}', -\omega') n^I(\underline{k}_I, \omega_I) \quad (10)$$

where we have used $V(\underline{k}, \underline{k}', \omega, \omega') = V(\underline{k}, -\underline{k}', \omega, -\omega')$. Substituting eq. (10) into eq. (9) we find the nonlinear dispersion relation to lowest order

$$\tilde{D}^I(\underline{k}_I, \omega_I) n^I(\underline{k}_I, \omega_I) = 0 \quad (11)$$

where $\tilde{D}^I(\underline{k}_I, \omega_I) = D^I(\underline{k}_I, \omega_I) + \delta D^I(\underline{k}_I, \omega_I)$. The nonlinear part of eq. (11) can be written

$$\delta D^I(\underline{k}_I, \omega_I) = - \int d^2 \underline{k}' d\omega' \frac{V(\underline{k}_I, \underline{k}', \omega_I, \omega) V(\underline{k}_I - \underline{k}', \underline{k}', \omega_I - \omega', \omega')}{D^{I-II}(\underline{k}_I - \underline{k}', \omega_I - \omega')} n^{II}(\underline{k}', \omega') \quad (11a)$$

where $I^{II}(\underline{k}, \omega) = \langle n^{II}(\underline{k}, \omega) n^{II}(-\underline{k}, -\omega) / n_0^2 \rangle$ is the power spectrum of the $\underline{E} \times \underline{B}$ turbulence. It has been previously shown¹⁰ that $I^{II}(\underline{k}, \omega)$ can be calculated using the direct-interaction approximation of Kraichnan¹¹

$$|\omega - \omega(\underline{k}) + \Gamma^{II}(\underline{k}, \omega)|^2 I^{II}(\underline{k}, \omega) = \frac{1}{2} \int d^2 \underline{k}' d\omega' |w(\underline{k}, \underline{k}')|^2 I^{II}(\underline{k}', \omega') I^{II}(\underline{k} - \underline{k}', \omega - \omega') \quad (12)$$

$$\Gamma^{II}(\underline{k}, \omega) = - \int d^2 \underline{k}' d\omega' \frac{w(\underline{k}, \underline{k} - \underline{k}') w(\underline{k} - \underline{k}', \underline{k}) I^{II}(\underline{k}', \omega')}{\omega - \omega' - \omega(\underline{k} - \underline{k}') + \Gamma^{II}(\underline{k} - \underline{k}', \omega - \omega')} \quad (13)$$

with $w(\underline{k}, \underline{k}') = V(\underline{k}, \underline{k}') + V(\underline{k}, \underline{k} - \underline{k}')$ and $\omega(\underline{k}) = \omega_{kr} + i\gamma(\underline{k})$ where $\gamma(\underline{k}) = (\psi/1 + \psi) [(\Omega_e/\nu_e)(V_d/L)\cos^2\theta - \underline{k}^2 c_s^2/\nu_i]$. In eq. (12) $\Gamma^{II}(\underline{k}, \omega)$ is the self-damping of the long wavelength $\underline{E} \times \underline{B}$ fluctuations $(\underline{k}_{II}, \omega_{II})$. We now proceed to solve equations (11)-(13).

First, we note that since the interacting waves considered here are non-dispersive (see eq. (6)) the beat wave dielectric $D^{I-II}(\underline{k}_I - \underline{k}_{II}, \omega_I - \omega_{II}) \approx 0$ and the right hand side of eq. (11a) diverges. As a result we replace $D^{I-II}(\underline{k}_I - \underline{k}', \omega_I - \omega')$ by its renormalized value $D^{I-II}(\underline{k}_I - \underline{k}', \omega_I - \omega') + \delta D^{I-II}(\underline{k}_I - \underline{k}', \omega_I - \omega') \approx \delta D^{I-II}(\underline{k}_I - \underline{k}', \omega_I - \omega')$ in eq. (11a). In order to study the nonlinear saturation we solve eq. (11a) for $\text{Im } \delta D^I \equiv -i\Gamma(\underline{k}_I, \omega_I(\underline{k}_I))$. From previous studies^{4, 10} the steady state solution of eqs. (12)-(13) can be written

$$I^{II}(\underline{k}, \omega) = I^{II}(\underline{k}) (2\pi)^{-\frac{1}{2}} (\Gamma^{II}(\underline{k}))^{-1} \exp[-(\omega - \omega(\underline{k}))^2 / 2(\Gamma^{II}(\underline{k}))^2] \quad (14)$$

where $\omega(\underline{k}) = \underline{k} \cdot \underline{v}_d / (1 + \psi)$, $\Gamma^{II}(\underline{k}) = 3.4 n^{-\frac{1}{2}} (\nu_i / \Omega_i) k^2 V_d (I^{II}(\underline{k}))^{\frac{1}{2}}$, and $I^{II}(\underline{k}) = I k^{-n}$ is isotropic^{4, 12} with $n \approx 3-4$ while¹³ $I \propto V_d^m$, $m \approx 2$. Substituting eq. (14) into (11a) and assuming that $k' \approx k_{II} < k_I \equiv k$, $\omega' \approx \omega_{II} < \omega_I \equiv \omega$, $D^{I-II}(\underline{k}_I - \underline{k}', \omega_I - \omega') \approx D^I(\underline{k}_I, \omega_I) + O(k'/k_I) \approx \delta D^I(\underline{k}_I, \omega_I)$ we find

$$\begin{aligned}
[\Gamma(\underline{k})]^2 &= \int d^2k' V(\underline{k}, \underline{k}') V(\underline{k}-\underline{k}', \underline{k}') I^{II}(\underline{k}') \\
&\approx (v_i/\Omega_i)^2 \int d^2k' (\hat{x} \times \underline{k}' \cdot \underline{k})^2 (\underline{k}' \cdot \underline{v}_d)^2 |\underline{k}'|^{-4} I^{II}(\underline{k}') \\
&= (v_i/\Omega_i)^2 k^2 v_d^2 \int d\theta' k' dk' \sin^2(\theta-\theta') \cos^2\theta' I^{II}(\underline{k}') \quad (15)
\end{aligned}$$

where $\Gamma(\underline{k}) \gg \gamma(\underline{k})$ has been assumed and θ' and θ are the angles made by \underline{k}' and \underline{k} , respectively, with \underline{v}_d . In evaluating the quantity $V(\underline{k}, \underline{k}') V(\underline{k}-\underline{k}', \underline{k})$ in eq. (15) we have kept only terms proportional to $(v_i/\Omega_i)^2$ where $v_i/\Omega_i \gg 1$. Experimental studies¹⁴ as shown in Figure 2(a) indicate that $\Gamma(\underline{k})$ is approximately independent of angle θ . This allows the replacement of $\Gamma(\underline{k})$ in eq. (15) by its angle averaged result $(2\pi)^{-1} \int d\theta \Gamma(\underline{k}) = \Gamma(k)$ giving

$$\Gamma(k) = (v_i/\Omega_i) \left(\frac{1}{2}\right) k v_d < |n/n_o|^2_{II} >^{\frac{1}{2}} \quad (16)$$

where $< |n/n_o|^2_{II} > = \int d\theta k dk I^{II}(\underline{k}) = 2\pi \int dk k I^{II}(k)$. Using $v_i/\Omega_i \approx 22$, $2\pi/k \approx 5m$, $v_d = 4 \times 10^2 m/sec$, $c_s \approx 3.6 \times 10^2 m/sec$, and $< |n/n_o|^2_{II} >^{\frac{1}{2}} \approx 0.01$, eq. (16) gives for the spectral width $\Gamma(k)/2\pi \approx 10$ Hz (cf. Fig. 1). Figure 2(b) shows that the scaling of $\Gamma(k) \propto k$ from eq. (16) is in reasonable agreement with experimental results^{13,14} which indicate that the spectral broadening $\Delta\omega_k \propto \Gamma(k) \propto k^{0.7}$.

The nonlinear dispersion relation in eq. (11) for the short wavelength Farley-Buneman instability in the long wavelength $\underline{E} \times \underline{B}$ turbulent background can then be written explicitly

$$\omega - \underline{k} \cdot \underline{v}_d (1 + \psi)^{-1} - i\psi\omega^2 (1 + \psi)^{-1} v_i^{-1} + i\psi k^2 c_s^2 / v_i + i \frac{1}{2} (v_i/\Omega_i) k v_d < |n/n_o|^2_{II} >^{\frac{1}{2}} = 0. \quad (17)$$

Separating $\omega = \omega_{kr} + i\gamma(\underline{k})$ we find

$$\gamma(\underline{k}) = [\psi/(1 + \psi)] \left\{ (\underline{k} \cdot \underline{v}_d)^2 / v_i - k^2 c_s^2 / v_i \right\} - \frac{1}{2} (v_i/\Omega_i) k v_d < |n/n_o|^2_{II} >^{\frac{1}{2}} \quad (17a)$$

For complete stabilization of the fastest growing linear mode ($\theta \approx 0$) the long wavelength background fluctuation level

$$\langle |n/n_o|_{II}^2 \rangle^{1/2} \geq 2 v_e \Omega_e^{-1} (1 + \psi)^{-1} (kV_d/v_i - k^2 c_s^2 / v_i kV_d).$$

with $kV_d/v_i \approx 0.1$, $k^2 c_s^2 / v_i kV_d \approx 0.05$ we find $n/n_o \geq 0.002$ which is consistent with available experimental estimates.

Part of this work was completed at the Centre de Physique Théorique, Ecole Polytechnique, Palaiseau, France. We wish to thank the Centre National de la Recherche Scientifique for their invitation to visit the Ecole Polytechnique. Useful discussions with R. Pellat of the Ecole Polytechnique and M. Crochet and C. Hanuise of the Université' de Toulon et du Var are acknowledged.




Fig. 1 — Simultaneous equatorial electrojet ionospheric plasma density fluctuation power spectra (vertical axis $I(k, \omega)$ vs. frequency ω in Hz (horizontal axis) from Ref. 14 for several radar backscatter observation frequencies; the radar frequencies 29.0 MHz, 9.0 MHz, 6.6 MHz, 5.6 MHz correspond to wavelengths 5.1m, 10.5m, 16.6m, 22.7m, and 26.7m, respectively.




Fig. 2 - (a) Power spectrum width vs. wave vector angle made by \underline{k} and \underline{V}_d , the electrojet electron drift velocity, at several wavelengths from Ref. 14; note approximate isotropy in angle. (b) Power spectrum width vs. wave number k at $\theta = 15^\circ$ from Ref. 14; solid line is best fit $\propto k^{0.7}$.

REFERENCES

1. O. Buneman, Phys. Rev. 115, 503 (1959).
2. D.T. Farley, J. Geophys. Res. 68, 6083 (1963); O. Buneman, Phys. Rev. Lett., 10, 285 (1963).
3. A. Simon, Phys. Fluids 6, 382 (1963); F.C. Hoh, Phys. Fluids 6, 1184 (1963).
4. M.J. Keskinen, R.N. Sudan, and R.L. Ferch, J. Geophys. Res. 84, 1419 (1979); R.N. Sudan, J. Akinrimisi, and D.T. Farley, J. Geophys. Res. 78, 240 (1973); B.E. McDonald, T.P. Coffey, S.L. Ossakow, and R.N. Sudan, J. Geophys. Res. 79, 2557 (1974); R.L. Ferch and R.N. Sudan, J. Geophys. Res. 82, 2283 (1977).
5. T. Sato, Phys. Rev. Lett. 28, 732 (1972).
6. J. Weinstock and A. Sleeper, J. Geophys. Res. 77, 3721 (1972).
7. A. Rogister and E. Jamin, J. Geophys. Res. 80, 1820 (1975).
8. A. Rogister and N. D'Angelo, J. Geophys. Res. 75, 3879 (1970).
9. M.J. Schmidt and S.P. Gary, J. Geophys. Res. 78, 8261 (1973).
10. R.N. Sudan and M.J. Keskinen, Phys. Rev. Lett. 38, 966 (1977); R.N. Sudan and M.J. Keskinen, Phys. Fluids 22, 2305 (1979).
11. R.H. Kraichnan, J. Fluid Mech. 5, 497 (1959); B.B. Kadomtsev, Plasma Turbulence (Academic, New York, 1965), Chap. III.
12. H.M. Ierkić, B.G. Fejer, and D.T. Farley, Geophys. Res. Lett. 7, 497 (1980).
13. B.B. Balsley and D.T. Farley, J. Geophys. Res. 76, 8341 (1971); B.B. Balsley and D.T. Farley, J. Geophys. Res. 78, 7471 (1973); B.B. Balsley, J. Geophys. Res. 74, 2333 (1969).
14. C. Hanuise and M. Crochet, J. Geophys. Res. 86, 3567 (1981).

IONOSPHERIC MODELING DISTRIBUTION LIST

Please distribute one copy to each of the following people (unless otherwise noted).

Naval Research Laboratory
Washington, D.C. 20375

Dr. T. Coffey - Code 4700 (25 cys)
Jack D. Brown - Code 4701
Dr. S. Ossakow - Code 4780 (150 cys)
Dr. P. Mange - Code 4101
Dr. E. Szuszczewicz - Code 4127
Dr. J. Goodman - Code 7560

Science Applications, Inc.
1150 Prospect Plaza
La Jolla, CA 92037

Dr. D. A. Hamlin
Dr. L. Linson
Dr. D. Sachs

Director of Space and Environmental
Laboratory, NOAA
Boulder, CO 80302

Dr. A. Glenn Jean
Dr. G. W. Adams
Dr. D. N. Anderson
Dr. K. Davies
Dr. R. F. Donnelly

A. F. Geophysics Laboratory
L. G. Hansom Field
Bedford, MA 01730

Dr. T. Elkins
Dr. W. Swider
Mrs. R. Sagalyn
Dr. J. M. Forbes
Dr. T. J. Keneshea
Dr. J. Aarons

Office of Naval Research
800 North Quincy Street
Arlington, VA 22217

Dr. H. Mullaney

Commander
Naval Ocean Systems Center
San Diego, CA 92151

Mr. R. Rose - Code 5321

U.S. Army Aberdeen Research
and Development Center
Ballistic Research Laboratory
Aberdeen, MD

Dr. J. Heimerl

Commander
Naval Air Systems Command
Department of the Navy
Washington, D.C. 20360

Dr. T. Czuba

Harvard University
Harvard Square
Cambridge, MA 02138

Dr. M. B. McElroy
Dr. R. Lindzen

Pennsylvania State
University
University Park, PA 16802

Dr. J. S. Nisbet
Dr. P. R. Rohrbaugh
Dr. L. A. Carpenter
Dr. M. Lee
Dr. R. Divany
Dr. P. Bennett
Dr. F. Klevans

University of California,
Los Angeles
405 Hillgard Avenue
Los Angeles, CA 90024

Dr. F. V. Coroniti
Dr. C. Kennel

University of California,
Berkeley
Berkeley, CA 94720

Dr. M. Hudson

Utah State University
4th N. and 8th Streets
Logan, Utah 84322

Dr. P. M. Banks
Dr. R. Harris
Dr. K. Baker

Cornell University
Ithaca, NY 14850

Dr. W. E. Swartz
Dr. R. Sudan
Dr. D. Farley
Dr. M. Kelley

NASA
Goddard Space Flight Center
Greenbelt, MD 20771

Dr. S. Chandra
Dr. K. Maeda
Dr. R. F. Benson

Princeton University
Plasma Physics Laboratory
Princeton, NJ 08540

Dr. F. Perkins
Dr. E. Frieman

Institute for Defense Analysis
400 Army/Navy Drive
Arlington, VA 22202

Dr. E. Bauer

University of Maryland
College Park, MD 20740

Dr. K. Papadopoulos
Dr. E. Ott

University of Pittsburgh
Pittsburgh, PA 15213

Dr. N. Zabusky
Dr. M. Biondi

Defense Documentation Center
Cameron Station
Alexandria, VA 22314

(12 copies if open publica-
tion, otherwise 2 copies)

12CY Attn TC

University of California
Los Alamos Scientific Laboratory
J-10, MS-664
Los Alamos, NM 87545

M. Pongratz
D. Simons
G. Barasch
L. Duncan

Massachusetts Institute of
Technology
Plasma Fusion Center
Library, NW16-262
Cambridge, MA 02139

**DAT
FILM**