UNCLAS		18-9	قيهرهم	_	_	_	_			_	
 	07   1.344										
								END DATE FILMED 103-182			



REPOR OCUMENTATION PAGE	EAD INSTRUCTIONS
1. REPORT NUMBER 2. GOVT ACCESSION Technical Report No. 9	NO. 3. HEL. NT'S CATALOG NUMBER
HD-HII	344
4. TITLE (and Subtitle)	S. TYPE OF REPORT & PERIOD COVER
Synthesis of the Tube Silicate Litidionite and	Interim Technical Report
Other Silicates	6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(a)	8. CONTRACT OR GRANT NUMBER(*)
Jesse Hefter and Malcolm E. Kenney	N00014-75-C-0693
9. PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TAS
Department of Chemistry	
Clase Western Keserve University Cleveland, OH 44106	NR 356-590
11. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE
Department of the Navy	February 17, 1982
Office of Naval Research	13. NUMBER OF PAGES
ALLINGLON, VA 4441/ 14. MONITORING AGENCY NAME & ADDRESS(II different from Controlling Offic	<ul> <li>b) 15. SECURITY CLASS. (of this report)</li> </ul>
	linclassified
	UNC1439111EU
	154. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of the Report) Distribution Unlimited 17. DISTRIBUTION ST, 4ENT (of the abstract entered in Block 20, if differen	from Report)
16. DISTRIBUTION STATEMENT (of the Report) Distribution Unlimited 17. DISTRIBUTION ST, 4ENT (of the abstract entered in Block 20, if different	FEB 2 4 1982
16. DISTRIBUTION STATEMENT (of the Report) Distribution Unlimited 17. DISTRIBUTION ST, 4ENT (of the abstract entered in Block 20, if different 18. SUPPLEMENTARY 15 TES	FEB 2 4 1982
16. DISTRIBUTION STATEMENT (of the Report) Distribution Unlimited 17. DISTRIBUTION ST. 4ENT (of the abstract entered in Block 20, if differen 18. SUPPLEMENTARY (TES Accepted for Publication in <u>Inorganic Chemistr</u>	FEB 2 4 1982
<ul> <li>16. DISTRIBUTION STATEMENT (of the Report)</li> <li>Distribution Unlimited</li> <li>17. DISTRIBUTION ST. (of the abstract entered in Block 20, if differen</li> <li>18. SUPPLEMENTARY (TES</li> <li>Accepted for Publication in <u>Inorganic Chemistr</u></li> </ul>	FEB 2 4 1982
<ul> <li>16. DISTRIBUTION STATEMENT (of the Report)</li> <li>Distribution Unlimited</li> <li>17. DISTRIBUTION ST. 4ENT (of the abstract entered in Block 20, if differen</li> <li>18. SUPPLEMENTARY 12 TES</li> <li>Accepted for Publication in <u>Inorganic Chemistr</u></li> <li>19. KEY WORDS (Continue on reverse elde if necessary and identify by block num</li> </ul>	from Report) FEB 2 4 1982 A Y
<ul> <li>16. DISTRIBUTION STATEMENT (of the Report) Distribution Unlimited</li> <li>17. DISTRIBUTION ST. 4ENT (of the abstract entered in Block 20, if differen</li> <li>18. SUPPLEMENTARY (TES Accepted for Publication in <u>Inorganic Chemistr</u></li> <li>19. KEY WORDS (Continue on reverse side if necessary and identify by block num Litidionite, Silicates, Copper Silicate, Tube S Sheet Silicates, Latticework Tubes</li> </ul>	from Report) ELLE FEB 2 4 1982 A Y ber) ilicates, Ladder Silicates,
<ul> <li>16. DISTRIBUTION STATEMENT (of the Report) Distribution Unlimited</li> <li>17. DISTRIBUTION ST. (of the abstract entered in Block 20, if differen <ul> <li>18. SUPPLEMENTARY</li> <li>18. SUPPLEMENTARY</li> <li>19. KEY WORDS (Continue on reverse eide if necessary and identify by block num Litidionite, Silicates, Copper Silicate, Tube S Sheet Silicates, Latticework Tubes</li> </ul> </li> <li>20. ABSTRACT (Continue on reverse eide if necessary and identify by block num </li> </ul>	Trom Report) ELL FEB 2 4 1982 A Y ber) illicates, Ladder Silicates,
<ul> <li>16. DISTRIBUTION STATEMENT (of the Report)</li> <li>Distribution Unlimited</li> <li>17. DISTRIBUTION ST. (ENT (of the abstract entered in Block 20, if differen</li> <li>18. SUPPLEMENTARY (TES)</li> <li>Accepted for Publication in <u>Inorganic Chemistr</u></li> <li>19. KEY WORDS (Continue on reverse side if necessary and identify by block num</li> <li>Litidionite, Silicates, Copper Silicate, Tube S Sheet Silicates, Latticework Tubes</li> <li>20. ABSTRACT (Continue on reverse side if necessary and identify by block num</li> <li>Two good syntheses of the rare tube silicate reported. In one a 2:1:1:8 mole-ratio mixture SiO, is sintered and in the other a glass made in the same mole ratios is devitrified. The s silicate and those in other known tube silicate</li> </ul>	Trom Report) FEB 2 4 1982 FEB 2 4 1982 A Y ber) filicates, Ladder Silicates, and filicates, Ladder Silicates, and from a mixture of these reagen tructures of the ion in this s are discussed in terms of
<ul> <li>16. DISTRIBUTION STATEMENT (of the Report)</li> <li>Distribution Unlimited</li> <li>17. DISTRIBUTION ST. (ENT (of the abstract entered in Block 20, if differen</li> <li>18. SUPPLEMENTARY (TES)</li> <li>Accepted for Publication in <u>Inorganic Chemistr</u></li> <li>19. KEY WORDS (Continue on reverse side if necessary and identify by block num Litidionite, Silicates, Copper Silicate, Tube S Sheet Silicates, Latticework Tubes</li> <li>20. ABSTRACT (Continue on reverse side if necessary and identify by block num Two good syntheses of the rare tube silicat reported. In one a 2:1:1:8 mole-ratio mixture SiQ is sintered and in the other a glass made in the same mole ratios is devitrified. The s silicate and those in other known tube silicate their relationships to each other and to the io pointed out that many latticework tubes like the</li> </ul>	Trom Report) FEB 2 4 1982 FEB 2 4 1982 A Y ber) ilicates, Ladder Silicates, er) T 1/7 te litidionite, NaKCuSi 0,0 of CuO, Na CO3, K CO3, and from a mixture of these reagen tructures of the ion in this s are discussed in terms of ns in other silicates. It is page which are the parents of
<ul> <li>16. DISTRIBUTION STATEMENT (of the Report)</li> <li>Distribution Unlimited</li> <li>17. DISTRIBUTION ST. (ENT (of the abstract entered in Block 20, if differen</li> <li>18. SUPPLEMENTARY TES</li> <li>Accepted for Publication in <u>Inorganic Chemistr</u></li> <li>19. KEY WORDS (Continue on reverse elde if necessary and identify by block num Litidionite, Silicates, Copper Silicate, Tube S Sheet Silicates, Latticework Tubes</li> <li>20. ABSTRACT (Continue on reverse elde if necessary and identify by block num Two good syntheses of the rare tube silicat reported. In one a 2:1:1:8 mole-ratio mixture SiO, is sintered and in the other a glass made in the same mole ratios is devitrified. The s silicate and those in other known tube silicate their relationships to each other and to the io pointed out that many latticework tubes like th</li> </ul>	FEB 2 4 1982 FEB 2 4 1982 A Y ber) filicates, Ladder Silicates,

f

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

 $\boldsymbol{\boldsymbol{\mathcal{F}}}$ 

these tube silicate ions exist and it is concluded that silicate ions corresponding to some of them will be found.

≻

### SECURITY CLASSIFICATION OF THIS PAGE(When Date Entered)

in any start and the

1

THE FIL MON

OFFICE OF NAVAL RESEARCH Contract N00014-75-C-0693 Task No. NR 356-590 TECHNICAL REPORT No. 9

Synthesis of the Tube Silicate Litidionite and Structural Relationships between It and Some Other Silicates

Ъy

Jesse Hefter and Malcolm E. Kenney Accepted for Publication in <u>Inorganic Chemistry</u>

Case Western Reserve University Department of Chemistry Cleveland, Ohio 44106

February 17, 1982

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited. Synthesis of the Tube Silicate Litidionite<sup>1</sup> and Structural Relationships between It and Some Other Silicates

Jesse Hefter and Malcolm E. Kenney

1

Contribution from the Department of Chemistry Case Western Reserve University Cleveland, Ohio 44106

4

## Abstract

Two good syntheses of the rare tube silicate litidionite,  $NaKCuSi_4O_{10}$ , are reported. In one a 2:1:1:8 mole-ratio mixture of CuO,  $Na_2CO_3$ ,  $K_2CO_3$ , and  $SiO_2$  is sintered and in the other a glass made from a mixture of these reagents in the same mole ratios is devitrified. The structures of the ion in this silicate and those in other known tube silicates are discussed in terms of their relationships to each other and to the ions in other silicates. It is pointed out that many latticework tubes like those which are the parents of these tube silicate ions exist and it is concluded that silicate ions corresponding to some of them will be found.

Section and the

While the scroll silicate chrysotile,  $Mg_3Si_2O_5(OH)_4^{4,5}$  is a familiar item of commerce (it is one of the silicates commonly called asbestos) and has been extensively studied,<sup>6</sup> tube silicates and aluminosilicates are unfamiliar and have been investigated relatively little. They are interesting, however, because tube structures in which the tube bonds are primary bonds have received little attention.

The most studied of the tube silicates and aluminosilicates is the aluminosilicate imogolite,  $Al_2(OH)_3SiO_3OH$ .<sup>7</sup> It is known to be composed of neutral tubes (the details of the structure of these tubes are, however, not understood).<sup>8</sup> Another aluminosilicate which can be viewed as belonging to the tube group, since it has an aluminosilicate ion that can be considered to be tubular, is bavenite,  $Ca_4Be_2Al_2Si_9O_{26}(OH)_2$ .<sup>9-11</sup>

Three silicates which have tube ions are litidionite,  $NaKCuSi_4O_{10}$ ,<sup>13</sup> fenaksite,  $NaKFeSi_4O_{10}$ ,<sup>14</sup> and the synthetic species,  $Na_2CuSi_4O_{10}$ .<sup>15</sup> Others are agrellite,  $NaCa_2Si_4O_{10}F$ ,<sup>16</sup> narsarsukite,  $Na_2TiSi_4O_{11}$ ,<sup>17</sup> miserite,  $KCa_5Si_2O_7Si_6O_{15}(OH)F$ ,<sup>18</sup> and probably canasite,  $Na_4K_2Ca_5Si_{12}O_{30}(OH,F)_4$ .<sup>19</sup>

As implied all but  $Na_2CuSi_4O_{10}$  have been found naturally.<sup>20-25</sup> So far as is known fenaksite, agrellite, narsarsukite, miserite, and canasite are rare. Litidionite is apparently very rare, the only reported occurrence of it being in the crater of Mt. Vesuvius.

Both litidionite<sup>15,26</sup> and narsarsukite<sup>27</sup> have been made in the laboratory using hydrothermal techniques. Hydrothermal techniques have also been used to make  $Na_2CuSi_4O_{10}$ .<sup>28-31</sup>

Since compounds containing tube ions are uncommon, these silicates are of interest and good synthetic routes to them are of value. In this paper we report two good new syntheses for litidionite and point out the advantages of each.

-2-

In addition we discuss the structure of its silicate ion, Figure 1, and those of other known tube silicates in terms of their relationships to each other and to the ions in other silicates. We also point out the fact that many latticework tubes like those which are the parents of these tube silicate ions exist and conclude that silicate ions corresponding to some of them will be found.

## Experimental

Synthesis of NaKCuSi $_40_{10}$ . Sintering Method. A mixture of CuO(3.82g), Na $_2CO_3(2.54g)$ , K $_2CO_3(3.32g)$ , and SiO $_2(11.5g)$  (a 2:1:1:8 mole-ratio mixture) in a platinum crucible was heated in a muffle furnace to 725°C over  $\sim$ 1h, held at this temperature for 1d, heated to 765°C over  $\sim$ 1h, and held at this temperature for 6d. The product, a hard, porous, blue solid with black impurities dispersed throughout it (18.8g) was carefully broken out of the crucible (17.3g).

This material was crushed with a diamond mortar to a powder which was almost all <100 mesh. The very small fraction of it which was >100 mesh was rejected. A mixture of some (3.0g) of the remaining powder and distilled water buffered to a pH of 5 with potassium acid phthalate and sodium hydroxide ( $\sim$ 200mL) was refluxed with stirring for  $\sim$ 10 min and allowed to stand for  $\sim$ 4 min. The portion which settled out was isolated by decantation. This partially purified material was further purified by a succession of five like decantation steps. The resulting solid was recovered by filtration, washed with water and acetone, and dried (1.8g). The product, a blue powder in which was interspersed small amount of a dark powder, gave the expected powder X-ray diffraction lines<sup>13,26</sup> and three impurity lines. The use of water buffered to a pH of 4 with potassium acid phthalate in the purification procedure led to a material grossly contaminated with decomposition products.

Modified Sintering Method. A mixture of CuO(1.27g), Na<sub>2</sub>CO<sub>3</sub>(0.848g),  $K_2CO_3(1.106g)$ , SiO<sub>2</sub>(3.85g), and Na<sub>2</sub>B<sub>4</sub>O<sub>7</sub>·1OH<sub>2</sub>O(0.707g) (a 2:1:1:8 mole-ratio mixture with 10% by weight of borax) was prepared. This was placed in a platinum crucible and heated to 675°C over  $\sim$ 1h, held at this temperature for 1d, heated to 725°C over  $\sim$ 1h, held at this temperature for 1d, heated to 760°C over  $\sim$ 1h, and held at this temperature for 5d. The product, a hard, porous, blue solid with a shiny surface and black particles dispersed throughout (6.67g) was carefully broken out of the crucible (5.02g).

This solid was crushed to a powder that was virtually all <100 mesh. The few particles >100 mesh were rejected. A portion (2.00g) of the rest was purified using a decantation process like that described above (12 cycles). The product, a blue powder in which was interspersed a small amount of a dark powder, gave the expected powder X-ray diffraction lines and one low intensity impurity line (0.92g).

Because of the excess of sodium in the reaction mixture, this product could have contained  $Na_2CuSi_4O_{10}$ . However, the X-ray data did not give evidence for its presence.<sup>15,29,32</sup>

Devitrification Method. A 2:1:1:8 mole-ratio mixture of CuO,  $Na_2CO_3$ ,  $K_2CO_3$ , and SiO<sub>2</sub> was heated in a platinum crucible to  $\sim 1200^{\circ}C$ . Part of the resulting glass was poured on a steel marver and flattened into a disc (1.31g). This was placed on a bed of  $NaKCuSi_4O_{10}$  (6mg, <100 mesh) which was supported by a fused silica plate, and heated at 765°C for 12d. The product, a disc approximately the size of the original disc, was gently

-4-

pried from the silica plate (1.33g). It was composed mainly of a mass of blue needle-like crystals oriented approximately perpendicularly to its surfaces. Along with these were a small amount of a gray material, a small silica shard, much of the powdered  $NaKCuSi_4O_{10}$ , and some blue glass. The gray material was present as a thin layer on the top and sides of the disc while the blue glass was present between the crystals. A powder of a portion of the disc gave the expected diffraction lines and two impurity lines.

As is to be expected, the time required for substantial devitrification of the glass employed under the general conditions used is variable. In an experiment which was similar to that described except for the use of a thicker disc of glass, 14d was insufficient but 28d was sufficient.

Instrumentation. The X-ray data were collected with a General Electric XRD-5 spectrometer using cobalt radiation.

## Discussion

Synthetic Work. The two sintering routes described for the synthesis of litidionite are parallel to the two routes reported by Pabst for making the chemically and structurally related silicate cuprorivaite (Egyptian blue),  $CaCuSi_4O_{10}$ .<sup>33,34</sup> The devitrification route is probably parallel to the route by which natural litidionite is formed since natural litidionite occurs in association with a blue glass.<sup>35</sup>

Both the non-borax sintering route and the devitrification route provide satisfactory methods for making the compound. The borax sintering route is less satisfactory because it is experimentally more cumbersome and may give a product which is less pure.

-5-

Structural Considerations. As already indicated fenaksite, agrellite, Na<sub>2</sub>CuSi<sub>4</sub>0<sub>10</sub>, narsarsukite, miserite, and probably canasite contain, like lititionite, tube silicate ions.<sup>13-19</sup> The first three contain ions that are the same as that in litidionite while the fourth contains a different type of ion, Figure 2. Still another is found in miserite, Figure 3, and probably also in canasite.

Each of these three types of ions is structurally related to other silicate ions. Thus, as pointed out earlier,<sup>36</sup> the ion in narsarsukite is related to a well-known sheet ion found in, for example, kaolinite,  $Al_2Si_2O_5(OH)_4$ .<sup>37</sup>

The basic relationship between these two ions is the same as that between their parent latticeworks. This relationship arises as is evident, Figure 4, because the pattern of the latticework tube taken circumferentially from a lengthwise line on its surface is the same as that in a strip of the latticework sheet.<sup>38,39</sup>

Similarly but not quite so directly, the ion in litidionite is related to a less common ion found in such silicates as apophyllite,  $KCa_4Si_8O_{20}F\cdot 8H_2O$ ,<sup>40</sup> and cuprorivaite.<sup>33</sup> Here, again making reference to the parent latticeworks, the arrangement of the latticework tube is like that of a strip of the latticework sheet except that all the pendent units project from one surface, Figure 5. In the same manner the parent latticework of the ion in miserite is related to that of the ion in dalyite, (K, Na)<sub>2</sub>ZrSi<sub>6</sub>O<sub>15</sub>,<sup>41</sup> Figure 6.<sup>42</sup>

All three of these latticework tubes are similar in that each is achiral and has a shortest circumferential circuit in which the repeat pattern occurs twice. However, in the first tube the shortest circuit does not incorporate a sequence of in-line links (i.e., links whose counterparts in

-6-

the corresponding strip are in a straight line). In the second and third tubes, in contrast, the shortest circuits do incorporate unbroken in-line sequences of links. Similarly, in the first tube no links in the circuit are parallel to a plane perpendicular to the tube axis while in the second and third tubes links in the circuits are parallel to planes perpendicular to the tube axes.

Additional latticework tubes associated with the kaolinite, apophyllite, and dalyite latticeworks are shown in Figure 7-9 (a variant of the third tube is associated with the dalyite latticework when arranged so as to outline squares and regular octagons). Each of these particular latticework tubes, like the first three latticework tubes, is achiral. Further, like the second and third of these tubes, each has a shortest circuit which contains two repeat patterns, incorporates an unbroken in-line sequence of links, and includes links that are parallel to a plane perpendicular to the tube axis.

Two latticework tubes associated with the latticework sheet corresponding to the ion in yet another sheet silicate, manganpyrosmalite,  $(Mn, Fe)Si_6O_{15}$ - $(OH, Cl)_{10}$ , <sup>43</sup> are shown in Figures 10 and 11.<sup>44</sup> These tubes too are achiral and have shortest circuits of the same nature as those in all but the first tube.

Many additional simple latticework tubes are also associated with the kaolinite, apophyllite, dalyite, and manganpyrosmalite latticework sheets. An example is the tube which is like the larger of the two tubes associated with the manganpyrosmalite sheet already discussed except for having a short-est circuit with one repeat, Figure 12. Another is the tube which is like the litidionite latticework tube except for having a shortest circuit with three repeats.

-7-

Not surprisingly many of these latticework tubes are chiral. A tube of this type associated with the apophyllite latticework sheet is shown in Figure 13. This tube, as is seen, has a shortest circuit which contains one repeat pattern, does not incorporate an unbroken in-line sequence of links, and does not include links parallel to a plane perpendicular to the tube axis.<sup>45</sup>

Still more latticework tubes similar to those already discussed exist. These are associated with additional latticework sheets.<sup>47</sup> Such a tube is shown in Figure 14.

Since there are so many closely related latticework tubes a simple systematic method for describing them is needed. This need can be met by utilizing descriptors containing the following elements: a c if the tube is chiral, a number giving the number of links in the shortest circuit, a subscript number giving the number of repeats in the circuit, and one or more numbers giving the number or numbers of links in the shortest circuit or circuits in the surface of the tube. Thus using this system the litidionite tube is the  $6_2$ :4,8 tube.

In connection with the latticework tubes discussed, it is interesting to note that (using the system just outlined) the  $8_2$ :4,8 tube, Figure 8, and the  $10_1$ :4,6,12 tube, Figure 12, are isomers. It is also interesting to note that half the latticework strip associated with the  $12_2$ :4,6,12 tube, Figure 10, has a drum-ladder counterpart, Figure 15. Obviously this is just one of a family of such latticeworks. So far no silicate based on a latticework of this type nor of the analogous chain-cage type is known.

As might be expected the known and potential silicate tubes have significant relationships with not just silicate sheets but also with other types

-8--

of silicate structures as well. One of these is the linear or chainband type.

Among the silicates that contain chains or bands with which the eight known and potential silicate tubes discussed first have significant relationships are: diopside,  $MgCaSi_2O_6$ , <sup>48</sup> howieite,  $Na(Fe^{2+},Mg,Fe^{3+},Al)_{12}(Si_6O_{17})_2^{-}$  $(0,0H)_{10}$ , <sup>49</sup> aenigmatite,  $Na_2Fe_5TiSi_6O_{20}$ , <sup>50</sup> astrophyllite,  $(K,Na)_3(Fe,Mn)_7^{-}$  $Ti_2Si_8(0,0H)_{31}$ , <sup>51</sup> tremolite,  $Mg_5Ca_2Si_8O_{22}(0H,F)_2$ , <sup>52</sup> jimthompsonite,  $(Mg,Fe)_5Si_6O_{16}(0H)_2$ , <sup>53</sup> synthetic  $Ba_5Si_8O_{21}$ , <sup>54</sup> synthetic  $Ba_6Si_{10}O_{26}$ , <sup>54</sup> vlasovite,  $Na_2ZrSi_4O_{11}$ , <sup>55</sup> xonotlite,  $Ca_6Si_6O_{17}(0H)_2$ , <sup>56</sup> tinaksite,  $K_2NaCa_2TiSi_7O_{19}(0H)$ , <sup>57</sup> and nordite,  $Na_2(Na,Mn)(Ca,Sr)RE(Zn,Mg,Fe,Mn)Si_6O_{17}$ . <sup>58</sup> The nature of some of these relationships is indicated in Figures 16 and 17. Again the relationships are complicated in some cases, because of the positioning of the pendent oxygens. <sup>59,60,62</sup>

As is obvious, the known silicate tubes and the potential small to moderate sized silicate tubes have (assuming reasonable bond parameters for the latter) diameters covering a wide range. The lower end of this range is  $\sim 4$ Å. Figures 1-3.<sup>63</sup>

Using past experience in similar situations as a guide, it appears highly likely that silicate tubes of the narsarsukite, litidionite, and miserite types will be found in silicates other than those in which they are now known. Since the litidionite tube has been found in several silicates, and since the relationships between it and several linear silicates are strong, it is particularly likely that this tube will be found in other silicates.

It also appears likely that new types of silicate tubes will be found soon. Among the most likely to be found are those having the  $8_2:6$ ,  $8_2:4,8$ , and  $6_2:4,6,8$  latticework tubes as parents since these latticework tubes

-9-

are like those which are parents of the litidionite and miserite tubes. The probability of finding the silicate tube whose parent is the  $8_2$ :6 tube is further enhanced because of the close relationships that exist between it and known silicate bands.

As is evident latticework tubes of the types described cannot simultaneously have continuous single-link members running lengthwise and circumferentially. Tubes of this type are, however, found among those having four, five, and six members radiating from a junction, Figure 18.

Finally, it is probable that in some cases organic tubes having carbon or carbon-heteratom frameworks like those discussed can be made. Thus, since norpinane can be made,<sup>64</sup>it seems likely that tricyclo[3.1.1.1.<sup>2,4</sup>]octane can be made and, hence, that narsarsukite-type and perhaps litidionitetype tubes can be made, Figure 19.<sup>65</sup> Such polymeric systems should have very interesting properties.

Ł

## Acknowledgment

We wish to thank Dr. Kenneth Wynne and Dr. Kerro Knox for helpful discussions. We also gratefully acknowledge support of this work by the Dow-Corning Corporation, the B. F. Goodrich Corporation, and the Office of Naval Research.

-10-

### References

- 1. This name is sometimes alternatively given as lithidionite. The name litidionite is employed here because it has been used in the most recent edition of Fleischer's glossary,<sup>2</sup> and because it is being used by the Chemical Abstracts Service.<sup>3</sup>
- Fleischer, M. "Glossary of Mineral Species"; Mineralogical Record, Tucson, AZ, 1980.
- 3. Private communication, Dennis Cipollo, Chemical Abstracts Service.
- 4. Yada, K. Acta Crystallogr. 1967, 23, 704.
- 5. Natural silicates commonly have variable compositions and the formulas given for them are thus of necessity generally idealized.
- Zussman, J. In "Asbestos Properties, Applications, and Hazards", Michaels, L. and Chissick, S. S., Ed.; Wiley: New York, 1979; Vol. I, Chapter 2.
- 7. Yoshinaga, N.; Aomine, S. Soil Sci. Plant Nutr. 1962, 8, 6.
- 8. Cradwick, P. D. G.; Farmer, V. C.; Russell, J. D.; Masson, C. R.; Wada, K.; Yoshinaga, N. Nature (London) Phys. Sci. 1972, 240, 187.
- 9. Cannillo, E.; Coda, A.; Fagnani, G. Acta Crystallogr. 1966, 20, 301.
- 10. Belov, N. V. Mineral. Sb. (Lvov) 1969, 23, 115.
- 11. The ions in bavenite can alternatively be viewed as belonging to the band group.
- 12. Liebau, F. Am. Mineral. 1978, 63, 918.
- 13. Martin Pozas, J. M.; Rossi, G.; Tazzoli, V. Am. Mineral. 1975, 60, 471.
- Golovachev, V. P.; Drozdov, Yu. N.; Kuz'min, E. A.; Belov, N. V. Sov. Phys. Dokl. (Engl. Transl.) 1971, 15, 902; Dokl. Akad. Nauk SSSR 1970, 194, 818.
- 15. Kawamura, K.; Kawahara, A. Acta Crystallogr. Sect. B, 1977, 33, 1071.

-11-

- 16. Ghose, S.; Wan, C. Am. Mineral. 1979, 64, 563.
- Pyatenko, Yu. A.; Pudovkina, Z. V.; Sov. Phys. Crystallogr. (Eng. Transl.)
   1961, 5, 540; Kristallografiya 1960, 5, 563.
- 18. Scott, J. D. Can. Mineral. 1976, 14, 515.
- 19. Chiragov, M. I.; Mamedov, Kh. S.; Belov, N. V. Dokl. Acad. Sci. USSR, Earth Sci. Sect. (Engl. Transl.) 1969, 185, 96; Dokl. Acad. Nauk. SSSR 1969, 185, 672.
- 20. Scacchi, E. Rend. Accad. Sci. Fis. Mat., Naples 1880, 19, 175.
- Dorfman, M. D.; Rogachev, D. L.; Goroshchenko, Z. I.; Mokretsova, A. V.
   Tr. Mineral. Muz., Akad. Nauk. SSSR 1959, No. 9, 152.
- 22. Gittens, J.; Bown, M. G.; Sturman, D. Can. Mineral. 1976, 14, 120.
- 23. Flink, G. Medd. Groen1. 1901, 24, 154.
- 24. Shaller, W. T. Am. Mineral. 1950, 35, 911.
- Dorfman, M. D.; Rogachev, D. L.; Goroshchenko, Z. I.; Vspenskaya, E. I.
   Tr. Mineral, Muz., Akad. Nauk. SSSR 1959, No. 9, 158.
- 26. Guth, J.-L.; Kalt, A.; Perati, B.; Wey, R.; C. R. Hebd. Seances Acad. Sci., Ser. D 1977, 285, 1221.
- 27. Blinov, V. A.; Mel'nikov, O. K.; Ilyukhin, V. V.; Voronkov, A. A.; Belov, N. V. Inorg. Mater. (Engl. Transl.) 1973, 9, 480; Izv. Akad. Nauk. SSSR Neorg. Mater. 1973, 9, 530.
- 28. Kornev, A. N.; Maksimov, B. A.; Lider, V. V.; Ilyukhin, V. V.; Belov, N. V. Sov. Phys. Dokl. (Engl. Transl.) 1973, 17, 735; Dokl. Akad. Nauk. SSSR 1972, 205, 831.
- Hubert, Y.; Guth, J.-L.; Perati, B.; Wey, R. C. R. Hebd. Seances Acad.
   Sci., Ser. D. 1976, 283, 291.

-12-

- 30. Kawamura, K.; Kawahara, A.; Henmi, K. Kobutsugku Zasshi 1976, 12, 403.
- 31. A report claiming the synthesis of canasite by wet methods has appeared; Melkonyan, G. S.; Finkel'shtein, B. I.; Kagramanyan, I. S. Tr. Nauchno.-Issled. Inst. Kamnya Silikatov 1970, No. 5, 233; Chem. Abstr. 1972, 77, 9098k.
- 32. Hubert, Y.; Jordan, D.; Guth, J.-L.; Kalt, A. C. R. Hebd. Seances Acad. Sci., Ser. D 1977, 284, 329.
- 33. Pabst, A. Acta Crystallogr. 1959, 12, 733.
- 34. Mazzi, F.; Pabst, A. Am. Mineral. 1962, 47, 409.
- 35. Zambonini, F. "Mineralogia Vesuviana", 2nd ed.; S. I. E. M.: Naples, 1935; p. 435.
- 36. Peacor, D. R.; Buerger, M. J.; Am. Mineral. 1962, 47, 539.
- 37. Pauling, L. Proc. Natl. Acad. Sci. U.S.A. 1930, 16, 578.
- 38. For clarity the orientations of the tubes in Figures 2 and 4 are matched.
- 39. The latticework tubes which are the true counterparts of the latticework sheets are, of course, those in which the links that are not parallel to the tube axes are curved. Here the discussion is not based on such tubes but rather on the corresponding tubes having appropriately shorter straight links in place of the curved links. The variations in the lengths of the links are not taken into account. This is acceptable because it appears, on the basis of the tubes examined, that often variants of these straightlink tubes have links of equal lengths (for example, a variant of the straight-link narsarsukite latticework tube has links with equal lengths). Further, the silicon-silicon distance in the Si-O-Si system is variable because the Si-O-Si angle is variable.

- 40. Taylor, W. H.; Náray-Szabó, St. Z. Kristallogr. Kristallgeom. Kristallphys. Kristallchem. 1931, 77, 146.
- 41. Fleet, S. G. Z. Kristallogr. Kristallgeom. Kristallphys. Kristallchem. 1965, 121, 349.
- 42. Again for clarity the orientations or the tubes in Figures 1 and 5 and Figures 3 and 6 are matched.
- 43. Takéuchi, Y.; Kawada, I.; Irmaziri, S.; Sadanaga, R. Mineral. J. 1969, 5,
  450; and Takéuchi, Y. private communication.
- 44. As examination shows, the latticework tubes in Figures 10 and 4 and Figures 11 and 7 are close structural analogues.
- 45. The kaolinite, apophyllite, dalyite, and manganpyrosmalite parent latticework sheets are, as is apparent, based on three-connected networks. It is of interest to note that the first of these latticeworks is based on the only regular three-connected network and that the second and fourth are based on two of the three semiregular three-connected networks.<sup>46</sup> It is also of interest that the dalyite parent latticework can be derived formally from the kaolinite parent latticework by appropriate insertion of sets of links arranged in U's and the apophyllite parent latticework similarly can be derived from the dalyite parent latticework.
- 46. Wells, A. F., "Structural Inorganic Chemistry", 4th ed.; Oxford, New York, 1975; Chapter 3.
- 47. These latticework sheets are also based, as they must be, on three-connected networks.
- 48. Warren, B.; Bragg, W. L. Z. Kristallogr. Kristallgeom. Kristallphys. Kristallchem. 1928, 69, 168.

-14-

- 49. Wenk, H. R. Am. Mineral. 1974, 59, 86.
- Cannillo, E.; Mazzi, F.; Fang, J. H.; Robinson, P. D.; Ohya, Y. Am. Mineral.
   1971, 56, 427.
- 51. Woodrow, P. J. Acta Crystallogr. 1967, 22, 673.
- 52. Warren, B. E. Z. Kristallogr. Kristallgeom. Kristallphys. Kristallchem. 1929, 72, 42.
- 53. Veblen, D. R.; Burnham, C. W. Am. Mineral. 1978, 63, 1058.
- 54. Hesse, K.-F.; Liebau, F. Z. Kristallogr. 1980, 153, 3.
- Voronkov, A. A.; Pyatenko, Yu. A.; Sov. Phys. Crystallogr. (Engl. Transl.)
   1962, 6, 755; Kristallograffya 1961, 6, 937.
- 56. Mamedov, Kh. S.; Belov, N. V. Zap. Vses. Mineral. Ova. 1956, 85, 13.
- Petrunina, A. A., Ilyukhin, V. V., Belov, N. V. Sov. Phys. Dokl. (Engl. Transl.) 1971, 16, 338; Dokl. Akad. Nauk. SSSR 1971, 198, 575.
- Bakakin, V. V.; Belov, N. V.; Borisov, S. V.; Solovyeva, L. P. Am. Mineral.
   1970, 55, 1167.
- 59. Some of these tube-chain and tube-band relationships have been discussed before, e.g., the litidionite-vlasovite,<sup>13</sup> fenaksite-vlasovite,<sup>14</sup> Na<sub>2</sub>CuSi<sub>4</sub>0<sub>10</sub>-vlasovite,<sup>28</sup> agrellite-vlasovite,<sup>16</sup> miserite-wallastonite(diopside),<sup>18</sup> miserite-xonotlite,<sup>18</sup> canasite-wallastonite(diopside),<sup>19</sup> and canasite-xonotlite<sup>19</sup> relationships (in the case of the miserite-xonotlite relationship the dissimilarities are focused on).
- 60. As is clear, a close formal relationship exists between the chain in diopside and the bands in tremolite, jimthompsonite, and the synthetic barium silicates. It is thus interesting to note that the tremolite-type band in nephrite  $Ca_2(Mg_Fe)_5Si_8O_{22}(OH,F)_2$ , is occasionally replaced by a triple-strand band.<sup>61</sup>

- Hutchison, J. L.; Jefferson, D. A.; Mallison, L. G.; Thomas, J. M. Mater. Res. Bull. 1976, 11, 1557.
- 62. A real relationship may exist between the band in howieite, and its counterpart sheet since it has been suggested that the band is formed from the sheet by a shearing process.<sup>49</sup>
- 63. It should not be assumed, of course, that sizable channels or even any channels will exist in silicates having moderate sized tubes since the tubes may contain ions (as does the tube in miserite) or may not be fully extended (or both).
- Musso, H.; Nauman, K. Angew. Chem. Int. Ed. Engl. 1966, 5, 127; Angew.
   Chem. 1966, 78, 116.
- 65. The litidionite-type tube having a carbon framework would be strained to an appreciable extent.

### Figures

- Figure 1. An ORTEP drawing of the tube ion in litidionite made with data gathered by Martin Pozas, Rossi, and Tazzoli.<sup>13</sup> The junctions indicate silicon atoms and the open circles oxygen atoms. The  $O_1-O_2$  distance is 4.85Å.
- Figure 2. An ORTEP drawing of the tube ion in narsarsufkite based on data gathered by Peacor and Buerger.<sup>36</sup> The  $0_1^{-0} 0_2$  distance is 4.32Å.
- Figure 3. An ORTEP drawing of the tube ion in miserite made with data gathered by Scott.<sup>18</sup> The  $0_1 0_2$  distance is 7.53Å.
- Figure 4. The parent latticework of the narsarsukite tube, this latticework opened along a lengthwise line and flattened, and the parent latticework of the kaolinite sheet with a strip emphasized that is the same as the opened and flattened tube latticework. The filled circles represent pendent units pointing outwards in the case of the tube and upwards in the case of the flat latticeworks. Like junctions have like numbers.
- Figure 5. The parent latticework of the litidionite tube, this latticework opened, and the parent latticework of the apophyllite sheet with a strip emphasized that is the same as the opened-tube latticework except for the arrangement of the pendent units. The filled circles represent pendent units pointing outwards or upwards and the open circles pendent units pointing downwards.
- Figure 6. The parent latticework of the miserite tube, this latticework opened, and the parent latticework of the dalyite sheet with a strip emphasized that is the same as the opened-tube latticework except for the arrangement of the pendent units.

- Figure 7. A second latticework tube related to the parent latticework of the kaolinite sheet, this latticework opened, and the kaolinite latticework sheet with a strip emphasized that is the same as the opened-tube latticework.
- Figure 8. A second latticework tube related to the parent latticework of the apophyllite sheet, this latticework opened, and the apophyllite latticework sheet with a strip emphasized that is the same as the opened-tube latticework except for the arrangement of the pendent units.
- Figure 9. A second latticework tube related to the parent latticework of the dalyite sheet, this latticework opened, and the dalyite latticework sheet with a strip emphasized that is the same as the opened-tube latticework except for the arrangement of the pendent units.
- Figure 10. A latticework tube related to the parent latticework of the manganpyrosmalite sheet, this latticework opened, and the manganpyrosmalite latticework sheet with a strip emphasized that is the same as the opened-tube latticework except for the arrangement of the pendent units.
- Figure 11. A second latticework tube related to the parent latticework of the manganpyrosmalite sheet, this latticework opened, and the manganpyrosmalite latticework sheet with a strip emphasized that is the same as the opened-tube latticework except for the arrangement of the pendent units.
- Figure 12. A latticework tube related to the larger of the two latticework tubes associated with the manganpyrosmalite latticework sheet.

-18-

- Figure 13. A chiral latticework tube related to the apophyllite latticework sheet. A part of a series of links following a generally helical path is emphasized.
- Figure 14. A latticework tube related to a pentagon-octagon latticework, this latticework opened, and the pentagon-octagon latticework with a strip emphasized that is the same as the opened-tube latticework.
- Figure 15. The latticework drum-ladder structure related to the 122:4,6,12latticework tube.
- Figure 16. Relationships between known and potential silicate tubes and known silicate linear structures. The frameworks are those of strips corresponding to the tubes. Superimposed on these are the latticeworks of the linear systems. These are altered, as necessary, by minor or significant amounts. The latticework-framework pairs shown are: (a) the diopside- and howieite-narsarsukite pairs, (b) the diopside-, aenigmatite-, astrophyllite-, tremolite-, jimthompsonite-,  $Ba_5Si_8O_{21}$ -, and  $Ba_6Si_{10}O_{26}$ -8<sub>2</sub>:6 pairs, (c) the diopside-, vlasovite-, xonotlite-, and tinaksite-litidionite pairs, and (d) the diopsideand aenigmatite-8,:4,8 pairs. In those cases where the latticeworks are substantially unaltered the directions of the pendent units are indicated: short lines, filled circles, and open circles representing respectively units pointing sideways, upwards, and downwards; and triangle-short line combinations representing pairs of units pointing at an angle. In the remaining cases only the presence of the pendent units is indicated: square-short line combinations representing groups of two or three units.

-19-

- Figure 17. Additional relationships between known and potential silicate tubes and known silicate linear structures. The latticework-framework pairs shown are: (a) the diopside-, and xonotlite-miserite pairs, (b) the diopside, aenigmatite-, and howieite- $6_2$ :4,6,8 pairs, (c) the diopside and nordite- $12_2$ :4,6,12 pairs, and (d) the diopsideand vlasovite- $20_2$ :4,6,12 pairs (for economy only half the  $12_2$ :4,6,12 and  $20_2$ :4,6,12 frameworks are shown).
- Figure 18. Tubes with single-link members running lengthwise and circumferentially: (a) a tube with four units at each junction, (b) a tube with five units at each junction, and (c) a tube with six units at each junction.
- Figure 19. The structures of: (a) norpinane, (b) tricyclo[3.1.1.1<sup>2,4</sup>]octane, (c) the hypothetical narsarsukite-type polymer, and (d) the hypothetical litidionite-type polymer.



Fig 3

4







ł

Fig 4



Fig 5



AND STREET





-22-



Fig 7







ł

Fig 8













Fig 10







Į

Fig 11



Fig 12



Fig 13







1

Fig 14





Martine William



-26-

Fig 16

-

1

100

•

![](_page_32_Figure_0.jpeg)

![](_page_33_Figure_0.jpeg)

![](_page_33_Figure_1.jpeg)

Fig 18

t

![](_page_33_Figure_3.jpeg)

![](_page_33_Figure_4.jpeg)

-

. •

-28-

# SP472-3/A1

-----

472:GAN:716:enj 78u472-608

# TECHNICAL REPORT DISTRIBUTION LIST, GEN

	No. Copies		<u>No</u> . <u>Copies</u>
Office of Naval Research		U.S. Army Research Office	
Attn: Code 472		Attn: CRD-AA-IP	
800 North Quincy Street		P.O. Box 1211	
Arlington, Virginia 22217	2	Research Triangle Park, N.C. 27709	1
ONR Western Regional Office		Naval Ocean Systems Center	
Attn: Dr. R. J. Marcus		Attn: Mr. Joe McCartney	
1030 East Green Street		San Diego, California 92152	1
Pasadena, California 91106	1		
		Naval Weapons Center	
ONR Eastern Regional Office		Attn: Dr. A. B. Amster,	
Attn: Dr. L. H. Peebles		Chemistry Division	
Building 114, Section D		China Lake, California 93555	1
666 Summer Street			•
Boston, Massachusetts 02210	t	Naval Civil Engineering Laboratory	
	-	Attn: Dr. R. W. Drisko	
Director Nevel Research Laborstory		Port Hueneme California 93401	1
Attn: Code 6100		iore addressed, ourselvente spave	4
Washington D C 20390	1	Department of Physics & Chemistry	
Washington, D.C. 20070	•	Nevel Doctoreducte School	
The Accietant Secretary		Monterey California 93940	1
of the News (DELS)	•	Muterey, Galiloidia 33340	1
Department of the News		Salantific Advisor	
Department of the Navy		Sciencific Advisor	-
Koom 41/30, Pentagon	•	Commandant of the Marine Corps	
Washington, D.C. 20350	L	(Code KD-1)	
Constant March Adm Constants Constants		wasnington, D.C. 20380	1
Commander, Naval Air Systems Command			
Attn: Code JIOC (H. Rosenwasser)		Naval Ship Research and Development	
Department of the Navy		Center	
Washington, D.C. 20360	1	Attn: Dr. G. Bosmajian, Applied Chemistry Division	
Defense Technical Information Center		Annapolis, Maryland 21401	1
Building 5, Cameron Station			
Alexandria, Virginia 22314	12	Naval Ocean Systems Center	
		Attn: Dr. S. Yamamoto, Marine	
Dr. Fred Saalfeld	•	Sciences Division	
Chemistry Division. Code 6100		San Diego, California 91232	1
Naval Research Laboratory			•
Washington, D.C. 20375	1	Mr. John Boyle	
	-	Materials Branch	
		Naval Ship Engineering Center	
		Philadelphia, Pennewlwania 19112	1
		encedections courses a 7112	1

SP472-3/A3

472:GAN:716:enj 78u472-608

## TECHNICAL REPORT DISTRIBUTION LIST, GEN

## <u>No</u>. Copies

1

Mr. James Kelley DTNSRDC Code 2803 Annapolis, Maryland 21402

Mr. A. M. Anzalone Administrative Librarian PLASTEC/ARRADCOM Bldg 3401 Dover, New Jersey 07801

. 1

# SP472-3/A6

· • •

....

472:GAN:716:enj 78u472-608 *и* •

## TECHNICAL REPORT DISTRIBUTION LIST, 356B

. . . . . .

. <b>.</b>	<u>No</u> . Copies		<u>No</u> . Copies
Dr. C. L. Shilling		Dr. E. Fischer, Code 2853	
Union Carbide Corporation		Naval Ship Research and	
Chemical and Plastics		Development Center	
Tarrytown Technical Center Tarrytown New York	1	Annapolis Division	•
Tallytowd, New IOLK	•	Annapolis, Maryland 21402	1
Dr. R. Soulen		Dr. Martin H. Kaufman, Head	
Contract Research Department		Materials Research Branch (Code 4542)	
Pennwalt Corporation		Naval Weapons Center	
900 First Avenue	-	China Lake, California 93555	1
King of Prussia, Pennsylvania 19406	1		
		Dr. C. Allen	
Dr. A. G. MacDiarmid		University of Vermont	
Department of Charistry		Department of Chemistry	•
Objladelphia Deprevivania 19174	1	Burlington, vermont 00401	1
Initalerphia, remistrante 17174	*	Professor R. Brago	
Dr. H. Allcock		Department of Chemistry	
Pennsylvania State University		University of Illinois	
Department of Chemistry		Urbana, Illinois 61801	1
University Park, Pennsylvania 16802	1		-
	1		
	*	COL R. W. Bowles, Code 100M	
Dr. R. Lenz		Office of Naval Research	
University of Massachusetts		800 N. Quincy Street	
Department of Chemistry		Arlington, Virginia 22217	1
Amherst, Massachusetts 01002	1		
		Professor T. Katz	
Dr. M. David Curtis		Department of Chemistry	
University of Michigan	•	Columbia University	
Department of Chemistry	•	New York, New York 10027	1
Ann Arbor, Michigan 48105	T		
		Professor Ismas Chien	
NASI-Taule Research Conter		Professor James Chien	
NASA-Lewis Research Center Artn: Dr. T. T. Serafini, MS 49-1		Professor James Chien Department of Chemistry University of Massachusetts	
NASA-Lewis Research Center Artn: Dr. T. T. Serafini, MS 49-1 21000 Brookpark Road		Professor James Chien Department of Chemistry University of Massachusetts Amhérst, Massachusetts 01002	1
NASA-Lewis Research Center Artn: Dr. T. T. Serafini, MS 49-1 21000 Brookpark Road Cleveland, Ohio 44135	1	Professor James Chien Department of Chemistry University of Massachusetts Amhérst, Massachusetts 01002	1
NASA-Lewis Research Center Artn: Dr. T. T. Serafini, MS 49-1 21000 Brookpark Road Cleveland, Ohio 44135	1	Professor James Chien Department of Chemistry University of Massachusetts Amhérst, Massachusetts 01002 Professor Malcolm B. Polk	1
NASA-Lewis Research Center Artn: Dr. T. T. Serafini, MS 49-1 21000 Brookpark Road Cleveland, Ohio 44135 Dr. J. Griffith	1	Professor James Chien Department of Chemistry University of Massachusetts Amhérst, Massachusetts 01002 Professor Malcolm B. Polk Department of Chemistry	1
NASA-Lewis Research Center Artn: Dr. T. T. Serafini, MS 49-1 21000 Brookpark Road Cleveland, Ohio 44135 Dr. J. Griffith Naval Research Laboratory	1	Professor James Chien Department of Chemistry University of Massachusetts Amhérst, Massachusetts 01002 Professor Malcolm B. Polk Department of Chemistry Atlanta University	1
NASA-Lewis Research Center Attn: Dr. T. T. Serafini, MS 49-1 21000 Brookpark Road Cleveland, Ohio 44135 Dr. J. Griffith Naval Research Laboratory Chemistry Section, Code 6120	1	Professor James Chien Department of Chemistry University of Massachusetts Amhérst, Massachusetts 01002 Professor Malcolm B. Polk Department of Chemistry Atlanta University Atlanta, Georgia 30314	1
NASA-Lewis Research Center Attn: Dr. T. T. Serafini, MS 49-1 21000 Brookpark Road Cleveland, Ohio 44135 Dr. J. Griffith Naval Research Laboratory Chemistry Section, Code 6120 Washington, D.C. 20375	1	Professor James Chien Department of Chemistry University of Massachusetts Amhérst, Massachusetts 01002 Professor Malcolm B. Polk Department of Chemistry Atlanta University Atlanta, Georgia 30314	1
NASA-Lewis Research Center Attn: Dr. T. T. Serafini, MS 49-1 21000 Brookpark Road Cleveland, Ohio 44135 Dr. J. Griffith Naval Research Laboratory Chemistry Section, Code 6120 Washington, D.C. 20375	1	Professor James Chien Department of Chemistry University of Massachusetts Amhérst, Massachusetts 01002 Professor Malcolm B. Polk Department of Chemistry Atlanta University Atlanta, Georgia 30314 Dr. G. Bryan Street	1
NASA-Lewis Research Center Attn: Dr. T. T. Serafini, MS 49-1 21000 Brookpark Road Cleveland, Ohio 44135 Dr. J. Griffith Naval Research Laboratory Chemistry Section, Code 6120 Washington, D.C. 20375 Dr. G. Gocdman	1	Professor James Chien Department of Chemistry University of Massachusetts Amhérst, Massachusetts 01002 Professor Malcolm B. Polk Department of Chemistry Atlanta University Atlanta, Georgia 30314 Dr. G. Bryan Street IBM Research Laboratory, K32/281 San Jose California 95193	1
NASA-Lewis Research Center Attn: Dr. T. T. Serafini, MS 49-1 21000 Brookpark Road Cleveland, Ohio 44135 Dr. J. Griffith Naval Research Laboratory Chemistry Section, Code 6120 Washington, D.C. 20375 Dr. G. Gocdman Globe-Union Incorporated 5757 North Green Bay Avenue	1	Professor James Chien Department of Chemistry University of Massachusetts Amhérst, Massachusetts 01002 Professor Malcolm B. Polk Department of Chemistry Atlanta University Atlanta University Atlanta, Georgia 30314 Dr. G. Bryan Street IBM Research Laboratory, K32/281 San Jose, California 95193	1 1 1

![](_page_37_Picture_0.jpeg)