





MERCING FOR PUBLIC MELANE, DISTRIBUTION CALINETED

ALL PORCE CROPING LOS LABORADORY ALL PORCE STRUCTURE COMMUNE CONTRACT PROVIDE COMMUNE CONTRACT PROVIDENTS 10731

88 01 14 004

Qualified requestors may obtain additional copies from the Defense Technical Information Center. All others should apply to the National Technical Information Service.

DEDORT DOCUMENTATION DAG	READ INSTRUCTIONS
REPURT DUCUMENTATION PAGE	BEFORE COMPLETING FORM
	VT ACCESSION NOL 3 RECIPIENT'S CATALOG NUMBER
AFGL-TR-82-0309	12 9 98 9
. TITLE (and Subtitle)	5. TYPE OF REPORT & PERIOD COVER
Derivation and Implementation of	f Expo- Scientific Report No.
nential Functions to Model Avia	
Scattoring Sportromotor Data Di	$\frac{1}{6} = \frac{1}{6} = \frac{1}$
tions	SCITOR IN FERFORMING DIG. REPORT NUMBER
	B. CONTRACT OF CRANT NUMPERCO
	S. CONTINUET ON ONANT NOMBER(S)
Lawrence E. Belsky	AF19628-81-C-0141
James P. Lally	
Peter L. Miller	
PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TAS AREA & WORK UNIT NUMBERS
Digital Programming Services, In	nc.
60 Hickory Drive	62101F
Waltham, MA 02154	23106580
1. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE
Air Force Compusice Laboratory	30 50 00
Hance AFR Macaachusatta A17	
Manshem (Dambarra) Main (TVC)	
MONITORING AGENCY NAME & ADDESS(1) different from (Controlling Office) 15, SECURITY CLASS (of this report)
A MONTONING ACCIVE AND A ADDRESSIT UTTERED INC.	
	Inclassified
	Unerabbilitea
	154. DECLASSIFICATION/DOWNGRADING
Approved for public release; dis • 7. DISTRIBUTION STATEMENT (of the abstract entered in Bloc	stribution unlimited. k 20, 11 dillerent from Report)
Approved for public release; dis • 7. DISTRIBUTION STATEMENT (of the abstract entered in Bloc	stribution unlimited. k 20, 11 different from Report)
Approved for public release; dis • 7. DISTRIBUTION STATEMENT (of the abstract entered in Bloc 9. SUPPLEMENTARY NOTES	stribution unlimited.
Approved for public release; dis • • • • • • • • • • • • • • • • • • •	stribution unlimited. k 20, 11 different from Report) Ity by block number) tribution functions, mathematica
Approved for public release; dis • • • • • • • • • • • • • • • • • • •	stribution unlimited. k 20, 11 different from Report) lify by block number) tribution functions, mathematica method, method of least squares,
Approved for public release; dis • • • • • • • • • • • • • • • • • • •	stribution unlimited. k 20, 11 different from Report) Ify by block number) tribution functions, mathematica method, method of least squares, fy by block number)
Approved for public release; dis Approved for public release; dis Approved for public release; dis Approved for public release; dis Approved for public release; and identified Supplementary notes Approved in Bloc Approved	<pre>stribution unlimited. k 20, If different from Report) Ify by block number) tribution functions, mathematical method, method of least squares, fy by block number) tical derivation of the Weibull s for the purpose of approxi- ta via a mathematical model, h function to model the data</pre>
Approved for public release; dis Approved for public release; dis DISTRIBUTION STATEMENT (of the abstract entered in Bloc SUPPLEMENTARY NOTES . SUPPLEMENTARY NOTES . SUPPLEMENTARY NOTES . SUPPLEMENTARY NOTES . KEY WORDS (Continue on reverse side if necessary and identify cloud droplet spectra data, dist derivation, maximum likelihood r Newton-Raphson method ABSTRACT (Continue on reverse side if necessary and identify this report details the mathemation and gamma distribution functions mating cloud droplet spectra data and analyzes the ability of each distributions. D FORM 1473 EDITION OF 1 NOV 65 IS OBSOLETE	<pre>stribution unlimited. k 20, 11 dillerent from Report) lty by block number) tribution functions, mathematica. method, method of least squares, // by block number) tical derivation of the Weibull s for the purpose of approxi- ta via a mathematical model, h function to model the data</pre>

LAND DEAL INTERPORT INTERACTION OF A L

•

Acknowledgements

The authors wish to thank Ms. Susan Sadofsky for her technical review. Special thanks is given to Mr. Morton Glass of AFGL for his suggestion of the work, for his guidance and for his many constructive criticisms.

	Accession For N2IS GRAAI DTIC ZAB Unannounced Justification		
6	<pre>Bistribution/ Availability Codes</pre>		
y	Dint Special		

TABLE OF CONTENTS

.

South and a standard and a standard

WOOD CONT

Availability Codes

Spectal

1810

I.	INTRODUCTION	3
II.	WEIBULL DISTRIBUTION FUNCTION	1
III.	THE KHRGIAN-MAZIN FORM OF THE GAMMA	
	DISTRIBUTION FUNCTION	ł
IV.	ANALYSIS	1
v.	CONCLUSIONS	ł
VI.	REFERENCES	;
	AT DATEABASA	
	NTTS GRALE ITLC TAN INMEMERIASI JUSTESCELLUI	
	89	
	/uo12ng122era	

I. Introduction

In the course of analyzing data under contract AF19628-81-C-0141, an attempt was made to fit certain exponential functions to number density data obtained from axial scattering spectrometer probes.

These data were collected in supercooled clouds associated with aircraft icing. It was noted that the distribution of the number density spectra in the size range of 2 to 30 microns diameter seemed to be describable by the Weibull function or the Khrgian-Mazin¹ form of the gamma distribution. The Weibull function has been used in time-to-failure probability analysis. These functions follow:

Weibull Distribution

$$f(x) = \int_{-\infty}^{\infty} \frac{x^{b-1} e^{-ax^{b}}}{x^{c}} x \ge 0 \quad a \ge 0, b \ge 0 \quad (1)$$

(2)

where N is the distribution function

Further discussion of this Weibull function can be found in Bethea et al.²

Gamma Distribution of degree 2 (Khrgian-Mazin)¹

$$f(x) = \begin{cases} N\frac{a}{2}^{3}x^{2}e^{-ax} & x \ge 0 \\ 0 & x < 0 \end{cases}$$

where N is the distribution function

The method of least squares analysis can be used to determine values of a and b in these functions. It is noted that this effort will require extensive use of numerical analysis to solve the resulting simultaneous non-linear equations. In order to save computation costs it was decided to utilize the maximum likelihood method (MLM) which has the advantage of reducting most solutions to a closed form. The maximum likelihood method (MLM) is described by Breiman³.

II. Weibull Distribution Function

In order to fit a function, f(x), to a data distribution (y), one mathematical method which can be used is the MLM. This method requires that, for the entire data distribution, the sum of the products of each observed value and the corresponding value of the function

$$\sum_{i=1}^{n} y_i f(x_i)$$

must be maximized.

Now, from equation 1, the Weibull is defined as:

$$f(x) = Nabx^{b-1}e^{-ax^{b}}$$

In order to fit this function to (x_i, y_i) , where,

x_i = channel i diameter
y_i = channel i normalized number density

"a" and "b" must be solved. Since the ln[f(x)] and f(x)
are maximized for the same values of "a" and "b", and since
f(x) is a product of functions of "a" and "b", it is easier
to solve for those values of "a" and "b" which maximize ln[f(x)].

Therefore, since the MLM implies that the partial derivatives equal zero, "a" and "b" can be solved for by the following equations:

$$\frac{\partial}{\partial a} \sum_{i=1}^{n} y_i \ln(Nabx_i^{b-1} e^{-ax_i^{b}}) = 0$$
(3)

and

$$\frac{\partial}{\partial b} = \sum_{i=1}^{n} y_i \ln(Nabx_i^{b-1}e^{-ax_i^b}) = 0$$
 (4)

Solving equation (1):

 $\frac{\partial}{\partial a} \sum_{i=1}^{n} y_{i} \ln (Nabx_{i}^{b-1}e^{-ax_{i}^{b}}) = 0$ $\sum_{i=1}^{n} (y_{i}) \frac{\partial}{\partial a} (\ln (Nabx_{i}^{b-1}e^{-ax_{i}^{b}})) = 0$ $\sum_{i=1}^{n} (y_{i}) \frac{\partial}{\partial a} (\ln N + \ln a + \ln b + (b-1) \ln x_{i}^{-ax_{i}^{b}}) = 0$ $\sum_{i=1}^{n} y_{i} (\frac{1}{a} - x_{i}^{b}) = 0$

Solving for "a" yields:

$$a = \frac{\underset{i=1}{\overset{n}{\underset{i=1}{\sum}} y_{i}}}{\underset{i=1}{\overset{n}{\underset{i=1}{\sum}} y_{i} x_{i}^{b}}}$$
(5)

Similarly solving equation (4):

1

$$\frac{\partial}{\partial b} \sum_{i=1}^{n} y_i \ln(\text{Nabx}_i^{b-1} e^{-ax_i^b}) = 0$$

 $\sum_{i=1}^{n} (y_i) \frac{\partial}{\partial b} (\ln(\operatorname{Nabx}_{i}^{b-1} e^{-ax_i^b})) = 0$

$$\sum_{i=1}^{n} (y_i) \frac{\partial}{\partial b} (\ln N + \ln a + \ln b + (b-1) \ln x_i - a x_i^b) = 0$$

$$\sum_{i=1}^{n} (y_i) \left(\frac{1}{b} + \ln x_i - a x_i^b \ln x_i\right) = 0$$

$$1 \qquad n \qquad n \qquad b$$

$$\frac{1}{b} \sum_{i=1}^{n} y_i + \sum_{i=1}^{n} y_i \ln x_i - a \sum_{i=1}^{n} y_i x_i^b \ln x_i = 0$$
(6)

Now, substituting for "a" (from equation (5)) in equation (6) and realizing that $\begin{array}{c}n\\\Sigma\end{array}$ $y_{i} \neq 0$, yields: i=1

$$\frac{1}{b} + \frac{\frac{\sum_{i=1}^{n} y_i \ln x_i}{\sum_{i=1}^{n} y_i} - \frac{\sum_{i=1}^{n} y_i x_i^{b} \ln x_i}{\sum_{i=1}^{n} y_i} = 0$$
(7)

Since b cannot be solved for in closed form, a numerical method must be used. One such method is the Newton-Raphson method as described by Scarborough⁴. In applying this method, let equation (7) equal g(b). Since the expression



is a constant (K) for a given distribution, and independent of the choice of a, b:

$$g(b) = \frac{1}{b} + \kappa - \frac{\sum_{i=1}^{n} y_i x_i^{b} \ln x_i}{\sum_{i=1}^{n} y_i x_i^{b}}$$

Now, to solve for the root, b, such that g(b) = 0, an initial guess for $g(b_0)$ is made. Next, solve for b_{n+1} where:

$$b_{n+1} = b_n - \frac{g(b_n)}{g'(b_n)} \quad \text{and where} \qquad (8)$$

 $|\mathbf{b}_{n+1}-\mathbf{b}_n| < \varepsilon$. The object now is to define ε small enough

so that $b \doteq b_{n+1}$. Since the Newton-Raphson method is repetitive by nature, g'(b_n) can be approximated without significant loss of accuracy:

$$g'(b_n) = \lim_{E \to 0} \frac{g(b_n + E) - g(b_n)}{E}$$

Therefore, substituting this approximation for $g'(b_n)$ in equation 8:

$$\mathbf{b}_{n+1} = \mathbf{b}_n - \frac{g(\mathbf{b}_n)E}{g(\mathbf{b}_n+E)-g(\mathbf{b}_n)}$$

Letting $E = .01(b_n)$ yields:

$$b_{n+1} = b_n - \frac{g(b_n)(.01)(b_n)}{g(b_n + .01b_n) - g(b_n)}$$
$$b_{n+1} = b_n - \frac{.01(b_n)}{\frac{g(1.01b_n) - g(b_n)}{g(b_n)}}$$

or

$$b_{n+1} = b_n - \frac{.01(b_n)}{\frac{g(1.01b_n)}{g(b_n)} - 1}$$
 (9)

With "a" and "b" defined, equations (5) and (9) respectively, only the independent factor N remains to be solved. Now, it can be shown that determining the value of N by way of the MLM, as was done for "a" and "b" in equation 3 and 4, would result in the expression:

$$\frac{1}{N} \sum_{i=1}^{n} y_i = 0$$

Since this expression is undefined for N, another method must be utilized to determine N. Therefore, applying the method of least squares to equation 1 and solving for N using partial derivatives yields:

$$\frac{\partial}{\partial N} \sum_{i=1}^{n} (y_i - Nabx_i^{b-1} e^{-ax_i^b})^2 = 0$$

$$\sum_{i=1}^{n} 2(y_i - Nabx_i^{b-1} e^{-ax_i^b}) (-abx_i^{b-1} e^{-ax_i^b}) = 0$$

$$\sum_{i=1}^{n} -2y_i abx_i^{b-1} e^{-ax_i^b} + \sum_{i=1}^{n} 2N(abx_i^{b-1} e^{-ax_i^b})^2 = 0$$

Factoring out 2ab yields:

$$\sum_{i=1}^{n} y_i x_i^{b-1} e^{-ax_i^b} = N(ab) \sum_{i=1}^{n} (x_i^{b-1} e^{-ax_i^b})^2,$$

and solving for N yields:

$$N = \frac{\sum_{i=1}^{n} y_{i} x_{i}^{b-1} e^{-ax_{i}^{b}}}{ab \sum_{i=1}^{n} (x_{i}^{b-1} e^{-ax_{i}^{b}})^{2}}$$
(10)

-.- III. The Khrgian-Mazin form of the Gamma Distribution Function

The gamma distribution derivation is accomplished in a manner similar to the Weibull function by fitting f(x) to (x_i, y_i) by way of the MLM. Thus, given that:

$$f(x) = \begin{cases} \frac{Na^{3}}{2} x_{i}^{2} e^{-ax_{i}} & x \ge 0 \\ 0 & x < 0 \end{cases}$$
(2)

solving for "a" via partial derivatives yields:

 $\frac{\partial}{\partial a} \sum_{i=1}^{n} (y_i \ln(N\frac{a^3}{2}x_i^2 e^{-ax_i}) = 0)$ $\sum_{i=1}^{n} (y_i \frac{\partial}{\partial a} (\ln N + \ln \frac{a^3}{2} + \ln x_i^2 - ax_i) = 0$ $\sum_{i=1}^{n} y_i ((\frac{2}{a^3}) (\frac{3a^2}{2}) - x_i) = 0$ $\sum_{i=1}^{n} y_i (\frac{3}{a} - x_i) = 0$ $\frac{\partial}{a} \sum_{i=1}^{n} y_i (\frac{3}{a} - x_i) = 0$

Solving for "a" yields:

$$\mathbf{a} = \frac{3 \sum_{i=1}^{k} y_i}{n}$$

$$\sum_{i=1}^{k} y_i x_i$$
(9)

Again, to avoid $\Sigma y_i = 0$ the method of least squares analysis is used to solve for N. Thus:

$$\frac{\partial}{\partial N} \sum_{i=1}^{n} (y_i - N(\frac{a^3}{2}) x_i^2 e^{-ax_i})^2 = 0$$

$$\sum_{i=1}^{n} 2(y_i - N(\frac{a^3}{2}) x_i^2 e^{-ax_i}) (-\frac{a^3}{2}) (x_i^2 e^{-ax_i}) = 0$$

$$\sum_{i=1}^{n} -2 y_i (\frac{a^3}{2}) x_i^2 e^{-ax_i} + \sum_{i=1}^{n} 2N((\frac{a^3}{2}) (x_i^2 e^{-ax_i}))^2 = 0$$
Factoring out and cancelling $\frac{2a^3}{2}$ yields:

$$\sum_{i=1}^{n} y_{i} x_{i}^{2} e^{-ax_{i}} = N \frac{a^{3}}{2} \sum_{i=1}^{n} (x_{i}^{2} e^{-ax_{i}})^{2}$$

and solving for N:

$$N = \frac{2 \sum_{i=1}^{n} y_i x_i^2 e^{-ax_i}}{a^3 \sum_{i=1}^{n} (x_i^2 e^{-ax_i})^2}$$

(12)

IV. Analysis

The Weibull and gamma distribution functions were applied to axial scattering spectrometer probe data obtained from 26 discrete data gathering flight passes made by the Air Force Geophysics Laboratory's instrumental Cl30-E aircraft during the 1979-1980 winter. The instrument probe was designed to collect data in the 2-30 micron range at 2 micron intervals. The resultant 15 data values (channels) for each flight pass comprised the raw data sets. However, due to the excessively noisy data signal present in channel 1, these data were eliminated from the analyses. Thus data channels 2-15 were used as input to the Weibull and gamma functions.

In order to demonstrate the ability of each function to approximate the number density distribution of each flight pass, Figures 1-3 are presented to compare the actual number density distribution for channels 2-15 to the modeled distributions. In addition, the pertinent meteorological data (liquid water content and temperature) for each flight pass are indicated on each figure.

The distributions from the three flight passes illustrated in Figures 1-3 were chosen to typify the range of liquid water content (LWC) and temperature values encountered during the 26 flights. As the figures show, both functions approximate the number density distributions for channels 2-6 with similar accuracy, but the Weibull generally fits the data better than the gamma function for channels 7-9. However, as the number density data levels off for channels 11-15, only the gamma function consistently models this behavior in an acceptable



•







manner. The Weibull's repeated tendency to approach zero at a rapid rate for channels 13-15, 11-15, and 9-15 respectively, necessarily eliminated it as the modeling function of choice. (Note: number density values which are less than the lower limit of the Y axis (1.0E-1) were not plotted.)

The impact of this behavior on the overall performance of the Weibull function is demonstrated in Figure 4, which depicts each functions' log RMS value for each of the 26 flight passes. As can be seen from the figure, 23 of the 26 (88%) gamma RMS values are less than or equal to 2.0 cm⁻³ μ^{-1} ; whereas only seven of the 26 (27%) Weibull RMS values are in this category.

This dissimilarity in the performance of the two functions prompted several analyses of the data to determine if some relationship or correlation factor could be discerned between a given data distribution and/or the meteorological conditions (LWC and temperature), during which the data distribution was obtained. Among these analyses was the comparison of the ratio of the number of LWC values above the median LWC value to the number of LWC values below the median for each function. The same comparison of ratios was made using temperature as the variate, with neither comparison evidencing a correlation between the magnitude of LWC or temperature values and the performance of the Weibull function. This lack of correlation between flight pass variables and the Weibull performance held true for log RMS values both above and below the 2.0 cm⁻³ \mu⁻¹ delimiter.

In addition, since the Weibull performed poorest for data channels 11-15 (usually values < 100), an attempt was

made to determine if the Weibull was sensitive to a particular range of data values less than 100. Despite the fact that the Weibull attained log RMS values very close to those of the gamma function for seven flight passes where channels 11-15 were in the 7-70 range (see Figure 4, passes 17-23), there were an equal number of instances where the RMS values deteriorated for flight passes where channels 11-15 were in this same range. (Note: the two flight passes (2 and 14) for which the Weibull attained the worst RMS values were considered an inadequate sample on which to perform analyses. However, it can be mentioned that the gamma function also attained its worst RMS values for the same two passes, albeit an order of magnitude difference exists between these values for the respective functions.)

V. Conclusions

These observations and analyses indicate that for this application, the gamma distribution function is a more accurate and consistent modeling method with which to approximate the data distributions. Subsequence analysis is planned to develop a weighting scheme to refine these modeling techniques.

VI. References

- Khrgian, A. Kh., Mazin, I.P., <u>Analysis of Methods for</u> <u>Describing Cloud Droplet Distribution Spectra</u>, Trudy TSAO, No. 17, 1956.
- Bethea, R. M., Duran, B. S., Boullion, T. L., <u>Statisti-</u> <u>cal Methods for Engineers and Scientists</u>, Marcel Dekker, Inc., 1975, p. 66.
- 3. Breiman, L., <u>Statistics: With a View Toward Applications</u>, Houghton Mifflin, 1973, p. 194 (6.12).

an and adding the second states of states where the second states of the second states of

 Scarborough, J. B., <u>Numerical Mathematical Analysis</u>, Johns Hopkins Press, 1962, p. 199.

