

AD-A123 779


DISTRIBUTIONAL ANALYSIS OF INVENTORY DEMAND OVER  
LEADTIME(U) NAVAL POSTGRADUATE SCHOOL MONTEREY CA  
M L YOUNT JUN 82.


1/2

UNCLASSIFIED

F/G 15/5

NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

(2)

ADA 123779

# NAVAL POSTGRADUATE SCHOOL

## Monterey, California



# THESIS

DISTRIBUTIONAL ANALYSIS  
OF INVENTORY  
DEMAND OVER LEADTIME

by

Mark Lee Yount

June 1982

Thesis Advisor: Charles F. Taylor, Jr.

Approved for public release, distribution unlimited

2  
S DTIC ELECTED JAN 26 1983 E

## Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
		ADA123 779
4. TITLE (and Subtitle) Distributional Analysis of Inventory Demand Over Leadtime		5. TYPE OF REPORT & PERIOD COVERED Master's Thesis June 1982
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s)  Mark Lee Yount		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Postgraduate School Monterey, California 93940		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Postgraduate School Monterey, California 93940		12. REPORT DATE June 1982
14. MONITORING AGENCY NAME & ADDRESS/If different from Controlling Office)		13. NUMBER OF PAGES 125
		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of info Report)  Approved for public release, distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Inventory demand distributions, Mixed Bernoulli distributions, Bootstrap, Data analysis, Resampling, Probabilistic models, Statistical models		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The distribution of inventory demand over procurement leadtime is modeled using mixed probability distributions that explicitly account for the high incidence of zero demands observed in Inventory Control Point Demand History Files. Analysis was limited to the right-hand tail area of the distribution on the assumption that that area is of critical importance in determining the effectiveness of an inventory system. Probabilistic		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 68 IS OBSOLETE  
S/N 0102-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE/When Data Entered

#20 - ABSTRACT - (Continued)

models studied were: 1) Bernoulli-exponential, 2) Bernoulli-lognormal, and 3) Bernoulli-logistic. These compound distributions were compared to several standard distributions including the Poisson, negative binomial and normal distributions using a resampling procedure appropriate in cases such as this where a paucity of data exists. Fits obtained from the 75th to 95th percentiles indicated the mixed distributions may be superior as a class to the standard distributions for high-demand items.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	



Approved for public release, distribution unlimited

Distributional Analysis  
of Inventory  
Demand Over Leadtime

by

Mark Lee Yount  
Lieutenant Commander, Supply Corps  
United States Navy  
A.B., Earlham College, 1968  
M.A., University of Colorado, 1971

Submitted in partial fulfillment of the  
requirements for the degree of

MASTER OF SCIENCE IN OPERATIONS RESEARCH

from the

NAVAL POSTGRADUATE SCHOOL  
June, 1982

Author:

Mark L. Yount

Approved by:

Charles F. Taylor, Jr.

Thesis Advisor

Donald R. Gaver

Second Reader

O.R. Pearl (acting)  
Chairman, Department of Operations Research

H. M. Woods  
Dean of Information and Policy Sciences

## ABSTRACT

The distribution of inventory demand over procurement leadtime is modeled using mixed probability distributions that explicitly account for the high incidence of zero demands observed in Inventory Control Point Demand History Files. Analysis was limited to the right-hand tail area of the distribution on the assumption that that area is of critical importance in determining the effectiveness of an inventory system. Probabilistic models studied were: 1) Bernoulli-exponential, 2) Bernoulli-lognormal, and 3) Bernoulli-logistic. These compound distributions were compared to several standard distributions including the Poisson, negative binomial and normal distributions using a resampling procedure appropriate in cases such as this where a paucity of data exists. Fits obtained from the 75th to 95th percentiles indicated the mixed distributions may be superior as a class to the standard distributions for high-demand items.

## TABLE OF CONTENTS

I.	INTRODUCTION	9
II.	BACKGROUND	12
A.	CURRENT MODEL SELECTION TECHNIQUES	12
B.	PREVIOUS RESEARCH	14
1.	Bernoulli-exponential Distribution	14
2.	Logarithmic-Poisson-Gamma Distribution	16
III.	PROPOSED MODELS	20
A.	COMPOUND BERNouLLI MODELS IN GENERAL	20
B.	MODEL DESCRIPTIONS	21
1.	Bernoulli-exponential Distribution	21
2.	Bernoulli-lognormal Distribution	23
3.	Bernoulli-logistic Distribution	25
IV.	EVALUATION PROCEDURE	28
A.	THE DATA	28
B.	THE RESAMPLING PROCEDURE	30
C.	MEASURES OF EFFECTIVENESS	33
D.	COMPUTER VALIDATION	34
V.	RESULTS	36
A.	LOW-DEMAND ITEMS	37
B.	MEDIUM-DEMAND ITEMS	38
C.	HIGH-DEMAND ITEMS	39
VI.	CONCLUSIONS AND RECOMMENDATIONS	41
A.	SUMMARY	41
B.	RECOMMENDATIONS	42

C. ADDITIONAL RESEARCH - - - - -	42
APPENDIX A: GLOSSARY OF TERMS - - - - -	44
APPENDIX B: CONSUMABLE ITEM TABULATED RESULTS - - - - -	46
APPENDIX C: REPAIRABLE ITEM TABULATED RESULTS - - - - -	58
APPENDIX D: ESTIMATION OF LOGNORMAL MEAN AND VARIANCE FROM UNTRANSFORMED DATA - - - - -	70
APPENDIX E: DATA RECORD LAYOUT - - - - -	72
APPENDIX F: FORTRAN PROGRAMS - - - - -	78
LIST OF REFERENCES - - - - -	122
BIBLIOGRAPHY - - - - -	123
INITIAL DISTRIBUTION LIST - - - - -	124

## LIST OF TABLES

1.	Current Inventory Control Point Probabilistic Demand Over Leadtime Models	10
2.	Required Data for Inverse Computations	31
3.	Low-Demand Items, Total Mean Squared Error ( $10^{-4}$ )	38
4.	Medium-Demand Items, Total Mean Squared Error ( $10^{-4}$ )	39
5.	High-Demand Items, Total Mean Squared Error ( $10^{-4}$ )	40

LIST OF FIGURES

1. Compound Cumulative Distribution Function,  
Bernoulli-lognormal Distribution - - - - - 22

## I. INTRODUCTION

Inventory system performance<sup>1</sup> in a situation with random demand depends upon reorder point<sup>2</sup> computations. Reorder point computations depend upon the probabilistic model chosen to represent the inventory system in that the reorder point is composed of two parts: 1) The expected demand during a procurement leadtime and 2) The safety level<sup>3</sup> determined from the probabilistic model of demand over leadtime. Probabilistic models are generally only utilized to represent demand over procurement leadtime since that is the only time period when stockouts<sup>4</sup> potentially occur. The chosen distribution is then a conditional distribution given procurement leadtime. The Navy has historically used three distributions of demand for models utilized at the Inventory Control Point (ICP) echelon of the Navy Supply System. The current probabilistic models and the average annual demand that is used to determine which model to utilize are displayed in Table 1.<sup>5</sup>

---

<sup>1</sup>See Appendix A for definition of "System Performance".

<sup>2</sup>See Appendix A for definition of "Reorder Point".

<sup>3</sup>See Appendix A for definition of "Safety Level".

<sup>4</sup>See Appendix A for definition of "Stockouts".

<sup>5</sup>The Aviation Supply Office (ASO) has recently changed to the use of the normal distribution as the model of choice for

Table 1: Current Inventory Control Point  
Probabilistic Demand Over Leadtime Models

<u>Distribution</u>	<u>Average Annual Demand Range</u>	
Poisson	0 - 1	Low
Negative Binomial	1 - 20	Medium
Normal	> 20	High

If the actual quarterly demand records for an ICP are examined, one striking characteristic that is present in nearly all records is the high incidence of zero observations. This is not surprising when one considers the echeloning of the supply system and the fact that users only place a demand on the wholesale system as their reorder points are reached. Since most activities stock at least one quarter's expected demand for an item, reorders are expected at most four times a year. By the natural phase differences of each activity's reorder actions, the ICP may experience zero demands during a given quarter for even high-demand items, and will surely often experience quarters of zero demands for medium- and low-demand items. A previous study of Pacific Fleet Combat Stores Ships' demand data [Ref. 1] found that the frequency of zero observations in those demand

---

all demand categories. Ships Parts Control Center (SPCC) still uses all the distributions in Table 1.

records appeared to have a low correlation with the overall demand level (i.e. high, medium or low).\*

A desirable model for an inventory system should compute reorder levels that accurately correspond to a specified stockout risk based on the level of actual or anticipated demand for each item. If such a model were available, the inventory system could avoid the added expense of either having too much or too little stock. One problem with the distributions in Table 1 is that none can account for the probability mass at zero; therefore, none can accurately compute reorder levels.

This study explores the use of compound probability models and their potential to more accurately achieve the goals presented above by explicitly accounting for the probability mass at zero. The results obtained should be considered as a first step in the exploration of a class of models which have heretofore received little attention in the context of modeling inventory demand.

---

\*See Section II.B.1 for a more detailed discussion of this study.

## II. BACKGROUND

### A. CURRENT MODEL SELECTION TECHNIQUES

In the past, the usual approach of choosing the probabilistic model for lead time demand was based on matching the empirical cumulative distribution function determined from the actual demand to various theoretical cumulative distribution functions. The shapes of the two curves are compared using the Kolmogorov-Smirnov Statistical Goodness of Fit test.<sup>7</sup> The Chi-Squared test has historically not been used because it is not very powerful when the number of observations is small (supply demand data is retained for at most twelve quarters and usually only eight quarters at the ICP level). The Kolmogorov-Smirnov test provides only a relative measure of goodness of fit, but it is usable with very small sample sizes. The goal then is to select the model with the best agreement between the empirical and theoretical cumulative distribution function curves.

The Navy Fleet Material Support Office (FMSO) has utilized the above Kolmogorov-Smirnov test to evaluate several theoretical distributions. Most recently [Ref. 2] FMSO

-----

<sup>7</sup>A good description of the Kolmogorov-Smirnov one and two-sample tests may be found in Siegel, S., Non Parametric Statistics, p.47-52 and 127-136, McGraw Hill, 1956.

evaluated seven distributions\* at the 90% confidence level using demand data from NAS Brunswick. The use of demand data from a stock point vice an ICP assumes the underlying demand behavior at the stock point and ICP are similar. Though this was never validated directly, the NAS Brunswick demand data were deemed to have "unique qualities better suited for demand analysis" as follows:

1. Twelve quarters of demand and demand frequency data were available per item, 50% more than available in ICP files,
2. New item identifiers that permit the selection of only steady state demand items were available,
3. Reliability of the data was established from prior PMSO studies.

The PMSO analysis showed that Navy demand patterns are very poorly modeled by any of the seven distributions studied. No new models were proposed and no attributes of the demand patterns were indicated as the most significant factor in the failure of the standard distributions to model Navy demand patterns.

There are several problems with the way the Kolmogorov-Smirnov test was used in the PMSO study:

---

\*Distributions tested were: Normal, Negative-Binomial, Poisson, Logistic, LaPlace, Gamma and Uniform.

1. The procedure is not strictly applicable in the case of discrete distributions such as the Poisson and negative binomial.
2. The test assumes that the parameters of the distribution being tested are completely specified in advance, i.e. not estimated from the data. (Lilliefors [Ref. 3] developed alternate tables for use in testing the exponential distribution when parameters are estimated from the data. This procedure was used in Reference 1.)
3. The Kolmogorov-Smirnov test evaluates goodness-of-fit over the entire range of the distribution. For inventory problems, the region of greatest interest is the right-hand tail of the distribution.

## B. PREVIOUS RESEARCH

Research into distributions which have properties that more closely model actual demand patterns has been very limited and, when conducted, often did not explore anything but the usual standard distributions. Two projects though stand out because of their innovative use of non-standard distributions and the good fit achieved.

### 1. Bernoulli-exponential Distribution

In Reference 1, Pacific Fleet Combat Stores Ship demand was modeled using the mixed Bernoulli-exponential

distribution. This distribution was chosen because it was observed that nearly all line items had an unusually high occurrence of demands of zero. Analysis of the data indicated that with high probability, the number of non-zero observations for a given item was unrelated to the average value of those non-zero observations. The conclusion was that the demand process could be thought of as two independent subprocesses with one process determining whether a demand would occur or not and the other determining the quantity of the demand, given the demand did occur. The former process was modeled as a Bernoulli process with parameter  $p$ ,  $p$  being the probability that a demand did occur. The latter process was modeled as a continuous exponential process based on exploratory data analysis. The resulting probability distribution function for the number of units demanded in a leadtime for the Bernoulli-exponential distribution is given by:

$$h(x) = \begin{cases} 1 - p & , x = 0 \\ p\lambda e^{-\lambda x}, & x > 0, \end{cases} \quad (1)$$

where:

$1/\lambda$  = the expected value of demand, given that the demand is greater than zero.

$p$  = probability that demand does occur.

The complementary cumulative distribution function is given by:

$$H(x) = p \exp(-\lambda x), \quad x \geq 0, \quad (2)$$

where  $H(x)$  is the probability that demand will not exceed  $x$ ; this is equivalent to the complementary cumulative distribution function used by Hadley and Whitin [Ref. 4].

The hypothesis that the demand of Combat Stores Ships could be modeled according to the above probability distribution was tested using the Kolmogorov-Smirnov test described earlier plus tests on the theoretical risk and the observed risk for values of risk in the upper right hand tail area of the distribution. The results of these tests on five independent samples from the demand data available provided strong evidence that the Bernoulli-exponential distribution describes the demand for any given stock item very well.

## 2. Logarithmic-Poisson-Gamma Distribution

A model for the distribution of demand during procurement leadtime [Ref. 5] was derived under the following assumptions:

- a. Requisitions occur according to a stationary Poisson Process,
- b. Requisition sizes follow a logarithmic distribution,

c. Leadtime is a random variable with the gamma distribution.

The resulting probability function for the number of units demanded in a leadtime is:

$$h(x) = \left( \frac{\theta}{\lambda + \theta} \right)^x \frac{\theta^x}{x!} \sum_{k=1}^x \left( \frac{\theta}{\lambda + \theta} \right)^k s_{x,k} \sum_{j=1}^k s_{k,j} \quad (3)$$

for  $x = 1, 2, 3, \dots$

and

$$h(0) = \left( \frac{\theta}{\lambda + \theta} \right)^0$$

where:

$\alpha, \beta$  = parameters of the Gamma distribution defining leadtime,

$\theta$  = parameter of the logarithmic distribution solved for by interval bisection from

$$E(X) = -\theta / (1 - \theta) \ln(1 - \theta),$$

$\lambda$  = requisition arrival rate,

$$c = -\lambda / \ln(1 - \theta)$$

$s_{x,k}$  = Stirling numbers of the first kind computed from the recursion

$$s_{x,k} = s_{x-1,k-1} + (x - 1)s_{x-1,k}$$

for  $k = 1, 2, \dots, x$  and  $x = 1, 2, \dots$

with  $s_{x,0} = 0$  for all  $x$

This distribution is called the Logarithmic-Poisson-Gamma (LPG). It is defined from four parameters  $\alpha$ ,  $\beta$ ,  $\theta$ , and  $\lambda$  ( $\theta$  and  $\lambda$  determine  $c$ ).

For more efficient computations, a recursion equation for integer  $\alpha$  was developed.

$$h(x) = \sum_{k=1}^x T_{x,k} \quad (4)$$

where:

$$T_{x,k} = \frac{\theta}{x} [C_2(\alpha + k - 1) T_{x-1,k-1} + (x - 1) T_{x-1,k}]$$

$$C_2 = c / (\lambda + \beta)$$

and  $\theta$ ,  $c$ ,  $\alpha$ ,  $\beta$  and  $\lambda$  defined as in (3).

Computing reorder points from fractiles of the LPG distribution, even using the simplified version of (4), is generally too time consuming for many real applications. As a result, an approximation was developed and tested that used a scaled version of the Poisson distribution to approximate the negative binomial distribution. The resulting approximation is:

$$p\{Z(t) = kx\} \approx \frac{(\alpha + x - 1)!}{x! (\alpha - 1)!} \left( \frac{\beta}{\mu + \beta} \right)^{\alpha} \left( \frac{\mu}{\mu + \beta} \right)^x \quad (5)$$

for  $x = 0, 1, 2, \dots$

where:

$Z(t)$  = number of units demanded in time  $t$

$$\mu = c\theta$$

$\alpha$ ,  $\beta$ ,  $c$  and  $\theta$  defined as in (3).

The authors conducted subjective tests of goodness of fit to their LPG model and its derivatives using Air Force consumable items and concluded, using as an example one item, that there was a "very close agreement between the observed and the predicted cumulative distribution functions for this item."

### III. PROPOSED MODELS

#### A. COMPOUND BERNoulli MODELS IN GENERAL

The results of the previous research indicate that a potential exists for improving inventory performance by the use of slightly more complex models. The focus of this study is on the compound Bernoulli models, leaving further analysis of the LPG model and its derivatives to others. The use of compound Bernoulli models has several advantages over standard simple distributions:

1. The model, by necessity, has an added parameter whose sole function is to estimate the probability of a non-zero demand.
2. For simple models defined only over the positive real numbers, the addition of the Bernoulli parameter simply results in the scaling of the cumulative distribution, thus adding little additional complexity to the model.
3. By explicitly accounting for zero demands, compound models account for more of the observed variance than do simple models.

Compound models are not without disadvantages:

1. Each parameter added to the model must be estimated for each individual stock item. For a large inventory

system such as the Navy's, this equates to considerable additional computational requirements.

2. Additional parameters must be stored for future retrieval or else computed from stored data when required, thus either extending computational time or else requiring additional online storage space.

The Bernoulli compound models all have similarly shaped cumulative distribution functions. Figure 1 is a sample distribution function for the compound Bernoulli-lognormal distribution with a Bernoulli factor of  $p$  equal to 0.6, a mean of 1.0 and a standard deviation of 0.5. Note that the distribution has a mass of zero for negative observations, a mass of  $1-p$  at zero and the usual cumulative distribution function shape for positive observations but of mass  $p$  vice mass one.

#### B. MODEL DESCRIPTIONS

Three compound Bernoulli models were formulated for evaluation. Each model was derived from the base distribution after compounding with the Bernoulli process. An additional requirement was made that the inverse of the compound model must be computationally relatively simple.

##### 1. Bernoulli-exponential Distribution

This distribution was derived in Reference 1 and the derivation will not be repeated here. Equations (1) and (2)




Figure 1: Compound Cumulative Distribution Function,  
Bernoulli-lognormal Distribution

are the probability density and complementary cumulative distribution functions respectively. The Bernoulli-exponential inverse complementary cumulative distribution function is computed as:

$$x = \begin{cases} 0 & , H(x) \geq p \\ \mu[\ln(p) - \ln(H(x))] & , H(x) < p \end{cases} \quad (6)$$

The parameters of this distribution are estimated from the sample data as follows:

$\hat{p}$  = number of non-zero demand observations in sample  
divided by sample size,

$\hat{\mu} = \bar{x} / \hat{p}$ , where  $\bar{x}$  is the entire sample mean,

or

$\hat{\mu} = \bar{x}$ , where  $\bar{x}$  is the sample mean of only the non-zero  
demand observations.

## 2. Bernoulli-lognormal Distribution

The lognormal distribution was chosen as a candidate for testing because of several desirable properties:

- a. It is defined for all real numbers greater than zero,
- b. Its inverse is readily computed using inverse normal approximations that are well documented for accuracy,
- c. Alone, the lognormal distribution does not allow the case of zero demand, but when combined with the Bernoulli distribution, the compound Bernoulli-lognormal distribution defines all the expected demand values greater than or equal to zero.

Utilizing the Bernoulli parameter  $p$  as a scaling factor for the lognormal density function, the compound Bernoulli-lognormal density function may be expressed as:

$$h(x) = \begin{cases} 1 - p & , x = 0 \\ p\phi\left(\frac{\ln(x) - \mu}{\sigma}\right), & x > 0 \end{cases} \quad (7)$$

where:

$p$  = probability of a non-zero demand,

$\mu$  = mean of the natural log of all demands greater than zero,

$\sigma$  = Standard deviation of the natural log of all demands greater than zero,

$\phi$  = Standard normal density function.

Integrating (7) results in the complementary cumulative distribution function given by:

$$p, x = 0 \quad (8)$$

$$H(x) = p \left[ 1 - \phi \left( \frac{\ln(x) - \mu}{\sigma} \right) \right], x > 0$$

where:

$\phi$  = Standard normal cumulative distribution function.

The Bernoulli-lognormal inverse cumulative distribution function is easily derived from (8) as:

$$x = \begin{cases} 0 & , H(x) \geq p \\ \exp[\mu + \sigma \phi^{-1}(1 - H(x)/p)], & H(x) < p \end{cases} \quad (9)$$

The parameters are estimated from the sample data in the usual manner:

$\hat{p}$  = number of non-zero demand observations in sample divided by sample size,

$\hat{\mu} = \bar{x}$ , where  $\bar{x}$  is the sample mean of the natural log of all non-zero demand observations,  
 $\hat{\sigma} = s$ , where  $s$  is the sample standard deviation of the natural log of all non-zero demand observations. An alternate method for estimating the mean and standard deviation of the lognormal distribution is to use the method of moments. The procedure in effect, transforms the mean and standard deviation obtained from the untransformed sample data vice transforming the data first and then computing the sample mean and standard deviation. This method involves fewer logarithms and will run faster on a computer. The procedure is described in Appendix D.

### 3. Bernoulli-logistic Distribution

The logistic distribution is a pseudo-normal distribution that is similar in shape to the normal, but is easier to handle mathematically. Its density function is defined as [Ref. 6]:

$$f(x) = \frac{\pi}{4\sqrt{3}\sigma} \operatorname{sech}^2\left(\frac{\pi(x - \mu)}{2\sqrt{3}\sigma}\right), -\infty < x < \infty, \quad (10)$$

while its cumulative distribution function is defined as:

$$F(x) = \frac{1}{2} \left[ 1 + \tanh\left(\frac{\pi(x - \mu)}{2\sqrt{3}\sigma}\right) \right], -\infty < x < \infty, \quad (11)$$

where:

$\mu$  = mean of the entire sample,

$\sigma$  = standard deviation of the entire sample.

With the logistic distribution, it is no longer a simple matter of scaling the distribution by the Bernoulli  $p$  parameter, since this distribution is defined over the entire real line and not just on the positive half. Consequently, the distribution requires a new constant of integration to replace the  $1/2$  used in (11). Integrating (10) from 0 to  $\infty$  provides the new constant as:

$$\int_0^\infty \operatorname{sech}^2\left(\frac{\pi(x - \mu)}{2\sqrt{3}\sigma}\right) dx = 1 + \tanh\left(\frac{\pi\mu}{2\sqrt{3}\sigma}\right) \quad (12)$$

which, when compounded with the Bernoulli distribution, gives the Bernculli-logistic distribution density function as:

$$h(x) = \begin{cases} 1 - p & , x = 0 \\ p \frac{1}{\left[1 + \tanh\left(\frac{\pi\mu}{2\sqrt{3}\sigma}\right)\right]} \operatorname{sech}^2\left(\frac{\pi(x - \mu)}{2\sqrt{3}\sigma}\right) & , x > 0 \\ 0 & , \text{otherwise,} \end{cases} \quad (13)$$

where:

$\mu$  = mean of all non-zero demand observations,

$\sigma$  = standard deviation of all non-zero demand observations.

After integrating (13), the complementary cumulative distribution function may be written as:

$$H(x) = p \left\{ 1 - \frac{1}{\left[ 1 + \tanh\left(\frac{\pi\mu}{2\sqrt{3}\sigma}\right) \right]} \right\} \left[ \tanh\left(\frac{\pi\mu}{2\sqrt{3}\sigma}\right) + \tanh\left(\frac{\pi(x-\mu)}{2\sqrt{3}\sigma}\right) \right] \quad (14)$$

for  $x \geq 0$

The inverse cumulative distribution is:

$$x = \begin{cases} 0 & , H(x) \geq p \\ \mu + \frac{\sqrt{3}\sigma}{\pi} \ln \left\{ \frac{2p}{H(x) \left[ 1 + \tanh\left(\frac{\pi\mu}{2\sqrt{3}\sigma}\right) \right] - 1} \right\} & , H(x) < p \end{cases} \quad (15)$$

The parameters are estimated from the sample data in the usual manner:

$\hat{p}$  = number of non-zero demand observations in sample divided by sample size,

$\hat{\mu} = \bar{x}$ , where  $\bar{x}$  is the sample mean of all non-zero demand observations,

$\hat{\sigma} = s$ , where  $s$  is the sample standard deviation of all non-zero demand observations.

#### IV. EVALUATION PROCEDURE

##### A. THE DATA

In order to test the validity of the proposed models, samples of actual demand were obtained from the Operations Analysis Department at FMSO. The data, accumulated from the demand history files of the Aviation Supply Office, was originally used as input data for the 5A (Aviation Afloat and Ashore Allowance Analyzer) [Ref. 7]. The data consists of 1587 consumable 1R cog items and 2892 non-program-related\* repairable 2R cog items. The information for each item is contained in one master record and several subrecords. The master record contains identifying information on each item such as the national stock number, replacement price, etc. Each subrecord contains up to forty-six demand records, each including the demand quantity and the day of the demand. A complete record layout is contained in Appendix E. To ease processing, the Julian Dates of the original demands were replaced by FMSO with a sequential date ranging from 1 to 1500 representing approximately four years of available history.

-----

\*See Appendix A for definition of "program related" items.

For this application, additional screening was conducted to group the demands into thirty-day buckets, thus providing a demand time series of forty-eight observations. Further editing was required prior to analysis to remove the negative demands noted for many of the items. It was assumed that the negative observations were the result of cancellations of previous demands which were never filled. Therefore, whenever a negative demand was encountered, the demand series was searched backwards for the first positive demand equal to or larger than the negative demand. If such a demand was found, the positive demand quantity was reduced by the absolute value of the negative quantity and the negative quantity was set equal to zero. If no offsetting positive demand was found, the negative demand quantity was still set equal to zero. The resulting edited demand time series was used as the base for all further analysis. Two additional demand series were created from the edited series:

1. A demand series of only the non-zero demands,
2. A demand series of the log of the non-zero demands.

The sample mean and sample standard deviation for each of the three demand series were computed<sup>10</sup> and the group, representing one line item, was accepted for further processing if:

-----

<sup>10</sup>Stationarity was assumed for the demand time series under study. If a trend were in fact present, the resampling technique effectively would eliminate it by shuffling the observed demands into random order.

1. There were at least two non-zero demands, and
2. The standard deviation of all three series was non-zero.

Once accepted, the sample mean and/or sample standard deviation of the appropriate series was used to estimate the value of  $x$ , such that for a given probability, demand will not exceed that value of  $x$ . The values of  $x$  were computed for each model tested at several probabilities from the inverse cumulative probability functions. Table 2 displays the sample mean and/or sample standard deviation required for each model investigated.

#### B. THE RESAMPLING PROCEDURE

The resampling procedure used was chosen over the more traditional methods discussed in Chapter II because it provides a method of comparing a theoretical distribution to the sample data at specified probabilities and is applicable in cases such as this where there is a paucity of data available for analysis. This procedure is similar to the "bootstrap" procedure used by Efron [Ref. 8]. The idea behind this procedure is to randomly sample with replacement the series of available data to create additional pseudo-samples that possess the same statistical properties as the original sample. The desired statistical property can then be estimated from

Table 2: Required Data for Inverse Computations

<u>MODEL</u>	<u>MEAN</u>	<u>STANDARD DEVIATION</u>
1. Exponential	A	N/R
2. Normal	A	A
3. Poisson	A	N/R
4. Negative Binomial	A	A
5. Lognormal	C	C
6. Logistic	A	A
7. La Place	A	A
8. Bernoulli-exponential	B	N/R
9. Bernoulli-lognormal	C	C
10. Bernoulli-logistic	B	B

Where:

A is from edited demand series

B is from non-zero demand series

C is from log of non-zero demand series

N/R indicates parameter Not Required for this model

each pseudo sample. With a sufficient number of repetitions, the distribution of the estimated property is known to be normal from the central limit theorem with mean equal to the population mean and standard deviation equal to the population standard deviation divided by the number of repetitions.

Thus by the use of the resampling procedure, it is possible to study theoretical distributions at points in their

right hand tail area, which is the region of most interest in computing safety levels in inventory models. The resampling procedure, as applied in this study, is a relatively straightforward application of the following technique:

1. The edited series for each sample of  $n$  demand observations is treated as the population from which the random samples are drawn.
2. A random sample of size  $n$  is drawn from this population, creating a pseudo sample. The pseudo sample is not a permutation of the original population since the sample values are selected with replacement from the original population. This is easily accomplished in a computer by generating  $n$  uniform random integers in the range from 1 to  $n$ , and using these numbers as subscripts to select the sample from the original population.
3. The pseudo sample percentile is computed for each repetition from (16) and is compared to the theoretical percentile.

$$\hat{p} = (\text{number of demand observations } \leq x) / n \quad (16)$$

The various values of  $x$  are computed from the inverse cumulative probability functions for each model evaluated.

Under the null hypothesis that the sample demands come from a particular probability distribution, the expected value of  $\hat{\rho}$  should equal  $\rho$ , the theoretical percentile:

$$E[\hat{\rho}] = \rho \text{ under } H_0 \quad (17)$$

and the distribution of  $\rho$  should be  $\text{Normal}(\rho, \sigma^2/n)$ . The percentile  $\rho$  is estimated as above and  $\sigma^2$  is estimated from standard binomial results as:

$$\text{Var}[\hat{\rho}] = \hat{\rho}(1 - \hat{\rho}) / n. \quad (18)$$

### C. MEASURES OF EFFECTIVENESS

When the above procedure is executed at several theoretical percentiles, the fit of the sample data may be compared at several locations in the theoretical distribution and the overall fit evaluated. Several common measures of effectiveness are available to provide a criteria on which to judge the success or failure of a model. Some of those available are:

1. Maximum absolute deviation,
2. Mean squared error,
3. Algebraic sum of errors.

Method three was eliminated since errors may offset one another resulting in a seemingly good fit, but in reality a

very poor fit. Method one is potentially a good way to evaluate error, but is generally not as sensitive as method two. Therefore, the mean squared error of the pseudo sample percentile estimates was chosen as the measure of effectiveness of choice and was accumulated as:

$$MSE = \frac{1}{n} \sum_{i=1}^n (\hat{\rho}_i - \rho_i)^2. \quad (19)$$

#### D. COMPUTER VALIDATION

The computer programs for this analysis were written in FORTRAN IV for execution on the IBM 3033 attached processor (System 370) computer installed at the Naval Postgraduate School. The programs are batch oriented since the volume of data available prevented storage directly on the user's private disk space. The data tapes received from FMSO were loaded onto the system mass storage device and when required, all or portions of the data were transferred to the Virtual Machine (VM) facility for processing. Two programs were written, the first to copy the data to make it accessible to the VM system and the second to conduct the actual analysis. A complete listing of the FORTRAN source code is available in Appendix F. Each program or subprogram contains a documentation block at the top that provides a complete description of

the program's purpose, a definition of variables used and a list of user-written subroutines and functions required.

The random number generator used is a part of the LLRANDOM II series [Ref. 9] developed at the Naval Postgraduate School. The programs and subprograms were thoroughly tested both independently and as a unit using known data. The results were verified by hand held calculator and in all cases agreed with the program output.

## V. RESULTS

The results of the distributional analysis conducted on the ten distribution models listed in Table 2 are tabulated in Appendix B for 1R consumable items. Appendix B is divided into the three demand classes listed in Table 1, and then each class is divided by distribution and percentile within each distribution. The results for the 2R repairable items are presented in a similar format in Appendix C.

The consistency of the resampling method with varying data was evaluated by testing independent sections of the consumable and repairable data sets and comparing the results. In both cases, independent subsamples were created by sampling every third item but changing the initial item sampled from the first to the third item. The analysis, not presented here, showed that the resampling method produced consistent results over all of the subsamples tested. The total mean square error<sup>11</sup> of the distributions in each subsample varied by no more than 20% from the total mean square error values obtained from the entire sample population as listed in Appendix B and Appendix C. The relative standing

---

<sup>11</sup>Total mean square error is the sum of the mean square errors computed at the 75th, 80th, 85th, 90th and 95th percentiles.

of each distribution was not altered in any of the subsamples.

The effect of the number of pseudo sample repetitions was also studied. The results proved similar while varying the number of repetitions from twenty to fifty. The standard deviation of the percentile estimates decreased as the number of repetitions increased. The total mean square error for each trial varied randomly but remained within 10% of the overall values for each distribution with more repetitions producing consistently smaller variations. For the data analysis production runs, forty was chosen as the number of resampling repetitions as a trade off between computer run time and minimizing variance.

#### A. LOW-DEMAND ITEMS

A summary of the total mean squared error figures for the models providing the best fit or the smallest total mean squared error is provided in Table 3 for the low-demand items. The best models for the low demand items all had very small total mean squared errors. This can be attributed to the very low probability of a demand ever being greater than one and to the integer sampling plan for accumulating the Bernoulli trials. For the integer results of the Poisson and negative binomial distributions, the estimated percentile

TABLE 3: LOW-DEMAND ITEMS,  
Total Mean Squared Error ( $10^{-4}$ )

	<u>Poisson</u>	<u>Negative binomial</u>	<u>Compound</u>
Consumable 1R	5.64	7.86	10.77
Repairable 2R	5.86	7.57	12.91

will closely match the theoretical percentile unless the number of Bernoulli successes is less than the expected number of successes for a given theoretical percentile. This produces a conservative result that will tend to overstate the quantity of stock required by one unit to provide a stated level of protection, but in the case of the small quantities involved, is not an undesirable property. The compound distributions, as a group, gave the same total mean squared error for much the same reasons. The probability of a non-zero demand is very small for the low-demand items, generally less than 0.05, thus these models tend to predict a quantity of zero for all percentiles. The resampling procedure and the sampling plan again provide a conservative result.

#### B. MEDIUM-DEMAND ITEMS

A summary of the total mean squared error figures for the models providing the best fit or the smallest total mean

squared error is provided in Table 4 for the medium-demand items. Here again the predicted demand for the various percentiles is small. The resampling technique still gives the

TABLE 4: MEDIUM-DEMAND ITEMS,  
Total Mean Squared Error ( $10^{-4}$ )

	<u>Negative binomial</u>	<u>Poisson</u>	<u>Bernoulli- lognormal</u>
Consumable 1R	28.55	48.28	80.60
Repairable 2R	29.14	44.84	112.44

edge to the integer distributions. The compound distributions have begun to show some differences among themselves, but as a group are clearly better than any of the continuous simple distributions. The magnitude of the total mean squared error is significantly larger than for the low-demand class primarily due to the larger allowable total demand range defining the medium-demand class.

#### C. HIGH-DEMAND ITEMS

The summary total mean squared error data for the high-demand items are listed in Table 5. With the unlimited range of mean demands in this class, the higher total mean squared errors are expected. The compound models have come into

TABLE 5: HIGH-DEMAND ITEMS,  
Total Mean Squared Error ( $10^{-4}$ )

	<u>Bernoulli-</u> <u>lognormal</u>	<u>Bernoulli-</u> <u>exponential</u>	<u>Exponential</u>
Consumable 1R	168.25	188.90	298.21
Repairable 2R	183.77	221.80	289.01

their own in this class with the Bernoulli-lognormal yielding a total mean squared error less than one half that of the normal distribution. The second best distribution, the Bernoulli-exponential, gives a total mean squared error only slightly higher than the Bernoulli-lognormal.

## VI. CONCLUSIONS AND RECOMMENDATIONS

### A. SUMMARY

This distributional analysis, conducted using a completely different experimental procedure than prior analyses, provides satisfactorily similar results for the low and medium demand categories in showing that the Poisson distribution gives the best fit for the low-demand category of items and that the negative binomial distribution gives the best fit for the medium-demand category of items. The class of compound distributions, as a whole, gave good fits in the low- and medium-demand categories, though the total mean squared error was two to four times as large as that of the Poisson or negative binomial distribution.

The analysis indicated that there are several distributions which provide better results than the normal distribution for high-demand items. The Bernoulli-lognormal distribution consistently provided total mean squared errors approximately one half that of the normal distribution, indicating a superior fit in the right-hand tail area of the distribution. Other distributions giving good fits were the Bernoulli-exponential and the standard exponential, both yielding total mean squared errors less than that of the normal distribution.

## B. RECOMMENDATIONS

The results of this study indicate the Navy should continue to use the Poisson and negative binomial distributions as the models for the low- and medium-demand classes respectively. Specifically, the Aviation Supply Office should re-evaluate its position on the use of the normal distribution for all demand classes and revert to the Poisson and negative binomial distributions for low- and medium-demand classes as before. The normal distribution tends to inflate the stock required for low- and medium-demand items for a specified level of protection resulting in increased safety levels and excessive dollar investment.

The Navy should consider replacing the normal distribution model used for high-demand items with the compound Bernoulli-lognormal distribution. If the Bernoulli-lognormal distribution is considered computationally too difficult for a large inventory system, the Bernoulli-exponential could be used in its place with little loss of effectiveness.

## C. ADDITIONAL RESEARCH

Follow on research in the following areas may improve and expand upon the results presented above:

1. Optimize the divisions of demand categories (i.e. low, medium, high) with the possible use of the probability

- of a non-zero demand as an element in the determination,
2. Implement the Bernoulli-lognormal model in FMSO's SA simulator to establish the effect of the change on the entire supply system,
  3. Investigate the implications of the assumption that the demand process is stationary,
  4. Evaluate other compound distributions, such as the Bernoulli-log-logistic, which is particularly appealing analytically.

## APPENDIX A

### GLOSSARY OF TERMS

**Program Related Item:** An item of stock whose demand can be predicted from the value of a specific Navy program, i.e., flying hours, steaming hours, etc.

**Reorder Point:** The on hand stock quantity that when reached, triggers an order for replenishment of stock material. The reorder point is the expected demand during procurement lead-time plus the safety level.

**Risk:** Probability of a stockout during leadtime.

**Safety Level:** The quantity of material which is required to be on hand to permit continued operation in the event of minor interruptions on normal replenishment or unpredictable fluctuations in demand. The safety level determined is structured so as to minimize time-weighted, essentiality-weighted requisitions short.

**Stockout:** A condition that exists when the on hand inventory is insufficient to fill the current demand requirements.

**System Performance:** A subjective measure maximized when the time-weighted, essentiality-weighted requisitions short is minimized. Time weighting is the consideration of the average number of days delay in the availability of material, essentiality-weighting is the consideration of the relative essentiality of each item. Requisitions short are requisitions for which material is not available.

## APPENDIX\_B

### CONSUMABLE ITEM TABULATED RESULTS

40 REPETITIONS FOR EACH OF 61 PSEUDO SAMPLES

LOW DEMAND RANGE:  $0.0 < D < 1.0$  PER YEAR

-----  
DISTRIBUTION: POISSON

#### THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.750000	0.812500	0.854006	0.913924	0.956069
STD DEV	0.008766	0.007902	0.007148	0.005678	0.004149
MSE	0.0	0.000156	0.000018	0.000290	0.000100
TOTAL MSE:	5.64E-04				

-----

DISTRIBUTION: NEGATIVE BINOMIAL

#### THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.750000	0.812500	0.854006	0.913924	0.948994
STD DEV	0.008766	0.007902	0.007148	0.005678	0.004454
MSE	0.0	0.000156	0.000018	0.000290	0.000322
TOTAL MSE:	7.86E-04				

40 REPETITIONS FOR EACH OF 61 PSEUDO SAMPLES

LOW DEMAND RANGE:  $0.0 < D < 1.0$  PER YEAR

---

DISTRIBUTION: NORMAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.953221	0.953221	0.953221	0.953221	0.953221
STD DEV	0.004275	0.004275	0.004275	0.004275	0.004275
MSE	0.042319	0.024488	0.011662	0.003835	0.001006
TOTAL MSE:	833.09E-04				

---

DISTRIBUTION: EXPONENTIAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.953221	0.953221	0.953221	0.953221	0.953221
STD DEV	0.004275	0.004275	0.004275	0.004275	0.004275
MSE	0.042319	0.024488	0.011662	0.003835	0.001006
TOTAL MSE:	833.09E-04				

---

DISTRIBUTION: LOGNORMAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.978404	0.995246	0.995246	1.000000	1.000000
STD DEV	0.002943	0.001392	0.001392	0.0	0.0
MSE	0.052629	0.038292	0.021267	0.009998	0.002500
TOTAL MSE:	1246.85E-04				

40 REPETITIONS FOR EACH OF 61 PSEUDO SAMPLES

LOW DEMAND RANGE:  $0.0 < D < 1.0$  PER YEAR

---

DISTRIBUTION: LAPLACE

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.953221	0.953221	0.953221	0.953221	0.953221
STD DEV	0.004275	0.004275	0.004275	0.004275	0.004275
MSE	0.042319	0.024488	0.011662	0.003835	0.001006
TOTAL MSE:	833.09E-04				

---

DISTRIBUTION: LOGISTIC

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.953221	0.953221	0.953221	0.953221	0.953221
STD DEV	0.004275	0.004275	0.004275	0.004275	0.004275
MSE	0.042319	0.024488	0.011662	0.003835	0.001006
TOTAL MSE:	833.09E-04				

---

DISTRIBUTION: BERNOULLI-EXPONENTIAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.750000	0.812500	0.854006	0.913924	0.944939
STD DEV	0.008766	0.007902	0.007148	0.005678	0.004618
MSE	0.0	0.000156	0.000018	0.000290	0.000613
TOTAL MSE:	10.77E-04				

40 REPETITIONS FOR EACH OF 61 PSEUDO SAMPLES

LOW DEMAND RANGE: 0.0 < D < 1.0 PER YEAR

---

DISTRIBUTION: BERNOULLI-LOGNORMAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.750000	0.812500	0.854006	0.913924	0.944939
STD DEV	0.008766	0.007902	0.007148	0.005678	0.004618
MSE	0.0	0.000156	0.000018	0.000290	0.000613

TOTAL MSE: 10.77E-04

DISTRIBUTION: BERNOULLI-LOGISTIC

---

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.750000	0.812500	0.854006	0.913924	0.944939
STD DEV	0.008766	0.007902	0.007148	0.005678	0.004618
MSE	0.0	0.000156	0.000018	0.000290	0.000613

TOTAL MSE: 10.77E-04

40 REPETITIONS FOR EACH OF 753 PSEUDO SAMPLES

MEDIUM DEMAND RANGE:  $1.0 < D < 20.0$  PER YEAR

---

DISTRIBUTION: POISSON

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.748131	0.807719	0.843932	0.893923	0.925303
STD DEV	0.002501	0.002271	0.002091	0.001774	0.001515
MSE	0.000138	0.000398	0.000603	0.001423	0.002266
TOTAL MSE:	48.28E-04				

---

DISTRIBUTION: NEGATIVE BINOMIAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.746099	0.805430	0.843400	0.900210	0.943628
STD DEV	0.002508	0.002281	0.002094	0.001727	0.001329
MSE	0.000332	0.000522	0.000589	0.000807	0.000606
TOTAL MSE:	28.55E-04				

---

DISTRIBUTION: NORMAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.888030	0.896699	0.907882	0.920152	0.935573
STD DEV	0.001817	0.001754	0.001666	0.001562	0.001415
MSE	0.024771	0.014168	0.007327	0.003509	0.002400
TOTAL MSE:	521.74E-04				

40 REPETITIONS FOR EACH OF 753 PSEUDO SAMPLES

MEDIUM DEMAND RANGE:  $1.0 < D < 20.0$  PER YEAR

---

DISTRIBUTION: EXPONENTIAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.852981	0.861003	0.871591	0.884311	0.898352
STD DEV	0.002040	0.001993	0.001928	0.001843	0.001741
MSE	0.020357	0.011779	0.006819	0.004953	0.006210
TOTAL MSE:	501.19E-04				

---

DISTRIBUTION: LOGNORMAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.953018	0.962552	0.970638	0.980623	0.990731
STD DEV	0.001219	0.001094	0.000973	0.000794	0.000552
MSE	0.043846	0.028367	0.015977	0.007411	0.002040
TOTAL MSE:	976.41E-04				

---

DISTRIBUTION: LAPLACE

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.884254	0.893767	0.904733	0.918110	0.936078
STD DEV	0.001843	0.001775	0.001692	0.001580	0.001409
MSE	0.024079	0.013865	0.007191	0.003573	0.002363
TOTAL MSE:	510.72E-04				

40 REPETITIONS FOR EACH OF 753 PSEUDO SAMPLES

MEDIUM DEMAND RANGE:  $1.0 < D < 20.0$  PER YEAR

---

DISTRIBUTION: LOGISTIC

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.875036	0.886687	0.896908	0.913037	0.934834
STD DEV	0.001905	0.001826	0.001752	0.001624	0.001422
MSE	0.022871	0.013193	0.006955	0.003690	0.002447
TOTAL MSE:	491.56E-04				

---

DISTRIBUTION: BERNOULLI-EXPONENTIAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.738072	0.795951	0.840430	0.895563	0.950685
STD DEV	0.002533	0.002322	0.002110	0.001755	0.001248
MSE	0.002132	0.002338	0.002267	0.002224	0.001327
TOTAL MSE:	102.88E-04				

---

DISTRIBUTION: BERNOULLI-LOGNORMAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.749344	0.809159	0.849391	0.903886	0.948263
STD DEV	0.002497	0.002264	0.002061	0.001698	0.001276
MSE	0.001456	0.001751	0.001730	0.001866	0.001258
TOTAL MSE:	80.60E-04				

40 REPETITIONS FOR EACH OF 753 PSEUDO SAMPLES

MEDIUM DEMAND RANGE:  $1.0 < D < 20.0$  PER YEAR

---

DISTRIBUTION: BERNOULLI-LOGISTIC

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.760006	0.821542	0.864176	0.916219	0.955416
STD DEV	0.002461	0.002206	0.001974	0.001596	0.001189
MSE	0.002256	0.002499	0.002304	0.002030	0.001241
TOTAL MSE:	103.30E-04				

40 REPETITIONS FOR EACH OF 429 PSEUDO SAMPLES

HIGH DEMAND RANGE:  $20.0 < D$  PER YEAR

---

DISTRIBUTION: POISSON

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.689018	0.730300	0.753320	0.780038	0.800697
STD DEV	0.003534	0.003388	0.003291	0.003162	0.003050
MSE	0.024843	0.027146	0.033935	0.042154	0.051388
TOTAL MSE:	1794.66E-04				

---

DISTRIBUTION: NEGATIVE BINOMIAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.724173	0.747130	0.791313	0.848041	0.896526
STD DEV	0.003412	0.003318	0.003102	0.002740	0.002325
MSE	0.003047	0.024647	0.026816	0.029773	0.031637
TOTAL MSE:	1159.20E-04				

---

DISTRIBUTION: NORMAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.850264	0.871100	0.890154	0.910670	0.934248
STD DEV	0.002724	0.002558	0.002387	0.002177	0.001892
MSE	0.016622	0.010312	0.005815	0.003155	0.002227
TOTAL MSE:	381.32E-04				

40 REPETITIONS FOR EACH OF 429 PSEUDO SAMPLES

HIGH DEMAND RANGE: 20.0 < D PER YEAR

---

DISTRIBUTION: EXPONENTIAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.784610	0.814275	0.846685	0.881207	0.918702
STD DEV	0.003138	0.002969	0.002750	0.002470	0.002086
MSE	0.009534	0.006885	0.005219	0.004384	0.003800
TOTAL MSE:	298.21E-04				

---

DISTRIBUTION: LOGNORMAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.855867	0.886448	0.918614	0.949144	0.978288
STD DEV	0.002681	0.002422	0.002087	0.001677	0.001113
MSE	0.019028	0.013031	0.008113	0.004389	0.001639
TOTAL MSE:	462.00E-04				

---

DISTRIBUTION: LAPLACE

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.843416	0.863600	0.885612	0.907473	0.934956
STD DEV	0.002774	0.002620	0.002430	0.002212	0.001883
MSE	0.015775	0.009709	0.005680	0.003248	0.002167
TOTAL MSE:	365.79E-04				

40 REPETITIONS FOR EACH OF 429 PSEUDO SAMPLES

HIGH DEMAND RANGE:  $20.0 < D$  PER YEAR

---

DISTRIBUTION: LOGISTIC

THEORETICAL PERCENTILES (P)

ESTIMATES:	0.75	0.80	0.85	0.90	0.95
MEAN	0.822132	0.846791	0.871709	0.900110	0.933755
STD DEV	0.002919	0.002750	0.002553	0.002289	0.001899
MSE	0.013721	0.008841	0.005633	0.003523	0.002259
TOTAL MSE:	339.76E-04				

---

DISTRIBUTION: BERNOULLI-EXPONENTIAL

THEORETICAL PERCENTILES (P)

ESTIMATES:	0.75	0.80	0.85	0.90	0.95
MEAN	0.759754	0.816600	0.867492	0.914225	0.955101
STD DEV	0.003261	0.002954	0.002588	0.002138	0.001581
MSE	0.005828	0.005034	0.003977	0.002618	0.001432
TOTAL MSE:	188.90E-04				

---

DISTRIBUTION: BERNOULLI-LOGNORMAL

THEORETICAL PERCENTILES (P)

ESTIMATES:	0.75	0.80	0.85	0.90	0.95
MEAN	0.745078	0.797614	0.847684	0.901963	0.953670
STD DEV	0.003327	0.003067	0.002743	0.002270	0.001605
MSE	0.004768	0.004403	0.003632	0.002613	0.001409
TOTAL MSE:	168.25E-04				

40 REPETITIONS FOR EACH OF 429 PSEUDO SAMPLES

HIGH DEMAND RANGE: 20.0 < D PER YEAR

---

DISTRIBUTION: BERNOULLI-LOGISTIC

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.828539	0.864722	0.892571	0.921326	0.947036
STD DEV	0.002877	0.002611	0.002364	0.002055	0.001710
MSE	0.012517	0.008862	0.005485	0.003020	0.001538
TOTAL MSE:	314.23E-04				

## APPENDIX C

### REPAIRABLE ITEM TABULATED RESULTS

40 REPETITIONS FOR EACH OF 137 PSEUDO SAMPLES

LOW DEMAND RANGE:  $0.0 < D < 1.0$  PER YEAR

---

DISTRIBUTION: POISSON

#### THEORETICAL PERCENTILES (P)

ESTIMATES:	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.750000	0.812496	0.853613	0.912594	0.955936
STD DEV	0.005849	0.005273	0.004775	0.003815	0.002772
MSE	0.0	0.000156	0.000019	0.000311	0.000099
TOTAL MSE:	5.86E-04				

---

DISTRIBUTION: NEGATIVE BINOMIAL

#### THEORETICAL PERCENTILES (P)

ESTIMATES:	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.750000	0.812496	0.853613	0.912594	0.950753
STD DEV	0.005849	0.005273	0.004775	0.003815	0.002923
MSE	0.0	0.000156	0.000019	0.000311	0.000270
TOTAL MSE:	7.57E-04				

40 REPETITIONS FOR EACH OF 137 PSEUDO SAMPLES

LOW DEMAND RANGE:  $0.0 < D < 1.0$  PER YEAR

---

DISTRIBUTION: NORMAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.949164	0.949164	0.949164	0.949164	0.949164
STD DEV	0.002967	0.002967	0.002967	0.002967	0.002967
MSE	0.040918	0.023465	0.011016	0.003568	0.001118
TOTAL MSE:	800.84E-04				

---

DISTRIBUTION: EXPONENTIAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.949164	0.949164	0.949164	0.949164	0.949164
STD DEV	0.002967	0.002967	0.002967	0.002967	0.002967
MSE	0.040918	0.023465	0.011016	0.003568	0.001118
TOTAL MSE:	800.84E-04				

---

DISTRIBUTION: LOGNORMAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.978874	0.991028	0.991028	1.000000	1.000000
STD DEV	0.001943	0.001274	0.001274	0.0	0.0
MSE	0.052955	0.036815	0.020201	0.009996	0.002500
TOTAL MSE:	1224.66E-04				

40 REPETITIONS FOR EACH OF 137 PSEUDO SAMPLES

LOW DEMAND RANGE:  $0.0 < D < 1.0$  PER YEAR

---

DISTRIBUTION: LAPLACE

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.949164	0.949164	0.949164	0.949164	0.949164
STD DEV	0.002967	0.002967	0.002967	0.002967	0.002967
MSE	0.040918	0.023465	0.011016	0.003568	0.001118
TOTAL MSE:	800.84E-04				

---

DISTRIBUTION: LOGISTIC

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.949164	0.949164	0.949164	0.949164	0.949164
STD DEV	0.002967	0.002967	0.002967	0.002967	0.002967
MSE	0.040918	0.023465	0.011016	0.003568	0.001118
TOTAL MSE:	800.84E-04				

---

DISTRIBUTION: BERNOULLI-EXPONENTIAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.750000	0.812496	0.853613	0.912594	0.942635
STD DEV	0.005849	0.005273	0.004775	0.003815	0.003141
MSE	0.0	0.000156	0.000019	0.000311	0.000804
TOTAL MSE:	12.91E-04				

40 REPETITIONS FOR EACH OF 137 PSEUDO SAMPLES

LOW DEMAND RANGE:  $0.0 < D < 1.0$  PER YEAR

---

DISTRIBUTION: BERNOULLI-LOGNORMAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.750000	0.812496	0.853613	0.912594	0.942635
STD DEV	0.005849	0.005273	0.004775	0.003815	0.003141
MSE	0.0	0.000156	0.000019	0.000311	0.000804
TOTAL MSE:	12.91E-04				

---

DISTRIBUTION: BERNOULLI-LOGISTIC

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.750000	0.812496	0.853613	0.912594	0.942635
STD DEV	0.005849	0.005273	0.004775	0.003815	0.003141
MSE	0.0	0.000156	0.000019	0.000311	0.000804
TOTAL MSE:	12.91E-04				

40 REPETITIONS FOR EACH OF 1487 PSEUDO SAMPLES

MEDIUM DEMAND RANGE:  $1.0 < D < 20.0$  PER YEAR

---

DISTRIBUTION: POISSON

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.747034	0.806323	0.843928	0.894732	0.929457
STD DEV	0.001782	0.001620	0.001488	0.001258	0.001050
MSE	0.000238	0.000507	0.000582	0.001321	0.001836
TOTAL MSE:	44.84E-04				

---

DISTRIBUTION: NEGATIVE BINOMIAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.744688	0.804450	0.843869	0.900645	0.944953
STD DEV	0.001788	0.001626	0.001488	0.001227	0.000935
MSE	0.000444	0.000598	0.000533	0.000788	0.000551
TOTAL MSE:	29.14E-04				

---

DISTRIBUTION: NORMAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.861474	0.875334	0.890316	0.905783	0.927155
STD DEV	0.001416	0.001354	0.001281	0.001198	0.001066
MSE	0.019444	0.011363	0.006084	0.003600	0.002944
TOTAL MSE:	434.35E-04				

40 REPETITIONS FOR EACH OF 1487 PSEUDO SAMPLES

MEDIUM DEMAND RANGE:  $1.0 < D < 20.0$  PER YEAR

---

DISTRIBUTION: EXPONENTIAL

THEORETICAL PERCENTILES (P)

ESTIMATES:	0.75	0.80	0.85	0.90	0.95
MEAN	0.823217	0.834022	0.845365	0.864271	0.886389
STD DEV	0.001564	0.001526	0.001482	0.001404	0.001301
MSE	0.016435	0.010140	0.007542	0.006691	0.007993
TOTAL MSE:	488.02E-04				

---

DISTRIBUTION: LOGNORMAL

THEORETICAL PERCENTILES (P)

ESTIMATES:	0.75	0.80	0.85	0.90	0.95
MEAN	0.946618	0.955351	0.964288	0.975138	0.987135
STD DEV	0.000922	0.000847	0.000761	0.000638	0.000462
MSE	0.041761	0.026454	0.014803	0.006692	0.001890
TOTAL MSE:	916.00E-04				

---

DISTRIBUTION: LAPLACE

THEORETICAL PERCENTILES (P)

ESTIMATES:	0.75	0.80	0.85	0.90	0.95
MEAN	0.856232	0.870161	0.885855	0.902481	0.928262
STD DEV	0.001439	0.001378	0.001304	0.001216	0.001058
MSE	0.018823	0.011026	0.006113	0.003736	0.002851
TOTAL MSE:	425.48E-04				

40 REPETITIONS FOR EACH OF 1487 PSEUDO SAMPLES

MEDIUM DEMAND RANGE:  $1.0 < D < 20.0$  PER YEAR

---

DISTRIBUTION: LOGISTIC

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.845409	0.858922	0.875792	0.896303	0.925823
STD DEV	0.001482	0.001427	0.001352	0.001250	0.001075
MSE	0.017690	0.010603	0.006269	0.004086	0.003093
TOTAL MSE:	417.42E-04				

---

DISTRIBUTION: BERNOULLI-EXPONENTIAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.730065	0.787917	0.834508	0.893026	0.954267
STD DEV	0.001820	0.001676	0.001524	0.001267	0.000857
MSE	0.003437	0.003608	0.003449	0.003021	0.001465
TOTAL MSE:	149.79E-04				

---

DISTRIBUTION: BERNOULLI-LOGNORMAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.750179	0.808666	0.852134	0.904194	0.949477
STD DEV	0.001775	0.001613	0.001455	0.001207	0.000898
MSE	0.002226	0.002681	0.002600	0.002327	0.001410
TOTAL MSE:	112.44E-04				

40 REPETITIONS FOR EACH OF 1487 PSEUDO SAMPLES

MEDIUM DEMAND RANGE:  $1.0 < D < 20.0$  PER YEAR

---

DISTRIBUTION: BERNOULLI-LOGISTIC

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.762406	0.822694	0.866844	0.915851	0.953988
STD DEV	0.001745	0.001566	0.001393	0.001138	0.000859
MSE	0.003183	0.003523	0.003107	0.002468	0.001379
TOTAL MSE:	136.61E-04				

40 REPETITIONS FOR EACH OF 356 PSEUDO SAMPLES

HIGH DEMAND RANGE: 20.0 < D PER YEAR

---

DISTRIBUTION: POISSON

THEORETICAL PERCENTILES (P)

ESTIMATES:	0.75	0.80	0.85	0.90	0.95
MEAN	0.695829	0.735554	0.762330	0.791447	0.821025
STD DEV	0.003855	0.003696	0.003567	0.003405	0.003212
MSE	0.022812	0.026569	0.032141	0.038638	0.044655
TOTAL MSE:	1648.16E-04				

---

DISTRIBUTION: NEGATIVE BINOMIAL

THEORETICAL PERCENTILES (P)

ESTIMATES:	0.75	0.80	0.85	0.90	0.95
MEAN	0.722861	0.755975	0.799852	0.857560	0.904930
STD DEV	0.003751	0.003599	0.003353	0.002929	0.002458
MSE	0.005285	0.024724	0.026795	0.028699	0.030113
TOTAL MSE:	1156.16E-04				

---

DISTRIBUTION: NORMAL

THEORETICAL PERCENTILES (P)

ESTIMATES:	0.75	0.80	0.85	0.90	0.95
MEAN	0.836334	0.859810	0.883941	0.905503	0.932035
STD DEV	0.003100	0.002909	0.002684	0.002451	0.002109
MSE	0.014669	0.009278	0.005429	0.003190	0.002319
TOTAL MSE:	348.85E-04				

40 REPETITIONS FOR EACH OF 356 PSEUDO SAMPLES

HIGH DEMAND RANGE:  $20.0 < D \text{ PER YEAR}$

---

DISTRIBUTION: EXPONENTIAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.780940	0.818187	0.856974	0.898153	0.935255
STD DEV	0.003466	0.003232	0.002934	0.002535	0.002062
MSE	0.009578	0.007296	0.005229	0.003830	0.002968
TOTAL MSE:	289.01E-04				

---

DISTRIBUTION: LOGNORMAL

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.819119	0.855337	0.895522	0.933281	0.971445
STD DEV	0.003226	0.002948	0.002563	0.002091	0.001396
MSE	0.013003	0.008704	0.005985	0.003420	0.001454
TOTAL MSE:	325.65E-04				

---

DISTRIBUTION: LAPLACE

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.827652	0.852088	0.877334	0.902360	0.932870
STD DEV	0.003165	0.002975	0.002749	0.002487	0.002097
MSE	0.013659	0.008798	0.005346	0.003302	0.002267
TOTAL MSE:	333.71E-04				

40 REPETITIONS FOR EACH OF 356 PSEUDO SAMPLES

HIGH DEMAND RANGE: 20.0 < D PER YEAR

---

DISTRIBUTION: LOGISTIC

THEORETICAL PERCENTILES (P)

ESTIMATES:	0.75	0.80	0.85	0.90	0.95
MEAN	0.805812	0.833134	0.860882	0.892308	0.930748
STD DEV	0.003315	0.003125	0.002900	0.002598	0.002128
MSE	0.012232	0.008347	0.005666	0.003803	0.002397
TOTAL MSE:	324.44E-04				

---

DISTRIBUTION: BERNOULLI-EXPONENTIAL

THEORETICAL PERCENTILES (P)

ESTIMATES:	0.75	0.80	0.85	0.90	0.95
MEAN	0.770942	0.829521	0.877192	0.921468	0.960362
STD DEV	0.003522	0.003151	0.002750	0.002254	0.001635
MSE	0.007108	0.005916	0.004548	0.003092	0.001516
TOTAL MSE:	221.80E-04				

---

DISTRIBUTION: BERNOULLI-LOGNORMAL

THEORETICAL PERCENTILES (P)

ESTIMATES:	0.75	0.80	0.85	0.90	0.95
MEAN	0.746108	0.797150	0.847197	0.901039	0.953617
STD DEV	0.003647	0.003370	0.003015	0.002502	0.001762
MSE	0.005655	0.004764	0.003680	0.002809	0.001469
TOTAL MSE:	183.77E-04				

40 REPETITIONS FOR EACH OF 356 PSEUDO SAMPLES

HIGH DEMAND RANGE:  $20.0 < D$  PER YEAR

---

DISTRIBUTION: BERNOULLI-LOGISTIC

THEORETICAL PERCENTILES (P)

<u>ESTIMATES:</u>	<u>0.75</u>	<u>0.80</u>	<u>0.85</u>	<u>0.90</u>	<u>0.95</u>
MEAN	0.829834	0.861163	0.886755	0.914062	0.941856
STD DEV	0.003149	0.002898	0.002656	0.002349	0.001961
MSE	0.013074	0.008963	0.005495	0.003143	0.001722
TOTAL MSE:	323.98E-04				

## APPENDIX D

### ESTIMATION OF LOGNORMAL MEAN AND VARIANCE FROM UNTRANSFORMED DATA

The usual method of estimating the sample mean and variance of a random sample assumed to come from a lognormal distribution is to take the log transform of the data and then compute the standard method of moments estimates. This is time consuming on a computer as the logarithm must be taken for each observation in the sample. Formulae to compute the mean and variance of the untransformed data from the mean and variance of the transformed data can be derived for the lognormal distribution.

If  $Y$  is distributed normal( $\mu_y, \sigma_y^2$ ), then  $X$  is distributed lognormal if  $Y = \ln(X)$ . The mean of  $X$  [Ref. 10] is expressed as:

$$\mu_x = e^{\mu_y + \frac{1}{2}\sigma_y^2} \quad (20)$$

and the variance as:

$$\sigma_x^2 = e^{2(\mu_y + \frac{1}{2}\sigma_y^2)} \left( e^{\sigma_y^2} - 1 \right). \quad (21)$$

Equations (20) and (21) are the reverse of that required in this case since  $\mu_y$  and  $\sigma_y^2$  are obtained from the transformed

data. What is needed then, is to solve for  $\mu_y$  and  $\sigma_y^2$  in terms of  $\mu_x$  and  $\sigma_x^2$ . Substituting (20) into (21) results in:

$$\sigma_x^2 = \mu_x^2 \left( e^{\sigma_y^2} - 1 \right) \quad (22)$$

which when solved for  $e^{\sigma_y^2}$ , taking logs and simplifying, gives the solution of  $\sigma_y^2$  as:

$$\sigma_y^2 = \ln \left( \frac{\sigma_x^2 + \mu_x^2}{\mu_x^2} \right). \quad (23)$$

Substituting (23) back into the log of (20), solving for  $\mu_y$  and simplifying gives the solution of  $\mu_y$  as:

$$\mu_y = \ln \left( \frac{\mu_x^2}{\sqrt{\mu_x^2 + \sigma_x^2}} \right). \quad (24)$$

Thus the mean and variance of the transformed data may be estimated from the sample mean and variance of the untransformed data reducing significantly the computational requirements.

APPENDIX E

DATA-RECORD-LAYOUT

5A Master Input File; 600 Characters Per Record, 10 Records Per Block, Fixed Length

CH	POS	1	2-3 4-12	13-22	23-32	33-42	43-52	53	54
D	RECORD	COGININ	REPLACE-	COST	SET	REPAIR	PROCUREMENT	NEWLY	PROVISIONED
A	TYPE	"1"	MENT	TO	UP	SET UP	METHOD	INDICATOR	INDICATOR
T			PRICE	REPAIR	COST	COST	CODE		(B067A)
A			(B055)	(B055A)	(B058)	(B05A)	(D025E)		
PICTURE	1	2	9	1	8.2	8.2	8.2	1	1

CH	POS	55	56	57	58	59	60
D	DEMAND	HIGH DOLLAR	HIGH REPLACE-	PROGRAM	REPAIRABLE	AUTOMATIC	
A	INDICATOR	DEMAND	MENT PRICE	RELATED	ITEM	REPAIR CYCLE	
T	(B067B)	INDICATOR	INDICATOR	INDICATOR	INDICATOR	INDICATOR	
A		(B067C)	(B067D)	(B067E)	(B067F)	(B067G)	
PICTURE	1	1	1	1	1	1	1

CH POS	61	62-65	1	66-68	1	69-71	1	72-74	1	75-77	1	78-80
D	NAVY REPORTING	PROCURE-		REPAIR-		REPAIR		REPAIR		WEAR-		ITEM
A	REPAIRABLE	MENT	LEAD-	SURVIVAL		SURVIVAL		OUT		ESSEN-		TIALITY
T	INDICATOR	LEADTIME	TIME M.A.D.	RATE		RATE M.A.D.		RATE		(F007)		(C008C)
A	(B067H)	(B011A)	(B011B)	(P009)		(F009A)		(F009)				
PICTURE	1	2.2	2.1	1.2		1.2		1.2		1.2		.3

CH POS	81-83	84-86	1	87-96	1	97-106	1	107-116	1	117-126	1	127-136
D	SHLF	OBSO-	SYSTEM	SYSTEM	SYSTEM	OVER-		OVER-		SYSTEM CAR-		
A	LIFE	LESSENCE	DEMAND	DEMAND	HAUL	DEMAND		DEMAND		CASS RETURN		
T	(C028)	RATE	AVERAGE	M.A.D.	AVERAGE	M.A.D.		M.A.D.		AVERAGE		
A	(B057)	(B02)	(A019)	(B022A)	(A019)	(B022A)		(A019A)		(B022B)		
PICTURE	1.2	1.2	5.5	5.5	5.5	5.5		5.5		5.5		5.5

CH POS	137-146	147-152	1	153-157	1	158-167	1	168-177	1	178-185	1	186-193
D	SYSTEM CAR-	SYSTEM	UNIT	ACQUISITION	QUARTERLY	REPAIR		REPAIR				
A	CASS RETURN	REQUISITION	PACK	WAR	DEMAND	LEVEL	QUANTITY	DEMAND	LEVEL	QUANTITY		
T	M. A. D.	AVERAGE	(C021B)	RESERVE	FORECAST	(B019B)	(B021A)	(B019B)	(B021B)	(B021A)		
A	(A019B)	(A023B)		(B028C)	(B074)							
PICTURE	5.5	4.2	5	5	10	8.2	8	8	8	8		

CH POS	194-192	197-199	200-202	203-205	206 - 208
D	NUMBER	NAVY NONREPORT-	NAVY REPORTING	NAVY REPORTING	
A	POLICY	ING/COMMERCIAL	REPAIR IN PRO-	REPAIR IN PRO-	
T	RECEIVER	REPAIR FAIR	REPAIR H.A.D.	PROCESS TIME	CESS TIME M.A.D.
A	ACTIVITIES	(B012)	(B012B)	(B012C)	(B012D)
PICTURE	3	1.2	2.1	1.2	2.1

CH	POS	387-390	391-397	398-404	405-411	412-418	419-425	426-432	433-439
D	MAXIMUM	ON-HAND	ON-HAND	ON-HAND	ON-HAND	ON-HAND	ON-HAND	NOT-FIT	M
A	INDUCTION	WHOLESALE	NAS	RETAIL	WAS	RETAIL	FOR	CONDITION	
T	QUANTITY	CONUS	ALAMEDA	CONUS	CECIL	OVERSEAS	ISSUE	ON-HAND	
A	(B095)	TIRS		TIRS	FIELD	TIRS	ON-HAND		
PICTURE	3	7	7	7	7	7	7	7	7

CH	POS	440-446	447-453	454-460	461-467	468-474
D	TOTAL RESERVA-	ZG5 RESERVATIONS		ZG5 RESERVA-	TOTAL RESER-	
A	TIONS AT WHOLE-	AT WHOLESALE	RESERVATIONS	TIONS AT NAS	VATIONS AT	
T	SALE CONUS TIRS	CONUS TIRS	AT NAS ALAMEDA	ALAMEDA	RETAIL CONUS	
A	(A013A)	(A013)	(A013A)	(A013)	TIRS (A013A)	(A013)
PICTURE	7	7	7	7	7	7

CH	POS	475-481	482-488	489-495	496-502	503-509
D	ZG5 RESERVA-	TOTAL RESERVA-	ZG5 RESERVA-	TOTAL RESERVA-	ZG5 RESERVATIONS-	
A	TIONS AT RE-	TIONS AT NAS	TIONS AT NAS	TIONS AT RETAIL	AT RETAIL OVER-	
T	TAIL CONUS	CECIL FIELD	CECIL FIELD	OVERSEAS TIRS	SEAS TIRS	
A	TIRS (A013)	(A013A)	(A013)	(A013A)	(A013)	
PICTURE	7	7	7	7	7	7

CH POS	510-512	513	514-520	521-527	528-534	535-544	545-554
D	NAS	TYPE	PROTECTED	POOL	OUTPUT-	ERROR TERM	
A	ALAMEDA	REPAIRABLE	(MOBILIZA-	RESER-	TING	OF THE	
T	LOCATION	INDICATOR	TION REQ-	ATIONS	RESER-	FIRST QUAR-	SECOND QUARTER
A	FACTOR		UIREMENTS		VATIONS	TER - e1	e2
PICTURE	3	1	7	7	7	55.5	55.5

CH POS	555-564	565-574	575-585	586-595	596	597-600
D	EXPECTED	VARIANCE		STANDARD		FAMILY
A	ERROR	- SIGMA	BLANKS	PRICE	BLANK	GROUP
T	VALUE	- SQUARED		(B053)		CODE
A	E(X)					
PICTURE	55.5	5.5	11	8.2	1	4

5A Demand Trailers; 600 Characters Per Record, 10 Records Per Block, Fixed Length

CH POS	1	2	3	4	5	6-9	10-13	14	15	16	17
D	RECORD	ENTRY	TYPE	RECURRING				BO	ENTRY	TYPE	RECUR-
A	TYPE	POINT	DEMAND	(PRIORITY)	DAY	AMOUNT	(LEVEL)	POINT	DEMAND	RING	
T	("2")	CODE	CODE	CODE	CODE	CODE	CODE	CODE	CODE	CODE	CODE
A											
PICTURE	1	1	1	1	1	1	1	1	1	1	1

CH POS	18	19-20	23-26	27		587	588	589	590		
D	PRIORITY	DAY	AMOUNT	LEVEL	BO	ENTRY	TYPE	RECUR-		PRIORITY	
A	T	A			...	POINT	DEMAND	RING	CODE	DEMAND	
PICTURE	1	4	4	1		1	1	1	1	1	1

CH POS	591-594	595-598	599	600	
D	DAY	AMOUNT	LEVEL	"0"	
A	T	A			
PICTURE	4	4	1	1	1

## APPENDIX E

### FORTRAN PROGRAMS

```
//YOUNTSSA JOB (1642,0045), 'MARK YOUNT', CLASS=B, MSGCLASS=Z  
// EXEC FORTICLG  
//PORT.SYSIN DD *
```

```
C**  
C** MVSTOWN PROGRAM READS THE RAW DATA FROM THE MASS STORAGE  
C** DEVICE AND COPIES THE INPUT RECORDS TO A TRANSFER DISK THAT  
C** IS ACCESSIBLE TO THE VM SYSTEM. THE MAIN DATA RECORDS AND  
C** SUBRECORDS ARE TREATED DIFFERENTLY IN THAT FOR THE MAIN  
C** RECORDS, ONLY THE RECORD/SUBRECORD INDICATOR IS COPIED. THE  
C** REMAINING DATA ELEMENTS REPRESENTING INFORMATION NOT USED IN  
C** THIS ANALYSIS ARE DISCARDED. ALL DEMAND SUBRECORDS FOR EACH  
C** MAIN RECORD ARE COPIED UNALTERED. BY SETTING THE VALUES OF  
C** THE VARIABLES START AND FREQ, THE USER IS ABLE TO SELECT  
C** ONLY A PORTION OF THE RAW DATA FOR TRANSFER. IN THIS WAY,  
C** INDEPENDENT SETS OF DATA COULD BE EVALUATED TO VERIFY THE  
C** POWER OF THE PROCEDURE.  
C**  
C** THIS PROGRAM IS DESIGNED TO TAKE A 1 IN FREQ SAMPLE FROM THE  
C** PMSO 5A SIMULATOR ASO DEMAND DATA STORED ON  
C** MSS.S1642.THESES1 FOR 1R DATA AND MSS.S1642.THESES2 FOR 2R  
C** DATA. THE FILEDEF FOR FT01FO01 MUST BE CHANGED FOR  
C** WHICHEVER DATA SET IS REQUIRED. USE MVSGET EXEC TO RETRIEVE  
C** THE SAMPLE FROM MVS004.  
C**  
C** DEFINITION OF VARIABLES:  
C**  
C** A(599) -- CHARACTER ARRAY USED TO TRANSFER DATA
```

```

*** EOP      -- WRITTEN TO OUTPUT FILE AS END OF FILE INDICATOR
*** FREQ     -- SAMPLING FREQUENCY
*** IN       -- COUNT OF RECORDS READ
*** M        -- 1 = RECORD/2 = SUBRECORD INDICATOR
*** OUT      -- COUNT OF RECORDS WRITTEN
*** START    -- RECORD NUMBER OF FIRST RECORD TO READ

*** NOTE: TO COPY ENTIRE FILE TO MVS004, SET START = 0 AND FREQ = 1

*** INTEGER4
*   A(599) ,
*   EOF/0/,
               N,
               IN/0/,
               FREQ,
               OUT/0/,
               START.

C      SET FREQ TO DESIRED SAMPLING FREQUENCY.
C      FREQ = 1
C      SET START TO DESIRED FIRST LINE ITEM.
C      START = 0
C      READ(1,800,END=80) N
C      CONTINUE
10     IF (.NOT.((N.EQ.1) .AND. (MOD(IN+1,FREQ) .EQ. START))) GOTO 40
C      READ AND TRANSFER TO MVS004 DESIRED RECORDS AND SUBRECORDS
C      IN = IN + 1
C      OUT = OUT + 1
C      WRITE(2,800) N,(A(I),I=1,599)
C      READ(1,810,END=80) N,(A(I),I=1,599)
C      CONTINUE
20     IF (.NOT.(IN - EQ. 2)) GOTO 30
          WRITE(2,810) N,(A(I),I=1,599)
          READ(1,810,END=80) N,(A(I),I=1,599)
GOTO 20

```

```

      30    CONTINUE
      GOTO 70
      CONTINUE
C
C   SKIP BY UNWANTED RECORDS AND SUBRECORDS
C
C   IN = IN + 1
C   READ(1,800,END=80) N
C   CONTINUE
C   IF ( .NOT. (N .EQ. 2) ) GOTO 60
C   READ(1,800,END=80) N
C   GOTO 50
C   CONTINUE
      60    CONTINUE
      GOTO 10
      CONTINUE
C
C   WRITE END OF FILE MARKER
C
C   WRITE(2,800) EOF
C   FREQ = FREQ - 1
C   WRITE(2,820) START,FREQ,IN,OUT
C
      70    CONTINUE
      GOTO 10
      CONTINUE
C
C   WRITE END OF FILE MARKER
C
C   WRITE(2,800) EOF
C   FREQ = FREQ - 1
C   WRITE(2,820) START,FREQ,IN,OUT
C
      80    CONTINUE
      STOP
      F1 FORMAT(1I1)
      F10 FORMAT(1I,8 (75A1))
      F20 FORMAT('STARTING AT RECORD ',I3,' AND SKIPPING EVERY ',I3,
      *          ' RECORDS',/,I5,' RECORDS WERE READ IN AND ',I5,' RECORDS ',
      *          ' WERE OUTPUT.')
      END
      /*
//GO.FT01P001 DD DISP=SHR,DSN=MSS.S1642.THESTIS1
//GO.FT02F001 DD UNIT=3 350, VOL=SER=MVS004,DISP=(NEW,KEEP),
// DCB=(RECFM=FB,LRECL=600,BLKSIZE=6000),SPACE=6000
// DSN=S1642.OUTPUT
/*

```

C001 THESIS IS THE MAIN PROGRAM FOR THE RESAMPLING PROCEDURE.  
 C002 THAT FUNCTIONS AS A CONTROL OVER THE DATA  
 C003 BEING EVALUATED BY CHECKING PARAMETERS RETURNED BY  
 C004 SUBROUTINES. IF THE PARAMETERS ARE OUTSIDE ACCEPTABLE  
 C005 LIMITS, THE ENTIRE SAMPLE IS DISCARDED BEFORE FURTHER  
 C006 PROCESSING. THE PARAMETERS CHECKED ARE:  
 C007 1. NUMBER OF NON-ZERO OBSERVATIONS GREATER THAN ONE.  
 C008 2. STANDARD DEVIATION OF ANY DEMAND SERIES EQUALS ZERO.  
 C009 THE SAMPLES ARE ALSO CATEGORIZED INTO LOW, MEDIUM AND HIGH  
 C010 DEMAND CATEGORIES AND COUNTS KEPT FOR THE NUMBER IN EACH  
 C011 CATEGORY.  
 C012  
 C013 DEFINITIONS OF VARIABLES:  
 C014  
 C015 AN -- FLOAT OF NBR  
 C016 CAT -- BREAKPOINTS OF DEMAND CATEGORIES FOR LOW,  
 C017 MEDIUM AND HIGH DEMANDS  
 C018 DY -- NUMERIC CALENDAR DAY OF MONTH  
 C019 ERROR -- LOGICAL TEST FOR ERRORS IN STATISTICAL VALUES  
 C020 FIRST -- LOGICAL VARIABLE TRUE ONLY FOR FIRST SAMPLE  
 C021 FX -- DISTRIBUTION PERCENTILES TO TEST  
 C022 HR -- HOUR  
 C023 I -- DEMAND CATEGORY FOR SAMPLE UNDER TEST  
 C024 IC -- COUNT OF SAMPLES IN EACH DEMAND CATEGORY  
 C025 ICNOUT -- DSRN FOR TERMINAL OUTPUT  
 C026 IDSK2 -- DSRN OF PROGRAM CONTROL PARAMETERS  
 C027 IER -- RETURN CODE FROM STATS SUBROUTINE  
 C028 IPRINT -- DSRN OF PRINTER DISK  
 C029 IPRT -- CONTROL PARAMETER FOR PRINTING DIAGNOSTICS  
 C030 ISEED -- SEED FOR RANDOM NUMBER GENERATOR USED IN SAMPLE  
 C031 J -- DO INDEX FOR GENERAL USE  
 C032 MI -- MINUTES  
 C033 MO -- NUMERIC MONTH OF YEAR  
 C034 MODELS -- NUMBER OF MODELS TO STUDY

```

C48 MONTH -- CHARACTER MONTHS OF THE YEAR
C48 MSE -- MEAN SQUARED ERROR OF PREDICTED VS. ACTUAL
C48
C48 C48 HU -- SAMPLE MEAN OF EACH GROUP OF OBSERVATIONS
C48 NBR -- NUMBER OF DEMAND OBSERVATIONS TO TEST IS 48
C48 NBR0BS -- NUMBER OF DEMAND OBSERVATIONS IN EACH GROUP,
C48
C48 NPTILE -- NUMBER OF SAMPLER PERCENTILES TO TEST
C48 NR ECD -- MAXIMUM NUMBER OF LINE ITEMS TO PROCESS
C48 NREPS -- NUMBER OF PSEUDO SAMPLE REPETITIONS
C48 OBS -- ARRAY OF DEMAND OBSERVATIONS WHERE COLUMNS ARE
C48
C48 1. RAW OBSERVATIONS AFTER REMOVING NEGATIVES
C48 2. NON-ZERO DEMAND OBSERVATIONS
C48 3. LOG OF NON-ZERO DEMAND OBSERVATIONS
C48
C48 P -- PROBABILITY OF 1. NON-ZERO DEMAND
C48 PHAT -- COUNTER FOR ACCUMULATING PERCENTILE ESTIMATES
C48 SC -- SECONDS
C48 STDV -- SAMPLE STANDARD DEVIATION OF EACH DEMAND GROUP
C48 X -- ABSISSA VALUE OF EACH PROBABILITY MODEL UNDER
C48
C48 YR -- TEST DETERMINED BY SUBROUTINE XINV
C48
C48 SUBROUTINES REQUIRED:
C48
C48 OPEN -- SETS UP INITIAL FILE DEFINITIONS
C48 INITIAL -- READS RAW DATA FROM DISK AND CREATES AN ARRAY
C48
C48
C48 C48 1. RAW DATA
C48 2. NON-ZERO OBSERVATIONS ONLY
C48 3. LOG OF NON-ZERO OBSERVATIONS
C48
C48 TSTATS -- COMPUTES MEAN AND STANDARD DEVIATIONS OF ABOVE
C48 XINV -- GIVEN A PROBABILITY, COMPUTES THE ABSISSA X
C48
C48 SAMPLE -- GENERATES PSEUDO SAMPLES AND TESTS AGAINST
C48
C48 C48 THE X VALUE COMPUTED IN XINV. ACCUMULATES
C48
C48 BERNoulli SUCCESS AND AVERAGE MEAN SQUARE ERROR

```

```

C**      DATINE    -- GETS TIME AND MONTH, DAY AND YEAR FROM SYSTEM
C**      OUTPUT   -- GENERATES OUTPUT REPORT
C**      CFOR
C**      *****
C**      INTEGER#4
*      YR,          MO,
*      HR,          MI,
*      IDSK,        I,
*      IPRINT,      IPRT,
*      NPITLE/5/,  NREPS,
*      ISEED/392746/, MODELS/10/,
*      ICNOUT,      NBRNEG/0/,
*      'APR', 'MAY', 'JUN', 'JUL', 'AUG', 'SEP', 'OCT', 'NOV', 'DEC'/
REAL#4
*      AN,
*      MU(3),
*      X(10,5),
*      MSE(3,10,5)/150*0.0/,
*      CAT(4)/0.0,1.0,20.0,9999.99/, FX(5)/0.75,0.80,0.85,0.90,0.95/
LOGICAL#1
*      ERROR,      FIRST/.TRUE./
COMMON /FILES/
*      IPRT,       ICONIN,
*      IDSK(30),    ICNOUT,
*      CALL OPEN('THESSIS')
IDSK2 = IDSK(2)
READ(IDSK2,*) NRECD,NREPS
CAT(2) = CAT(2) / 12.0
CAT(3) = CAT(3) / 12.0
AN = FLOAT(NBR)
CONTINUE
CALL INITAL(FIRST,NBR,OBS,NBROBS,NBRNEG,680)
FIRST = .FALSE.
P = FLOAT(NBROBS(2)) / AN
C      RETURN BASIC STATISTICS FOR EACH GROUP OF OBSERVATIONS

```

```

C      ERROR = .FALSE.
DO 20 J=1,3
  CALL TSTATS(OBS(1,J),NBRJBS(J),MU(J),STDV(J),IER)
  IF (IER .NE. 0) ERROR = .TRUE.
CONTINUE
C
C      SKIP ENTIRE SAMPLE IF TOO SMALL OR STANDARD DEVIATION EQUALS 0
C
C      IF (ERROR) NBRDEL = NBRDEL + 1
C      IF (ERROR) GOTO 70
C      IF (IPRT .GE. 5) WRITE(IPRINT,600) MU(1),STDV(1)
C      IF (IPRT .GE. 5) WRITE(IPRINT,610) MU(2),STDV(2),P
C      IF (IPRT .GE. 5) WRITE(IPRINT,620) MU(3),STDV(3),NBRBOS(3)
C
C      DIVIDE INTO DEMAND CATEGORIES BY OVERALL MEAN
C
C      IF (MU(1) .GT. CAT(2)) GOTO 30
C      I = 1
C      GOTO 50
C      CONTINUE
C      IF (MU(1) .GT. CAT(3)) GOTO 40
C      I = 2
C      GOTO 50
C      CONTINUE
C      I = 3
C      CONTINUE
C      IC(I) = IC(I) + 1
C
C      COMPUTE LOCATION OF EACH PERCENTILE FOR ALL MODELS.
C
C      CALL XINV(P,MU,STDV,FX,MODELS,NPTILE,X,IER)
C
C      CONDUCT RESAMPLING PROCEDURE RETURNING THE NUMBER OF
C      C      BERNOULLI SUCCESSES AND THE AVERAGE MEAN SQUARED ERROR OF THE
C      C      MODEL.

```

```

C      DO 60 J=1,NREPS
C          CALL SAMPLE(ISEED,OBS,NBR,I,MODELS,NPTILE,X,PX,PHAT,MSE)
60      CONTINUE
CONTINUE
IF (IC(1)+IC(2)+IC(3) .LT. NRECD) GOTO 10
70      CONTINUE
C
C      GENERATE OUTPUT REPORT
C
CALL DATIME(YR,MO,DY,JD,HR,MI,SC,HD)
WRITE(ICNOUT,630) NBRDEL,NBRNEG,(IC(I),I=1,3)
WRITE(IPRINT,630) NBRDEL,NBRNEG,(IC(I),I=1,3)
WRITE(ICNOUT,640) HR,HI,SC,DY,MONTH(MO),YR
WRITE(IPRINT,640) HR,HI,SC,DY,MONTH(MO),YR
CALL OUTPUT(MODELS,NPTILE,NREPS,PX,IC,PHAT,MSE,CAT)
STOP
FORMAT('0SAMPLE MEAN: ',F10.4,' SAMPLE STANDARD DEVIATION:','
     * F10.4)
600      FORMAT('0SAMPLE MEAN: ',F10.4,' SAMPLE STANDARD DEVIATION:','
     * F10.4,' PROBABILITY OF A NON-ZERO OBSERVATION: ',F10.8)
610      FORMAT('0SAMPLE MEAN: ',F10.4,' SAMPLE STANDARD DEVIATION:','
     * F10.4,' NUMBER OF NON-ZERO OBSERVATIONS: ',I10)
620      FORMAT('1',20X,'SUMMARY RUN STATISTICS',/,
     * '-NUMBER OF SAMPLES DELETED BY POOR STATISTICS: ',I3,/,
     * 'NUMBER OF SAMPLES WITH NEGATIVE DEMANDS: ',I3,/,
     * 'PROCESSED ',I4,' RECORDS ASSIGNED AS LOW DEMAND',/,
     * 'PROCESSED ',I4,' RECORDS ASSIGNED AS MEDIUM DEMAND',/,
     * 'PROCESSED ',I4,' RECORDS ASSIGNED AS HIGH DEMAND')
630      FORMAT('REPORT GENERATED AT ',I2,':',I2,':',I2,' ON ',
     * I2,'.',A3,' 19',I2)
640      END

```

SUBROUTINE OPEN ( FNAME\$ )

C\*\*  
C\*\* OPEN IS A GENERAL PURPOSE SUBROUTINE DESIGNED TO HANDLE ALL  
C\*\* THE OPERATING SYSTEM FILE DEFINITIONS REQUIRED TO INPUT AND  
C\*\* OUTPUT DATA TO THE REQUIRED DEVICE. OPEN IS CALLED AS THE  
C\*\* FIRST EXECUTABLE STATEMENT OF THE MAIN PROGRAM AND READS A  
C\*\* DATA FILE TO DETERMINE THE DATA SET REFERENCE NUMBERS (DSRN)  
C\*\* REQUIRED BY THE MAIN PROGRAM. OPEN THEN CALLS THE SYSTEM  
C\*\* SUBROUTINE PRTCMS TO EXECUTE THE CMS FILEDEF COMMANDS WITH  
C\*\* ARGUMENTS SPECIFIED IN THE DATA FILE. OPEN ALSO INITIALIZES  
C\*\* A COMMON BLOCK WITH THE DSRN'S ASSIGNED TO MAKE THEM  
C\*\* ACCESSIBLE TO OTHER PROGRAM UNITS.  
C\*\*  
C\*\* TO OPERATE, THE USER MUST HAVE A FILE NAMED 'FNAME\$'  
C\*\* FILES'. OPEN READS THE FILE AS FOLLOWS:  
C\*\*  
C\*\* LINE 1. FORMAT(4I5,5X, A8) ASSIGNED TO OUTPUT CONTROL  
C\*\* VARIABLE IPRT AND I/O DEVICE NUMBERS FOR  
C\*\* TERMINAL INPUT, TERMINAL OUTPUT, PRINTER DISK,  
C\*\* AND FILENAME FOR PRINT FILE '<PN> OUTPUT A1'.  
C\*\*  
C\*\* LINE 2. FORMAT(I5,5X,7(1A,2X)) ASSIGNED TO I/O DEVICE  
C\*\* NUMBERS FOR OTHER THAN PRINTER DISK I/O AND THE  
C\*\* FILE NAME AND FILETYPE. FILEMODE DEFAULTS TO A1.  
C\*\* REMAINING VALUES ARE FIVE OR LESS OPTIONAL  
C\*\* FILEDEF PARAMETERS, IE. LRECL, BLOCK, ETC.  
C\*\*  
C\*\* LINE 2 IS OPTIONAL AND MAY BE LEFT OUT OR REPEATED UP TO  
C\*\* THIRTY TIMES AS NECESSARY.  
C\*\*  
C\*\* OTHER SUBROUTINES CALLED: STR\$  
C\*\*  
C\*\* WRITTEN BY MARK YOUNT AUG 1981  
C\*\*  
C\*\* C:\mark\src\open.sas  
C\*\*  
C\*\* IMPLICIT REAL\*8 ( A-H, O-Z )

```

COMMON /FILE$/ IPRT, ICONIN, ICNOUT, IPRINT, IDSK(30)
REALS8 PAR(14)
* /'FILEDEF ',' TERMINAL', 'DISK ', 'OUTPUT '
* 'A1 ',' ( ',' RECPH ', 'PA '
* 'BLOCK ',' 133 ',' PERM ', 'FILES '
* ') ',' /'
DIMENSION OPT$(5)

C IPRT = 0
DNUM$ = STR$(99)
CALL PRTCM$ (PAR(1) , DNUM$ , PAR(3) , FNAME$ , PAR(12) , PAR(5) ,
* PAR(6) , PAR(11) )

C READ DATA FOR TERMINAL I/O AND PRINTER DESTINED OUTPUT
C
C READ(99,100) ITMP , ICONIN, ICNOUT, IPRINT, PNAME$
C FORMAT(4I5,5X,A8)
100 CONIN$ = STR$(ICONIN)
CALL PRTCM$ (PAR(1) , CONIN$ , PAR(2) , PAR(6) , PAR(11))
COUT$ = STR$(ICNOUT)
CALL PRTCM$ (PAR(1) , COUT$ , PAR(2) , PAR(6) , PAR(11))
PRINT$ = STR$(IPRINT)
IF (IPRINT .EQ. 6) GOTO 10
CALL PRTCM$ (PAR(1) , PRINT$ , PAR(3) , PNAME$ , PAR(4) , PAR(5) , PAR(6) ,
* PAR(7) , PAR(8) , PAR(9) , PAR(10) , PAR(11) )
CONTINUE
10
C READ DATA FOR DISK FILE OPERATIONS
C
C FORMAT(15,5X,7(A8,2X))
110 IPRT = ITMP
DO 20 I=1,30
  READ(99,110,END=30) IDSK(I) , DNAMES$, DTYPES$ , (OPT$(J) , J=1,5)
DO 15 J=1,5
  IF (OPT$(J) .NE. PAR(14)) GOTO 15
  OPT$(J) = PAR(13),

```

```

      GOTO 16
15    CONTINUE
16    CONTINUE
      DNUM$ = STR$(IDSK(I))
      IF (IPRT .GE. 20) WRITE(ICNOUT,200) PAR(1),DNUM$,PAR(3),DNAMES$,
      DTYPE$,PAR(6),PAR(11),(OPT$(J),J=1,5)
      *   FORMAT(' ',7A10/10X,5A10)
      CALL FRTCMS(PAR(1),DNUM$,PAR(3),DNAMES$,DTYPE$,PAR(6),PAR(11),
      *   OPT$(1),OPT$(2),OPT$(3),OPT$(4),OPT$(5))

20    CONTINUE
30    CONTINUE
      IF (IPRT .GE. 10) CALL PRTCMS(PAR(1))
      REWIND 99
      RETURN
      END

```

DOUBLE PRECISION FUNCTION STR\$(IVAL)

Character

STR\$ IS A GENERAL PURPOSE FUNCTION INVOKED BY OPEN DESIGNED  
TO ENHANCE THE FLEXIBILITY OF THE FRTCMS SUBROUTINE BY  
CONVERTING INTEGER DSRN'S INTO REAL<sup>8</sup> CHARACTER STRINGS  
SUITABLE AS ARGUMENTS FOR FRTCMS. (FRTCMS USES ONLY  
CHARACTER STRING ARGUMENTS. ALPHABETIC ARGUMENTS MAY BE FROM  
1 TO 8 CHARACTERS LEFT JUSTIFIED, BUT NUMERIC ARGUMENTS MUST  
BE EIGHT CHARACTERS LEFT JUSTIFIED AND PADDED ON THE RIGHT  
WITH BLANKS IF NECESSARY.)  
STR\$ CONVERTS INTEGER VALUES IN THE RANGE OF +99,999,999 TO  
-9,999,999 INTO ALPHANUMERIC REPRESENTATION AS THE PUNCTUATION  
OUTPUT. STR\$ MUST BE TYPED REAL<sup>8</sup> IN THE CALLING PROGRAM.  
STR\$ RETURNS A CHARACTER VALUE OF 0 IF CONVERSION UNSUCCESSFUL.  
WRITTEN BY MARK YOUNT SEP 1981  
REAL<sup>8</sup> BF,BLINK/  
COMMON /FILES/IPT,IPRINT,ICONIN,ICONOUT,ICNOUT,IDS<sup>1</sup>SK(30)  
LOGICAL<sup>1</sup> NUM(8),ONUM(8),MINUS/-/  
LOGICAL<sup>1</sup> DIGIT(10)/\*0\*,\*1\*,\*2\*,\*3\*,\*4\*,\*5\*,\*6\*,\*7\*,\*8\*,\*9\*/  
INTEGER<sup>4</sup> FUNC\$/STR\$/  
EQUIVALENCE (ONUM(1),B\$)  
IF (IPT .GE. 20) WRITE(ICNOUT,201) IVAL  
201 FORMAT(' INPUT INTEGER = ',I10)  
IM = 8  
B\$ = BLINK  
IF ((IVAL .GE. 10\*\*8) .OR. (IVAL .LE. -1\*10\*\*7)) GOTO 6000  
ITEMP = IABS(IVAL)  
C EXTRACT CHARACTERS ONE AT A TIME FROM IVAL AND STORE AS  
C ALPHANUMERIC CHARACTERS IN ARRAY NUM

```

C      DO 10 I=1,8
      IF (ITEMP .EQ. 0) GOTO 20
      N = ITEMP
      ITEMP = ITEMP / 10
      N = N - ITEMP * 10
      NUM (IM) = DIGIT (N+1)
      IF (IPRT .GE. 40) WRITE (ICNOUT, 200) N, NUM (IM)
      FORMAT (' CCMV XITED NUMERIC ', I1, ' TO CHARACTER ', A1)
      IM = IM - 1
10    CONTINUE
20    CONTINUE
C      LEFT JUSTIFY CHARACTERS IN BS (1)
C      DO 60 I=1,8
          IF (I .GT. 1) GOTO 50
          IF (IVAL) 30, 40, 50
          CONTINUE
          ONUM (I) = MINUS
          IM = IM - 1
          IF (IPRT .GE. 40) WRITE (ICNOUT, 210)
          FORMAT (' ASSIGNED NEGATIVE IVALUE TO STR$')
          GOTO 60
60    CONTINUE
          ONUM (I) = DIGIT (1)
          IM = IM - 1
          IF (IPRT .GE. 40) WRITE (ICNOUT, 200) IVAL, ONUM (I)
          GOTO 60
210   CONTINUE
          J = I + IM
          IF (J .GT. 8) GOTO 60
          ONUM (I) = NUM (J)
          CONTINUE
60    CREATE OUTPUT ARRAY

```

```
C      STR$ = B$  
C      IF (IPRT .GE. 30) WRITE (ICNOUT,220) IVAL,STR$  
220  FORMAT (' NUMERIC VALUE = ',18.0 ALPHANUMERIC VALUE = ',A8)  
RETURN  
  
C      ERROR HANDLING SECTION FOLLOWS  
C  
6000  CONTINUE  
      WRITE (ICNOUT,6010) FUNCS  
6010  FORMAT (' *** STRING LENGTH ERROR *** ',A4)  
      STR$ = DIGIT(1)  
      RETURN  
END
```

```

SUBROUTINE INITIAL (FIRST,M,OBS,VBROBS,NBRNEG,")
C*#
C*# INITIAL FIRST CALLS SUBROUTINE DATARD TO READ ONE SAMPLE
C*# FROM THE SPECIFIED DISK WHEN CALLED FROM THE MAIN PROGRAM.
C*# THE STRING OF FORTY-EIGHT DEMAND OBSERVATIONS THUS READ IN
C*# ARE FIRST PROCESSED TO REMOVE ANY NEGATIVE DEMAND
C*# OBSERVATIONS BY BACKING OUT THE PRIOR POSITIVE DEMAND THAT
C*# IS EQUAL OR LARGER THAN THE NEGATIVE VALUE. WHETHER A
C*# LARGER VALUE IS FOUND OR NOT, THE NEGATIVE DEMAND IS SET TO
C*# ZERO. THE RESULTING DEMAND SERIES IS THE EDITED DEMAND
C*# SERIES USED FOR FURTHER PROCESSING. TWO MORE STEPS ARE
C*# REQUIRED PRIOR TO SUBROUTINE RETURN, FIRST A SECOND DEMAND
C*# SERIES IS CONSTRUCTED OF ONLY THE NON-ZERO DEMANDS AND THE
C*# COUNT STORED, SECOND A THIRD DEMAND SERIES IS CONSTRUCTED OF
C*# THE NATURAL LOG OF THE NON-ZERO DEMAND SERIES JUST CREATED.
C*# THUS WHEN INITIAL RETURNS CONTROL TO THE MAIN PROGRAM, THREE
C*# DEMAND SERIES HAVE BEEN CONSTRUCTED
C*#
C*# DEFINITION OF VARIABLES:
C*#
C*#   COUNT    -- TEMPORARY COUNTER EVENTUALLY ASSIGNED TO
C*#             N BROBS (2) AND NBROBS (3)
C*#   FIRST    -- LOGICAL VARIABLE TO INDICATE FIRST USE OF INITIAL
C*#   I,J,K    -- DO LOOP INDICIES
C*#   IDSK1   -- I INPUT DSRN FOR DEMAND DATA
C*#   IPRINT  -- OUTPUT DSRN FOR LOCAL PRINTER
C*#   IPRT    -- CONTROLS LEVEL OF DIAGNOSTICS PRINTED
C*#   IREV    -- METHOD TO COUNT BACKWARD IN OBS ARRAY
C*#   M       -- I INPUT COUNT OF OBSERVATIONS TO INPUT,
C*#             N BROBS (1) ALWAYS EQUALS M
C*#   NBRNEQ  -- COUNTS NUMBER OF SAMPLES WITH NEGATIVE OBS.
C*#   NBROBS  -- COUNTS OF OBSERVATIONS IN EACH OF ABOVE
C*#   NEGOBS  -- LOGICAL VARIABLE, TRUE IF SAMPLE HAS NEG OBS.
C*#   OBS (M,1) -- ORIGINAL OBSERVATIONS WITH NEGATIVES REMOVED
C*#   OBS (M,2) -- ALL ZERO OBSERVATIONS REMOVED FROM DATA

```

```

C      OBS(M,3) -- NATURAL LOG OF NON-ZERO OBSERVATIONS
C      OTHER SUBROUTINES REQUIRED BY INITIAL:
C      DATARD -- DOES ACTUAL READING OF DATA FROM RAW DATA SETS
C
C      INTEGER*4
C      IPRT,           ICONIN,          ICNOUT,          IPRINT,
C      IDSK(30),       IREV,            IDSK1,           NBR OBS(3),
C      COUNT,          I,                J,                K,
C      M,              NBRNEG
C      REAL*4
C      OBS(M,3)
C      LOGICAL*1
C      NEGOMS,         FIRST
C      COMMON /FILES/
C      IPRT,           ICONIN,          ICNOUT,          IPRINT,
C      IDSK
C      IDSK1 = IDSK(1)
C      NBR OBS(1) = M
C      CALL DATARD(OBS(1,1), M, FIRST, 640)

C      ECHO PRINT INPUT VALUES (CONTROLLED BY IPRT VALUE)
C
C      IF (IPRT .GE. 5) WRITE(IPRT,600) (OBS(K,1), K=1,M)
C      NEGOMS = .FALSE.
C      DO 20 I=1,M
C      IP (OBS(I,1) .GE. 0.0) GOTO 20

C      C      EDIT BY BACKING OUT AND REMOVING NEGATIVE OBSERVATIONS
C
C      NEGOMS = .TRUE.
C      DO 10 J = 2,I
C      IREV = I - J + 1
C      IF (OBS(IREV,1) .LT. -OBS(I,1)) GOTO 10

```

```

OBS(IREV,1) = OBS(IREV,1) + OBS(I,1)
OBS(I,1) = 0.0
GOTO 20

10   CONTINUE
      OBS(I,1) = 0.0
CONTINUE
      IF (NEG(OBS) NBRNEG = NBRNEG + 1

C   PRINT EDITED INPUT VALUES (CONTROLLED BY IPRT)
C
C   IF (IPRT .GE. 5) WRITE(IPRT,610) (OBS(K,1),K=1,M)

C   CREATE ARRAY OF ONLY NON-ZERO OBSERVATIONS
C   AND ARRAY OF LOG OF THESE NON-ZERO OBSERVATIONS
C
C   COUNT = 0
DO 30 I=1,M
      IF (OBS(I,1) .EQ. 0.0) GOTO 30
      COUNT = COUNT + 1
      OBS(COUNT,2) = OBS(I,1)
      OBS(COUNT,3) = LOG(OBS(I,1))

30   CONTINUE
      NBROBS(2) = COUNT
      NBROBS(3) = COUNT

C   PRINT NON-ZERO INPUT VALUES (CONTROLLED BY IPRT)
C
C   IF (IPRT .GE. 5) WRITE(IPRT,620) (OBS(K,2),K=1,COUNT)

C   PRINT LOG OF NON-ZERO INPUT VALUES (CONTROLLED BY IPRT)
C
C   IF (IPRT .GE. 5) WRITE(IPRT,630) (OBS(K,3),K=1,COUNT)

40   CONTINUE
      RETURN 1
      FORMAT('1RAW OBSERVATIONS ',3(T20,20F5.0,/))
600

```

```
610 FORMAT('ONEGATIVE REMOVED ',3(T20,20F5.0,'/'))
620 FORMAT('NON ZERO OBS ONLY ',3(T20,20F5.0,'/'))
630 FORMAT('LOG NON-ZERO ONLY ',3(T20,20F5.0,'/'))
800 FORMAT(50P4.0)
END
```

RD-A123 779

DISTRIBUTIONAL ANALYSIS OF INVENTORY DEMAND OVER  
LEADTIME(U) NAVAL POSTGRADUATE SCHOOL MONTEREY CA  
M L YOUNT JUN 82


2/2

UNCLASSIFIED

F/G 15/5

NL

END  
FILED  
DTB



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

SUBROUTINE DATARD (MONTH,NBR,TIME,\*)

```

C*#
C*# DATARD DOES THE ACTUAL READING OF DATA FROM THE VM MINIDISK
C*# AS TWO PARALLEL ARRAYS OF DATA. THE FIRST THE SEQUENTIAL DAY
C*# OF DEMAND AND THE SECOND THE QUANTITY DEMANDED. ALL THE
C*# INDIVIDUAL DEMANDS FOR EACH DAY ARE TOTLED AND THEN GROUPED
C*# INTO THIRTY DAY BUCKETS REPRESENTING THE ACCUMULATED DEMAND
C*# FOR THIS SAMPLE FOR ONE MONTH. THE DATA RETURNED FROM
C*# DATARD IS THE RAW MONTHLY DEMAND DATA TO BE PROCESSED BY
C*# SUBROUTINE INITIAL.
C*#
C*# DEFINITION OF VARIABLES:
C*#
C*# DATE      -- SEQUENTIAL DATE OF DEMAND, DAY 1 TO DAY 1440
C*#           -- ONE ELEMENT FOR EACH ELEMENT OF INPUT DATA
C*# DEM       -- ACCUMULATES DAILY DEMANDS FOR LATER AGGREGATION
C*#           -- DO LOOP INDICES
C*# IDSK1    -- INPUT DSRN FOR DEMAND DATA
C*# FIRST    -- STARTING INDEX OF CURRENT 30 DAY MONTH BLOCK
C*# LAST     -- ENDING INDEX OF CURRENT 30 DAY MONTH BLOCK
C*# MONTH   -- OUTPUT DEMAND VALUES AGGRAGATED IN 30 DAY BLOCKS
C*# NBR     -- NUMBER OF OBSERVATIONS TO CREATE
C*# QUANT   -- QUANTITY DEMANDED ON SEQUENTIAL DATE
C*# TIME    -- LOGICAL VARIABLE TRUE ONLY ON FIRST CALL
C*#
C*# INTEGER#4 FIRST,      LAST,
C*#           IDSK1,
C*#           J,
C*# REAL#4   DATE(46),   N
C*#           DEM(1550),
C*#           LOGICAL#1 QUANT(46)
C*#           TIME
C*#           COMMON /FILES/
C*#           IPRT,      ICONIN,      IPRINT,

```

```

* IDSK(30)
IDSK1 = IDSK(1)
IF (TIME) READ(IDSK1, 800) N
C
C   INITIALIZE COUNTERS USED TO TOTAL DEMAND
C
C   DO 20 I=1,1550
      DEM(I) = 0.0
20   CONTINUE
      DO 30 I=1,NBR
          MONT(I) = 0.0
30   CONTINUE

C   READ IN DEMANDS AND ACCUMULATE IN DAILY BUCKETS
C
C   READ(1,810) N, (DATE(I), QUANT(I), I=1,46)
40   CONTINUE
      IF (.NOT. (N .EQ. 2)) GOTO 70
      DO 60 I=1,46
          IF (.NOT. (DATE(I) .NE. 0)) GOTO 50
              DEM(DATE(I)) = DEM(DATE(I)) + QUANT(I)
60   CONTINUE
      READ(1,810) N, (DATE(I), QUANT(I), I=1,46)
      GOTO 40
70   CONTINUE

C   AGGREGATE ACCUMULATED DEMANDS INTO MONTHLY BUCKETS
C
C   FIRST = 31
LAST = 60
DO 90 I=1,NBR
    DO 80 J=FIRST, LAST
        MONT(I) = MONT(I) + DEM(J)
80   CONTINUE
    FIRST = LAST + 1
90   CONTINUE

```

```
      LAST = LAST + 30
      CONTINUE
      IF (N .EQ. 0) RETURN 1
      RETURN
      FORMAT(I1)
      FORMAT(I1,46(4X,I4,F4.0,1X)) )
      END
```

SUBROUTINE TSTATS (X,N,XMEAN,STDV,IER)

C\*#  
C\*# COMPUTES THE SAMPLE MEAN AND SAMPLE STANDARD

C\*# DEVIATION OF AN INPUT DATA ARRAY WHEN CALLED FROM THE MAIN

C\*# PROGRAM. VARIOUS ERROR RETURN CODES SIGNIFY WHETHER THE

C\*# SAMPLE SIZE IS TOO SMALL FOR PROCESSING OR THE STANDARD

C\*# DEVIATION EQUALS ZERO.

C\*#  
C\*# DEFINITION OF VARIABLES:

C*#	AN	--	FLOATING POINT VALUE OF N
C*#	DEV	--	INTERMEDIATE VALUE OF STDV CALCULATION
C*#	I	--	GENERAL DO INDEX
C*#	IER	--	ERROR RETURN CODE EQUAL TO:
C*#		0	-- NORMAL RETURN
C*#		-1	-- INSUFFICIENT DATA (IE. 0 OR 1)
C*#		-3	-- STDV EQUALS 0
C*#	N	--	NUMBER OF DATA ELEMENTS IN THE SAMPLE
C*#	STDV	--	SAMPLE STANDARD DEVIATION OF DATA VECTOR X (N-1)
C*#	SUM	--	DOUBLE PRECISION SUM OF OBSERVATIONS
C*#	X	--	INPUT DATA VECTOR OF LENGTH N
C*#	XMEAN	--	SAMPLE MEAN OF DATA VECTOR X

```

INTEGERS4
*   I,
REALS4
*   X(N),
      XMEAN,
      SUM,
      DEV
REALS8
*   AN = FLOAT(N)
IER = 0
IF(N .GE. 2) GO TO 10
IER = -1

```

```
      GOTO 40
10    CONTINUE
C      COMPUTE MEAN AND MOMENT ESTIMATES
C      FIND THE MEAN
C
      SUM = 0.D0
      DO 20 I=1,N
          SUM = SUM + X(I)
20    CONTINUE
      XMEAN = SNGL(SUM) /AN
      SUM = 0.D0
      DO 30 I=1,N
          DEV = X(I) - XMEAN
          SUM = SUM + DEV * DEV
30    CONTINUE
      STDV = SQRT(SNGL(SUM) / (AN - 1.0))
      IF (STDV .EQ. 0.0) TERR = -3
      CONTINUE
      RETURN
END
```

10

SUBROUTINE XINV(P,MU,STDV,PX,MODEL,NPTILE,X,IER)

C\*\*  
C\*\* XINV IS CALLED BY THE MAIN PROGRAM TO COMPUTE THE ABSISSA  
C\*\* VALUES FOR EACH MODEL BEING EVALUATED AT THE FIVE  
C\*\* PERCENTILES OF INTEREST, CURRENTLY THE 75TH, 60TH, 85TH,  
C\*\* 90TH AND 95TH PERCENTILES.  
C\*\* DISTRIBUTIONS, (NORMAL, LOGNORMAL AND BERNOLLI-LOGNORMAL)  
C\*\* XINV CALLS THE IMSL SINGLE PRECISION SUBROUTINE MDN RIS TO  
C\*\* FIND THE NORMALIZED ABSISSA VALUE AND THEN CONVERTS THAT  
C\*\* SAMPLE MEAN AND STANDARD DEVIATION CONVERSIONS. FOR THE  
C\*\* EXPONENTIAL, BERNOLLI-EXPONENTIAL AND BERNOLLI-LOGISTIC  
C\*\* DISTRIBUTIONS, XINV COMPUTES THE ABSISSA VALUE DIRECTLY FROM  
C\*\* THE DATA SUPPLIED. THE REMAINING DISTRIBUTIONS, POISSON,  
C\*\* NEGATIVE BINOMIAL, LOGISTIC AND LA PLACE, XINV CALLS THE  
C\*\* SUBROUTINES POISSN, NEGBIN, LOGIST AND LAPLCE RESPECTIVELY  
C\*\* TO COMPUTE THE ABSISSA VALUES.  
C\*\* UPON RETURN TO THE MAIN PROGRAM, XINV HAS COMPUTED THE  
C\*\* INVERSE FOR EACH SAMPLE AT EVERY PERCENTILE FOR EVERY MODEL.  
C\*\*  
C\*\* COMPUTES THE INVERSE CUMULATIVE PROBABILITY FUNCTION FOR THE  
C\*\* FOLLOWING DISTRIBUTIONS SELECTED BY THE VALUE OF THE PARAMETER  
C\*\* MODEL:  
C\*\*  
C\*\* 1. POISSON  
C\*\* 2. NEGATIVE BINOMIAL  
C\*\* 3. STANDARD NORMAL  
C\*\* 4. EXPONENTIAL  
C\*\* 5. LOGNORMAL  
C\*\* 6. LOGISTIC  
C\*\* 7. LAPLACE  
C\*\* 8. BERNOLLI- EXPONENTIAL  
C\*\* 9. BERNOLLI- LOGNORMAL  
C\*\* 10. BERNOLLI- LOGISTIC

```

C#* DEFINITION OF VARIABLES:
C#* PI      -- PROBABILITY VALUE
C#* HX      -- COMPLEMENTARY PROBABILITY VALUE
C#* IER      -- ERROR PARAMETER INDICATING
C#*          0 NORMAL RETURN
C#*          NE 0 RETURN CODE FROM SUBROUTINE "MDNRIS"
C#*          OTHER CODE RETURNED FROM ROUTINE "MDNRIS"
C#*          SELECTS DISTRIBUTION
C#* MODEL    -- DETERMINES WHICH MU & STDV EACH MODEL USES
C#* MTYPE   -- SAMPLE MEAN OF EACH GROUP OF DATA
C#* MU      -- ARRAY OF PERCENTILES FOR TESTING
C#* NPTILE  -- BERNoulli PROBABILITY PARAMETER
C#* P       --
C#* PI      -- CONSTANT PI
C#* STDV   -- SAMPLE STANDARD DEVIATION OF EACH GROUP OF DATA
C#* X       -- ABSCISSA VALUE RESULT OF INVERSE FUNCTION
C#* Z       -- NORMALIZED RESULT OF INVERSE FUNCTIONS
C#*          OTHER SUBROUTINES REQUIRED BY XINV:
C#*          LAPLCE -- DISTRIBUTION, DENSITY OR INVERSE LAPLACE VALUES
C#*          LOGIST -- DISTRIBUTION, DENSITY OR INVERSE LOGISTIC VALUES
C#*          MDNRIS -- IMSL INVERSE NORMAL SUBROUTINE
C#*          NEGBIN -- DISTRIBUTION, DENSITY OR INVERSE NEGATIVE
C#*          BINOMIAL VALUES
C#*          POISSN -- DISTRIBUTION, DENSITY OR INVERSE POISSON VALUES
C#*          THE MEAN AND STANDARD DEVIATION ARE ESTIMATED BY THE METHOD OF
C#*          MOMENTS. FOR DISTRIBUTIONS SUCH AS THE LAPLACE AND LOGISTIC,
C#*          THESE ESTIMATORS ARE BIASED, BUT REMAIN THE CHOICE FOR USE IN
C#*          THIS ANALYSIS DUE TO EASE OF COMPUTATION.
C#*          REAL4
C#*          * FACT,      FX(NPTILE), MU(3),
C#*          P,

```

```

C   PR,          HX,
C   X(MODEL,NPTILE),
C   INTEGER*4
C   * IER,      MODEL,    NPTILE,    MTYPE(10)/1,1,1,3,1,1,2,3,2/,
C   * J,          K
C   IER = 0
DO 130 J=1,MODEL
DO 120 K=1,NPTILE
HX = 1.0 - FX(K)

C   RETURN X = 0.0 FOR COMPOUND DISTRIBUTIONS IF P <= HX
C

C   X(J,K) = 0.0
IF ((J .GE. 8) .AND. (P .LE. HX)) GOTO 110

C   SELECT THE DISTRIBUTION
C
GOTO (10,20,30,40,50,60,70,80,90,100),J
CONTINUE

10   C   INVERSE POISSON FUNCTION
C
CALL POISSN(X(J,K),MU(MTYPE(J)),3,FX(K),IER)
IF (IER .NE. 0) GOTO 130
GOTO 110
CONTINUE

20   C   INVERSE NEGATIVE BINOMIAL FUNCTION
C
CALL NEGBIN(X(J,K),MU(MTYPE(J)),
IF (STDV(MTYPE(J)),3,FX(K),IER)
IF (IER .NE. 0) GOTO 130
GOTO 110
CONTINUE

30   C   INVERSE STANDARD NORMAL FUNCTION

```

```

C          CALL MDNRIS(FX(K),Z,IER)
C          IF (IER .NE. 0) GOTO 130
C          X(J,K) = MU(MTYPE(J)) + STDV(MTYPE(J)) * Z
C          GOTO 110
C          CONTINUE
40
C          C          INVERSE EXPONENTIAL FUNCTION
C          C          X(J,K) = -MU(MTYPE(J)) + LOG(HX)
C          GOTO 110
C          CONTINUE
50
C          C          INVERSE LOG NORMAL FUNCTION
C          C          CALL MDNRIS(FX(K),Z,IER)
C          IF (IER .NE. 0) GOTO 140
C          X(J,K) = EXP(MU(MTYPE(J)) + STDV(MTYPE(J)) * Z)
C          GOTO 110
C          CONTINUE
60
C          C          INVERSE LOGISTIC FUNCTION
C          C          CALL LOGIST(X(J,K),MU(MTYPE(J)),STDV(MTYPE(J)),
C          3,FX(K),IER)
C          IF (IER .NE. 0) GOTO 140
C          GOTO 110
C          CONTINUE
70
C          C          INVERSE LAPLACE FUNCTION
C          C          CALL LAPLCE(X(J,K),MU(MTYPE(J)),STDV(MTYPE(J)),
C          3,FX(K),IER)
C          IF (IER .NE. 0) GOTO 140
C          GOTO 110
C          CONTINUE
80

```

```

C      INVERSE MIXED BERNOULLI-EXPONENTIAL FUNCTION
C
C      X(J,K) = MU(MTYPE(J)) * LOG(P / HX)
C
C      GOTO 110
C      CONTINUE
C
C      INVERSE BERNOULLI-LOGNORMAL FUNCTION
C
C      PR = (P - HX) / P
C      CALL MDNRIS(PR,Z,IER)
C      IF (IER .NE. 0) GOTO 140
C      X(J,K) = EXP(MU(MTYPE(J)) + STDV(MTYPE(J)) * Z)
C
C      GOTO 110
C      CONTINUE
C
C      INVERSE BERNOULLI-LOGISTIC FUNCTION
C
C      FACT = SQRT(3.0) * STDV(MTYPE(J)) / PI
C      Z = HX & (1.0 + TANH(MU(MTYPE(J)) / (2.0 * FACT)))
C      X(J,K) = MU(MTYPE(J)) + FACT * LOG(2.0 * P / Z - 1.0)
C
C      CONTINUE
C      CONTINUE
C      CONTINUE
C      RETURN
C      END
C
C
C      100
C      110
C      120
C      130
C      140

```

```

SUBROUTINE POISSN (X,MU,TYPE,FX,IER)
C*#
C*# POISSN IS A GENERAL PURPOSE POISSON DISTRIBUTION FUNCTION
C*# THAT RETURNS EITHER THE DENSITY OR CUMULATIVE DISTRIBUTION
C*# VALUE GIVEN THE ABSISSA VALUE OR ALTERNATIVELY, RETURNS THE
C*# ABSISSA VALUE GIVEN THE PROBABILITY.
C*# COMPUTES POISSON DENSITY AND DISTRIBUTION FUNCTIONS BY USE OF
C*# RECURSION EQUATIONS.
C*#
C*# X - INPUT PARAMETER REPRESENTING ABSISSA VALUES OF THE
C*# POISSON DISTRIBUTION. FOR ITYPE = 3, X IS AN OUTPUT
C*# PARAMETER.
C*#
C*# MU - MEAN OF THE DESIRED POISSON DISTRIBUTION.
C*#
C*# TYPE = 1 RETURNS PDF IN FX GIVEN X
C*#        2 RETURNS CDF IN FX GIVEN X
C*#        3 RETURNS INVERSE CDF VALUES IN X GIVEN FX AS CDF
C*#
C*# FX - OUTPUT PARAMETER OF PROBABILITY (X<=X). FOR ITYPE = 3,
C*# FX IS AN INPUT PARAMETER.
C*#
C*# IER - ERROR PARAMETER RETURN CODE EQUAL TO:
C*#      0 - NORMAL RETURN
C*#      -1 - INVALID SELECTION OF TYPE
C*#      -3 - ALGORITHM DID NOT CONVERGE WITHIN 1000 ITERATIONS.
C*#
C*# INTEGER I4
C*#      I4,          IER,          TYPE
C*#      REAL4,        PDF,          FX,
C*#      CDF,          X,           MU,
C*#      A,
C*#      PDF = 0.0
C*#      IF (MU .LT. 170.0) PDF = EXP (-MU)

```

```

CDF = PDF
A = 0.0
DO 50 I=1,1000
  IF (TYPE .LT. 1) GOTO 60
  IF (TYPE .GT. 3) GOTO 60
  GOTO (10,20,30),TYPE
CONTINUE
  IF (A .LT. X) GOTO 40
    FX = PDF
    GOTO 99
  CONTINUE
  IF (A .LT. X) GOTO 40
    FX = CDF
    GOTO 99
  CONTINUE
  IF (CDF .LT. FX) GOTO 40
    X = A
    GOTO 99
  CONTINUE
  A = A + 1.0
  IF (PDF .GT. ZL) PDF = MU * PDF / A
  IF (PDF .LE. ZL) PDF = 0.0
  CDF = CDF + PDF
CONTINUE
IER = -3
GOTO 99
CONTINUE
IER = -1
CONTINUE
RETURN
END

```

```

SUBROUTINE NEGBIN (X,MU,STDV,TYPE,FX,IER)
C
C   NEGBIN IS A GENERAL PURPOSE NEGATIVE BINOMIAL DISTRIBUTION
C   FUNCTION THAT RETURNS EITHER THE DENSITY OR CUMULATIVE
C   DISTRIBUTION VALUE GIVEN THE ABSCISSA VALUE, OR ALTERNATIVELY
C   RETURNS THE ABSCISSA VALUE GIVEN THE PROBABILITY.
C
C   COMPUTES NEGATIVE BINOMIAL DENSITY AND DISTRIBUTION FUNCTIONS
C   BY USE OF RECURSION EQUATIONS.
C
C   X - INPUT PARAMETER REPRESENTING ABSCISSA VALUES OF THE
C   NEGATIVE BINOMIAL DISTRIBUTION. FOR TYPE = 3, X IS
C   AN OUTPUT PARAMETER.
C
C   MU - MEAN OF THE NEGATIVE BINOMIAL DISTRIBUTION.
C
C   STDV - STANDARD DEVIATION OF THE NEGATIVE BINOMIAL
C   DISTRIBUTION.
C
C   TYPE = 1 RETURNS PDF IN FX GIVEN X
C         2 RETURNS CDF IN FX GIVEN X
C         3 RETURNS INVERSE CDF VALUES IN X GIVEN FX AS CDF
C
C   FX - OUTPUT PARAMETER OF PROBABILITY (X<=X) . FOR TYPE = 3,
C   FX IS AN INPUT PARAMETER.
C
C   IER - ERROR PARAMETER RETURN CODE EQUAL TO:
C         0 - NORMAL RETURN
C         -1 - INVALID SELECTION OF TYPE
C         -3 - ALGORITHM DID NOT CONVERGE WITHIN 1000 ITERATIONS.
C
C   INTEGER I4
C   REAL I,
C          TYPE,
C          FX,
C          CDF,
C          MU,
C          X,
C          Y,
C          Z,
C          T,
C          S,
C          R,
C          Q,
C          P,
C          O,
C          N,
C          M,
C          L,
C          K,
C          J,
C          I,
C          H,
C          G,
C          F,
C          E,
C          D,
C          C,
C          B,
C          A,
C          B1,
C          B2,
C          B3,
C          B4,
C          B5,
C          B6,
C          B7,
C          B8,
C          B9,
C          B10,
C          B11,
C          B12,
C          B13,
C          B14,
C          B15,
C          B16,
C          B17,
C          B18,
C          B19,
C          B20,
C          B21,
C          B22,
C          B23,
C          B24,
C          B25,
C          B26,
C          B27,
C          B28,
C          B29,
C          B30,
C          B31,
C          B32,
C          B33,
C          B34,
C          B35,
C          B36,
C          B37,
C          B38,
C          B39,
C          B40,
C          B41,
C          B42,
C          B43,
C          B44,
C          B45,
C          B46,
C          B47,
C          B48,
C          B49,
C          B50,
C          B51,
C          B52,
C          B53,
C          B54,
C          B55,
C          B56,
C          B57,
C          B58,
C          B59,
C          B60,
C          B61,
C          B62,
C          B63,
C          B64,
C          B65,
C          B66,
C          B67,
C          B68,
C          B69,
C          B70,
C          B71,
C          B72,
C          B73,
C          B74,
C          B75,
C          B76,
C          B77,
C          B78,
C          B79,
C          B80,
C          B81,
C          B82,
C          B83,
C          B84,
C          B85,
C          B86,
C          B87,
C          B88,
C          B89,
C          B90,
C          B91,
C          B92,
C          B93,
C          B94,
C          B95,
C          B96,
C          B97,
C          B98,
C          B99,
C          B100,
C          B101,
C          B102,
C          B103,
C          B104,
C          B105,
C          B106,
C          B107,
C          B108,
C          B109,
C          B110,
C          B111,
C          B112,
C          B113,
C          B114,
C          B115,
C          B116,
C          B117,
C          B118,
C          B119,
C          B120,
C          B121,
C          B122,
C          B123,
C          B124,
C          B125,
C          B126,
C          B127,
C          B128,
C          B129,
C          B130,
C          B131,
C          B132,
C          B133,
C          B134,
C          B135,
C          B136,
C          B137,
C          B138,
C          B139,
C          B140,
C          B141,
C          B142,
C          B143,
C          B144,
C          B145,
C          B146,
C          B147,
C          B148,
C          B149,
C          B150,
C          B151,
C          B152,
C          B153,
C          B154,
C          B155,
C          B156,
C          B157,
C          B158,
C          B159,
C          B160,
C          B161,
C          B162,
C          B163,
C          B164,
C          B165,
C          B166,
C          B167,
C          B168,
C          B169,
C          B170,
C          B171,
C          B172,
C          B173,
C          B174,
C          B175,
C          B176,
C          B177,
C          B178,
C          B179,
C          B180,
C          B181,
C          B182,
C          B183,
C          B184,
C          B185,
C          B186,
C          B187,
C          B188,
C          B189,
C          B190,
C          B191,
C          B192,
C          B193,
C          B194,
C          B195,
C          B196,
C          B197,
C          B198,
C          B199,
C          B200,
C          B201,
C          B202,
C          B203,
C          B204,
C          B205,
C          B206,
C          B207,
C          B208,
C          B209,
C          B210,
C          B211,
C          B212,
C          B213,
C          B214,
C          B215,
C          B216,
C          B217,
C          B218,
C          B219,
C          B220,
C          B221,
C          B222,
C          B223,
C          B224,
C          B225,
C          B226,
C          B227,
C          B228,
C          B229,
C          B230,
C          B231,
C          B232,
C          B233,
C          B234,
C          B235,
C          B236,
C          B237,
C          B238,
C          B239,
C          B240,
C          B241,
C          B242,
C          B243,
C          B244,
C          B245,
C          B246,
C          B247,
C          B248,
C          B249,
C          B250,
C          B251,
C          B252,
C          B253,
C          B254,
C          B255,
C          B256,
C          B257,
C          B258,
C          B259,
C          B260,
C          B261,
C          B262,
C          B263,
C          B264,
C          B265,
C          B266,
C          B267,
C          B268,
C          B269,
C          B270,
C          B271,
C          B272,
C          B273,
C          B274,
C          B275,
C          B276,
C          B277,
C          B278,
C          B279,
C          B280,
C          B281,
C          B282,
C          B283,
C          B284,
C          B285,
C          B286,
C          B287,
C          B288,
C          B289,
C          B290,
C          B291,
C          B292,
C          B293,
C          B294,
C          B295,
C          B296,
C          B297,
C          B298,
C          B299,
C          B300,
C          B301,
C          B302,
C          B303,
C          B304,
C          B305,
C          B306,
C          B307,
C          B308,
C          B309,
C          B310,
C          B311,
C          B312,
C          B313,
C          B314,
C          B315,
C          B316,
C          B317,
C          B318,
C          B319,
C          B320,
C          B321,
C          B322,
C          B323,
C          B324,
C          B325,
C          B326,
C          B327,
C          B328,
C          B329,
C          B330,
C          B331,
C          B332,
C          B333,
C          B334,
C          B335,
C          B336,
C          B337,
C          B338,
C          B339,
C          B340,
C          B341,
C          B342,
C          B343,
C          B344,
C          B345,
C          B346,
C          B347,
C          B348,
C          B349,
C          B350,
C          B351,
C          B352,
C          B353,
C          B354,
C          B355,
C          B356,
C          B357,
C          B358,
C          B359,
C          B360,
C          B361,
C          B362,
C          B363,
C          B364,
C          B365,
C          B366,
C          B367,
C          B368,
C          B369,
C          B370,
C          B371,
C          B372,
C          B373,
C          B374,
C          B375,
C          B376,
C          B377,
C          B378,
C          B379,
C          B380,
C          B381,
C          B382,
C          B383,
C          B384,
C          B385,
C          B386,
C          B387,
C          B388,
C          B389,
C          B390,
C          B391,
C          B392,
C          B393,
C          B394,
C          B395,
C          B396,
C          B397,
C          B398,
C          B399,
C          B400,
C          B401,
C          B402,
C          B403,
C          B404,
C          B405,
C          B406,
C          B407,
C          B408,
C          B409,
C          B410,
C          B411,
C          B412,
C          B413,
C          B414,
C          B415,
C          B416,
C          B417,
C          B418,
C          B419,
C          B420,
C          B421,
C          B422,
C          B423,
C          B424,
C          B425,
C          B426,
C          B427,
C          B428,
C          B429,
C          B430,
C          B431,
C          B432,
C          B433,
C          B434,
C          B435,
C          B436,
C          B437,
C          B438,
C          B439,
C          B440,
C          B441,
C          B442,
C          B443,
C          B444,
C          B445,
C          B446,
C          B447,
C          B448,
C          B449,
C          B450,
C          B451,
C          B452,
C          B453,
C          B454,
C          B455,
C          B456,
C          B457,
C          B458,
C          B459,
C          B460,
C          B461,
C          B462,
C          B463,
C          B464,
C          B465,
C          B466,
C          B467,
C          B468,
C          B469,
C          B470,
C          B471,
C          B472,
C          B473,
C          B474,
C          B475,
C          B476,
C          B477,
C          B478,
C          B479,
C          B480,
C          B481,
C          B482,
C          B483,
C          B484,
C          B485,
C          B486,
C          B487,
C          B488,
C          B489,
C          B490,
C          B491,
C          B492,
C          B493,
C          B494,
C          B495,
C          B496,
C          B497,
C          B498,
C          B499,
C          B500,
C          B501,
C          B502,
C          B503,
C          B504,
C          B505,
C          B506,
C          B507,
C          B508,
C          B509,
C          B510,
C          B511,
C          B512,
C          B513,
C          B514,
C          B515,
C          B516,
C          B517,
C          B518,
C          B519,
C          B520,
C          B521,
C          B522,
C          B523,
C          B524,
C          B525,
C          B526,
C          B527,
C          B528,
C          B529,
C          B530,
C          B531,
C          B532,
C          B533,
C          B534,
C          B535,
C          B536,
C          B537,
C          B538,
C          B539,
C          B540,
C          B541,
C          B542,
C          B543,
C          B544,
C          B545,
C          B546,
C          B547,
C          B548,
C          B549,
C          B550,
C          B551,
C          B552,
C          B553,
C          B554,
C          B555,
C          B556,
C          B557,
C          B558,
C          B559,
C          B560,
C          B561,
C          B562,
C          B563,
C          B564,
C          B565,
C          B566,
C          B567,
C          B568,
C          B569,
C          B570,
C          B571,
C          B572,
C          B573,
C          B574,
C          B575,
C          B576,
C          B577,
C          B578,
C          B579,
C          B580,
C          B581,
C          B582,
C          B583,
C          B584,
C          B585,
C          B586,
C          B587,
C          B588,
C          B589,
C          B590,
C          B591,
C          B592,
C          B593,
C          B594,
C          B595,
C          B596,
C          B597,
C          B598,
C          B599,
C          B600,
C          B601,
C          B602,
C          B603,
C          B604,
C          B605,
C          B606,
C          B607,
C          B608,
C          B609,
C          B610,
C          B611,
C          B612,
C          B613,
C          B614,
C          B615,
C          B616,
C          B617,
C          B618,
C          B619,
C          B620,
C          B621,
C          B622,
C          B623,
C          B624,
C          B625,
C          B626,
C          B627,
C          B628,
C          B629,
C          B630,
C          B631,
C          B632,
C          B633,
C          B634,
C          B635,
C          B636,
C          B637,
C          B638,
C          B639,
C          B640,
C          B641,
C          B642,
C          B643,
C          B644,
C          B645,
C          B646,
C          B647,
C          B648,
C          B649,
C          B650,
C          B651,
C          B652,
C          B653,
C          B654,
C          B655,
C          B656,
C          B657,
C          B658,
C          B659,
C          B660,
C          B661,
C          B662,
C          B663,
C          B664,
C          B665,
C          B666,
C          B667,
C          B668,
C          B669,
C          B670,
C          B671,
C          B672,
C          B673,
C          B674,
C          B675,
C          B676,
C          B677,
C          B678,
C          B679,
C          B680,
C          B681,
C          B682,
C          B683,
C          B684,
C          B685,
C          B686,
C          B687,
C          B688,
C          B689,
C          B690,
C          B691,
C          B692,
C          B693,
C          B694,
C          B695,
C          B696,
C          B697,
C          B698,
C          B699,
C          B700,
C          B701,
C          B702,
C          B703,
C          B704,
C          B705,
C          B706,
C          B707,
C          B708,
C          B709,
C          B710,
C          B711,
C          B712,
C          B713,
C          B714,
C          B715,
C          B716,
C          B717,
C          B718,
C          B719,
C          B720,
C          B721,
C          B722,
C          B723,
C          B724,
C          B725,
C          B726,
C          B727,
C          B728,
C          B729,
C          B730,
C          B731,
C          B732,
C          B733,
C          B734,
C          B735,
C          B736,
C          B737,
C          B738,
C          B739,
C          B740,
C          B741,
C          B742,
C          B743,
C          B744,
C          B745,
C          B746,
C          B747,
C          B748,
C          B749,
C          B750,
C          B751,
C          B752,
C          B753,
C          B754,
C          B755,
C          B756,
C          B757,
C          B758,
C          B759,
C          B760,
C          B761,
C          B762,
C          B763,
C          B764,
C          B765,
C          B766,
C          B767,
C          B768,
C          B769,
C          B770,
C          B771,
C          B772,
C          B773,
C          B774,
C          B775,
C          B776,
C          B777,
C          B778,
C          B779,
C          B780,
C          B781,
C          B782,
C          B783,
C          B784,
C          B785,
C          B786,
C          B787,
C          B788,
C          B789,
C          B790,
C          B791,
C          B792,
C          B793,
C          B794,
C          B795,
C          B796,
C          B797,
C          B798,
C          B799,
C          B800,
C          B801,
C          B802,
C          B803,
C          B804,
C          B805,
C          B806,
C          B807,
C          B808,
C          B809,
C          B810,
C          B811,
C          B812,
C          B813,
C          B814,
C          B815,
C          B816,
C          B817,
C          B818,
C          B819,
C          B820,
C          B821,
C          B822,
C          B823,
C          B824,
C          B825,
C          B826,
C          B827,
C          B828,
C          B829,
C          B830,
C          B831,
C          B832,
C          B833,
C          B834,
C          B835,
C          B836,
C          B837,
C          B838,
C          B839,
C          B840,
C          B841,
C          B842,
C          B843,
C          B844,
C          B845,
C          B846,
C          B847,
C          B848,
C          B849,
C          B850,
C          B851,
C          B852,
C          B853,
C          B854,
C          B855,
C          B856,
C          B857,
C          B858,
C          B859,
C          B860,
C          B861,
C          B862,
C          B863,
C          B864,
C          B865,
C          B866,
C          B867,
C          B868,
C          B869,
C          B870,
C          B871,
C          B872,
C          B873,
C          B874,
C          B875,
C          B876,
C          B877,
C          B878,
C          B879,
C          B880,
C          B881,
C          B882,
C          B883,
C          B884,
C          B885,
C          B886,
C          B887,
C          B888,
C          B889,
C          B890,
C          B891,
C          B892,
C          B893,
C          B894,
C          B895,
C          B896,
C          B897,
C          B898,
C          B899,
C          B900,
C          B901,
C          B902,
C          B903,
C          B904,
C          B905,
C          B906,
C          B907,
C          B908,
C          B909,
C          B910,
C          B911,
C          B912,
C          B913,
C          B914,
C          B915,
C          B916,
C          B917,
C          B918,
C          B919,
C          B920,
C          B921,
C          B922,
C          B923,
C          B924,
C          B925,
C          B926,
C          B927,
C          B928,
C          B929,
C          B930,
C          B931,
C          B932,
C          B933,
C          B934,
C          B935,
C          B936,
C          B937,
C          B938,
C          B939,
C          B940,
C          B941,
C          B942,
C          B943,
C          B944,
C          B945,
C          B946,
C          B947,
C          B948,
C          B949,
C          B950,
C          B951,
C          B952,
C          B953,
C          B954,
C          B955,
C          B956,
C          B957,
C          B958,
C          B959,
C          B960,
C          B961,
C          B962,
C          B963,
C          B964,
C          B965,
C          B966,
C          B967,
C          B968,
C          B969,
C          B970,
C          B971,
C          B972,
C          B973,
C          B974,
C          B975,
C          B976,
C          B977,
C          B978,
C          B979,
C          B980,
C          B981,
C          B982,
C          B983,
C          B984,
C          B985,
C          B986,
C          B987,
C          B988,
C          B989,
C          B990,
C          B991,
C          B992,
C          B993,
C          B994,
C          B995,
C          B996,
C          B997,
C          B998,
C          B999,
C          B9999

```

```

* STDV,      VAR,      A,      X,
* RHO,      ONE/1.0/,   K,      ZL/1.0E-20/
* VAR = STDV*2
RHO = MU / VAR
K = MU*2 / (VAR - MU)
PDF = RHO*K
CDF = PDF
A = 0.0
DO 50 I=1,1000
  IF (TYPE .LT. 1) GOTO 60
  IF (TYPE .GT. 3) GOTO 60
  GOTO (10,20,30),TYPE
10  CONTINUE
  IF (A .LT. X) GOTO 40
    FX = PDF
    GOTO 99
20  CONTINUE
  IF (A .LT. X) GOTO 40
    FX = CDF
    GOTO 99
30  CONTINUE
  IF (CDF .LT. FX) GOTO 40
    X = A
    GOTO 99
40  CONTINUE
  A = A + ONE
  IF (PDF .GT. ZL) PDF = (A + K - ONE) * (ONE - RHO) * PDF / A
  IF (PDF .LE. ZL) PDF = 0.0
  CDF = CDF + PDF
50  CONTINUE
  IER = -3
  GOTO 99

```

60      CONTINUE  
      IER = -1  
99      CONTINUE  
      RETURN  
      END

```

SUBROUTINE LOGIST (X,MU,STDV,TYPE,FX,IER)
C
C LOGIST IS A GENERAL PURPOSE LOGISTIC DISTRIBUTION FUNCTION
C THAT RETURNS EITHER THE DENSITY OR CUMULATIVE DISTRIBUTION
C VALUE GIVEN THE ABSISSA VALUE OR ALTERNATIVELY, RETURNS THE
C ABSISSA VALUE GIVEN THE PROBABILITY.
C
C X - INPUT PARAMETER REPRESENTING THE ABSISSA VALUE OF
C      THE LOGISTIC DISTRIBUTION. FOR TYPE = 3, X IS AN
C      OUTPUT PARAMETER.
C
C MU - MEAN OF THE LOGISTIC DISTRIBUTION OR METHOD OF MOMENTS
C      OR MLE ESTIMATE.
C
C STDV- STANDARD DEVIATION OF THE LOGISTIC DISTRIBUTION OR
C      METHOD OF MOMENTS OR MLE ESTIMATE.
C
C TYPE = 1 RETURNS PDF IN FX GIVEN X
C        2 RETURNS CDF IN FX GIVEN X
C        3 RETURNS INVERSE CDF VALUES IN X GIVEN FX AS CDF
C
C FX- OUTPUT PARAMETER OF PROBABILITY (X<=X). FOR TYPE = 3,
C      FX IS AN INPUT PARAMETER.
C
C IER - ERROR PARAMETER RETURN CODE EQUAL TO:
C        0 - NORMAL RETURN
C        -1 - INVALID SELECTION OF TYPE
C
C INTEGER
C * IER,          TYPE
C * REAL4        ARG,          FX,
C * PI/3,1415926/, SECNX,
C * MU,          SQRPI,
C * STDV

```

```

C   INITIALIZE CONSTANTS
C
C   SQRT3 = SQRT(3.0)
C   ARG = HALF * PI * (X - MU) / (SQRT3 * STDV)
C
C   IF (TYPE .LT. 1) GOTO 40
C   IF (TYPE .GT. 3) GOTO 40
C   GOTO (10,20,30), TYPE
10  CONTINUE

C   COMPUTE DISTRIBUTION PDF
C
C   SECHX = ONE / COSH(ARG)
C   PX = PI * SECHX**2 / (4.0 * SQRT3 * STDV)
C   GOTO 50
C
C   CONTINUE

20  COMPUTE DISTRIBUTION CDF
C
C   PX = HALF * (ONE + TANH(ARG))
C   GOTO 50
C
C   CONTINUE

30  COMPUTE X VALUES FROM PROBABILITY INTEGRAL TRANSFORM
C
C   X = MU + STDV * SQRT3 * LOG(PX / (ONE - PX)) / PI
C   GOTO 50
C
C   CONTINUE
C   IER = -1
40  CONTINUE
C
C   RETURN
50  END

```

```

SUBROUTINE LAPLCE (X,MU,STDV,TYPE,FX,IER)
C*   LAPLCE IS A GENERAL PURPOSE LAPLACE DISTRIBUTION FUNCTION
C*   THAT RETURNS EITHER THE DENSITY OR CUMULATIVE DISTRIBUTION
C*   VALUE GIVEN THE ABSISSA VALUE OR ALTERNATIVELY, RETURNS THE
C*   ABSISSA VALUE GIVEN THE PROBABILITY.
C*
C*   X - INPUT PARAMETER REPRESENTING THE ABSISSA VALUE OF
C*   THE LAPLACE DISTRIBUTION. FOR TYPE = 3, X IS AN
C*   OUTPUT PARAMETER.
C*
C*   MU - MEAN OF THE LAPLACE DISTRIBUTION OR METHOD OF MOMENTS
C*   OR MLE ESTIMATE.
C*
C*   STDV - STANDARD DEVIATION OF THE LAPLACE DISTRIBUTION OR
C*   METHOD OF MOMENTS OR MLE ESTIMATE.
C*
C*   TYPE = 1 RETURNS PDF IN FX GIVEN X
C*         2 RETURNS CDF IN FX GIVEN X
C*         3 RETURNS INVERSE CDF VALUES IN X GIVEN FX AS CDF
C*
C*   FX - OUTPUT PARAMETER OF PROBABILITY (X<=X) . FOR TYPE = 3,
C*   FX IS AN INPUT PARAMETER.
C*
C*   IER - ERROR PARAMETER RETURN CODE EQUAL TO:
C*         0 - NORMAL RETURN
C*         -1 - INVALID SELECTION OF TYPE
C*
C*   INTEGER
C*     IER,
C*     REAL*4
C*     ARG,
C*     TWO/2.0/,
C*     STDV
C*     ONE/1.0/,
C*     ZERO/0.0/,
C*     SQR 2/1.4142135/.

```

```

C      INITIALIZE CONSTANTS
C
C      ARG = SQR2 * ABS(X - MU) / STDV
C      IF (TYPE .LT. 1) GOTO 40
C      IF (TYPE .GT. 3) GOTO 40
C      GOTO (10,20,30), TYPE
10    CONTINUE
C
C      COMPUTE DISTRIBUTION PDF
C
C      FX = EXP(-ARG) / (SQR2 * STDV)
C      GOTO 50
20    CONTINUE
C
C      COMPUTE DISTRIBUTION CDF
C
C      FX = EXP(-ARG) / TWO
C      IF (X .GE. ZERO) FX = ONE - FX
C      GOTO 50
30    CONTINUE
C
C      COMPUTE X VALUES FROM PROBABILITY INTEGRAL TRANSFORM
C
C      IF (FX .LT. 0.5) X = MU + STDV * LOG(TWO * FX) / SQR2
C      IF (FX .GE. 0.5) X = MU - STDV * LOG(TWO - TWO * FX) / SQR2
C      GOTO 50
40    CONTINUE
        IER = -1
50    CONTINUE
        RETURN
        END

```

SUBROUTINE SAMPLE (ISEED,JBS,NBR,I,MODEL,NPTILE,X,PX,PHAT,MSE)

```

C** SAMPLE IS THE HEART OF THE ANALYSIS AND IS CALLED REPEATEDLY
C** BY THE MAIN PROGRAM TO GENERATE SUCCESSIVE REPETITIONS OF
C** THE PSEUDO SAMPLES FOR THE PURPOSE OF AVERAGING THE RESULTS.
C** THE RANDOM INTEGERS FROM ONE TO FORTY-EIGHT ARE OBTAINED
C** EXPANDING FORTY-EIGHT ZERO TO ONE RANDOM NUMBERS BY
C** MULTIPLYING BY FORTY-EIGHT, ADDING ONE AND TRUNCATING TO
C** INTEGERS. THE PSEUDO SAMPLE IS THEN CREATED FROM THE EDITED
C** DEMAND OBSERVATIONS AS DESCRIBED IN SECTION IV.B.
C** EACH SAMPLE IS ANALYZED BY SUCCESSIVELY CHECKING EACH
C** OBSERVATION AGAINST THE DESIRED ABSCISSA VALUE COMPUTED
C** EARLIER BY XINV AND RECORDING A SUCCESS IF THE SAMPLE
C** OBSERVATION IS LESS THAN OR EQUAL TO THE CURRENT ABSCISSA
C** VALUE. SUCCESSES ARE REFLECT BY ADDING ONE TO THE VARIABLE
C** SUCCES WHILE FAILURES ADD ZERO TO THE VARIABLE. TO HANDLE
C** THE CASE OF INTEGER ABSCISSA VALUE, FOR THE POISSON AND
C** NEGATIVE BINOMIAL DISTRIBUTIONS AND ALSO FOR THE COMPOUND
C** DISTRIBUTIONS WHERE THE ABSCISSA VALUE IS ZERO, SUCCESSES
C** WERE RECORDED ONLY UNTIL THE NUMBER OF SUCCESSES WERE LESS
C** THAN THE EXPECTED NUMBER OF SUCCESSES (COMPUTED AS THE
C** PERCENTILE TIMES THE NUMBER OF OBSERVATIONS-- FORTY EIGHT).
C** AFTER THE ENTIRE PSEUDO SAMPLE HAS BEEN CHECKED, THE RESAMPLING
C** PROCEDURE ESTIMATE OF P IS COMPUTED AND ADDED TO THE
C** ACCUMULATOR VARIABLE PHAT FOR LATER OUTPUT. LIKEWISE, THE
C** MEAN SQUARED ERROR IS COMPUTED AND ACCUMULATED IN THE
C** VARIABLE MSE FOR LATER OUTPUT

C** VARIABLES USED ARE AS FOLLOWS:
C**          A      -- DUMMY ARRAY FOR RANDOM NUMBERS FOR SAMPLERAP
C**          AN     -- FLOAT OF NBR
C**          BPROB   -- RESAMPLING PROBABILITY OF DEMAND LT EXPECTED
C**          EXPS    -- EXPECTED NUMBER OF BERNOULLI SUCCESSES
C**          FX     -- PROBABILITIES OF INTEREST

```

```

C00      I          -- DEMAND CATEGORY
C00      IPBT     -- CONTROL PARAMETER FOR OUTPUT OF DIAGNOSTIC INFO
C00      IPRINT   -- DSRN FOR PRINTER DISK
C00      ISSEED   -- SEED FOR RANDOM NUMBER GENERATOR LRND
C00      J          -- DO INDEX SEQUENCING THROUGH MODELS
C00      K          -- DO INDEX SEQUENCING THROUGH PERCENTILES
C00      L          -- DO INDEX SEQUENCING THROUGH SAMPLERAP SAMPLES
C00      IND      -- INDEX TO SELECT RANDOM OBSERVATION FROM OBS
C00      MODEL    -- NUMBER OF MODELS TO TEST
C00      MSE      -- MEAN SQUARED ERROR OF PERCENTILES
C00      NBR      -- NUMBER OF OBSERVATIONS IN SAMPLE
C00      NPTILE   -- NUMBER OF PERCENTILES TO TEST
C00      OBS      -- ARRAY OF DEMAND OBSERVATIONS AND DERIVATIVES
C00      PHAT     -- CUMULATIVE PERCENTILE ESTIMATES
C00      SAMP     -- PSEUDO SAMPLE OF DEMAND OBSERVATIONS
C00      SUCCES   -- NUMBER OF BERNOULLI SUCCESSES EACH TRIAL
C00      X          -- EXPECTED DEMAND OBSERVATION FOR A GIVEN
C00                  PERCENTILE AND MODEL
C00
C00      INTEGER 4
C00      ISSEED,   NBR,
C00      I,        J,
C00      SUCCES,   IND,
C00      REAL 4
C00      AN,       SAMP(48),
C00      MSE(3,MODEL,NPTILE),
C00      OBS(NBR,3),
C00      A(48),
C00      COMMON /FILE/
C00      IPRT,
C00      IDSK(30)
C00      AN = FLOAT(NBR)
C00      DO 10 K=1,NPTILE
C00      EXP5(K) = FX(K) * AN
C00
C00      IPRINT,
C00      ICNIN,
C00      IPRINT,

```

```

10    CONTINUE
C   GENERATE PSEUDO SAMPLES
C
CALL LRND(ISEED,A,NBR,2,0)
DO 20 L=1,NBR
  IND = IPIX(A(L)+AN) + 1
  SAMP(L) = OBS(IND,1)
CONTINUE
IF (IPRT .GE. 5) WRITE(IPRINT,600) (SAMP(L),L=1,NBR)

C   COMPUTE SAMPLE PERCENTILES FOR EACH PSEUDO SAMPLE
C
DO 90 J=1,MODEL
  IF (IPRT .GE. 5) WRITE(IPRINT,610) J
  DO 80 K=1,NPTILE
    SUCCES = 0
    DO 70 L=1,NBR
      IF (.NOT.(SAMP(L) .LE. X(J,K))) GOTO 60
      IF (AMOD(X(J,K),1.0) .EQ. 0.0) GOTO 30
C   FOR NON-INTEGER VALUES OF EXPECTED DEMAND,
C   ACCUMULATE BERNOULLI SUCCESSES WHENEVER
C   ACTUAL DEMAND, REPRESENTED BY THE PSEUDO
C   SAMPLE, IS LESS THAN OR EQUAL TO EXPECTED
C   DEMAND.
      SUCCES = SUCCES + 1
    GOTO 50
  CONTINUE
30
C   IN THE CASE OF INTEGER VALUES OF EXPECTED
C   DEMAND, SUCCESSES ARE ACCUMULATED UNTIL
C   THE EXPECTED NUMBER OF SUCCESSES IS
C   REACHED.
C

```

```

        IF (FLOAT(SUCSES) .GE. EXPS(K)) GOTO 40
        SUCSES = SUCSES + 1
        CONTINUE
        CONTINUE
        CONTINUE
        BPROB = FLOAT(SUCSES) / AN
        IP (IPRT .GE. 5) WRITE(IPRINT,620) X(J,K),BPROB,FX(K)

C      ACCUHULATE SQUARED ERROR AND ESTIMATED PERCENTILE
C      FOR EACH DISTRIBUTION AT EACH THEORETICAL PERCENTILE
C

        PHAT(I,J,K) = PHAT(I,J,K) + BPROB
        MSE(I,J,K) = MSE(I,J,K) + (BPROB - FX(K)) ** 2
        IP(IPRT .GE. 5) WRITE(IPRINT,630) PHAT(I,J,K),MSE(I,J,K)
        CONTINUE
        CONTINUE
        RETURN
        FORMAT('OPS EUDO SAMPLE ',3(T20,20F5.0,/))
        FORMAT('OPERS MODEL ',I1,I1)
        FORMAT('OPROBABILTY PSEUDO SAMPLE .IE. ',F12.6,' IS ',F12.6,
     *      ' AT THE ',F4.2,' PERCENTILE')
        FORMAT('OCUMULATIVE PROBABILITY ',F5.1,' SQUARED ERROR: ',F12.6)
        END

```

```

SUBROUTINE OUTPUT (MODEL,NPTILE,NREPS,FX,IC,PHAT,MSE,CAT)
C48
C49      OUTPUT IS THE FINAL SUBROUTINE CALLED BY THE MAIN PROGRAM
C50      AND IS USED TO GENERATE THE OUTPUT REPORT. ONLY MINOR
C51      COMPUTATIONS ARE PERFORMED BY OUTPUT, THOSE BEING COMPUTING
C52      THE AVERAGE P VALUE AND ITS STANDARD DEVIATION, AVERAGING
C53      THE MEAN SQUARED ERROR AND TOTALING THE AVERAGE MEAN SQUARED
C54      ERROR FOR EACH DEMAND CLASS AND MODEL.
C55
C56      DEFINITION OF VARIABLES:
C57
C58      CAT      -- BREAKPOINTS OF DEMAND CATEGORIES FOR LOW,
C59      C58      MEDIUM AND HIGH DEMANDS
C60      DISTR     -- CHARACTER NAME OF EACH MODEL TESTED
C61      FX        -- DISTRIBUTION PERCENTILES TO TEST
C62      I          -- DEMAND CATEGORY FOR SAMPLE UNDER TEST
C63      IC         -- COUNT OF SAMPLES IN EACH DEMAND CATEGORY
C64      IPRINT    -- DSRN OF PRINTER DISK
C65      J          -- DO INDEX SEQUENCING THROUGH MODELS
C66      K          -- DO INDEX SEQUENCING THROUGH PERCENTILES
C67      MODEL      -- NUMBER OF MODELS TO STUDY
C68      MSE        -- MEAN SQUARED ERROR OF PREDICTED VS. ACTUAL
C69      PERCENTILE
C70      MU        -- BERNOULLI MEAN OF PSEUDO SAMPLE REPETITIONS
C71      C70      FOR EACH PERCENTILE AND MODEL TESTED
C72      NPTILE    -- NUMBER OF SAMPLE PERCENTILES TO TEST
C73      NREPS     -- NUMBER OF PSEUDO SAMPLE REPETITIONS
C74      PHAT      -- CUMULATIVE PERCENTILE ESTIMATES FOR EACH
C75      DEMAND CATEGORY, MODEL AND PERCENTILE
C76      STDV      -- BERNOULLI STANDARD DEVIATION OF PSEUDO SAMPLE
C77      REPETITIONS FOR EACH PERCENTILE AND
C78      MODEL TESTED
C79      TMSE      -- TOTAL MEAN SQUARE ERROR SUMMED OVER PERCENTILES
C80      C79      FOR EACH DEMAND CATEGORY AND MODEL
C81      TRIALS    -- TOTAL NUMBER OF REPETITIONS. USED TO NORMALIZE

```



```

      MSE(I,J,K) = MSE(I,J,K) / TRIALS
      TMSE = TMSE + MSE(I,J,K)
      CONTINUE
      WRITE(IPRINT,620) (MU(K),K=1,NPTILE)
      WRITE(IPRINT,630) (STDV(K),K=1,NPTILE)
      TMSE = TMSE + 1.0E04
      WRITE(IPRINT,640) (MSE(I,J,K),K=1,NPTILE),TMSE
      CONTINUE
      CONTINUE
      RETURN
      * FORMAT(6X,I2,' REPETITIONS FOR EACH OF ',I4,' PSEUDO SAMPLES',/,
      * 6X,A7,' DEMAND RANGE:',F5.1,' < D < ',F7.2,' PER YEAR')
      610  FORMAT(/, ' DISTRIBUTION: ',2A16,',/')
      * * ESTIMATES',22X,'PERCENTILES (P)',/,'12X,5F10.2)
      620  FORMAT(' MEAN      ',5F10.6)
      630  FORMAT(' STD DEV   ',5F10.6)
      640  FORMAT(' MSE       ',5F10.6,' TOTAL MSE: ',F8.2,'E-04')
      END

```

## LIST OF REFERENCES

1. Taylor, C.P., A Multi Item Inventory Model for Combat Stores Ships, Master's Thesis, Naval Postgraduate School, Monterey CA, 1975.
2. U.S. Navy Fleet Material Support Office Report 120, Probability Distributions for Leadtime Demand, by T. Tupper, November 1975.
3. Lilliefors, H. W., "On the Kolmogorov-Smirnov Test for the Exponential Distribution with Mean Unknown", American Statistical Association Journal, V. 64, pp. 387-389, March 1969.
4. Hadley, G. and Whitin, T. M., Analysis of Inventory Systems, p. 166, Prentice-Hall, 1963.
5. Nahmias, S. and Demmy, W.S., The Logarithmic Poisson Gamma Distribution: A Model for Leadtime Demand, paper presented at ORSA - FIMS Joint National Meeting, Houston TX, October 1981.
6. Johnson, N. L. and Kotz, S., Distributions in Statistics. Continuous Univariate Distributions, v. 2, pp. 1-7, Houghton Mifflin, 1970.
7. Navy Fleet Material Support Office Operations Analysis Report 128, User's Manual for SA (Aviation Afloat and Ashore Allowance Analyzer), by J. W. Sari, R. J. Gabriel, J. P. Shinskie and B. B. Sloan, 1 March 1977.
8. Efron, B., "Bootstrap Methods: Another Look at the Jackknife", The Annals of Statistics, V. 7, No. 1, pp. 1-26, 1979.
9. Naval Postgraduate School Report NPS55-81-005, The New Naval Postgraduate School Random Number Package-LIRANDOMI, by P.A.W. Lewis and L. Uribe, February 1981.
10. Johnson, N. L. and Kotz, S., Distributions in Statistics. Continuous Univariate Distributions, v. 1, pp. 112-120, Houghton Mifflin, 1970.

## BIBLIOGRAPHY

Croston, J. D., "Forecasting and Stock Control for Intermittent Demands" Operations Research Quarterly, V. 23, pp. 289-303, September 1972.

Ross, S. M., Applied Probability Models with Optimization Applications, Holden-Day, 1970.

Silver, E. A., "Operations Research in Inventory Management: A Review and Critique", Operations Research, V. 29, pp. 628-645, July-August 1981.

Sweet, A. L., "An Ad Hoc Method for Forecasting Series with Zero Values", AIEE Transactions, V. 12, pp. 97-103, March 1980.

U.S. Army Inventory Research Office Report 263, Integrated Forecasting Techniques for Secondary Item Classes - Part I - Active Items, by E. Gotwals III and D. Orr, September 1980.

U.S. Army Inventory Research Office Report 263, Integrated Forecasting Techniques for Secondary Item Classes - Part II - Inactive Items, by E. Gotwals III, September 1980.

INITIAL DISTRIBUTION LIST

	<u>No. Copies</u>
1. Defense Technical Information Center Cameron Station Alexandria, VA 22314	2
2. Defense Logistics Studies Information Exchange U. S. Army Logistics Management Center Fort Lee, VA 23807	2
3. Library, Code 0142 Naval Postgraduate School Monterey, CA 93940	2
4. Commander ATTN: Code 04A Naval Supply Systems Command Washington, DC 20376	1
5. Department Chairman, Code 55Mt Department of Operations Research Naval Postgraduate School Monterey, CA 93940	1
6. Professor Peter A. W. Lewis, Code 55Lw Department of Operations Research Naval Postgraduate School Monterey, CA 93940	1
7. Professor Alan W. McMasters, Code 54Mg Department of Administrative Sciences Naval Postgraduate School Monterey, CA 93940	1
8. Professor Donald P. Gaver Jr., Code 55Gv Department of Operations Research Naval Postgraduate School Monterey, CA 93940	1
9. Professor Russell Richards, Code 55Rh Department of Operations Research Naval Postgraduate School Monterey, CA 93940	1

10. Commanding Officer 1  
ATTN: Code 93  
Navy Fleet Material Support Office  
Mechanicsburg, PA 17055
11. Commanding Officer 1  
ATTN: Code 799  
Ships Parts Control Center  
Mechanicsburg, PA 17055
12. Mr. Bernard Rosenmann, Chief 1  
U. S. Army Inventory Research Office  
Room 800, Custom House  
2nd and Chestnut Sts.  
Philadelphia, PA 19106
13. Mr. V. J. Presutti Jr., Code XRS 1  
Operations Analysis Office  
Headquarters Air Force Logistic Command  
Wright-Patterson Air Force Base, OH 45433
14. Commanding Officer 1  
ATTN: LCDR Charles F. Taylor, Code 9RA  
Navy Fleet Material Support Office  
Mechanicsburg, PA 17055
15. Commanding Officer 1  
ATTN: LCDR Mark L. Yount, Code SDB4-A  
Aviation Supply Office  
700 Robbins Avenue  
Philadelphia, PA 19111