

MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A


PHOTOGRAPH THIS SHEET

AD A125037

DTIC ACCESSION NUMBER



LEVEL



INVENTORY

Terminal Forecast Reference Notebook
For Heidelberg A B, Germany
DOCUMENT IDENTIFICATION Final, 1 Oct. 82

DISTRIBUTION STATEMENT A


Approved for public release;
Distribution Unlimited

59

DISTRIBUTION STATEMENT

| | |
|--------------------|---|
| ACCESSION FOR | |
| NTIS | GRA&I <input checked="" type="checkbox"/> |
| DTIC | TAB <input type="checkbox"/> |
| UNANNOUNCED | <input type="checkbox"/> |
| JUSTIFICATION | |
| | |
| | |
| BY | |
| DISTRIBUTION / | |
| AVAILABILITY CODES | |
| DIST | AVAIL AND/OR SPECIAL |
| A | |

DISTRIBUTION STAMP



DATE ACCESSIONED




"Original contains color plates: All DTIC reproductions will be in black and white"

83 02 025 066

DATE RECEIVED IN DTIC

PHOTOGRAPH THIS SHEET AND RETURN TO DTIC-DDA-2

AD A125037



TERMINAL FORECAST REFERENCE NOTEBOOK

FOR

HEIDELBERG AB, FRG

APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED

PUBLISHED BY
DETACHMENT 3
7TH WEATHER SQUADRON
2ND WEATHER WING (MAC)
UNITED STATES AIR FORCE
1 OCTOBER 1982

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

| REPORT DOCUMENTATION PAGE | | READ INSTRUCTIONS BEFORE COMPLETING FORM |
|---|-----------------------|--|
| 1. REPORT NUMBER | 2. GOVT ACCESSION NO. | 3. RECIPIENT'S CATALOG NUMBER |
| 4. TITLE (and Subtitle) TERMINAL FORECAST REFERENCE NOTEBOOK FOR HEIDELBERG AB, GERMANY | | 5. TYPE OF REPORT & PERIOD COVERED FINAL |
| | | 6. PERFORMING ORG. REPORT NUMBER |
| 7. AUTHOR(s) | | 8. CONTRACT OR GRANT NUMBER(s) |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS DETACHMENT 3, 7TH WEATHER SQUADRON APO NEW YORK 09102 | | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS |
| 11. CONTROLLING OFFICE NAME AND ADDRESS Headquarters, 2nd Weather Wing Aerospace Sciences Division (2WW/DN) APO New York 09012 | | 12. REPORT DATE 1 Oct 82 |
| | | 13. NUMBER OF PAGES 59 (including covers) |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) | | 15. SECURITY CLASS. (of this report) Unclassified |
| | | 15a. DECLASSIFICATION/DOWNGRADING SCHEDULE |
| 16. DISTRIBUTION STATEMENT (of this Report) APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED | | |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) | | |
| 18. SUPPLEMENTARY NOTES | | |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) | | |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number) | | |

DD FORM 1473
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

TERMINAL FORECAST REFERENCE NOTEBOOK (TFRN)

SECTION I

LOCATION AND TOPOGRAPHY

DETACHMENT 3, 7TH WEATHER SQUADRON

1. Physical Features. In describing the European weather, we are apt to compare the European continent with the United States. Although there may be no marked differences in the properties of the air masses invading both continents, the weather observed is sometimes notably different than one might expect due to the interaction with the European physical environment. In contrast to the United States, there is no extensive south to north mountain barrier like the western highland upon which is superimposed a series of mountain ranges. Instead the principal mountain systems are arranged in a general west-east direction and are described as follows:

a. The Alps are the highest European mountain chain and form the southern boundary of Germany. They extend from southern France well into southeastern Austria. In the main ridge of the mountains are many peaks higher than 10000 ft.

b. The Pyrenees chain, separating France from Spain, has peaks extending above 9000 feet.

c. The Apennines stretch the entire length of Italy and have peaks which rise to 8500 feet in central Italy.

d. The Scandinavian mountains are orientated in a southwest to northeast direction. The highest peaks rarely exceed 7000 feet.

2. Besides the above mountain barriers, there are numerous smaller mountainous regions in Europe which have their characteristic effects upon invading air masses. In Germany, there are many areas where the elevation raises to 3000 feet above the surrounding valleys causing small scale weather variations. The most important of these local groups of smaller mountains and hills are:

a. The Black Forest in SW Germany with tops of over 4000 feet.

b. The Haardt and Hunsruck mountains covering areas west of the Rhine river.

c. The Schwabisch Alps between Stuttgart and Ulm.

d. The Odenwald to the east of the Rhine near Heidelberg and Mannheim.

e. The Taunus mountains to the north of Frankfurt.

f. The Eifels and Westerwalds in the Bonn and Koblenz area.

g. The Rothaar mountains to the SW of Kassel.

h. The Harz SE of Hannover.

i. The Vogelsberg between Giessen and Fulda.


j. The Spessarts east of Frankfurt.

k. The Rhon mountains near Fulda.

l. The Thuringerwald to the east of Fulda.

- m. The Frankenwald SW of Hof.
- n. The Frankenalbs S of Nurnberg.
- o. The Bohemian Forest along the Czechoslovakian border.
- p. The Bavarian Alps forming the boundary with Switzerland.

3. Influence of Topography. Because of the absence of the north-south mountain barrier, the relatively warm Atlantic maritime air masses can invade the entire land area. Therefore, there are no abrupt changes in climate across western and central Europe. In general, the mountain barriers extending west-east across southern Europe merely separate the warm southern region from the cooler areas to the north. With the numerous mountains and hills south of 52°N in Germany, winds from virtually any direction will produce upslope effects. The effects of upslope may produce extensive areas of low stratus, fog, and precipitation, or, in many cases, the effects may produce only very localized conditions.



ROTHAR

WESTERWALD

TAUUS

SPESARTS

VOGELGER

THURINGERWALD

FRANKENWALD

ODENWALD

HUNSRICK

HARDT

FRANKEN ALPS

BAVARIAN ALPS

**SCHARZWALD
(BLACK FOREST)**

SCHWABISCH ALPS

BAVARIAN ALPS

**FIGURE 1
1-3**

LOWE

AIRFIELD WEATHER BRIEF

HEIDELBERG ARMY AIRFIELD, GERMANY
49° 24'N, 008° 39'E
Elevation 357 feet

TERRAIN SUMMARY: Heidelberg AAF lies in the Rhine River Valley approximately two miles southwest of the city of Heidelberg, which lies astride the Neckar River as it flows into the Rhine Valley.

The topographic features that dominate the local area are relatively complex. The valley elevations average 300 to 500 feet above sea level and is described as gently rolling. The majority of the open land in the valley is devoted to farming. Forming the eastern edge of the Rhine Valley, approximately two nautical miles east of Heidelberg AAF rise the Odenwald Mountains. The Haardt Mountains form the western border of the Rhine Valley approximately 25 miles west of Heidelberg AAF and are oriented generally north-south. The elevations in these mountain chains average from 1500 to 2000 feet MSL.

AIRFIELD WEATHER: The predominant type of air mass effecting the Heidelberg area during all seasons is maritime polar which produces a typical maritime climate with mild winters and cool summers. The Haardt mountains influence the local weather by acting as a partial barrier to the predominant west-northwesterly flow of moist air. The Odenwald and Haardt mountains form the east and west walls of the natural Rhine basin in which air often stagnates causing poor visibilities.

CLOUDINESS AND VISIBILITY: The typical synoptic situation effecting the local area during autumn and winter is a high pressure cell centered over Western Europe. The air mass is generally moist and stable with wet, cloudy, and foggy weather prevalent. When the high is displaced northward and eastward, the source region of the air flow is continental polar and good flying conditions prevail for two to ten days. Haze and smoke, however, will reduce visibilities after the second day in the shallow cold air unless the high pressure system is centered over northern Germany which results in moderate easterly gradient, good visibility, and occasional gusty winds.


The autumn frontal activity is generally marked by a well-defined wind shift, wide band of clouds and steady precipitation with little temperature change at the surface. Frontal systems lifted over the dome of colder air takes on the characteristics of warm frontal occlusions eliminating much of the surface phenomena. Frequency of autumn and winter frontal passages averages 4 to 10 per month from October through March.

During late spring and early summer, the Azores high is displaced northward and becomes stronger. The prevailing circulation over the area is west to northwest. The moist maritime air is warmed as it moves over the land mass of Europe and combined with mechanical lift from the terrain features causes a relatively high frequency of convective showers and thunderstorms. The frequency of thunderstorms per month is 1 to 3 during April, May, and September; and 6 to 8 during June, July, and August. Most summertime fronts are relatively weak, characterized by narrow cloud and precipitation belts.

Fog, smoke and haze are the main causes of low visibilities throughout the year. During the spring and summer months nocturnal cooling will cause fog to form during the early morning hours. This phenomenon occurs 5 to 6 days a month from May through August, but seldom persists beyond 0300 to 0900 LST. During the fall and winter, fog occurs more frequently and persists for longer periods of time (averages 7-10 times per month from November through March). Low stratus ceilings are a frequent occurrence during the late fall and winter. When wind velocity is sufficient to preclude fog formation, the low ceilings have a tendency to form near sunset and intensify during the night hours. They tend to lift and burn off during the late morning and afternoon hours the following day. Stratus ceilings below 1000 feet have a frequency of 6 to 8 days per month during this period.

PRECIPITATION: Precipitation occurs frequently and quite uniformly throughout the year. Measurable precipitation is recorded on an average of 150 days a year (41%). During the late spring and summer, precipitation occurs most frequently as brief convective showers or thundershowers. In the fall and winter it occurs as either rain or snow (the frequency of occurrence of rain is approximately 75% while snow and freezing rain is approximately 25%).

WINDS: Surface winds are seldom a problem to flying at Heidelberg AAF. Normally gusty surface winds are associated with thunderstorm activity or with strong drainage winds down the western slopes of the Odenwald mountains at night. Gusts greater than 30 knots are infrequent.



514

EDZN
515

2710

EDEN

Main

642

50°N

2090

EDAM

EDOT

EDAW

EDDF

640

637

636

1778

EDEH

639

EDOF

657

1571

1864

2250

Pollution Source

EDOR

1891

EDAS

712

729

Neckar

EDIE 1857

734 K-Stuhl

2208

Rhine

724

723

3054

EDRS

708

EDAM

714

742

EDOP

745

49°N

1841

1568

1877

727

LFQP

186

EDAL

722

725

739

737

EDOS

738

LFST

190

EDAN

809

1-6

METEOROLOGICAL INSTRUMENTATION

Heidelberg AAF is equipped with the following meteorological instrumentation:

- a. ML-24 psychrometer located inside an instrument shelter, with aerator.
- b. RO₂GMO-20 Wind Recorder with wind vane located on west side of field, 3.2 meters above the ground near the helicopter pads. Wind recorder in observer section.
- c. GMO-11 Wind Indicator located in forecaster section but hooked into the RO₂GMO-20 system.
- d. GMO-10 Indicator located in observer section with transmissometer instrumentation located parallel to runway.
- e. GMO-13 Rotating Beam Ceilometer Readout in observer section with instruments located near runway. The RBC has a 400 ft baseline.
- f. ML-102-E Aneroid Barometer, ML-512 Mercurial Barometer, and ML-563A/UM Barograph for pressure.
- g. Also located in the observer area is a helium tank to fill 10 gram balloons for cloud height estimation.

Heidelberg Army Airfield
Weather Equipment




FIGURE 3

[GAS] 1-8 / 6

TERMINAL FORECAST REFERENCE NOTEBOOK (TFRN)

SECTION II

CLIMATIC AIDS

1. Operationally significant weather elements/thresholds at EDIE.

a. Winds: See Table 1.

b. Ceiling/Visibility:

- (1) Cat 5 \geq 3000'/3.0NM
- (2) 3000'/3.0NM $>$ Cat 4 \geq 1000'/2.0NM
- (3) 1000'/2.0NM $>$ Cat 3 \geq 500'/0.9NM
- (4) 500'/0.9NM $>$ Cat 2 \geq 300'/0.4NM
- (5) 300'/0.4NM $>$ Cat 1

c. Meteorological Watch Advisory (MVA) Criteria:

- (1) Snow accumulation \geq 0.5 inch on paved surfaces.
- (2) Thunderstorms within 5NM of EDIE.

d. Weather warning criteria are in accordance with USAREUR Reg 115-2 and include:

- (1) Tornadoes.
- (2) Surface winds \geq 49 knots.
- (3) Surface winds \geq 34 knots but less than 49 knots.
- (4) Hail \geq 3/4 inch.
- (5) Hail \geq 1/4 inch but less than 3/4 inch.
- (6) Freezing precipitation.
- (7) \geq 6 inches of snow accumulation in 12 hours.
- (8) \geq 2 inches but less than 6 inches of snow accumulation in 12 hours.
- (9) \geq 2 inches of rain in 12 hours.
- (10) 20°F (11°C) change in temperature from max-max/min-min.
- (11) Temperature of 0°F (-18°C) or less.

2. This section includes the following climatic aids:

a. Monthly wind roses for winds less than seven (7) knots and for winds equal to or greater than seven (7) knots.

b. Monthly climatological ceiling/visibility probabilities stratified in three hour blocks for the following criteria:

(1) Equal to or greater than 3000'/3.0NM.

(2) Less than 3000'/3.0NM.

(3) Less than 1000'/2.0NM.

(4) Less than 500'/0.9NM.

(5) Less than 300'/0.4NM.

WEATHER EFFECTS
US ARMY AIRCRAFT

TABLE 1. LIMITATIONS

(The final go/no decision is the pilots')

| | C-12/U-21 | UH-1 | AH-1 | OH-58 | CH-47 |
|------------------------|---|---|---|---|--|
| SUSTAINED SURFACE WIND | (1) 30 kts (start) | 30 kts (start) | 30 kts (start) | 20 kts (autorotation) 45 kts (start) | 30 kts (start) |
| GUST SPREAD | (1) 15 kts (start) | 15 kts (start) | 15 kts (start) | 10 kts (autorotation) 15 kts (start) | |
| TAIL WIND | (1) 30 kts (hover) | 30 kts (hover) | 30 kts (hover) | | 25 kts (start) 10 kts (start w/o auxillary power unit) |
| CROSS WIND | (1) 35 kts (hover) | 35 kts (hover) | 35 kts (hover) | 10 kts (autorotation) | |
| ICING | SEVERE LIGHT (CAN FLY IF CIG ≥ 010, TOPS ≤ 080 and IN/OUT) | SEVERE CAN'T FLY IN ANY ICING (NO IFR) | SEVERE CAN'T FLY IN ANY ICING (NO IFR) | SEVERE CAN'T FLY IN ANY ICING (NO IFR) | SEVERE |
| TURBU-LENCE | SEVERE (2) | SEVERE (2) | SEVERE (2) | SEVERE | SEVERE |

NOTE: A. Aircraft Dash-10 TM Used As Reference.

(1) Limitations for these aircraft vary according to the experience level of the pilot and various other factors. It is not possible to list all combinations.

(2) Moderate turbulence limitations are:

(a) Prohibited when the report or forecast is based on transport type aircraft.

(b) Permitted when the report or forecast is based on helicopters or light aircraft under 12,500 pounds gross weight.

B. AR Reg 95-1/USAREUR Sup 1 covers the weather minimums for uncontrolled airspace.



C. DOD FLIPS cover the weather minima at each airfield.

D. This information is provided only to inform the weather briefer of weather limitations to US Army aircraft, not to provide the weather briefer with go/no go decision responsibility.

SURFACE WINDS

LESS THAN 7 KTS (49.3%)

EQUAL TO/GREATER THAN 7 KTS (25.9%)



CALM (24.8%)

SCALE





ANNUAL

SURFACE WINDS

LESS THAN 7 KTS (49.3%)

EQUAL TO/GREATER THAN 7 KTS (28.5%)





CALM (22.3%)

MONTH: JANUARY

SURFACE WINDS

LESS THAN 7 KTS (46.8%)

EQUAL TO/GREATER THAN 7 KTS (28.0%)





CALM (25.2%)

MONTH: FEBRUARY


SURFACE WINDS

LESS THAN 7 KTS (46.5%)

EQUAL TO/GREATER THAN 7 KTS (31.3%)



SCALE





CALM (22.2%)

MONTH: MARCH


SURFACE WINDS

LESS THAN 7 KTS (46.9%)

EQUAL TO/GREATER THAN 7 KTS (33.7%)



SCALE






CALM (19.3%)

MONTH: APRIL

SURFACE WINDS

LESS THAN 7 KTS (49.2%)

EQUAL TO/GREATER THAN 7 KTS (36.7%)





CALM (23.9%)

MONTH: MAY

SURFACE WINDS

LESS THAN 7 KTS (50.3%)

EQUAL TO/GREATER THAN 7 KTS (22.4%)





CALM (27.3%)

MONTH: JUNE


SURFACE WINDS

LESS THAN 7 KTS (50.5%)

EQUAL TO/GREATER THAN 7 KTS (22.5%)



SCALE





CALM (27.0%)

MONTH: JULY


SURFACE WINDS

LESS THAN 7 KTS (51.1%)

EQUAL TO/GREATER THAN 7 KTS (23.2%)



SCALE






CALM (25.6%)

MONTH: AUGUST

SURFACE WINDS

LESS THAN 7 KTS (49.2%)

EQUAL TO/GREATER THAN 7 KTS (23.0%)






CALM (27.5%)

MONTH: SEPTEMBER

SURFACE WINDS

LESS THAN 7 KTS (50.1%)

EQUAL TO/GREATER THAN 7 KTS (20.1%)





CALM (29.7%)

MONTH: OCTOBER


SURFACE WINDS

LESS THAN 7 KTS (51.3%)

EQUAL TO/GREATER THAN 7 KTS (25.4%)



SCALE





CALM (23.2%)

MONTH: NOVEMBER


SURFACE WINDS

LESS THAN 7 KTS (49.2%)

EQUAL TO/GREATER THAN 7 KTS (27.2%)



SCALE



CALM (23.6%)

MONTH: DECEMBER

FREQUENCY OF OCCURRENCE %

JANUARY

| | ≥ 3000/3. | < 3000/3. | < 1000/2.0 | < 500/.3 | < 300/.4 |
|-------|-----------|-----------|------------|----------|----------|
| 00-02 | 46.7 | 53.3 | 28.1 | 11.8 | 6.8 |
| 03-05 | 44.4 | 55.6 | 30.1 | 12.1 | 7.1 |
| 06-08 | 39.3 | 60.7 | 34.8 | 14.1 | 6.9 |
| 09-11 | 35.2 | 64.8 | 40.2 | 16.0 | 7.7 |
| 12-14 | 41.0 | 59.0 | 33.1 | 12.4 | 6.1 |
| 15-17 | 42.3 | 57.7 | 33.0 | 12.9 | 5.9 |
| 18-20 | 44.7 | 55.3 | 30.6 | 11.8 | 5.3 |
| 21-23 | 46.4 | 53.6 | 29.7 | 11.7 | 6.5 |

FEBRUARY

| | ≥ 3000/3. | < 3000/3. | < 1000/2.0 | < 500/.9 | < 300/.4 |
|--|-----------|-----------|------------|----------|----------|
| | 50.8 | 49.2 | 29.6 | 12.3 | 7.8 |
| | 48.1 | 51.9 | 30.3 | 13.6 | 8.5 |
| | 40.5 | 59.5 | 38.2 | 18.9 | 11.1 |
| | 39.8 | 60.2 | 41.0 | 18.5 | 7.8 |
| | 44.9 | 55.1 | 31.9 | 12.2 | 4.1 |
| | 49.9 | 50.1 | 29.1 | 11.0 | 3.8 |
| | 51.5 | 48.5 | 29.1 | 11.6 | 4.4 |
| | 51.9 | 48.1 | 27.0 | 10.9 | 5.7 |

MARCH

| | | | | | |
|-------|------|------|------|-----|-----|
| 00-02 | 47.6 | 52.4 | 9.8 | 2.2 | 1.0 |
| 03-05 | 68.1 | 31.9 | 13.7 | 3.9 | 2.3 |
| 06-08 | 50.1 | 49.9 | 28.5 | 9.0 | 4.4 |
| 09-11 | 50.6 | 49.4 | 26.9 | 5.8 | 2.2 |
| 12-14 | 64.4 | 35.6 | 14.5 | 2.8 | .5 |
| 15-17 | 74.3 | 25.7 | 9.2 | 1.6 | .4 |
| 18-20 | 77.3 | 22.7 | 8.1 | 1.3 | .3 |
| 21-23 | 77.9 | 22.1 | 7.6 | 1.7 | .7 |

APRIL

| | | | | | |
|--|------|------|------|-----|-----|
| | 81.4 | 18.6 | 5.9 | 1.0 | .5 |
| | 72.7 | 27.3 | 11.2 | 2.9 | 1.3 |
| | 60.6 | 39.4 | 19.7 | 4.0 | 1.8 |
| | 62.9 | 37.1 | 13.5 | .9 | .1 |
| | 74.5 | 25.5 | 5.7 | .4 | .1 |
| | 84.1 | 15.9 | 3.0 | .5 | .1 |
| | 87.7 | 12.3 | 3.1 | 1.2 | .0 |
| | 80.6 | 19.4 | 4.0 | .9 | .2 |

FREQUENCY OF OCCURRENCE *

MAY

| | ≥ 3000/3. | < 3000/3. | < 1000/2.0 | < 500/.9 | < 300/.4 |
|-------|-----------|-----------|------------|----------|----------|
| 00-02 | 81.5 | 18.5 | 3.9 | .8 | .3 |
| 03-05 | 76.7 | 23.3 | 9.3 | 1.8 | .6 |
| 06-08 | 68.8 | 31.2 | 12.9 | .9 | .5 |
| 09-11 | 77.1 | 22.9 | 5.7 | .2 | .1 |
| 12-14 | 68.0 | 32.0 | 2.2 | .1 | .0 |
| 15-17 | 77.5 | 22.5 | 2.0 | .0 | .0 |
| 18-20 | 91.1 | 8.9 | 2.2 | .0 | .0 |
| 21-23 | 85.9 | 14.1 | 1.8 | .3 | .0 |

JULY

| | | | | | |
|-------|------|------|------|-----|-----|
| 00-02 | 86.0 | 14.0 | 4.3 | .7 | .3 |
| 03-05 | 76.3 | 23.7 | 10.7 | 2.9 | 1.7 |
| 06-08 | 69.2 | 30.8 | 12.6 | 1.4 | .6 |
| 09-11 | 79.3 | 20.7 | 4.8 | .3 | .0 |
| 12-14 | 86.8 | 13.2 | 1.9 | .0 | .0 |
| 15-17 | 91.9 | 8.1 | 1.1 | .1 | .0 |
| 18-20 | 92.1 | 7.9 | 2.5 | .3 | .0 |
| 21-23 | 75.8 | 24.2 | 3.2 | .5 | .1 |

JUNE

| | ≥ 3000/3. | < 3000/3. | < 1000/2.0 | < 500/.9 | < 300/.4 |
|--|-----------|-----------|------------|----------|----------|
| | 61.0 | 39.0 | 2.9 | 1.2 | .2 |
| | 72.4 | 27.6 | 13.7 | 2.8 | 1.2 |
| | 62.9 | 37.1 | 15.8 | 2.6 | 1.2 |
| | 73.8 | 26.2 | 7.5 | .5 | .2 |
| | 82.3 | 17.7 | 3.3 | .2 | .0 |
| | 87.8 | 12.2 | 2.5 | .1 | .1 |
| | 89.7 | 10.3 | 3.2 | .4 | .1 |
| | 88.2 | 11.8 | 3.1 | .5 | .0 |

AUGUST

| | | | | | |
|--|------|------|------|-----|-----|
| | 80.8 | 19.2 | 5.5 | 1.2 | .5 |
| | 74.2 | 25.8 | 11.5 | 3.2 | 1.5 |
| | 65.2 | 34.8 | 17.0 | 3.1 | 1.4 |
| | 73.9 | 26.1 | 7.9 | .4 | .0 |
| | 86.4 | 13.6 | 3.3 | .1 | .0 |
| | 90.9 | 9.1 | 1.9 | .1 | .0 |
| | 92.3 | 7.7 | 2.5 | .1 | .0 |
| | 89.6 | 10.4 | 3.0 | .2 | .2 |

FREQUENCY OF OCCURRENCE %

SEPTEMBER

| | ≥ 3000/3. | < 3000/3. | < 1000/2.0 | < 500/.9 | < 300/.4 |
|-------|-----------|-----------|------------|----------|----------|
| 00-02 | 77.0 | 23.0 | 10.3 | 2.2 | 1.3 |
| 03-05 | 62.9 | 37.1 | 20.7 | 8.5 | 5.8 |
| 06-08 | 50.2 | 49.8 | 30.9 | 12.2 | 7.6 |
| 09-11 | 61.9 | 38.1 | 17.6 | 3.5 | 1.2 |
| 12-14 | 79.7 | 20.3 | 6.2 | .7 | .1 |
| 15-17 | 86.6 | 13.4 | 3.9 | .3 | .1 |
| 18-20 | 86.4 | 13.6 | 4.7 | .4 | .0 |
| 21-23 | 84.4 | 15.6 | 5.6 | .5 | .3 |

NOVEMBER

| | | | | | |
|-------|------|------|------|------|------|
| 00-02 | 48.8 | 51.2 | 28.6 | 12.7 | 8.0 |
| 03-05 | 43.5 | 56.5 | 30.2 | 13.3 | 9.1 |
| 06-08 | 36.1 | 63.9 | 37.2 | 16.9 | 10.7 |
| 09-11 | 39.4 | 60.6 | 36.4 | 14.0 | 7.0 |
| 12-14 | 45.3 | 54.7 | 28.5 | 10.1 | 4.2 |
| 15-17 | 48.5 | 51.5 | 27.0 | 10.7 | 5.3 |
| 18-20 | 50.4 | 49.6 | 24.4 | 11.4 | 6.5 |
| 21-23 | 49.8 | 50.2 | 26.1 | 11.6 | 7.3 |

OCTOBER

| | ≥ 3000/3. | < 3000/3. | < 1000/2.0 | < 500/.9 | < 300/.4 |
|--|-----------|-----------|------------|----------|----------|
| | 57.8 | 42.2 | 23.3 | 12.8 | 9.9 |
| | 47.9 | 52.1 | 31.7 | 16.6 | 12.7 |
| | 37.4 | 62.6 | 44.1 | 12.9 | 16.7 |
| | 44.8 | 55.2 | 34.4 | 14.0 | 7.9 |
| | 61.8 | 38.2 | 19.3 | 5.6 | 1.2 |
| | 67.9 | 32.1 | 17.3 | 5.6 | 2.2 |
| | 67.9 | 32.1 | 17.9 | 6.8 | 3.8 |
| | 66.3 | 33.7 | 17.5 | 9.5 | 6.0 |

DECEMBER

| | | | | | |
|--|------|------|------|------|-----|
| | 39.6 | 60.4 | 30.1 | 12.8 | 8.1 |
| | 38.3 | 61.7 | 31.1 | 13.2 | 8.3 |
| | 33.2 | 66.8 | 36.6 | 15.1 | 8.3 |
| | 32.9 | 67.1 | 39.4 | 16.9 | 8.8 |
| | 36.2 | 63.8 | 32.9 | 12.0 | 5.8 |
| | 38.2 | 61.8 | 33.2 | 13.3 | 6.5 |
| | 41.0 | 59.0 | 31.1 | 11.7 | 6.2 |
| | 40.9 | 59.1 | 31.1 | 12.0 | 7.0 |

TERMINAL FORECAST REFERENCE NOTEBOOK (TFRN)

SECTION III

APPROVED LOCAL FORECAST STUDIES AND FORECASTER HINTS

APPROVED LOCAL FORECAST STUDIES

FORECASTER HINTS

SURFACE WINDS 35 KNOTS OR GREATER

1. Surface winds equal to or greater than thirty-five knots are not a major problem at Heidelberg (EDIE), but when they do occur the actual forecasting of these winds is difficult. Upstream winds at the higher elevations can significantly exceed 35 knots and yet EDIE will not experience these strong winds. EDIE is located on the eastern edge of the Rhine Valley, about three miles from the eastern ridge line. The east-west distance between the ridges is approximately 30 miles and the hill tops are in excess of 1800 feet. The regular windflow is deflected by lee side eddy and examining the circulation model of these eddies, the deflection of the wind explains the lower speeds at EDIE.

2. Winds outside of thunderstorms can be in excess of 35 knots with strong cold FROPA or TROF passage. The wind profile of these systems breaks down the eddy effect and allows the downrush from the cold air to reach the surface at EDIE.

a. General observation has indicated a positive correlation between the temperature/pressure change from Hahn AB (EDAH) to EDIE.

(1) There are three positive indicators needed prior to using this studv.

- (a) Strong cold FROPA or TROF passage.
- (b) Wasserkuppe (10544) winds greater than 44 knots.
- (c) Winds already meeting warning criteria at EDAH.

(2) If the above criteria is met, than compare the EDIE and EDAH temperature and ALSTG for the following criteria.

- (a) Temperature difference 3°C or more.
- (b) ALSTG difference is .08 inch or more.

(3) If all of the above criteria are met, then EDIE winds should equal or exceed 35 knots but the maximum gust should not exceed the Wasserkuppe's winds minus 10 knots.

b. The start of the warning should not be earlier than 1+30 from the time the above criteria is first observed and the maximum time of the warning should not exceed 5 hours.

3. If sustained winds are @ 25 knots and moderate RASH or CB/TSTMS occur, expect to exceed warning criteria for winds.

1. Fog:

a. Moderate easterly flow aloft (even though surface winds may be calm) either prevents dense fog formation at Heidelberg or keeps it one to two categories above potential visibility reduction for the conditions.

b. Downslope winds, even one to three knots, will keep visibility up in a similar manner. These winds are from 010 through 120 degrees.

c. Visibility, which is category 3 or less in fog during the night then improves markedly well before dawn with heavy dew, rarely will go below category 3 during the morning hours.

2. Ceiling: When EDIE has a low cloud ceiling, expect a one to two category drop in the ceiling with the approach and passage of an active trough or frontal system.

3. Freezing Precipitation: When surface temperature are below freezing due to radiational cooling and easterly cold advection from the Siberian High, beware of an apparently stalled frontal system in France that appears unable to breakdown the high pressure and move eastward. At the upper levels (700MB and above), high pressure will frequently breakdown very rapidly leaving a cold dome of air (surface to 850MB) over which warm air is overrunning. Freezing precipitation will begin within 12 hours of the initial movement of the front.

4. Fronts and Trofs. "Normal" frontal movement of north/south oriented fronts is approximately 12 hours from Paris to Heidelberg.

1

TERMINAL FORECAST REFERENCE NOTEBOOK (TFRN)

SECTION IV

WEATHER CONTROLS

PART I: SEASONAL DISCUSSIONS

SPRING AND SUMMER WEATHER

1. Pressure Systems: The summer circulation is characterized by the intensification and northward displacement of the Azores anticyclone together with the weakening of the Icelandic Low.
2. Air Masses: The principal European air masses are either polar or tropical, and they can have a maritime or continental source region and/or trajectory. Original characteristics obtained in the source region may be more or less transformed depending on the trajectory the air will follow. A detailed description of European air masses can be found in 2WWP 105-12, Chapter 3, Section E.
3. Fronts: During the summer, all frontal zones become somewhat indistinct because of the weaker temperature gradients and less favorable wind fields. The Atlantic polar front is located farther north than during the winter. It separates maritime tropical air masses from modified polar air masses and the mean position extends from Iceland to northern Scandinavia.
4. Temperatures: Rapid warming begins in March and continues until August. In summer, mean daily maximums are in the 70's, with overnight lows in the 50's. Hot weather occurs occasionally with easterly flow. In September, a distinct cooling trend begins.
5. Winds: The low level flow over central and southern Germany is predominantly northwesterly during the summer; however, many localized deviations are created by terrain. Due to cooling at night, drainage winds from the hills may cause local changes in the mean daytime wind direction and speed. Windspeeds generally are highest in the afternoon, in association with maximum solar heating, and lowest around sunrise. In summer, strong winds are uncommon but they can occur with thunderstorm activity.
6. Precipitation: Summertime precipitation over Europe is principally air mass type convective activity. Weak fronts or troughs and the associated lifting will often produce widespread shower activity. VFR usually cannot be maintained in summer rainshowers.
7. Thunderstorms: Summertime heating results in frequent shower activity over the interior region during May through late September. Thunderstorms, however, are generally associated with troughs or weak frontal systems. Although non-frontal thunderstorms are uncommon, afternoon cumulus may develop rapidly over ridge lines due to orographic lifting and cause shower activity with reduced ceilings. Although German thunderstorms do not build as high as the storms over the southern U.S., they are no less severe. During the transitional springtime, although the Icelandic Low is weakening and giving way to the Azores High, major polar outbreaks are not uncommon and can produce Heidelberg's most severe thunderstorms due to large air mass temperature contrasts.
8. Fog and Haze: Poor visibility is common in all seasons in southern Germany, although during May through August, visibility is greatly improved over the winter trends. The restriction to visibility is likely to be a thin layer of morning fog or haze that "burns off" rapidly by midmorning and lifts the visibility above VFR minimums.

9. Icing: In summer, icing in the low levels is generally not a problem, because of relatively high freezing levels. Summertime icing is principally associated with convective activity and is usually above the 5000 ft level.

10. Turbulence: Turbulent conditions exist about 30% of the time for aircraft flying below 5000 ft. Terrain influences are one of the prime considerations when trying to avoid turbulence. The leeside of ridge lines normal to the wind direction will be the likely place for turbulent air when the wind speed is 25 knots or greater.

AUTUMN AND WINTER WEATHER

1. Pressure System: In winter, the Azores high pressure cell is much less intense, while the high pressure system over Asia sometimes extends as far westward as France and Spain. The Icelandic low pressure system is quite deep and extensive.

2. Air Masses: Air masses, like those during the spring and summer, continue to have polar and tropical source regions. The polar air masses are of particular importance and some characteristics are illustrated in the following examples.

a. Maritime polar is the most common air mass found over Europe. The air mass properties will depend upon the over water trajectory; however, all maritime polar invasions are characterized by their relatively warm temperatures and high relative humidities. Since the air is usually warmer than the continent, stable conditions generally prevail. Stratus-type clouds, light rain or drizzle, and moderate to poor visibilities with fog and haze characterize the associated weather.

b. Continental polar air invades Germany during periods of easterly flow around an intense Russian or Scandinavian high pressure cell. Since continental polar air is cold and dry, one can expect clear skies and good visibilities during the first stages. As the anticyclonic circulation becomes more intense, temperature inversions will develop, and visibilities will be reduced in the stable air due to smoke and haze. If cyclonic activity is taking place in the Mediterranean, warm maritime air may overrun the cold polar air causing heavy snowfall over Germany.

3. Fronts: During the winter, most cold fronts which invade Europe originate over the North Atlantic and have a long over water trajectory. The cold air behind cold fronts is modified by the relatively warm Atlantic Ocean to such an extent that in most cases continuity is very difficult to maintain on surface analyses. Upon reaching the relative cold European continent, the air which travels behind the cold front is usually warmer than the air over Europe. The cold front loses its characteristics and can be analysed as a trough. Poorest frontal weather conditions are associated with pre-warm frontal situations. Ceilings less than 500 feet and a visibility less than 1 mile are common in such situations. Stationary cold fronts associated with a weak westerly flow may also produce low ceilings and visibilities.

4. Temperatures: During autumn and winter, differences between the mean

temperatures of the warmest and coldest month is about 20°F. A distinct cooling trend begins in September and by late October daily maximums are generally in the mid 50's with overnight lows in the mid 40's. During winter, the central core of Europe is cold while the north and western part is warm due to the warm oceans currents. The predominant westerly air flow produces relatively mild temperatures. Western Europe is occasionally affected by cold outbreaks from the large cold air mass source region to the east. During winter the daily maximum temperatures generally range from the mid 30's to the mid 40's and the minimums range from the high 20's to the mid 30's.

5. Winds: The low level flow over central and southern Germany is predominantly southwesterly in winter, however, local deviations are created by terrain. Valleys oriented within 20 degrees of the major axis of the air flow tend to channel the wind and may increase its speed to well above 35 knots. Infrequent occurrences of gale force winds are normally associated with winter time cold frontal passages. At elevations above 1500 feet, gale force winds occur on an average of two or three days a month. Diurnal variation in wind speed is less pronounced in winter at low levels. Wind speeds are generally highest in the afternoon, in association with maximum solar heating, and lowest around sunrise.

6. Precipitation: Winter precipitation is usually more widespread and persistent, but generally less intense than in summer. Precipitation, unless very light, will always lower the effective ceiling and reduce the horizontal and slant range visibility. Freezing rain, although not frequent, may occur a few times each year and has to be considered as a principal hazard to aircraft operations.

7. Thunderstorms: During the winter, thunderstorms do occur on rare occasions but are mainly confined to the months May through September.

8. Fog and Haze: Winter and autumn are the seasons having the most days with fog. During the period of October through March, frequent haze occurs on an average of 15 to 25 days per month, reducing the visibility to less than 5 miles. In winter the restriction to visibility is likely to be thick fog or dense haze coupled with extensive cloud cover. Due to the low sun angle, short days, and the heavy cloud/fog cover, little "burning off" of the fog or haze occurs during the day. It is not unusual for the poor visibility to persist over the entire area for several days during the winter season.

9. Icing: The high frequency of low cloudiness and freezing levels near the surface result in conditions favorable for icing on 15 to 20 days per winter month. About 3/4 of the icing is in the form of "rime" ice while the remainder is normally of the "mixed" variety.

10. Turbulence: The main causes of turbulence are the thermal discontinuities associated with fronts and tight pressure gradients and their interaction with the surrounding terrain. While the factors causing turbulence vary in intensity by season, the overall frequency remains about the same throughout the year. Terrain influences are one of the prime considerations when trying to avoid low level turbulence.

11. Cloudiness: Due to the rugged terrain over central Europe, the amount of cloudiness and ceilings have a wide variation. The average amount of cloudiness

is 70% or greater during the entire year. The elevation of the stations and the protection from the prevailing air flow effects the distribution of cloudiness and ceilings. The windward slopes have a greater amount of cloudiness and lower ceilings than the leesides, and lower ceilings are, of course, observed at stations with higher elevations. Maximum cloudiness and lowest ceilings occur in winter. During the morning cloudiness is greater than during the afternoon in winter. Over northern Germany there is a tendency for low clouds to persist into the afternoon. Stratocumulus associated with high pressure cells are quite persistent in the Ruhr Valley and other northern German industrial areas. Low ceilings and visibilities are often observed with the approach of a secondary trough. Ceilings below 1000 feet are frequently observed under such conditions. Low clouds are also observed with slow moving cold fronts approaching northern Germany and the lowlands from a northerly direction. When these fronts stagnate over this region, low ceilings will persist for days. Very low clouds with continuous precipitation are usually associated with an outbreak of cold moist air from a northerly direction, especially in the region between Brussels and Cologne. Southern Germany has more than 15 days per month with overcast skies during the winter. The mean cloud amounts are greatest in the morning. Low clouds are predominately stratiform or thick stratocumulus. At the majority of stations in southern Germany the ceiling is below 2000 feet on more than half of the winter mornings.

PART II: GENERAL WEATHER PHENOMENA

1. The seasonal cyclone tracks and seasonal anticyclone tracks are presented in Figures 1 and 2, respectively, pertinent tracks are referred to when they relate to operationally significant forecast problems. Forecast problems are categorized by month in Table 2. Discussion is topic oriented based upon forecast problems. A standard analysis package evolves from the problem discussion. Theory is not presented. Prior to going to Table 2 problems, a definition of the short wave trough is presented.

2. The Short Wave Trough: The short wave trough dominates the European weather forecasting problem. The short wave trough is defined via the following example: Consider the typical winter situation with the quasi-stationary Icelandic Low (a planetary scale feature) northwest of England. Frequently a synoptic scale low breaks out of the Icelandic Low moving along one of the cyclone tracks in Figure 1. This low is associated with a traveling, synoptic scale wave in the westerlies. This wave is not the short wave, since it can be easily tracked in upper air analysis and machine progs. However, there are short waves associated with the traveling low. These scales of motion are depicted in Figure 3. Most of the short waves will be surface features, but some waves will have increased amplitude detectable at gradient and 850MB levels as a wind shift or possibly a small-scale velocity maximum. This is a vertically deep short wave trough. To identify short waves, analyze surface isobaric patterns at 1 to 2 MB intervals and then analyze gradient and 850MB level flow lines to find which troughs are vertically deep. The short wave troughs give a spider web character to the traveling synoptic scale surface low. These short wave troughs have a wave length varying from a few to several hundred kilometers and the frequency of passage (at a point) varies anywhere from 90 minutes to 9 to 12 hours. The shorter the wave, the more rapidly it travels and the less the impact upon the weather. For example, the 90 minute waves may merely increase cloudiness and produce drizzle/rain while 9 to 12 hour waves may bring significant showers (Total-Totals values must be considered). On the 90 minute end of the scale, every 3rd or 4th wave may be detectable at the gradient level, while at the 9 to 12 hour end, they are all likely to be detectable at gradient level. The waves detectable at gradient and 850MB levels are important to identify and prog because the associated weather response is much more significant than that with the intervening shallow surface waves. This point will be stressed repeatedly as problem areas listed in Table 2 are systematically addressed.

3. The Siberian High: The Siberian High expands westward into Europe several times during the average winter. The associated cold front usually pushes through Germany to become stationary in central France. This nearly always occurs in conjunction with off shore blocking (Omega High) at 500MB and a deep cold core low at 500MB over Russia. The situation persists until the block breaks down (usually 7 to 10 days). Surface isotherms pack at the frontal boundary, but the leading edge of the front (area of Germany) is too shallow to analyze at 850MB. You can find the frontal inversion on a SKEW-T (usually between 500 to 1,200 feet AGL). The front comes into the central West Germany as a persistent cold easterly wind (8-12 knots). The 2nd day after frontal passage, daytime heating produces persistent stratus under the frontal inversion. In Bavaria, the front wedges up the mountains giving upslope fog/stratus below the frontal inversion and clear skies above. (See Figure 4). When the long wave pattern begins to shift, the Siberian High will recede eastward as a warm front. There will normally not be freezing precip associated with this withdrawal.

4. Persistent Stratus: Persistent stratus occurs with four synoptic situations. The stratus forms and breaks with passage of short wave troughs. The four situations are:

- a. The Siberian High, discussed above.
- b. In advance of a warm front (the stratus breaks are "sucker holes").
- c. Upslope Conditions: Stratus breaks occur when the wind direction changes with short wave trough passage. Following trough passage, stratus reforms if synoptic scale gradient level winds return to upslope direction.

- d. Post Cold Front Stratus (NW flow): Gradient level winds above 20 knots produce 3,000 to 4,000 foot ceilings overnight (often going scattered in late morning), and gradient winds below 20 knots produce 2,000 to 2,500 foot ceilings. Conditions are lower if a short wave trough passes. The dynamic mechanism is a combination of solar and turbulent mixing.

5. Fog (Radiation Versus Upslope): A general problem in central Germany is determination of the fog formation mechanism, radiation versus upslope. Upslope fog/stratus requires a surface inversion several hundred feet deep, moist air within this layer, and the proper wind direction. Use GMGO Upslope and Lee Effects Maps to infer proper wind direction, Upslope fog/stratus will not break with heating. Consider the following classic example in Figure 5. This model affects the Finthen AAF Terminal, but is of interest to all units which clear aircraft. A stationary trough sets up over the English Channel (through 500MB). This produces persistent SW flow. Given a stable air mass and ample moisture at low levels (a result of SW flow), you have potential upslope conditions (use a SKEW-T to evaluate stability, moisture content of the air, and depth of the surface inversion). The fog is in the ridge east of the channel trough. There is no front in the trough even though air mass characteristics vary substantially over a broad zone. The first fog day is frequently radiation fog. Radiation fog will break at 0900 to 1100L, but often visibility will not go above four to five miles. Temperatures are above 0°C (often four to five °C). Use SKEW-T (depth of moisture, etc. IAW AWS/TR-79/006) to forecast the breakout time. Winds at gradient level must be 10 knots or below for radiation fog onset, e.g., a short wave may come through and break the fog out of cycle. When the moisture reaches amount/depth needed for upslope fog (a surface inversion with air near saturation through 500 feet or more), the fog will not break following sunrise. It won't break until the wind direction or air mass changes. A typical upslope pattern is, Day 1: Ceiling 500/1 all day; Day 2: Ceiling 200/1/2 all day and moist column deepens; Day 3: Cooler temperatures with drizzle and continued 200/1/2 stratus/fog. Considering forecast specification categories (AWSR 105-27), upslope conditions are often above 200/1/2 but below 1000/1 before sunrise, but fall below 200/1/2 after sunrise, and then return to the below 1000/1 interval after sunset. Now consider the short wave trough as defined earlier. As a short wave moves through, wind direction and speed change and stability lessens. Fog reverts to radiation mechanism and the area normally breaks wide open. Trough timing is critical. When the trough arrives, the fog breaks. After the short wave passes, the synoptic scale fog reverts to the upslope pattern. Without an air mass change upslope fog will recur. Superficially, the day-to-day sequence of events appears to be a series of chaotic unforecastable variations wherein one-day fog/stratus conditions

break after sunrise and the next day they worsen after sunrise. However the sequences can be forecast. To do this, analyze/prog the short waves, and determine during each hour of the forecast whether the driving force is upslope or radiation. Note that the ends of the upslope season are not clear-cut; it depends on the synoptic pattern. See Table 2.

6. Freezing Precipitation: The primary seasons are late fall and early spring, but given the right synoptic situation, freezing precipitation can occur throughout the winter. Mixed precipitation (rain/snow/etc) tends to be associated with the retreat of the Siberian High. This precipitation is mixed or snow, dependent upon temperatures within the advancing air mass. Freezing precipitation usually comes ahead of the warm front associated with a cyclone on the southern track. The surface must be frozen (preferably for several days). Otherwise, the heat capacity of the soil will prevent freezing precipitation. There are two variations: widespread and spotty freezing precipitation.

a. Widespread Freezing Precipitation: If the 0°C isotherm at 850MB is trailing (behind) the warm front, snow will occur ahead of the warm front (or rain dependent on position of 0°C isotherm at the surface), and freezing precipitation will not occur (See Figure 6). Figure 7 is the model for freezing precipitation. The key is the position of the 850MB 0°C isotherm in relation to the warm front. As the warm front passes, freezing precipitation will continue if the surface air is close to or below freezing in the warm sector. The ice may be covered by snow as the cold front passes. Alternately, heavy rain in the warm sector will usually melt existing ice. When predicting freezing precipitation, start time is the arrival of the 0°C 850MB isotherm and stop time is the arrival of the warm front (except as noted above).

b. Spotty Freezing Precipitation: (See Figure 8). Here the 0°C isotherm only goes north of the warm front in the cold pockets associated with the meso-scale lows. There are two components of motion to predict. The short wave (meso-low) component (toward the northwest) and the warm frontal component. The vector resultant is usually toward the northeast. The areas of freezing precipitation are small. They may track continuously or they may jump in response to orographic and other effects. In this situation, you can make good area forecasts, but point forecasting is very difficult.

7. Air Mass Thunderstorms: The key ingredients are warm air advection, a deep layer of low-level moisture, and solar heating. These factors produce the instability. A typical pattern is a stationary, modifying air mass. After a few days, a few thunderstorms occur in late afternoon. The next day thunderstorms are widespread. The overturning resulting from the thunderstorms stabilizes the air mass, and the following day there are fewer storms again. The cycle repeats until the air mass changes. On any given day, analyze the TT and prog the maxima. Use Table 3 to make the forecast. Anywhere in the day to day sequence discussed above, the arrival of a short wave trough will trigger locally numerous thunderstorms day or night. Concurrent arrival of a vertically deep short wave trough (detectable at gradient or 850MB) and a TT maximum will occasionally trigger severe weather. However, the TT max and the short wave must be independently progged as they move at different rates, and the short wave will tend to outrun the instability maximum. Thunderstorms, of course, are always related to orography based on the extra lift from flow over a ridge or suppressive lee effects.

8. Thundershowers, Rainshowers, and Snowshowers: There are two models, the short wave trough and the closed (synoptic-scale) low.

a. Short Wave Trough: The short-wave problem is identical to that discussed with air mass thunderstorms. The only difference is the spotty showers, associated with air mass instability alone, are not present. Showers (rain or snow, dependent upon temperature) will be generated by the trough and move with it. Shower intensity depends on the TT values arriving at the time the trough arrives. The vertically deep trough will produce heavy showers (wintertime thunder/snow showers are even possible if TT are high enough). See Figure 9.

b. Closed Low: The closed surface low may or may not be associated with a frontal system. See Figures 10 and 11. The low usually follows the northern track. Analyze the 850MB isotherms to determine whether or not there are fronts and for frontal placement. When analyzing occlusions, sometimes the 700MB level is better than the 850MB level (the occlusion may be above 850MB). It is nearly impossible to identify a European front from surface data alone. Trough associated cloudiness can mask solar heating to give the appearance of a front when none exists. When fronts are not present, showers will be less widespread and will closely correlate to the passage of short wave troughs. Frontal lows will also be overtaken by short wave troughs, but the trough will merely intensify the showers on a background of rain. Use isallobaric analysis to predict the track and rate of movement of the low, extrapolate observed frontal movements, and pay special attention to the vertically deep short wave trough. Overlay a progged TT analysis on the progged pressure system analysis to determine event intensity.

9. Continuous Rain or Snow: These events are associated with the closed low (with and without fronts) just discussed. Continuous precipitation is a background condition related to the synoptic scale vertical motion pattern which the heavy showers come with passage of short wave troughs. Heavy snow is always associated with a warm or occluded warm front in winter. The retreat of the Siberian High will produce heavy snow in the southern and eastern areas of West Germany as it moves out as a warm front.

10. Trough Associated Winds: There are two models, those accompanying frontal systems and those not associated with frontal systems.

a. The frontal model is seen in Figure 12. This is the winter windstorm model. Once more you prog the low with isallobaric analysis and extrapolate the fronts. The key is to recognize that the strong winds are not associated with frontal pressure gradients, but with a vertically deep short wave trough passing through the system. The wind max will outrun the low and be dissipated in the ridge ahead of the low. Do an isotach analysis at gradient level and use extrapolation tools.

b. The channel trough discussed with upslope conditions is the fall and spring windstorm model. See the sequence in Figure 13. Note that there are no fronts and the wind does not change direction (at any one point) as the max moves through the major trough (the wind max is a compression wave). In the fall you tend to have more stable air (low TT) and showers are less likely to accompany the windstorm. The spring season can go either way but often when a TT max is associated, thunderstorms develop in the compression wave. The keys are: Identify the wave while it is still on the back side of the major trough, and

prog the wind isotach maximum using extrapolation.

11. Summary: From the foregoing, it is evident that local analysis is the key. It takes study, practice, and frequent forecast reviews/re-analysis to become a skilled European forecaster, but this is achievable in three to six months. The emphasis must be on:

a. Surface analysis (one to two MB analysis in key quadrants to identify short waves and analysis of closed isallobars).

b. Gradient level analysis (identify short waves, windstorm indicators, and upslope fog conditions).

c. 850MB to 700MB isothermal analysis (locate fronts and identify deep short waves).

d. Analysis and propping of the Total-Totals fields.

12. An important issue is whether or not a trough has a front (a boundary between two air masses) associated with it. Unfortunately, centralized products do not provide reliable guidance on frontal type or presence. Furthermore, centralized progs do not retain small scale synoptic features like the short wave trough and the compression wave wind maximum. Read the analysis models in the ECI 2570 course frequently, and view caramate follow-on-training products regularly. Finally, when long range forecasts are needed, the Baur Type catalog can be a useful source. Whenever a classic Baur pattern is observed, the situation will persist at least as long as the longer indicated time mode. The shorter time models are apparently the result of classifying mixed Baur Types (rather than discarding them).

PRIMARY FORECASTING PROBLEMS BY MONTH

SIBERIAN HIGH
CPSK

PERSISTENT STRATUS

RAD FOG

UPSLOPE/RAD FOG

RAD
FOG

AIRMASS
TRW

AIRMASS
TRW

4-10

A

S

N

D

J

F

M

A

M

J

J

A




TABLE 2


TOTAL-TOTALS INDEX VALUES GUIDE FOR FORECASTING THUNDERSTORMS

IN CENTRAL GERMANY

(MAY NEED ADJUSTING FOR SPECIFIC LOCATIONS IN GERMANY)


| FORECAST | VT | CT | TT |
|-----------|------|---------|---------|
| ISOLATED | 26 | 20 - 23 | 46 - 49 |
| FEW | ≥ 28 | 20 - 24 | 50 - 52 |
| SCATTERED | ≥ 28 | 22 - 26 | 53 - 55 |
| NUMEROUS | ≥ 28 | ≥ 26 | ≥ 55 |

TABLE 2



MEAN PATHS OF CYCLONES BY SEASON. —→ PRIMARY TRACK; - - -→ SECONDARY TRACK.

FIGURE 1



MEAN PATHS OF ANTICYCLONES BY SEASON. ———▶ PRIMARY TRACK; - - -▶ SECONDARY TRACK.

FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7

15 W 4-18




FIGURE 8




FIGURE 9

NO FRONTS




FIGURE 10

WITH FRONTS




FIGURE 11





FIGURE 12



TYPICAL URSLOPE FLORA

CURRENT LEVEL

FIGURE 136

60

50

4-24





FIGURE 13b



GRADIENT LEVEL AT
TIME - 776 HOURS

FIGURE 13c

15w
4-26

40





FIGURE 13d



GRADIENT LEVEL AT
TIME - T + 18 Hours

FIGURE 13e

15 W 4-28

60 + . . . +

50 + . . . +

40 + . . . +

35 N

