

AD-A125 430

ESTIMATION OF NOISY TELEGRAPH PROCESSES: NONLINEAR
FILTERING VERSUS NONLINEAR MASSACHUSETTS INST OF TECH
CAMBRIDGE STATISTICS CENTER Y YAO FEB 83 TR-29-ONR
N00014-75-C-0555

1/1


UNCLASSIFIED

F/G 20/1

NL



END
FILED
4 83
074



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A 125 430



ESTIMATION OF NOISY TELEGRAPH PROCESSES:
NONLINEAR FILTERING VS. NONLINEAR SMOOTHING

BY
YI-CHING YAO

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

TECHNICAL REPORT NO. ONR 29

FEBRUARY 1983

PREPARED UNDER CONTRACT
N00014-75-C-0555 (NR-609-001)
FOR THE OFFICE OF NAVAL RESEARCH



Reproduction in whole or in part is permitted for
any purpose of the United States Government

This document has been approved for public release
and sale; its distribution is unlimited

83 03 07 054

DTIC FILE COPY

AD A 125 430

Estimation of Noisy Telegraph Processes:
Nonlinear Filtering vs. Nonlinear Smoothing

Abstract

In the estimation problem of a two-state stationary Markov process with Gaussian white noise added, the optimal smoother is a two-filter smoother. In a special case, we compare analytically the optimal nonlinear filter and smoother and find that the latter is significantly better than the former when either the noise intensity or the rate of jump of the states is low.

Key words: Nonlinear filtering, nonlinear smoothing, telegraph process

MMS 1980 subject classification: Primary 93E14; Secondary 62M05, 93E11

Accession For	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
ERIC GRAI			
ERIC I/S			
Unannounced			
Justification			
By			
Distribution/			
Availability Codes			
Avail and/or			
Dist. Special			

A

1. Introduction

In a certain class of linear dynamic Gaussian systems, the optimal smoothing estimator of the states may be regarded as a two-filter smoother. See Wall, et al (1981) for a complete discussion. Several authors, e.g. Rauch, et al (1965) and Mehra and Bryson (1968) compared the performance of filters and smoothers in linear dynamic systems and found that in certain cases smoothers may be more preferable even at the expense of time delay.

In this paper, we consider the system of a telegraph process in the presence of additive Gaussian white noise, and study the relationship between the optimal filtering and smoothing estimators of the states. To be specific, define the signal process $(u_t)_{-\infty < t < \infty}$ to be a stationary two-state Markov process such that

$$\Pr(u_t = 1) = p = 1 - \Pr(u_t = -1)$$

$$(1.1) \Pr(u_{t+h} = 1 | u_t = 1) = 1 - \nu h + o(h)$$

$$\Pr(u_{t+h} = -1 | u_t = -1) = 1 - \nu h + o(h)$$

where $\nu = (1-p)\nu'$. Let the observed process

$$(1.2) Z_t = \int_0^t u_s ds + \alpha W_t, \quad t \in (-\infty, \infty)$$

where (W_t) is a standard Wiener process ($W_0 = 0$)



independent of $\{u_t\}$ and α is a positive constant to represent the intensity of the noise.

In the next section, we show that the optimal smoothing estimator of u_t is a function of the forward and backward filtering estimators of u_t . In Section 3, we compare analytically the performance of the optimal nonlinear filter and smoother in a special case.

2. Optimal Nonlinear Smoother as a Two-Filter Smoother

In this section, we assume the process $\{Z_t\}$ is observed from $t = 0$ on. Denote by \hat{u}_t^L , \hat{u}_t^R and \hat{u}_t the left-sided, right-sided and two-sided conditional expectations of u_t , i.e. $E(u_t | Z_0^t)$, $E(u_t | Z_t^T)$ and $E(u_t | Z_0^T)$, respectively, where T is a fixed time span (possibly ∞), $0 < t < T$, $Z_0^t \equiv \{Z_u; 0 \leq u \leq t\}$ and $Z_t^T \equiv \{Z_u; t \leq u < T\}$. Wonham (1965) showed that \hat{u}_t^L satisfies a stochastic differential equation. We can easily see that \hat{u}_t^R must satisfy a reversed-time stochastic differential equation of the same type. Now, the following proposition tells us that the smoothing estimate \hat{u}_t of u_t can be easily computed from knowing \hat{u}_t^L and \hat{u}_t^R .

Proposition 2.1

$$\hat{u}_t = \frac{(1 + \hat{u}_t^L)(1 + \hat{u}_t^R) - 2p(1 + \hat{u}_t^L \hat{u}_t^R)}{(1 + \hat{u}_t^L)(1 + \hat{u}_t^R) - 2p(\hat{u}_t^L \hat{u}_t^R)}$$

$$\text{i.e. } \tanh^{-1} \hat{u}_t = \tanh^{-1} \hat{u}_t^L + \tanh^{-1} \hat{u}_t^R + \frac{1}{2} \log \frac{1-p}{p}$$

In particular, when $p = 1/2$,

$$\hat{u}_t = (\hat{u}_t^L + \hat{u}_t^R) / (1 + \hat{u}_t^L \hat{u}_t^R)$$

$$\text{i.e. } \tanh^{-1} \hat{u}_t = \tanh^{-1} \hat{u}_t^L + \tanh^{-1} \hat{u}_t^R$$

Proof of Proposition 2.1

Denote by $f_{z_0^T}(\cdot)$ (or $f_{z_0^T}(\cdot | u_t = \pm 1)$) the Radon-

Nikodym derivative of the measure on $C[0, T]$ induced by z_0^T (or z_0^T conditional on $u_t = \pm 1$, respectively) with respect to the Wiener measure on $C[0, T]$. The existence of the derivatives is a consequence of Girsanov's Theorem (see [1], Theorem 7.2). Let $b = 1$ or -1 . Using the independence of Z_0^t and Z_t^T given u_t ,

$$\Pr(u_t = b | z_0^T = z_0^T) = f_{z_0^T}(z_0^T | u_t = b) \Pr(u_t = b) / f_{z_0^T}(z_0^T)$$

$$= f_{z_0^T}(z_0^T | u_t = b) \Pr(u_t = b)$$

$$= f_{z_0^T}(z_0^T | u_t = b) f_{z_t^T}(z_t^T | u_t = b) \Pr(u_t = b)$$

$$= \Pr(u_t = b | z_0^T = z_0^T) \Pr(u_t = b | z_t^T = z_t^T) / \Pr(u_t = b)$$

So,

$$\frac{\Pr(u_t=1|z_0^T=z_0^T)}{\Pr(u_t=-1|z_0^T=z_0^T)} = \frac{\Pr(u_t=1|z_0^L=z_0^L)}{\Pr(u_t=-1|z_0^L=z_0^L)} \frac{\Pr(u_t=1|z_t^T=z_t^T)}{\Pr(u_t=-1|z_t^T=z_t^T)} \frac{\Pr(u_t=-1)}{\Pr(u_t=1)}$$

Since $\hat{u}_t^L = \Pr(u_t=1|z_0^L=z_0^L) - \Pr(u_t=-1|z_0^L=z_0^L)$, etc.

$$\frac{1+\hat{u}_t}{1-\hat{u}_t} = \frac{1+\hat{u}_t^L}{1-\hat{u}_t^L} \frac{1+\hat{u}_t^R}{1-\hat{u}_t^R} \frac{1-p}{p}$$

Observing $2 \tanh^{-1} x = \log(1+x)(1-x)^{-1}$, we complete the proof. \square

3. Comparison Between Filtering and Smoothing

In this section, we study the performance of the optimal nonlinear filter and smoother. In order to derive some analytic results, we consider estimates based on an infinite time span and $p = 1/2$. Denote by q_t^L, q_t^R and q_t the left-sided, right-sided and two-sided conditional expectations of u_t , i.e. $E(u_t|z_{-}^L)$, $E(u_t|z_{+}^R)$ and $E(u_t|z_{-}^L, z_{+}^R)$, respectively. Define

$$(3.1) \text{MSE}(q_t^L) \equiv E\{E(u_t|z_{-}^L) - u_t\}^2 \equiv E(q_t^L - u_t)^2$$

and

$$(3.2) \text{MSE}(q_t) \equiv E\{E(u_t|z_{-}^L, z_{+}^R) - u_t\}^2 \equiv E(q_t - u_t)^2$$

It should be noted that $\text{MSE}(q_t)$ is constant for $t < (-\infty, \infty)$ but $\text{MSE}(q_t^L) < \text{MSE}(q_t^R) = \text{MSE}(q_t^L)$ for $t < 0 \leq u < v$, for when $t < 0$

$$(3.3) L(q_t^L|u_t) = L(E(u_t|z_{-}^L, z_{+}^R, \theta \leq t)|u_t) \\ = L(E(u_t|z_{-}^L, z_{+}^R, \theta > t)|u_t)$$

where $L(Y)$ is the distribution of random variable Y and $L(Y|X)$ is the conditional distribution of Y given X .

In the following, we only consider $t \geq 0$.

We can readily modify the proof of Proposition 2.1 to show

Proposition 3.1

$$q_t = \frac{q_t^L + q_t^R}{1 + q_t^L q_t^R}$$

Proposition 3.2 (Wonham (1965))

(A)

$$\Pr(q_t^L \in [q, q+dq] | u_t=1) = c(\gamma) (1+q^2)^{-2} \exp[-2\gamma(1-q^2)^{-1/2}dq]$$

where

$$c(\gamma) = \int_1^{\infty} z^{1/2} (z-1)^{-1/2} e^{-2\gamma z} dz^{-1}$$

and

$$\gamma = \alpha^2 v$$

(B)

$$\begin{aligned} \text{MSE}(q_t^L) &= \int_0^{\infty} z^{-1/2} (z+1)^{-1/2} e^{-2\gamma z} dz / \int_0^{\infty} z^{-1/2} (z+1)^{1/2} e^{-2\gamma z} dz \\ &= \begin{cases} -2\gamma \log \gamma + o(\gamma \log \gamma) & (\gamma \rightarrow 0^+) \\ 1 - (4\gamma)^{-1} + o(\gamma^{-2}) & (\gamma \rightarrow \infty) \end{cases} \end{aligned}$$

Proposition 3.3

$$\begin{aligned} \text{MSE}(q_t) &= c(\gamma)^2 \int_{-1}^1 \int_{-1}^1 \left[\frac{x+y}{1+xy} - 1 \right]^2 (1+x)(1-x^2)^{-2} (1+y)(1-y^2)^{-2} \\ &\quad \cdot \exp(-2\gamma[(1-x^2)^{-1} + (1-y^2)^{-1}]) dx dy \\ &= \begin{cases} 2\gamma + o(\gamma) & (\gamma \rightarrow 0^+) \\ 1 - \frac{1}{2}\gamma^{-1} + o(\gamma^{-2}) & (\gamma \rightarrow \infty) \end{cases} \end{aligned}$$

Proof of Proposition 3.3

Since $\{u_t\}$ is time reversible, $L(q_t^R | u_t) = L(q_t^L | u_t)$.

Also, q_t^R is independent of q_t^L given u_t . Therefore, by applying Propositions 3.1 and 3.2 (A), we can readily derive the formula for $\text{MSE}(q_t)$. The computation of its asymptotic behavior is given in the Appendix. \square

We may also compute the MSE for the Wiener filtering estimate of u_t and the best linear estimate based on Z_{t-}^u .

$\text{MSE}_W \equiv \text{MSE}$ for the Wiener filtering estimate of u_t

$$(3.4) \quad = 2\gamma[(1+\gamma^{-1})^{1/2} - 1]$$

$$= \begin{cases} 2\gamma^{1/2} + o(\gamma^{1/2}) & (\gamma \rightarrow 0^+) \\ 1 - (4\gamma)^{-1} + o(\gamma^{-2}) & (\gamma \rightarrow \infty) \end{cases}$$

$\text{MSE}_{BL} \equiv \text{MSE}$ for the best linear estimate of u_t based on Z_{t-}^u

$$(3.5) \quad = \left(\frac{\gamma}{1+\gamma} \right)^{1/2}$$

$$= \begin{cases} \gamma^{1/2} + o(\gamma^{1/2}) & (\gamma \rightarrow 0^+) \\ 1 - \frac{1}{2}\gamma^{-1} + o(\gamma^{-2}) & (\gamma \rightarrow \infty) \end{cases}$$

Now we are ready to compare the performance of the

various estimates. See Table 3.1 for the summary of the asymptotic results on their MSE.

Remark 1: As $\gamma \rightarrow 0^+$, the linear estimates are not efficient. It seems that in non-Gaussian systems linear estimates are rather inflexible and therefore can not perform well.

Remark 2: As $\gamma \rightarrow 0^+$, the optimal smoother is more efficient than the optimal filter by a factor $-\log \gamma$. This factor is about 6.9 when $\gamma = 0.001$ (e.g. $\alpha = 0.1$). Therefore, when either the noise intensity or the rate of jump of the states is low, the optimal smoother is significantly better than the optimal filter.

Remark 3: In finite-state processes, error probability is also an interesting criterion. In the following, we present the error probabilities for several optimal decision procedures. We consider decisions on whether u_0 is 1 or -1.

(i) Based on Z_{-m}^0 :

An optimal decision d_L is:

$$d_L(Z_{-m}^0) = 1 \text{ iff } \Pr(u_0=1 | Z_{-m}^0) \geq \frac{1}{2}.$$

The error probability is

$$e_L = c(\gamma) \int_{-1}^0 (1+q)(1-q^2)^{-2} e^{-2\gamma(1-q^2)^{-1}} dq \\ = \frac{1}{2} \left(1 - \frac{c(\gamma)}{\sqrt{\gamma}} e^{-2\gamma} \right)$$

$$(3.6) \quad = \begin{cases} -\frac{1}{2} \gamma \log \gamma + o(\gamma \log \gamma) & (\gamma \rightarrow 0^+) \\ \frac{1}{2} \left(1 - \frac{1}{\sqrt{2\gamma}} + o(\gamma^{-1/2}) \right) & (\gamma \rightarrow \infty) \end{cases}$$

(ii) Based on Z_{-m}^m :

An optimal decision d is

$$d(Z_{-m}^m) = 1 \text{ iff } \Pr(u_0=1 | Z_{-m}^m) \geq \frac{1}{2}$$

The error probability is

$$e = c(\gamma)^2 \int_{-1}^1 \int_{-y}^y (1-x)(1-x^2)^{-2} e^{-2\gamma(1-x^2)^{-1}} dx (1-y)(1-y^2)^{-2} \\ \cdot e^{-2\gamma(1-y^2)^{-1}} dy \\ = c(\gamma)^2 \left\{ \frac{1}{2} c(\gamma)^{-1} \gamma^{-1} e^{-2\gamma} \int_0^1 \left(1 + \frac{y}{v} \right)^{1/2} e^{-2\gamma v} dv \right. \\ \left. - \frac{1}{2} \gamma^{-2} e^{-4\gamma} \int_0^1 \left(1 + \frac{y}{v} \right)^{3/2} e^{-4\gamma v} dv + \frac{1}{8} \gamma^{-2} e^{-4\gamma} \int_0^1 \left(1 + \frac{y}{v} \right)^{1/2} \right. \\ \left. \cdot e^{-2\gamma v} dv \right\} \\ = \begin{cases} \gamma \log 2 + o(\gamma) & (\gamma \rightarrow 0^+) \\ \frac{1}{2} \left(1 - \frac{1}{\sqrt{\gamma}} + o(\gamma^{-1/2}) \right) & (\gamma \rightarrow \infty) \end{cases}$$

Table 3.1 Asymptotic Behavior of Mean Squared Errors for Various Estimates for the Noisy Two-State Markov Process

Estimate	Linear		Nonlinear	
	Filtering MSE _F	Smoothing MSE _{BL}	Filtering MSE(q _c)	Smoothing MSE(q _c)
$\gamma \rightarrow 0 +$	$2\gamma^{1/2}$	$\gamma^{1/2}$	$-2\gamma \log \gamma$	2γ
$\gamma \rightarrow \infty$	$1-(4\gamma)^{-1}$	$1-(2\gamma)^{-1}$	$1-(4\gamma)^{-1}$	$1-(2\gamma)^{-1}$

References

- [1] Liptser, R. S. and Shiriyayev, A.N. (1977). *Statistics of Random Processes. Vol. I: General Theory*. Springer-Verlag, New York.
- [2] Mehra, R.K. and Bryson, A. E. (1968). Linear smoothing using measurements containing correlated noise with an application to inertial navigation. *IEEE Trans. Aut. Control* AC-13, 496-503.
- [3] Rauch, H.E., Tung, F. and Striebel, C.T. (1965). Maximum likelihood estimates of linear dynamic systems. *AIAA J.* 3, 1445 - 1450.
- [4] Wall, J.E., Willsky, A.S. and Sandell, N.R. (1981). On the fixed-interval smoothing problem. *Stochastics* 5, 1-41.
- [5] Monham, W.N. (1965). Some applications of stochastic differential equations to optimal nonlinear filtering. *J. SIAM Control, Ser. A*, 2, 347-369.

Appendix

Proof of the Asymptotic Expansions for MSE(q.) in Proposition 3.3

In the following, c(y) is abbreviated to c. We use the notation A ~ B (y → y_0) to mean lim_{y → y_0} A/B = 1.

(I) The case of y → 0+.

$$\begin{aligned}
c^{-1} &= \int_0^1 z^{1/2} (z-1)^{-1/2} e^{-2yz} dz \\
&= \gamma^{-1} \int_0^1 u^{1/2} (u-\gamma)^{-1/2} e^{-2u} du \quad (\gamma z = u) \\
&= \gamma^{-1} \int_0^{\gamma} (v+\gamma)^{1/2} v^{-1/2} e^{-2(v+\gamma)} dv \quad (u=v+\gamma) \\
&= \gamma^{-1} \int_0^{\gamma} v^{1/2} v^{-1/2} e^{-2v} dv \\
(A.1) \quad &= \frac{1}{2} \gamma^{-1} \\
&\int_{-1}^1 \int_{-1}^1 \frac{(xy-1)^2 (1+x)(1+y)}{(1-x^2)^2 (1-y^2)^2} \exp\left(\frac{-2x}{1-x^2} + \frac{-2y}{1-y^2}\right) dx dy \\
&= \int_{-1}^1 \int_{-1}^1 \frac{1}{(1+xy)^2 (1+x)(1+y)} \exp\left(\frac{-2x}{1-x^2} + \frac{-2y}{1-y^2}\right) dx dy
\end{aligned}$$

$$(A.2) \quad \left\{ \int_{-1}^{-1+c} \int_{-1}^{-1+c} + 2 \int_{-1}^{-1+c} \int_{-1}^{-1+c} \right\} \frac{1}{(1+xy)^2 (1+x)(1+y)}$$

$$\cdot \exp\left(\frac{-2x}{1-x^2} + \frac{-2y}{1-y^2}\right) dx dy$$

(y → 0+)

Here ε is a small positive number.

$$\int_{-1}^{-1+c} \int_{-1}^{-1+c} \frac{1}{(1+xy)^2 (1+x)(1+y)} \exp\left(\frac{-2x}{1-x^2} + \frac{-2y}{1-y^2}\right) dx dy$$

$$\int_{-1}^{-1+c} \int_{-1}^{-1+c} \frac{1}{4(1+x)(1+y)} \exp\left(\frac{-2x}{1-x^2} + \frac{-2y}{1-y^2}\right) dx dy \quad (y \rightarrow 0+)$$

$$= \frac{1}{4} \left[\int_{-1}^{-1+c} \frac{1}{1+x} \exp\left(\frac{-2x}{1-x^2}\right) dx \right]^2$$

$$= \frac{1}{4} \left[\int_0^{\epsilon y^{-1}} \frac{1}{u} \exp\left[\frac{-2}{u(2-\gamma u)}\right] du \right]^2 \quad (1+x = \gamma u)$$

$$= \frac{1}{4} \left[\int_M^{\epsilon y^{-1}} \frac{1}{u} \exp\left[\frac{-2}{u(2-\gamma u)}\right] du \right]^2 \quad (M \text{ is a large number})$$

$$\frac{1}{4} \left[\int_M^{c/y} \frac{1}{u} du \right]^2$$

(A.3) $-\frac{1}{4} (\log y)^2$

$$\int_{-1}^{-1+c} \int_{1-c}^1 \frac{1}{(1+xy)^2 (1+x)(1+y)} \exp\left(\frac{-2y}{1-x} + \frac{-2x}{1-y}\right) dx dy$$

$$= \int_0^1 \int_0^1 \frac{1}{(u+v-uv)^2 (2-u)v} \exp\left(\frac{-2y}{u(2-u)} + \frac{-2x}{v(2-v)}\right) dudv$$

(u=1-x, v=1+y)

$$= \frac{1}{2} \int_0^1 \int_0^1 \frac{1}{(u+v-uv)^2 v} \exp\left(\frac{-2y}{u(2-u)} + \frac{-2x}{v(2-v)}\right) dudv$$

$$= \frac{1}{2} \int_0^1 \int_0^1 \frac{1}{(u+v)^2 v} \exp\left(\frac{-2y}{u(2-u)} + \frac{-2x}{v(2-v)}\right) dudv$$

$$= \frac{1}{2\gamma} \int_0^{c/\gamma} \int_0^{c/\gamma} \frac{1}{(x+y)^2 y} \exp\left(\frac{-2}{x(2-\gamma x)} + \frac{-2}{y(2-\gamma y)}\right) dx dy$$

(u = \gamma x, v = \gamma y)

$$= \frac{1}{4\gamma} \int_0^{c/\gamma} \int_0^{c/\gamma} \left[\frac{1}{(x+y)^2 y} + \frac{1}{(x+y)^2 x} \right] \exp\left[\frac{-2}{x(2-\gamma x)} + \frac{-2}{y(2-\gamma y)}\right] dx dy$$

dx dy

$$= \frac{1}{4\gamma} \int_0^{\pi/2} \int_0^{\pi/2} \frac{1}{xy(x+y)} \exp\left(-\frac{1}{x} - \frac{1}{y}\right) dx dy$$

$$= \frac{1}{4\gamma} \int_0^{\pi/2} \int_0^{\pi/2} \frac{1}{\cos \theta + \sin \theta} \exp[-r(\cos \theta + \sin \theta)] dr d\theta$$

(x^{-1} = r \cos \theta, y^{-1} = r \sin \theta)

(A.4) $= \frac{1}{4\gamma}$

Therefore, from (A.1), (A.2), (A.3), (A.4),

$$\text{MSE}(q_c) = 2\gamma (\gamma + 0^*)$$

(II) The case of $\gamma = \infty$.

$$c^{-1} = \int_1^{1+c} z^{1/2} (z-1)^{-1/2} e^{-2\gamma z} dz$$

$$= \int_1^{1+c} (1+(x-1))^{1/2} (x-1)^{-1/2} e^{-2\gamma x} dx \quad (c \text{ is small positive})$$

$$= \int_0^{1+y} \left[1 + \frac{1}{2}(z-1) + o((z-1)^2) \right] (z-1)^{-1/2} e^{-2yz} dz$$

$$(A.5) = e^{-2y} \left[\left(\frac{y}{2}\right)^{1/2} y^{-1/2} + 2^{-7/2} y^{1/2} y^{-3/2} + o(y^{-3/2}) \right]$$

$$\int_{-1}^1 \int_{-1}^1 \left(\frac{x+y}{1+xy} - 1 \right)^2 \frac{(1+x)(1+y)}{(1-x^2)^2(1-y^2)^2} \exp\left(\frac{-2y}{1-x^2} + \frac{-2y}{1-y^2}\right) dx dy$$

$$= \int_{-1}^1 \int_{-1}^1 \frac{1}{(1+xy)^2(1+x)(1+y)} \exp\left(\frac{-2y}{1-x^2} + \frac{-2y}{1-y^2}\right) dx dy$$

$$= \int_{-1}^1 \int_{-1}^1 (1-x-y-xy + x^2 + y^2 + o(x^2 + y^2))$$

$$\exp\left(\frac{-2y}{1-x^2} + \frac{-2y}{1-y^2}\right) dx dy$$

$$(A.6) = \left[\int_{-1}^1 \exp\left(\frac{-2y}{1-x^2}\right) dx \right]^2 + 2 \left[\int_{-1}^1 x^2 \exp\left(\frac{-2y}{1-x^2}\right) dx \right]$$

$$\cdot \left[\int_{-1}^1 \exp\left(\frac{-2y}{1-y^2}\right) dy \right] (1 + o(1))$$

$$\int_{-1}^1 \exp\left(\frac{-2y}{1-x^2}\right) dx = 2 \int_0^1 \exp\left(\frac{-2y}{1-x^2}\right) dx$$

$$= 2e^{-2y} \int_0^1 e^{-ys} s^{-1/2} (2+s)^{-1/2} ds \left(s = \frac{2x^2}{1-x^2} \right)$$

$$= 2e^{-2y} \int_0^1 e^{-ys} s^{-1/2} (2+s)^{-1/2} ds + o(s)$$

(A.7)

$$= 2e^{-2y} \left(2^{-3/2} y^{1/2} y^{-1/2} - 3 \cdot 2^{-9/2} y^{1/2} y^{-3/2} + o(y^{-3/2}) \right)$$

$$\int_{-1}^1 x^2 \exp\left(\frac{-2y}{1-x^2}\right) dx = 2 \int_0^1 x^2 \exp\left(\frac{-2y}{1-x^2}\right) dx$$

$$= 2e^{-2y} \int_0^1 e^{-ys} s^{1/2} (2+s)^{-5/2} ds \left(s = \frac{2x^2}{1-x^2} \right)$$

$$= 2e^{-2y} \int_0^1 e^{-ys} s^{1/2} (2+s)^{-5/2} + o(s) ds$$

-18-

$$(A.8) = 2 e^{-2Y} (2^{-7/2} Y^{1/2} Y^{-3/2} + o(Y^{-3/2}))$$

From (A.6), (A.7), (A.8),

$$\int_{-1}^1 \int_{-1}^1 \left(\frac{x+y}{1+xy} - 1 \right)^2 \frac{(1+x)(1+y)}{(1-x^2)^2(1-y^2)^2} \exp\left(\frac{-2Y}{1-x^2} + \frac{-2Y}{1-y^2}\right) dx dy$$

$$(A.9) = e^{-4Y} (2^{-1} Y^{-1} - 2^{-3} Y^{-2} + o(Y^{-2}))$$

So, from (A.5) and (A.9),

$$\begin{aligned} \text{MSE}(q_c) &= \frac{2^{-1} Y^{-1} - 2^{-3} Y^{-2} + o(Y^{-2})}{\left[\left(\frac{1}{2} \right) 1/2 Y^{-1/2} + 2^{-7/2} 1/2 Y^{-3/2} + o(Y^{-3/2}) \right]^2} \\ &= 1 - \frac{1}{2} Y^{-1} + o(Y^{-1}) \end{aligned}$$

This completes the proof. \square

END

DATE
FILMED

4-83

DTIC