

AD-A129 169

NONEXISTENCE OF A SHOCK LAYER IN GAS DYNAMICS WITH A
NONCONVEX EQUATION OF STATE(U) WISCONSIN UNIV-MADISON
MATHEMATICS RESEARCH CENTER R L PEGO APR 83

1/1


UNCLASSIFIED

MRC-TSR-2505 DAAG29-80-C-0041

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

MRC Technical Summary Report #2505

NONEXISTENCE OF A SHOCK LAYER
IN GAS DYNAMICS
WITH A NONCONVEX EQUATION OF STATE

ADA 129169

Robert L. Pego

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

April 1983

(Received March 10, 1983)

DTIC
SELECTE
JUN 9 1983
A

DTIC FILE COPY

Approved for public release
Distribution unlimited

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

National Science Foundation
Washington, DC 20550

83 06 07 076

- a -

UNIVERSITY OF WISCONSIN-MADISON
MATHEMATICS RESEARCH CENTER

NONEXISTENCE OF A SHOCK LAYER IN GAS DYNAMICS
WITH A NONCONVEX EQUATION OF STATE

Robert L. Pego

Technical Summary Report #2505

April 1983

ABSTRACT

A classical result of Gilbarg states that a simple shock wave solution of Euler's equations is compressive if and only if a corresponding shock layer solution of the Navier-Stokes equations exists, assuming, among other things, that the equation of state is convex. An "entropy condition" appropriate for weeding out "unphysical" shocks in the nonconvex case has been introduced by T.-P. Liu. For shocks satisfying his entropy condition, Liu showed that purely viscous shock layers exist (with zero heat conduction). Dropping the convexity assumption, but retaining many other reasonable restrictions on the equation of state, ^{Pego, the author} we construct an example of a (large amplitude) shock which satisfies Liu's entropy condition but for which a shock layer does not exist if heat conduction dominates viscosity. ^{Pego} We also give a simple restriction, weaker than convexity, which does guarantee that shocks which satisfy Liu's entropy condition always admit shock layers. ←

AMS (MOS) Subject Classifications: 76N10, 35Q10, 76L05, 35L65, 35B99

Key Words: shocks, shock layer, traveling waves, conservation laws,

admissibility, entropy condition, viscosity, heat conduction

Work Unit Number 2 - Physical Mathematics

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.
This material is based upon work supported by the National Science Foundation
under Grant No. MCS-7927062, Mod. 2.

SIGNIFICANCE AND EXPLANATION

A traditional method of determining which shocks (which are discontinuous solutions in gas dynamics or elasticity) are "physical" is to ask whether the discontinuity can be obtained in the limit of vanishing viscosity and heat conduction. For a constant profile shock, one then seeks corresponding constant profile "shock layer" solutions when viscosity and heat conduction are nonzero. If the equation of state is convex, a simple condition distinguishes those discontinuities which admit shock layers: The shock must be compressive, or entropy should increase across the jump. Convexity does not hold for some materials, however. In the nonconvex case, Liu proposed a revised "entropy condition" to select physical shocks. For small amplitude shocks, the present author has shown that a shock layer always exists if Liu's entropy condition is satisfied. Here we present an example to show that this is not the case for strong shocks, unless the equation of state satisfies some extra restrictions. We also supply some extra restrictions which are shown to be appropriate.

Distribution/	
Availability Codes	
Avail and/or	
Special	
A	

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

NONEXISTENCE OF A SHOCK LAYER IN GAS DYNAMICS
WITH A NONCONVEX EQUATION OF STATE

Robert L. Pego

§1. Introduction

In 1951, D. Gilbarg [2] showed that for each stationary shock wave solution of Euler's equations of gas dynamics in one space dimension, there exists a corresponding smooth solution (called a shock layer or shock profile) of the Navier-Stokes equations with viscosity and heat conduction, provided that the thermodynamic equation of state satisfies a short list of restrictions given by Weyl [11], including a convexity condition. The shock wave is required to satisfy the Rankine-Hugoniot jump relations, and the entropy condition, which ensures that entropy increases along particle paths in Euler's equations. Under Weyl's restrictions, the entropy condition simply requires shocks to be compressive, i.e., density must increase along particle paths.

More recently, interest has developed in equations of state which may be nonconvex ([5],[9],[10]). T.-P. Liu introduced an entropy condition appropriate for shocks in this situation. He showed that a shock satisfies his entropy condition if and only if a purely viscous shock layer exists, i.e., a shock layer for the equations with zero heat conduction [4].

The purpose of this note is to present an example concerning restrictions on the equation of state under which Liu's entropy condition does select just those shocks for which shock layers exist for any positive values of viscosity and heat conduction. We explicitly construct an equation of state such that

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.
This material is based upon work supported by the National Science Foundation under Grant No. MCS-7927062, Mod. 2.

for some shock wave which satisfies Liu's entropy condition, no corresponding shock layer can exist if the heat conduction dominates the viscosity. (A shock layer does exist if heat conduction is small compared to viscosity.) The equation of state satisfies all of Weyl's restrictions except the convexity condition, and satisfies other conditions which have appeared in recent work on the Riemann problem in gas dynamics ([5],[10]). We can give a simple restriction weaker than convexity (satisfied by van der Waals gases in regions of hyperbolicity, for example) under which all shocks satisfying Liu's entropy condition always admit corresponding shock layers. (See Theorem 1.)

An example such as we describe may be of interest in one dimensional thermo(visco)elasticity, where for quite general constitutive relations it is desirable to determine the smoothing effect of viscous and thermal dissipation (e.g. see [1]). Nonconvex equations of state also occur for materials exhibiting phase transitions. In this paper, however, we require that Euler's equations remain hyperbolic in the region of interest. Effects other than viscosity and heat conduction must be considered to determine "structure" in phase transition zones; see [8] and [9].

The Navier-Stokes equations in one space dimension, written in Lagrangian form, are

$$\begin{aligned}
 (1.1) \quad \tau_t - u_h &= 0 \\
 u_t + p_h &= (\mu u_h / \tau)_h \\
 \mathcal{E}_t + (pu)_h &= (\mu u u_h / \tau)_h + (\lambda \theta_h / \tau)_h
 \end{aligned}$$

Here h denotes the Lagrangian mass coordinate, t is time, τ is specific volume, u is velocity, p is pressure, θ is temperature, \mathcal{E} is energy density per unit mass, and μ and λ are the coefficients of viscosity and heat conduction, respectively. $\mathcal{E} = e + u^2/2$, where e is the internal energy density per unit

mass. We assume that τ and θ determine the thermodynamic state of the material, and that e , p , and the entropy S are given by sufficiently smooth equations of state $e = e(\tau, \theta)$, $p = p(\tau, \theta)$, $S = S(\tau, \theta)$. These functions are related through the Gibbs relation

$$(1.2) \quad \theta dS = de + p d\tau$$

The coefficients μ and λ may also depend smoothly on τ and θ . The quantities μ , λ , τ , θ , and p are positive. For convenience, we shall denote the triple (τ, u, ξ) by U , and the thermodynamic state by Z , specified by the pair (τ, θ) (or by (τ, p) , see assumption (1.10)).

Euler's equations are obtained from (1.1) by setting $\mu = \lambda = 0$. A shock wave traveling at speed s is a weak solution of these equations of the form

$$(1.3) \quad U(h, t) = \begin{cases} U_- & \text{for } h < st \\ U_+ & \text{for } h > st \end{cases}$$

The two end states must satisfy the Rankine-Hugoniot relations

$$(1.4) \quad \begin{aligned} 0 &= -s(\tau_+ - \tau_-) - (u_+ - u_-) \\ 0 &= -s(u_+ - u_-) + (p_+ - p_-) \\ 0 &= -s(\xi_+ - \xi_-) + (p_+ u_+ - p_- u_-) \end{aligned}$$

We call any solution (1.3) satisfying (1.4) a "simple jump solution", or "jump". It is customary to reserve the term "shock" for jumps which satisfy the entropy condition.

What determines a jump? The speed s may be determined up to sign from any suitable pair of thermodynamic states Z_+ and Z_- by the relation

$$(1.5) \quad 0 = s^2(\tau_+ - \tau_-) + (p_+ - p_-)$$

In our analysis it suffices to consider only back-facing shocks, for which $s < 0$. (Particle paths in (1.1) are vertical.) Now $u_+ - u_-$ is

determined. Imposing the third equation, we recall: The states Z_+ and Z_- determine a simple jump solution with $s < 0$ if and only if the Hugoniot function vanishes:

$$H(Z_+, Z_-) \equiv e_+ - e_- + \frac{1}{2}(p_+ + p_-)(\tau_+ - \tau_-) = 0$$

Fixing Z_0 , the set of states Z satisfying $H(Z, Z_0) = 0$ is typically a curve, called the Hugoniot curve with center Z_0 in the (τ, p) plane.

A shock layer for the jump (1.3) is a traveling wave solution $U(h-st)$ of (1.1) with $U(\xi) \rightarrow U_{\pm}$ as $\xi \rightarrow \pm\infty$. Plugging in and integrating once, $U(\xi)$ must satisfy

$$\begin{aligned} 0 &= -s(\tau - \tau_{\pm}) - (u - u_{\pm}) \\ \mu u_{\xi}/\tau &= -s(u - u_{\pm}) + (p - p_{\pm}) \\ \mu \mu u_{\xi}/\tau + \lambda \theta_{\xi}/\tau &= -s(\xi - \xi_{\pm}) + pu - p_{\pm}u_{\pm} \end{aligned}$$

The first equation determines $u(\xi)$ from $\tau(\xi)$, so we obtain

$$\begin{aligned} \lambda \theta_{\xi}/\tau &= -s(\xi - \xi_{\pm} + (\tau - \tau_{\pm})(p_{\pm} + su)) \equiv L(\tau, \theta) \\ \mu \tau_{\xi}/\tau &= \frac{1}{-s}(s^2(\tau - \tau_{\pm}) + p - p_{\pm}) \equiv M(\tau, \theta) \end{aligned} \quad (1.6)$$

These equations yield an autonomous system of ODEs in the (τ, θ) plane. It is clear that Z_+ and Z_- are rest points for this system. A shock layer exists for the jump (1.2) if and only if the system (1.5) admits a trajectory $Z(\xi)$ connecting Z_- to Z_+ , i.e., $Z(\xi) \rightarrow Z_{\pm}$ as $\xi \rightarrow \pm\infty$.

We remark that equations (1.6) are identical to those used by Gilbarg [2] for the stationary shock layer in Eulerian form, if we introduce an eulerian space variable $x = \int_{\xi}^{\infty} \tau(\eta) d\eta$ and recognize that $-s$ is the mass flux $pu = \text{const}$, where $\rho = 1/\tau$ is the density.

In order to motivate Liu's entropy condition for the jump (1.3) and examine the implications of nonconvexity, we briefly consider the simple

example of isothermal gas dynamics in Lagrangian form. The equations are

$$\begin{aligned}\tau_t - u_h &= 0 \\ u_t + p(\tau)_h &= (\mu_h/\tau)_h\end{aligned}$$

A simple jump with speed s connecting (τ_-, u_-) and (τ_+, u_+) for $\mu = 0$ satisfies

$$0 = s^2(\tau_+ - \tau_-) + p(\tau_+) - p(\tau_-)$$

A shock profile $(\tau, u)(\xi)$, with $\xi = h-st$, must satisfy (eliminating u)

$$(1.7) \quad \mu \tau_\xi / \tau = \frac{1}{-s} (s^2(\tau - \tau_\pm) + p(\tau) - p(\tau_\pm)) \equiv M(\tau)$$

Assuming that $p(\tau)$ is decreasing, but possibly nonconvex, any two states

τ_+, τ_- determine a jump with speed $s < 0$. A shock layer exists for that jump

if and only if: $\text{sgn } M(\tau) = \text{sgn } (\tau_+ - \tau_-)$ for τ between τ_+ and τ_- .

This is the "entropy condition" which selects those jumps which admit a shock

layer for these equations. The situation is pictured in Fig. 1, where the

phase portrait of (1.7) is indicated on the τ axis of the graph of $p(\tau)$.

Observe that since p is not convex, rarefaction shocks can exist admitting

shock layers, e.g. $\tau_- = \tau_2 < \tau_+ = \tau_3$. Also, a compressive jump need not

admit a shock layer, e.g. $\tau_- = \tau_4 > \tau_+ = \tau_1$.




Figure 1. Shock profiles in isothermal gas dynamics

For the nonisothermal Euler's equations, we now state Liu's strict entropy condition for a jump (1.3) determined by Z_+ , Z_- with $s < 0$:
the condition is that

$$\begin{aligned} \text{sgn } M(Z) &= \text{sgn}(\tau_+ - \tau_-) \text{ for all } Z = (\tau, p) \\ s(E) &\text{ between } Z_+ \text{ and } Z_- \text{ on the Hugoniot curve} \\ &\text{with center } Z_+ \text{ (or with center } Z_-). \end{aligned}$$

With $s < 0$, this condition simply means that the line $M(\tau, p) = s^2(\tau - \tau_{\pm}) + p - p_{\pm} = 0$ should lie above (below) the Hugoniot curve between Z_+ and Z_- if $\tau_- > \tau_+$ ($\tau_- < \tau_+$). The restrictions we shall impose on the equation of state imply that with some choice of center, the Hugoniot curve is regular between Z_+ and Z_- , so the condition $s(E)$ is well defined.

The restrictions we place on the equation of state are as follows:

$$(1.8) \quad e_\theta(\tau, \theta) > 0$$

This means that heat conduction is a dissipative effect in (1.1) for $\lambda > 0$.

$$(1.9) \quad p_\tau(\tau, S) \equiv dp/d\tau|_{S \text{ const}} < 0$$

Euler's equations are strictly hyperbolic just when this holds.

$$(1.10) \quad p_\theta(\tau, \theta) > 0$$

Thus the map $(\tau, \theta) \rightarrow (\tau, p(\tau, \theta))$ is one to one on the domain of states Ω_θ .

The image of this map we designate Ω_p . We assume

$$(1.11) \quad \Omega_p \text{ is convex.}$$

These restrictions above are the same as those imposed by Gilbarg [2], following Weyl [11], except that we omit the convexity assumption

$$d^2p/d\tau^2|_{S \text{ const}} > 0.$$

Under these restrictions, the Hugoniot curve is described by the following. The proof is deferred to §2.

Proposition 1. Fix $Z_0 = (\tau_0, p_0)$ in Ω_p . Then in the region $\tau > \tau_0$, the set $\{Z \in \Omega_p \mid H(Z, Z_0) = 0\}$ consists solely of a monotonically decreasing curve

$p = h_0(\tau)$ passing through Z_0 .

We now describe our positive results concerning existence and uniqueness of the shock layer for any given μ , λ , positive functions of τ and θ .

Theorem 1. Assume (1.8)-(1.11). Also assume

$$(1.12) \quad e_\tau(\tau, \theta) > 0 \quad \text{in } \Omega_\theta$$

Suppose Z_+ , Z_- in Ω_θ satisfy $H(Z_+, Z_-) = 0$ and (1.13) below. Then the simple jump solution determined by Z_+ , Z_- and $s < 0$ from (1.5) admits a unique shock layer if and only if the strict entropy condition $s(E)$ is satisfied.

In the absence of (1.12), compressive shocks ($\tau_- > \tau_+$ if $s < 0$) always admit a shock layer:

Theorem 2. Assume (1.8)-(1.11). Suppose Z_+ , Z_- in Ω_θ satisfy $H(Z_+, Z_-) = 0$ and (1.13) below, and suppose $\tau_- > \tau_+$. Then the conclusion of Theorem 1 holds.

Weak shocks ($|U_+ - U_-|$ small) always admit a shock layer, under the most basic of restrictions:

Theorem 3. Assume only (1.8), (1.9). Fix Z_0 in Ω_θ . Then if Z_+ and Z_- are sufficiently close to Z_0 and satisfy $H(Z_+, Z_-) = 0$, the simple jump solution determined by Z_+ , Z_- with $s < 0$ admits a shock layer (unique in a neighborhood of Z_0) if and only if the strict entropy condition $s(E)$ is satisfied.

Theorem 3 is proved in [7] as a consequence of a general theorem on singular viscosity matrices for systems of conservation laws. The proof of Theorems 1 and 2 is extremely similar to that of Gilbarg [2], and will be presented briefly in §2. The extra condition imposed is a technical "domain" condition,

$$(1.13) \quad \text{The line segment } [\tau_+, \tau_-] \times \{\theta_-\} \text{ lies in } \Omega_\theta,$$

needed so certain constructions are valid. (A similar assumption is made

tacitly by Gilbarg.)

Our main result is a "counterexample":

Theorem 4. There exists an equation of state, defined in a domain Ω_0 ,

satisfying (1.8)-(1.11), and a pair of states Z_+ , Z_- which satisfy

$H(Z_+, Z_-) = 0$ and (1.13), such that the jump solution determined by Z_+ , Z_- and $s < 0$ satisfies the strict entropy condition $s(E)$, but

- if λ/μ is sufficiently large, no corresponding shock layer exists,
- if λ/μ is sufficiently small, a unique shock layer exists.

From Theorem 2, this shock must be a rarefaction shock, $\tau_- < \tau_+$. We stress, however, that Liu's result shows that a purely viscous shock layer (with $\lambda = 0$) exists, so entropy does increase across this shock, i.e., $S(Z_-) < S(Z_+)$.

This equation of state must violate (1.12) and the convexity condition. However, the following conditions can be satisfied:

$$(1.14) \quad e_\tau(\tau, p) > 0$$

$$(1.15) \quad p_\tau(\tau, \theta) < 0$$

Condition (1.14) (along with (1.8)-(1.10), basically) was imposed by Liu [5] to guarantee that the Riemann problem for gas dynamics has a unique solution globally. Condition (1.15), along with (1.8), implies (1.9) and implies that constant states for the Navier-Stokes equations are linearly stable (see [6], [7]).

Finally, we briefly remark that although we have presented this example in the context of gas dynamics, it is valid in the context of 1D thermoviscoelasticity as well. One may redefine the equation of state outside a compact domain so that the restrictions imposed by Dafermos [1] are satisfied. The key step is to appropriately specify the function $p_0(\tau, \theta)$ globally (e.g. see section 3).

§2. Existence and uniqueness of the shock layer

We begin this section by gathering several useful thermodynamic calculations, then we proceed to prove Theorems 1 and 2. Take τ and θ as independent variables. Using (1.2), and equating mixed partials of $S(\tau, \theta)$, we obtain the standard identities

$$(2.1) \quad \theta S_{\theta}(\tau, \theta) = e_{\theta}(\tau, \theta)$$

$$(2.2) \quad \theta S_{\tau}(\tau, \theta) = e_{\tau}(\tau, \theta) + p = \theta p_{\theta}(\tau, \theta)$$

Considering τ and p independent, we get

$$(2.3) \quad \theta S_p(\tau, p) = e_p(\tau, p) = e_{\theta}(\tau, \theta) \theta_p(\tau, p)$$

$$(2.4) \quad \theta S_{\tau}(\tau, p) = e_{\tau}(\tau, p) + p$$

and also

$$(2.5) \quad p_{\tau}(\tau, p) = -S_{\tau}(\tau, p) / S_p(\tau, p)$$

The assumptions (1.8)-(1.10) imply that the derivatives

$S_{\theta}(\tau, \theta)$, $S_{\tau}(\tau, \theta)$, $S_p(\tau, p)$ and $S_{\tau}(\tau, p)$ are all positive.

The following identity helps relate the entropy condition $s(E)$ to the ODEs (1.6).

Proposition 2. Fix Z_+ and Z_- with $H(Z_+, Z_-) = 0$. Let $s < 0$ satisfy (1.5).

Now $L(\tau, \theta)$ and $M(\tau, \theta)$ in (1.6) are determined, and for any $Z = (\tau, \theta)$ we have

$$(2.6) \quad H(Z, Z_{\pm}) = \frac{1}{-s} L(Z) + \frac{1}{2} (\tau - \tau_{\pm}) (-sM)(Z)$$

It follows that

$$H(Z, Z_+) - H(Z, Z_-) = \frac{1}{2} (\tau_- - \tau_+) (-sM)(Z)$$

Proof.
$$\begin{aligned} H(Z, Z_{\pm}) + L(Z)/s &= \frac{1}{2} (u^2 - u_{\pm}^2) + (\tau - \tau_{\pm}) (\frac{1}{2} p + p_{\pm}) - p_{\pm} + su \\ &= (\tau - \tau_{\pm}) (\frac{1}{2} (u + u_{\pm}) + \frac{1}{2} p - p_{\pm}) - su = \frac{1}{2} (\tau - \tau_{\pm}) (-sM)(Z) \end{aligned}$$

(Recall that $u - u_{\pm} = -s(\tau - \tau_{\pm})$.)

Proof of Proposition 1. With Z_0 fixed, $H(Z, Z_0)$ is a function of $Z = (\tau, p)$.

Using (2.2), $H_{\tau}(\tau, p) = \theta S_{\tau}(\tau, p) + \frac{1}{2} p_0 - p > 0$ if $p < p_0$, and

$$H_p(\tau, p) = e_p(\tau, p) + \frac{1}{2}(\tau - \tau_0) > 0 \text{ if } \tau > \tau_0.$$

It is convenient now to calculate

$$(2.7) \quad \frac{1}{-s} L_\theta(\tau, \theta) = e_\theta(\tau, \theta) \quad \frac{1}{-s} L_\tau(\tau, \theta) = \theta p_\theta(\tau, \theta) + sM(\tau, \theta)$$

$$(2.8) \quad -sM_\theta(\tau, \theta) = p_\theta(\tau, \theta) \quad -sM_\tau(\tau, \theta) = s^2 + p_\tau(\tau, \theta)$$

To get L_τ , use Proposition 2 and (2.2).

Proof of Theorem 2. Let Z_+ , Z_- in Ω_0 satisfy $H(Z_+, Z_-) = 0$, (1.13), and

$\tau_- > \tau_+$, and let $s < 0$ satisfy (1.5). Suppose first that the entropy condition $s(E)$ is satisfied. Our proof, like Gilbarg's, is based on an analysis of the curves $(L = 0)$ and $(M = 0)$ which pass through Z_+ and Z_- . Since L_θ and M_θ are positive, θ is a given function of τ along each of these curves, $\theta = l(\tau)$, $\theta = m(\tau)$ respectively. We claim that $l(\tau)$ and $m(\tau)$ are defined for $\tau_+ < \tau < \tau_-$ and satisfy

$$i) \quad l(\tau) < m(\tau) \quad \text{for } \tau_+ < \tau < \tau_-$$

$$ii) \quad l'(\tau) < 0 \quad \text{for } \tau_+ < \tau < \tau_-$$

Indeed, the set $(M = 0)$ in Ω_p is a straight line segment, by (1.11). Pulling back to the (τ, θ) plane, $m(\tau)$ is defined as described. Using Proposition 2, the entropy condition $s(E)$ implies that between Z_+ and Z_- on the Hugoniot curve with center Z_+ , $L(Z) > 0 > M(Z)$. Property i) follows. Property ii) follows from i) using (2.7), since $M < 0$ on the curve $(L = 0)$. The "domain" condition (1.13), with (1.11), now implies $l(\tau)$ is defined as described.

The existence of the shock layer follows from i), ii) (see Fig. 2). The region $R = \{(\tau, \theta) \mid \tau_+ < \tau < \tau_- \text{ and } l(\tau) < \theta < m(\tau)\}$ is negatively invariant under the flow induced by (1.6). Also, any trajectory $Z(\xi)$ which starts in R must be monotone (θ increasing, τ decreasing) and tend to Z_- as $\xi \rightarrow -\infty$.

Consider a vertical line segment crossing R ,

$$\{(\tau, \theta) \mid \tau = \tau_0, l(\tau) < \theta < m(\tau)\}, \quad \tau_+ < \tau_0 < \tau_-.$$




Figure 2. Phase portrait of (1.6) for a compressive shock

A point on this segment may belong to one of three disjoint classes: the forward trajectory of (1.6) starting at the point may

- a) exit the region R on the curve ($L = 0$)
- b) exit the region R on the curve ($M = 0$)
- c) not exit the region R .

In the last case, the trajectory must tend toward Z_+ by monotonicity. The first two classes are open (continuity) and nonempty (endpoints). The third class is therefore also nonempty by connectedness, so some shock layer exists.

We turn to demonstrate the uniqueness of the shock layer. First we claim that any trajectory of (1.6) joining Z_+ and Z_- must lie entirely in the region R. Indeed, no trajectory can approach Z_+ within either of the regions ($M < 0$ and $L < 0$) or ($M > 0$ and $L > 0$). Also, the region

$$R_1 = \{(\tau, \theta) \in \Omega_0 \mid \tau < \tau_+ \text{ and } L > 0 \text{ or } M > 0\}$$

is negatively invariant. So our claim holds.

The characteristic equation for (1.6) at the critical point Z_+ always has one positive and one nonpositive root:

$$\begin{aligned}
 (2.9) \quad 0 &= \begin{vmatrix} M_T/\mu - \kappa & M_\theta/\mu \\ L_T/\lambda & L_\theta/\lambda - \kappa \end{vmatrix} \\
 &= \kappa^2 - (M_T/\mu + L_\theta/\lambda)\kappa + (\ell'(\tau_+) - m'(\tau_+))M_\theta L_\theta/\mu\lambda
 \end{aligned}$$

The constant term is nonpositive. If it is zero, then $M_T > 0$. In general, then, Z_+ is a saddle point, and the uniqueness of the trajectory approaching within R is easily established. In the degenerate case, one may construct a one dimensional center-stable manifold in a neighborhood of Z_+ , locally invariant under (1.6). A trajectory starting in this neighborhood which does not lie in the given center-stable manifold must eventually (for ξ large) leave any sufficiently small neighborhood of Z_+ (see Kelley [3]). Therefore, any two trajectories approaching Z_+ within R must lie on the same curve.

We have established the existence and uniqueness of the shock layer when $\tau_- > \tau_+$ and the entropy condition $s(E)$ is satisfied. If $\tau_- > \tau_+$ but the condition $s(E)$ fails to hold, arguments similar to those above show that no trajectory of (1.6) can connect Z_- to Z_+ . Theorem 2 (and Theorem 1 for compressive shocks) follows.

For a rarefaction shock Z_+ , Z_- with $\tau_- < \tau_+$, $s < 0$, satisfying the entropy condition $s(E)$, the situation is as follows: $L(Z) < 0 < M(Z)$ for Z between Z_+ and Z_- on the Hugoniot curve with center Z_- . Hence $m(\tau) < \ell(\tau)$ for $\tau_- < \tau < \tau_+$. If $\ell'(\tau) < 0$, then the region

$$R = \{(\tau, \theta) \mid \tau_- < \tau < \tau_+ \text{ and } m(\tau) < \theta < \ell(\tau)\}$$

is negatively invariant and the existence and uniqueness of a trajectory connecting Z_- to Z_+ is established as in the compressive case above. However, it no longer follows from (2.7) that $\ell'(\tau) < 0$ if $\ell(\tau) > m(\tau)$. But we may calculate, using (2.2),

$$\begin{aligned}
 \frac{1}{-s} L_T(\tau, \theta) &= e_T(\tau, \theta) + p(\tau, \theta) + sM(\tau, \theta) \\
 &= e_T(\tau, \theta) + p(\tau, m(\tau))
 \end{aligned}$$

So the condition $e_T(\tau, \theta) > 0$ ensures that $\ell'(\tau) < 0$ for all τ , and

Theorem 1 follows.

We conclude this section by remarking that it is easy to use Proposition 2, (2.7) and (2.8) to show that a simple jump solution admits a purely viscous shock layer ($\lambda = 0$) if and only if the entropy condition $s(E)$ is satisfied. This recovers the result of Liu [4], under the assumptions (1.8)-(1.10), which are rather different than those Liu imposed (see [5]).

§3. A shock without a shock layer

In this section we construct explicitly an equation of state satisfying assumptions (1.8)-(1.10), and (1.14), (1.15), defined in a suitable region of the (τ, θ) plane, but violating the restriction $e_\tau(\tau, \theta) > 0$, such that a (large amplitude) rarefaction shock exists, satisfying the entropy condition $s(E)$, for which no shock layer exists if λ/μ is sufficiently large. The shock layer does exist if λ/μ is small, and we stress that entropy does increase across this shock, i.e., $S(Z_-) < S(Z_+)$ with $s < 0$.

Our object is to construct Fig. 3, representing the phase portrait for (1.6). The idea is that if λ/μ is large, the vector field $(\tau M/\mu, \tau L/\lambda)$ for (1.6) is nearly horizontal, and a trajectory leaving Z_0 must hit the "hump" in the curve $(M = 0)$. Then it is easy to show that no trajectory connects Z_- to Z_+ . But using (2.6) and (2.7), one may easily check that the entropy condition $s(E)$ is satisfied for the jump determined by Z_+ , Z_- , and $s < 0$.




Figure 3. A shock without a profile

Our procedure is to specify selected data (in particular the curve $\theta = m(\tau)$ on which $M = 0$), using thermodynamic identities to consistently specify the equation of state. Fix $Z_0 = (\tau_0, \theta_0)$ in the first quadrant. We shall specify the curve ($M = 0$) as $\theta = m(\tau)$, where $m(\tau)$ satisfies conditions (m1)-(m3) below (so its graph appears as in Fig. 3). Later, we will specify $p_0 = p(\tau_0, \theta_0)$ and s . Then we will know p along the curve ($M = 0$):

$$(3.1) \quad p_m(\tau) \equiv p(\tau, m(\tau)) = p_0 - s^2(\tau - \tau_0)$$

For all (τ, θ) , we specify that

$$(3.2) \quad \theta p_\theta(\tau, \theta) = e_\tau(\tau, \theta) + p(\tau, \theta) = \frac{1}{\theta}$$

so that $p_\theta = 1/\theta^2$, and (1.10) holds. The equation of state for p is now determined:

$$(3.3) \quad p(\tau, \theta) = p_m(\tau) + (1/m(\tau) - 1/\theta)$$

Our construction will be complete once $e_\theta(\tau, \theta)$ is determined. For then, since p_0 , s , and $m(\tau)$ determine p , $e_\tau(\tau, \theta)$ is known from (3.2). Then e_τ and e_θ determine $e(\tau, \theta)$, and $S(\tau, \theta)$ is determined from the Gibbs relation (1.2). ((3.2) ensures that the expression obtained for dS can be integrated.)

We must have $(e_\theta)_\tau = (e_\tau)_\theta = -2/\theta^2$, so e_θ has the form

$$e_\theta(\tau, \theta) = e_\theta(\tau_0, \theta) - 2(\tau - \tau_0)/\theta^2$$

We specify $e_\theta(\tau_0, \theta)$ so that (1.8) holds in the region of interest. It suffices to define

$$e_\theta(\tau_0, \theta) = 80\tau_0(\theta^{-2} + \theta_0^{-2})$$

so that $e_\theta(\tau, \theta) > 0$ for $\theta > 0$, $0 < \tau < 40\tau_0$. An explicit formula for $e(\tau, \theta)$ is

$$(3.4) \quad \begin{aligned} e(\tau, \theta) = & 80\tau_0(\theta_0^{-2} - \theta^{-2})\theta - \int_{\tau_0}^{\tau} 1/m(\tau) d\tau \\ & + (\tau - \tau_0)(2/\theta - p_0 + 1/2^2(\tau - \tau_0)) + e_0 \end{aligned}$$

We must now show that $m(\tau)$, p_0 and s may be specified so that Fig. 3 is

valid and (1.9) holds, along with (1.11), (1.14), (1.15). Observe that

$-sM(\tau, \theta)$ is independent of p_0 and s :

$$-sM(\tau, \theta) = p(\tau, \theta) - p_m(\tau) = 1/m(\tau) - 1/\theta$$

Using (2.7), we must have

$$\frac{1}{-s}L_\tau(\tau, \theta) = \theta p_\theta - (-s)M = 2/\theta - 1/m(\tau)$$

Requiring $L(\tau_0, \theta_0) = 0$, the quantity $(1/-s)L(\tau, \theta)$ is independent of p_0 and s

(recall $(1/-s)L_\theta = e_\theta$):

$$(3.5) \quad \frac{1}{-s}L(\tau, \theta) = 80\tau_0(\theta_0^{-2} - \theta^{-2})\theta + 2(\tau - \tau_0) - \int_{\tau_0}^{\tau} 1/m(\tau) d\tau$$

We will use this formula to manipulate the curve $(L = 0)$ by choice of $m(\tau)$. Set $\tau_k = (k+1)\tau_0$. We shall show that $m(\tau)$ can be chosen so that:

- (3.6) i) $L(\tau, m(\tau)) > 0$ for $\tau - \tau_0 > 0$ small
- ii) $L(\tau, \theta_0 + \varepsilon_1) < 0$ for $\tau_3 < \tau < \tau_5$ for some $\varepsilon_1 > 0$,
while $\theta_0 < m(\tau) < \theta_0 + \varepsilon_1$ for $\tau_4 < \tau < \tau_5$
- iii) $L(\tau, m(\tau)) > 0$ for τ large

We ask that $m(\tau)$ be defined and smooth for $\frac{1}{2}\tau_0 < \tau < 40\tau_0$, and satisfy conditions (m1)-(m3) below:

$$(m1) \quad m(\tau_0) = \theta_0 \text{ and } m'(\tau_0) = 0$$

Since $(1/-s)L_\tau(\tau_0, \theta_0) = 1/\theta_0 > 0$, (3.6i) follows.

$$(m2) \quad \begin{aligned} m'(\tau) &< 0 & \text{for } \tau_0 < \tau < \tau_1, \\ m(\tau) &= \theta_0/6 & \text{for } \tau_1 < \tau < \tau_3, \\ m'(\tau) &> 0 & \text{for } \tau_3 < \tau < \tau_4, \text{ and} \end{aligned}$$

$$\max_{\tau_4 < \tau < \tau_5} m(\tau) = \theta_0 + \varepsilon_1, \quad \varepsilon_1 > 0 \text{ small, with } m(\tau_4) = \theta_0 = m(\tau_5).$$

To verify (3.6ii), we estimate, for $\tau_3 < \tau < \tau_5$,

$$\begin{aligned} \frac{1}{-s}L(\tau, \theta_0 + \varepsilon_1) &< C\varepsilon_1 + 2(\tau_5 - \tau_0)/\theta_0 - (\tau_3 - \tau_1)6/\theta_0 \\ &= C\varepsilon_1 - 2\tau_0/\theta_0 < 0 \quad \text{if } \varepsilon_1 \text{ is small.} \end{aligned}$$

$$(m3) \quad m(\tau) = \theta_0 \quad \text{for } \tau > \tau_5$$

To verify (3.6iii), we estimate, for $\tau > \tau_6$,

$$\begin{aligned} \frac{1}{-s} L(\tau, \theta_0) &> 2(\tau - \tau_0)/\theta_0 - ((\tau_4 - \tau_0)6/\theta_0 + (\tau - \tau_4)/\theta_0) \\ &= ((\tau - \tau_4) - 4(\tau_4 - \tau_0))/\theta_0 > 0 \quad \text{if } \tau > \tau_{20}. \end{aligned}$$

Now: Since $(1/-s)L_0(\tau, \theta) > c > 0$ for $1/\tau_0 < \tau < \tau_{40}$, the set $(L = 0)$ is a curve $\theta = l(\tau)$ which is bounded above. We claim that $l(\tau)$ is defined for $\tau_0 < \tau < \tau_5$. Indeed, $L(\tau, \theta_0) < 0$ for $\tau_3 < \tau < \tau_5$, and since $L_\tau(\tau, \theta_0/6) > 0$ for $\tau < \tau_3$ (use (2.7) with $M < 0$), we have $L(\tau, \theta_0/6) < 0$ for $\tau < \tau_3$, so $l(\tau)$ is bounded below. In fact, we have shown that $l(\tau) > m(\tau)$ for $\tau_1 < \tau < \tau_5$.

We now set

$$\begin{aligned} \tau_+ &= \min\{\tau \mid \tau > \tau_5 \text{ and } l(\tau) = m(\tau)\} \\ \tau_- &= \max\{\tau \mid \tau < \tau_1 \text{ and } l(\tau) = m(\tau)\} \end{aligned}$$

Then $\tau_5 < \tau_+ < \tau_{20}$, and since $m'(\tau_0) = 0 > l'(\tau_0)$, $\tau_0 < \tau_- < \tau_1$. We set $\theta_+ = m(\tau_+)$, $\theta_- = m(\tau_-)$, fixing Z_+ and Z_- , and completing the construction of Fig. 3.

It remains to choose p_0 and s so that $p(\tau, \theta) > 0$ and (1.9), (1.14),

(1.15) hold in a suitable domain. But

$$(3.7) \quad p_\tau(\tau, S) = p_\tau(\tau, \theta) - \theta p_\theta(\tau, \theta)^2 / e_\theta(\tau, \theta)$$

(One calculates $p(\tau, S(\tau, \theta))_\tau$ and $\theta_\tau(\tau, S) = -S_\tau/S_\theta(\tau, \theta)$ and uses (2.1), (2.2).)

So (1.15) implies (1.9). But from (3.2),

$$(3.8) \quad p_\tau(\tau, \theta) = -s^2 + (1/m(\tau))'$$

So we can ensure that $p_\tau(\tau, \theta) < 0$ for $0 < \tau < \tau_{40}$ if we choose s^2 sufficiently large.

As we choose p_0 , we seek to fix the domain of states Ω_p in the (τ, p) plane so that Ω_p is convex, $p > 0$ in Ω_p , (1.14) holds, and, pulled back to the (τ, θ) plane, the domain Ω_θ contains (say) the half strip

$\{(\tau, \theta) \mid 0 < \tau < \tau_{40}, \theta > \theta_0/10\}$ which contains all our constructions. It suffices to choose Ω_p of the form

$$\Omega_p = \{(\tau, p) \mid 0 < \tau < \tau_{40} \text{ and } p - p_0 + s^2(\tau - \tau_0) > -10/\theta_0\}$$

with

$$(3.9) \quad p_0 = 40\tau_0 s^2 + 10/\theta_0$$

To check (1.14), compute

$$e_\tau(\tau, p) = e_\tau(\tau, \theta) + e_\theta(\tau, \theta)(-p_\tau(\tau, \theta)/p_\theta(\tau, \theta))$$

Using (3.2) and (3.8) one may verify that $e_\tau(\tau, p) > 0$ in Ω_p if s^2 is sufficiently large.

Now the equation of state is completely specified, and (1.8)-(1.15) hold except for (1.12). We have already noted that the entropy condition $s(E)$ holds, since $l(\tau) > m(\tau)$ for $\tau_- < \tau < \tau_+$.

The penultimate stage of our analysis is to show that for λ/μ sufficiently large, there is a trajectory of (1.6) leaving Z_0 and intersecting the curve $(M = 0)$ before the "hump". Since $m'(\tau_0) > l'(\tau_0)$, Z_0 is a saddle point for (1.6) (see (2.9)). If κ is the positive characteristic value for (1.6) at Z_0 , the the characteristic vector $(\kappa - \tau L_\theta/\lambda, \tau L_\tau/\lambda)$ may be shown to have positive components. Let $Z(\xi)$ be the solution leaving Z_0 along this vector. So long as $Z(\xi)$ remains above the curve $(M = 0)$, τ is increasing on $Z(\xi)$ and we may write $\theta = z(\tau)$ along $Z(\xi)$. Then

$$\frac{dz}{d\tau} = \frac{\mu L(\tau, z(\tau))}{\lambda M(\tau, z(\tau))}$$

It is clear that $z(\tau)$ is defined and increasing for $\tau_0 < \tau < \tau_*$, where

$$\tau_* = \min\{\tau \mid \tau > \tau_0 \text{ and } L(\tau, \theta_0) = 0\}$$

Note that $\tau_* < \tau_3$. We will "channel" the curve $z(\tau)$ under the curve $(L = 0)$ by using a thin box with a corner cut out: Define

$$R^* = (\tau_0, \tau_3) \times (\theta_0 - \varepsilon_2, \theta_0 + \varepsilon_1) \setminus (\tau_0, \tau_*) \times (\theta_0 - \varepsilon_2, \theta_0 + \varepsilon_1/2)$$

Here $\varepsilon_2 > 0$ is chosen so that \bar{R}^* is contained in the region $(M > 0)$. Now, the

trajectory $Z(\xi)$ must enter R^* on the cut-out part of the boundary with $\theta > \theta_0$, i.e., it must enter at a point $(\tau, \theta_0 + \frac{1}{2}\epsilon_1)$ for $\tau_0 < \tau < \tau_*$, or at a point (τ_*, θ) for $\theta_0 < \theta < \theta_0 + \frac{1}{2}\epsilon_1$.

Now $\sup_{R^*} |L/M|$ is finite. If λ/μ is sufficiently large in R^* , we may obtain

$$\left| \frac{dz}{d\tau} \right| < \min\{\frac{1}{2}\epsilon_1/(\tau_3 - \tau_0), \epsilon_2/(\tau_3 - \tau_*)\}$$

so long as $Z(\xi)$ is in R^* . This implies that the trajectory $Z(\xi)$ must leave R^* on its right boundary, $\{\tau_3\} \times (\theta_0 - \epsilon_2, \theta_0 + \epsilon_1)$, which is contained in the region $(L < 0 \text{ and } M > 0)$. Past this point, θ decreases and τ increases along $Z(\xi)$, so since $m(\tau)$ achieves the value $\theta_0 + \epsilon_1$, $Z(\xi)$ must intersect the curve $(M = 0)$ at a point $(\tilde{\tau}, \tilde{\theta})$, where $\tau_5 > \tilde{\tau} > \tau_3 > \tau_-$.

We may now easily construct a positively invariant region containing Z_- and disjoint from Z_+ , so that no trajectory can connect Z_- to Z_+ . The region is bounded on the right by the line $\tau = \tilde{\tau}$ for $\theta < \tilde{\theta}$ ($M < 0$ here), from above by the curve $\theta = z(\tau)$ for $\tau_0 < \tau < \tilde{\tau}$ (trajectories cannot cross), and by the curve $\theta = \min\{l(\tau), m(\tau)\}$ for $\tau \leq \tau_0$ (the vector field either points down, or to the left at a point where $l'(\tau) < 0$, as in §2). Our example is complete.

ACKNOWLEDGEMENT

This work comprises one chapter of this author's Ph.D. dissertation, written with the guidance and encouragement of Professor Andrew Majda at the University of California, Berkeley.

REFERENCES

1. C. Dafermos, Global smooth solutions for the initial boundary value problem for the equations of one dimensional nonlinear thermoviscoelasticity, SIAM J. Math. Anal. 13(1982)397-408.
2. D. Gilbarg, The existence and limit behavior of the one dimensional shock layer, Amer. J. Math. 7(1951)256-274.
3. A. Kelley, Stability of the center-stable manifold, J. Math. Anal. Appl. 18(1967)336-344.
4. T.-P. Liu, The entropy condition and the admissibility of shocks, J. Math. Anal. Appl. 53(1976)78-88.
5. T.-P. Liu, Shock waves in the nonisentropic gas flow, J. Diff. Eqns. 22(1976)442-452.
6. A. Matsumura and T. Nishida, The initial value problem for the equations of motion of compressible viscous and heat-conductive fluids, Proc. Japan Acad 55, Ser. A(1979)337-342.
7. R. Pego, Stable viscosities and shock profiles for systems of conservation laws, (to appear).
8. J. Serrin, Phase transitions and interfacial layers for van der Waals fluids, in Recent Methods in Nonlinear Analysis and Applications, ed. A. Confora, et al., Instituto Matematico, Universita di Napoli, Naples(1980)169.
9. M. Slemrod, Admissibility criteria for propagating phase boundaries in a van der Waals fluid, to appear in Arch. Rat. Mech. Anal.
10. R. G. Smith, The Riemann problem in gas dynamics, Transactions of the AMS 249(1979)1-50.
11. H. Weyl, Shock waves in arbitrary fluids, Comm. Pure Appl. Math. 2(1949)103-122.

RLP/jvs

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER #2505	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Nonexistence of a Shock Layer in Gas Dynamics With a Nonconvex Equation of State		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Robert L. Pego		8. CONTRACT OR GRANT NUMBER(s) MCS-7927062, Mod. 2. DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 2 - Physical Mathematics
11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18 below		12. REPORT DATE April 1983
		13. NUMBER OF PAGES 20
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES U. S. Army Research Office P. O. Box 12211 Research Triangle Park North Carolina 27709 National Science Foundation Washington, DC 20550		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) shocks, shock layer, traveling waves, conservation laws, admissibility, entropy condition, viscosity, heat conduction.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A classical result of Gilbarg states that a simple shock wave solution of Euler's equations is compressive if and only if a corresponding shock layer solution of the Navier-Stokes equations exists, assuming, among other things, that the equation of state is convex. An "entropy condition" appropriate for weeding out "unphysical" shocks in the nonconvex case has been introduced by T.-P. Liu. For shocks satisfying his entropy condition. Liu showed that purely viscous shock layers exist (with zero heat conduction). Dropping the convexity assumption, but retaining many other reasonable restrictions on the equation		

ABSTRACT (continued)

of state, we construct an example of a (large amplitude) shock which satisfies Liu's entropy condition but for which a shock layer does not exist if heat conduction dominates viscosity. We also give a simple restriction, weaker than convexity, which does guarantee that shocks which satisfy Liu's entropy condition always admit shock layers.

END

DATE
FILMED

6-83

DTIC