


AD-A129 185

A VERY SMALL REMARK ON SMOOTH DENJOY MAPS OF THE CIRCLE 1/1
(U) WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER
G R HALL APR 83 MRC-TSR-2512 DAAG29-80-C-0041

UNCLASSIFIED

F/G 12/1 NL

END
DATE
FILMED
7-83
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

MRC Technical Summary Report #2512

A VERY SMALL REMARK ON
SMOOTH DENJOY MAPS OF THE CIRCLE

AD A129185

Glen Richard Hall

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

April 1983

(Received March 30, 1983)

DTIC
ELECTE
S JUN 9 1983
A

DTIC FILE COPY

Approved for public release
Distribution unlimited

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

and

National Science Foundation
Washington, DC 20550

83 06 07 06 9

UNIVERSITY OF WISCONSIN - MADISON
MATHEMATICS RESEARCH CENTER

A VERY SMALL REMARK ON
SMOOTH DENJOY MAPS OF THE CIRCLE

Glen Richard Hall

Technical Summary Report #2512

April 1983

ABSTRACT

In this note we show that if f is a smooth homeomorphism of the circle with one 'non degenerate' critical point such that f has an irrational rotation number which is poorly approximable by rationals (e.g., a quadratic irrational) and f is a Denjoy map, then the length of the associated wandering interval cannot tend to zero too rapidly.

AMS (MOS) Subject Classifications: 58F99, 54H20

Key Words: Denjoy Maps, wandering intervals

Work Unit Number 1 - Applied Analysis

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041 and National Science Foundation Postdoctoral Fellowship.

SIGNIFICANCE AND EXPLANATION

Maps of a circle into itself, when considered as discrete dynamical systems, have been observed to contain some of the qualitative behavior of complicated experimental situations. Much of this structure can be understood for homeomorphisms of the circle by considering the invariant called the rotation number. In fact, the homeomorphisms of the circle with irrational rotation number can be classified up to topological conjugacy, provided the map and its inverse are sufficiently smooth. Counter examples to this classification scheme exist which are C^1 diffeomorphisms or C^∞ homeomorphisms with non differentiable inverse. Whether C^∞ can be replaced with analytic in the second examples is an open question.

In this report ^{the author} shows that if certain assumptions are made on the rotation number and on the derivative of a homeomorphism from the circle onto itself, then even if the map does not fit into the classification scheme we can still obtain estimates on the behavior of its orbits under iteration.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<input type="checkbox"/>
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	



The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

A VERY SMALL REMARK ON
SMOOTH DENJOY MAPS OF THE CIRCLE

Glen Richard Hall

1. Introduction: Let $T = \mathbb{R}/\mathbb{Z}$ be the circle with unit circumference. We say a homeomorphism $f : T \rightarrow T$ is a Denjoy map if f has no periodic and no dense orbits. A theorem of Denjoy [2] states that if f is a C^2 diffeomorphism then f is not a Denjoy map. He also constructs C^1 diffeomorphisms which are Denjoy maps (see Herman [4]). Recently, examples of homeomorphisms $f : T \rightarrow T$ which are C^∞ Denjoy maps have been constructed ([3]). These maps fail to be diffeomorphisms because they have points where the derivative of f is zero. However, the question of the existence of analytic homeomorphisms which are Denjoy maps remains open.

It turns out that a homeomorphism $f : T \rightarrow T$ is a Denjoy map if and only if it has no periodic orbits and there exists a "wandering" interval $I \subseteq T$ such that $\{f^n(I)\}_{n=-\infty}^{\infty}$ forms a disjoint family of intervals. An interesting property of the examples of [3] is that they can be constructed so that $\text{length } f^n(I) \rightarrow 0$ arbitrarily rapidly. (This requires an addition to [3] which states that "case B" will not occur. This was shown to the author by J. C. Yoccoz (personal communication).)

In this note we consider homeomorphisms $f : T \rightarrow T$ with only a "non-degenerate" critical point (e.g., analytic homeomorphisms) and rotation number which is very poorly approximable by rationals (e.g., quadratic irrationals). If f is a Denjoy map with $I \subseteq T$ a wandering interval as described above, then we show $\text{length } f^n(I)$ cannot tend to zero arbitrarily rapidly.

As we will note in section 3, recent computer studies (Shenker, [5]) indicate that the estimates of this note might be significantly improved, yielding a corresponding improvement of the theorem.

2. Definitions and Notations: Fix a map $f_0 : \mathbb{R} \rightarrow \mathbb{R}$ satisfying the following conditions

- 1) f_0 is increasing, C^3 and $f_0(0) \in (0,1)$,
- 2) $\forall x \in \mathbb{R}, f_0(x+1) = f_0(x) + 1$, (i.e. f_0 is the lift of a homeomorphism of the circle),
- 3) there exists precisely one point $x_0 \in [0,1)$ such that $f_0'(x_0) = 0$ (hence $f_0''(x_0) = 0$),
- 4) $f_0'''(x_0) \neq 0$,
- 5) $f_0''(x) = 0$ for only finitely many $x \in [0,1)$.

We let $f(\cdot, \phi) : \mathbb{R} \rightarrow \mathbb{R}$ denote the one parameter family of maps given by

$$f(x, \phi) = f_0(x) + \phi.$$

Notation: Fix an interval $J \subseteq [0,1)$, $x_0 \in \text{interior } J$. Then let

$$M_1 = \sup_{x \in [0,1)} \frac{\partial f}{\partial x}(x, \phi),$$

$$M_2 = \sup_{x \in [0,1)} \left| \frac{\partial^2 f}{\partial x^2}(x, \phi) \right|,$$

$$M_3 = \inf_{x \in J} \left| \frac{\partial^3 f}{\partial x^3}(x, \phi) \right|,$$

$$M_4 = \inf_{x \in [0,1) - J} \frac{\partial f}{\partial x}(x, \phi).$$

We may choose J small so that $M_3 > 0$. (Clearly these constants are independent of ϕ .) We define $f^n(x, \phi) = f(f^{n-1}(x, \phi), \phi)$ for $n \in \mathbb{Z}$, i.e., $f^n(x, \phi)$ is the n^{th} iterate of $f(\cdot, \phi)$ with ϕ fixed.

Definition: The rotation number of $f(\cdot, \phi)$ denoted $\rho(\phi)$ is given by

$$\rho(\phi) = \lim_{n \rightarrow \infty} \frac{f^n(x, \phi) - x}{n}.$$

Remarks: 1) The rotation number $\rho(\phi)$ exists and is independent of the x in the definition. Moreover, $\rho(\phi)$ is a continuous increasing function of ϕ .

2) The function $\rho(\phi)$ is strictly increasing when its value is irrational, i.e., if $\rho(\bar{\phi}) \notin \mathbb{Q}$ and $\phi \neq \bar{\phi}$ then $\rho(\phi) \neq \rho(\bar{\phi})$ (see Herman [4] for proofs of these remarks).

Notation: For each $\beta \in \mathbb{R}$, let $A_\beta = \{\phi : \rho(\phi) = \beta\}$. Then for every β , A_β is a closed interval, which is a single point when β is irrational.

Finally, we fix $\alpha_0 \in [0, 1]$, α_0 an irrational such that there exists a constant $M_5 > 0$ so that if

$$\alpha_0 = \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{\ddots}}}, \quad a_i \in \mathbb{Z}.$$

Then $|a_i| < M_5$ for all i , (e.g., any quadratic irrational will satisfy this condition). We let $\frac{p_n}{q_n}$ denote the n^{th} convergent of α_0 , i.e.

$$\frac{p_n}{q_n} = \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{\ddots + \frac{1}{a_n}}}}.$$

Let $\phi_0 \in \mathbb{R}$ be the unique number such that

$$\rho(\phi_0) = \alpha_0.$$

3. Statement of the theorem: Suppose $f(\cdot, \phi_0)$ is a Denjoy map, i.e. $f(\cdot, \phi_0)$ is not continuously conjugate to rigid rotation by its rotation number α_0 . Then there exists a nontrivial (i.e., non empty, non singleton)

interval $I \subseteq [0,1)$ such that $\{f^n(x) + m : x \in I\}_{n,m \in \mathbb{Z}}$ form a family of disjoint intervals. Let $\mu_n = \text{length}(f^n(I))$, then $\mu_n \rightarrow 0$ as $n \rightarrow \infty$ (recall f is the lift of a circle map).

Theorem: There exists a constant $M_0 > 1$ such that

$$(3.1) \quad \mu_{\pm q_n} > 1/M_0^{q_n}.$$

infinitely often.

That is, the length of the wandering interval cannot decrease too fast.

The outline of the proof is as follows: First we derive an estimate on the length of the intervals A_{p_n/q_n} . Next we show that the orbit of the critical point must return to a neighborhood of the critical point in each set of q_n iterates, moreover, the size of this neighborhood is controlled by the size of μ_{q_n} . Hence, if μ_{q_n} is very small, a small change in ϕ will make a relatively large change in the rotation number, not leaving "enough room" for the A_{p_n/q_n} intervals.

Remark: It is important to note that the estimates we use on the derivatives of high iterates of $f(\cdot, \phi)$ are very crude (see lemma 1 below). Recent numerical studies (Shenker, [5]) indicate much better estimates may hold, and if so, much better bounds on μ_{q_n} would result.

4. The lemmas: In this section we give the necessary lemmas.

Lemma 1: For all $x \in \mathbb{R}$, $\phi \in \mathbb{R}$ and $n \in \mathbb{Z}^+$

$$(4.1) \quad \frac{\partial(f^n)}{\partial x}(x, \phi) < M_1^n,$$

$$(4.2) \quad \left| \frac{\partial^2(f^n)}{\partial x^2}(x, \phi) \right| < M_2 \cdot M_1^{2n-2}/(M_1-1),$$

$$(4.3) \quad \frac{\partial(f^n)}{\partial \phi}(x, \phi) < M_1^{n+1}/(M_1-1).$$

Proof: These estimates follow immediately from the chain rule and (2.1),

(2.2) (see remark, end of section 3). //

Lemma 2: For any $n > 0$,

$$\text{length}(A_{p_n/q_n}) > 1 / \left(\frac{4M_2 \cdot M_1^{3q_n}}{(M_1 - 1)} \right) .$$

Proof: Fix $n > 0$ and let $[\phi_1, \phi_2] = A_{p_n/q_n}$. Then

$$\forall x \in \mathbb{R}, f^{q_n}(x, \phi_1) < x + p_n ,$$

and there exists $x_1 \in (0, 1)$ with

$$f^{q_n}(x, \phi_1) = x_1 + p_n .$$

Now, $\frac{\partial f^{q_n}}{\partial x}(x_0, \phi) = 0$, so by (4.2) for $\delta = 1 / \left(\frac{2M_2 \cdot M_1^{2q_n - 1}}{(M_1 - 1)} \right)$ we have for any ϕ ,

$$\frac{\partial f^{q_n}}{\partial x}(x, \phi) < \frac{1}{2} \text{ whenever } |x_0 - x| < \delta .$$

But then it follows that for any ϕ ,

$$\begin{aligned} \max_{x \in [0, 1]} (x - f^{q_n}(x, \phi)) - \min_{x \in [0, 1]} (x - f^{q_n}(x, \phi)) \\ > 1 / \left(\frac{4M_2 \cdot M_1^{2q_n - 1}}{M_1 - 1} \right) . \end{aligned}$$

By (4.3) we then have that there exist $x_2, x_3 \in \mathbb{R}$ such that

$$f^{q_n}(x_2, \phi) < x_2 + p_n \text{ and } f^{q_n}(x_3, \phi) > x_3 + p_n$$

whenever $0 < \phi - \phi_1 < \frac{1}{\left(\frac{4M_1^{q_n+1} \cdot M_1^{2q_n-1}}{(M_1-1)^2} \right)}$ and the proof is complete. //

The following lemmas are standard results on the orbit structure for maps with irrational rotation number.

Lemma 3: The sets $\{f^i(x, \phi_0) + m : i, m \in \mathbb{Z}, 0 < i < q_n\}$ and $\{f^{-i}(x, \phi_0) + m : i, m \in \mathbb{Z}, 1 < i < q_n\}$ alternate on \mathbb{R} (i.e. between any two points of the first set there is a point of the second set and vice versa) for any fixed $x \in \mathbb{R}$ and q_n .

Lemma 4: For each $n > 0$, $x_1, x_2 \in \mathbb{R}$, if for some $q_n > i, j > 0, m_1, m_2 \in \mathbb{Z}$, $f^i(x_1, \phi_0) + m_1 < f^j(x_1, \phi_0) + m_2$ then there exists k and m_3 with $0 < k < q_{n+1}$ with

$$f^i(x_1, \phi_0) + m_1 < f^k(x_2, \phi_0) + m_3 < f^j(x_1, \phi_0) + m_2 .$$

The proofs of these lemmas follow from, for example, Theorem 4.1, page 409 of Coddington, Levinson [1] and II. 7.1 of Herman [4], respectively.

Lemma 5: For n sufficiently large, the set $\{f^i(y, \phi_0) - [f^i(y, \phi_0)] : i = 0, \dots, n\}$ (where $[\cdot]$ denotes greatest integer) contains more than two points in J for any $y \in [0, 1)$.

Proof: It suffices to show that the orbit of any point, say x_0 , contains the critical point in its closure. But if $\overline{\{f^i(x_0, \phi_0) - [f^i(x_0, \phi_0)] : i > 0\}}$ does not contain x_0 then by altering f in a neighborhood of x_0 we can make a C^3 diffeomorphism without altering the orbit of x_0 , (see Figure 1) which contradicts Denjoy's theorem and the proof is complete. //




Figure 1

Remark: In fact the orbit of x_0 must approach x_0 from "both sides" as was shown by J. C. Yoccoz (personal communication).

5. Proof of the theorem. Without loss of generality we may assume $\mu_0 = \max_{i \in \mathbb{Z}} \mu_i$. Now fix $n > 0$ and for $y, \hat{y} \in I$ let $y_i = f^i(y, \phi_0) - [f^i(y, \phi_0)]$, $\hat{y}_i = f^i(\hat{y}, \phi_0) - [f^i(\hat{y}, \phi_0)]$ where $[\cdot]$ denotes the greatest integer. Fix i_0 with $0 < i_0 < q_n$ and $|y_{i_0} - x_0| = \min\{|y_i - x_0| : i = 0, \dots, q_n - 1\}$. By condition (5) in the definition of f_0 there exist, say r , points in $(0, 1)$ such that $\frac{\partial^2 f}{\partial x^2}(x, \phi_0) = 0$. Hence by lemma 3 we see that taking n sufficiently large, so that the conclusion of lemma 5 holds,

$$\begin{aligned} \frac{\partial(f^n)}{\partial x}(y, \phi_0) &= \prod_{j=0}^{q_n-1} \frac{\partial f}{\partial x}(y_j, \phi_0) \\ &> \prod_{j=1}^{q_n} \frac{\partial f}{\partial x}(\hat{y}_{-j}, \phi_0) \cdot M_4^{r-1} \cdot \frac{\partial f}{\partial x}(y_{i_0}, \phi_0) \end{aligned}$$

This follows from lemma (3) since for each i , $0 < i < q_n$ there corresponds j , $0 > j > -q_n$ such that $\frac{\partial f}{\partial x}(y_j, \phi_0) < \frac{\partial f}{\partial x}(y_i, \phi_0)$ except for $i = i_0$ and the $r-1$ choices of $i \neq i_0$ where y_i is adjacent to a point with $\frac{\partial^2 f}{\partial x^2} = 0$. Hence, by an appropriate choice of $\hat{y} \in I$ we obtain

$$\begin{aligned} \frac{\partial f}{\partial x}^{q_n}(y, \phi_0) &> \frac{\mu_0}{\mu^{-q_n}} M_4^{r-1} \frac{\partial f}{\partial x}(y_{i_0}, \phi_1) \\ &> M_4^{r-1} \frac{\partial f}{\partial x}(y_{i_0}, \phi_1) . \end{aligned}$$

Noting that

$$\frac{\partial f}{\partial x}(y_{i_0}, \phi_1) > 6M_3 \cdot (y_{i_0} - x_0)^2$$

we see that

$$\frac{\partial f}{\partial x}^{q_n}(y, \phi_0) > 6M_4^{r-1} \cdot M_3 (y_{i_0} - x_0)^2$$

for all $y \in I$. Hence, for some $y \in I$

$$\mu_{q_n} > 6\mu_0 M_4^{r-1} \cdot M_3 (y_{i_0} - x_0)^2, \text{ i.e.,}$$

$$(5.1) \quad (\mu_{q_n} / (\mu_0 \cdot 6M_4^{r-1} \cdot M_3))^{1/2} > |y_{i_0} - x_0| .$$

Suppose $\mu_{q_n} < 1/c^{q_n}$ for all n , for some $c > 1$. Then, by (5.1), for each n there exists $y_n \in I$ such that if $\delta_n = \min_{i=0, \dots, q_n-1} |f^i(y_n, \phi_0) - [f^i(y_n, \phi_0)] - x_0|$ then

$$\delta_n < c_2 / c_1^{q_n}$$

where $c_1 = c^{1/2}$ and $c_2 = (6\mu_0 M_4^{r-1} M_3)^{-1/2}$. Fix m such that $\delta_{m+1} < \delta_m$.

Then by lemma 4 there exists t , $0 < t < q_{m+2}$ with

$$|f^t(x_0, \phi_0) - [f^t(x_0, \phi_0)] - x_0| < \delta_m .$$

But since $1 < \frac{\partial f^n}{\partial \phi}(x, \phi)$ for all x, ϕ, n we see that there exists ϵ such that $f^t(x_0, \phi_0 + \epsilon) - [f^t(x_0, \phi_0 + \epsilon)] = x_0$ and $|\epsilon| < \delta_m$. But then either $A_{P_{m+3}/q_{m+3}} \subseteq [\phi_0 - |\epsilon|, \phi_0 + |\epsilon|]$ or $A_{P_{m+4}/q_{m+4}} \subseteq [\phi_0 - |\epsilon|, \phi_0 + |\epsilon|]$ and hence by lemma 2

$$\delta_m > |\epsilon| > (M_1 - 1)^2 / (4M_2 \cdot M_1^{3q_{m+4}}) > (M_1 - 1)^2 / (4M_2 \cdot M_1^{3(M_4 + 1)^4 q_m})$$

i.e. $c_2/c_1^{q_m} > (M_1 - 1)^2 / (4M_2 \cdot M_1^{3(M_4 + 1)^4 q_m})$. Hence, we must have

$$c_1 < M_1^{3(M_4 + 1)^4}$$

which completes the proof of (3.1) for μ_{α_n} . To show $\mu_{-\alpha_n}$ also satisfies

the theorem for $M_0 = M_1^{9(M_4 + 1)^8} + 1$ is similar. //

6. Example: Let $f_0(x) = x + \frac{1}{2\pi} \sin(2\pi x)$. Then the family $f(x, \phi) =$

$x + \phi + \frac{1}{2\pi} \sin(2\pi x)$ satisfies the hypotheses, and if we let $\alpha_0 = \frac{\sqrt{5} - 1}{2}$

then we obtain $M_0 = 2^{2304} + 1$.

REFERENCES

- 1) E. A. Coddington, N. Levinson, Theory of Ordinary Differential Equations, McGraw-Hill Book Company, New York, 1955.
- 2) A. Denjoy, Sur les courbes definiées par les equation differentiables a la surface du tore. *J. de Math. Pure et Appliquees*, 11 (1932), 33-375.
- 3) G. R. Hall, A C^∞ Denjoy counter example, *Ergodic Theory and Dynamical Systems*, 1 (1981), 261-272.
- 4) M. R. Herman, Sur la conjugaison differentiable des diffeomorphismes du cercle a des rotation, *Publ. I.H.E.S.*, 49 (1979), 5-234.
- 5) S. J. Shenker, Scaling Behavior in a map of a circle onto itself: Empirical results, (preprint).

GRH/jvs

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER #2512	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A Very Small Remark on Smooth Denjoy Maps of the Circle		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Glen Richard Hall		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE April 1983
		13. NUMBER OF PAGES 10
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Denjoy Maps, wandering intervals		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this note we show that if f is a smooth homeomorphism of the circle with one 'non degenerate' critical point such that f has an irrational rotation number which is poorly approximable by rationals (e.g., a quadratic irrational) and f is a Denjoy map, then the length of the associated wandering interval cannot tend to zero too rapidly.		

DEFI