

AD-A129 553

IFR (INCREASING FAILURE RATE) FOR REPAIRABLE SYSTEMS

1/1

(U) CITY COLL NEW YORK DEPT OF MATHEMATICS

N R CHAGANTY APR 83 CUNY-MB4 AFOSR-TR-83-0511

UNCLASSIFIED


AFOSR-82-0024

F/G 12/1

NL



END  
DATE  
FILMED  
7 83  
DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

12

ADA 1 29553

IFR Results for Repairable Systems

by N.R. Chaganty

April 1983

City College, CUNY Report No. MB4  
AFOSR Technical Report No. 82-04  
AFOSR Grant No. 82-0024

DTIC  
ELECT  
S JUN 21 1983 D  
A

Approved for public release;  
distribution unlimited.

DTIC FILE COPY

83 06 20 118

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>AFOSR-TR- 83-0511</b>	2. GOVT ACCESSION NO. <b>AD-A129553</b>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) <b>IFR RESULTS FOR REPAIRABLE SYSTEMS</b>		5. TYPE OF REPORT & PERIOD COVERED <b>TECHNICAL</b>
7. AUTHOR(s) <b>N.R. Chaganty</b>		6. PERFORMING ORG. REPORT NUMBER <b>CUNY Report No. MB4</b>
9. PERFORMING ORGANIZATION NAME AND ADDRESS <b>Department of Mathematics The City College - CUNY New York NY 10031</b>		8. CONTRACT OR GRANT NUMBER(s) <b>AFOSR-82-0024</b>
11. CONTROLLING OFFICE NAME AND ADDRESS <b>Mathematical &amp; Information Sciences Directorate Air Force Office of Scientific Research Bolling AFB DC 20332</b>		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS <b>PE61102F; 2304/A5</b>
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE <b>APR 83</b>
		13. NUMBER OF PAGES <b>8</b>
		15. SECURITY CLASS. (of this report) <b>UNCLASSIFIED</b>
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) <b>Approved for public release; distribution unlimited.</b>		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) <b>IFR distributions; IFRA distributions; NBU distributions; repairable systems; k out of n systems; time to first failure; Markov processes.</b>		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  <b>Consider a k-out-of-n system with independent repairable components. Assume that the repair and failure distributions are exponential with parameters <math>\mu_1, \dots, \mu_n</math> and <math>\lambda_1, \dots, \lambda_n</math>. It is shown that if <math>\lambda_i - \mu_i = \Delta</math> for all i, then the distribution of system life is IFR.</b>		

DD FORM 1473 1 JAN 73

EDITION OF 1 NOV 68 IS OBSOLETE


83 06 20 118

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ABSTRACT

Consider a k-out-of-n system with independent repairable components. Assume that the repair and failure distributions are exponential with parameters  $\{\mu_1, \dots, \mu_n\}$  and  $\{\lambda_1, \dots, \lambda_n\}$  respectively. In this paper we show that if  $\lambda_i - \mu_i = \Delta$  for all i then the life distribution of the system is Increasing Failure Rate.



AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)  
NOTICE OF TRANSMITTAL TO DTIC  
This technical report has been reviewed and is approved for public release (AN APR 190-12).  
Distribution is unlimited.  
MATTHEW J. KENTER  
Chief, Technical Information Division

1. Introduction. A k-out-of-n system is one which functions if and only if at least k of the n components function. An n-out-of-n system is known as a series system and an l-out-of-n system is known as a parallel system. Consider a k-out-of-n system with independent repairable components, each of which has exponentially distributed failure times and repair times. Assume that the exponential failure time distributions have parameters  $\lambda_1, \dots, \lambda_n$  and the exponential repair times have parameters  $\mu_1, \dots, \mu_n$ . The study of the life distribution of these systems has received the attention of many authors. Birnbaum-Esary-Marshall [2] have shown that if the components are irreparable ( $\mu_i \equiv 0, 1 \leq i \leq n$ ), then the life distribution of the k-out-of-n system is Increasing Failure Rate on the Average (IFRA). This is true even for general coherent systems. In the special case where  $\lambda_i$ 's are all equal to  $\lambda$  and  $\mu_i$ 's are equal to  $\mu$ , the number of components not working at any time point forms a birth and death process. The results on birth and death processes of Keilson [7], Derman-Ross-Schechner [6], Brown and Chaganty [4] all imply that the life distribution of the k-out-of-n system is Increasing Failure Rate (IFR). More generally without any restrictions on the  $\lambda_i$ 's and  $\mu_i$ 's, it was shown by Ross [9], the life distribution of a coherent system is New Better than Used (NBU), from which it follows that the life distribution of the k-out-of-n system is NBU. The theorem of Ross [9] is actually a special case of the result in section 6.6 of Brown and Chaganty [4]. The question whether the

conclusion of Ross [9] can be strengthened to IFRA for a k-out-of-n system remains open. The case  $k=n=2$  failed to yield any counter example. In this paper we show that for a k-out-of-n system if  $\lambda_i - \mu_i$  is constant over  $i$ , the life distribution of the system is IFR. The motivation and importance for the study of the life distribution of the k-out-of-n system comes from the results of Esary and Proschan [5]. They have shown that the distribution function of the parallel and series systems provide lower and upper bounds respectively for the distribution function of an arbitrary coherent system. The Laplace transform and moments of the parallel system, without any restrictions on  $\lambda_i$ 's and  $\mu_i$ 's, and related results were derived in Brown [3].

2. Definitions and Preliminaries. A nonnegative random variable  $T$  with distribution  $F$  is said to be IFR if  $\bar{F}(x+t)/\bar{F}(t)$  is nonincreasing in  $-\infty < t < \infty$ , for each  $x \geq 0$ , where  $\bar{F} = 1-F$ . If the density  $f$  exists, this is equivalent to saying that  $r(x) = f(x)/\bar{F}(x)$  is nondecreasing in  $x \geq 0$ . The function  $r(x)$  is known as the failure rate. The random variable  $T$  is said to be IFRA if  $[\bar{F}(x)]^{1/x}$  is nonincreasing in  $x \geq 0$  and  $T$  is NBU if  $\bar{F}(x+y) \leq \bar{F}(x)\bar{F}(y)$  for all  $x, y \geq 0$ . It is well known that

$$\text{IFR} \rightarrow \text{IFRA} \rightarrow \text{NBU} ,$$

and none of the reverse implications are true. The properties and usefulness of these three classes of life distributions were well explained in Barlow and Proschan [1].

5

Consider a coherent system with  $n$  components. Let  $X_i(t) = 0$  if component  $i$  is functioning at time  $t$  and  $1$  if it is under repair,  $i=1, \dots, n$ . The vector  $X(t) = (X_1(t), \dots, X_n(t))$  determines the state of the system at time  $t$ . To start with we assume that all the components are working, that is,  $X(0) = (0, 0, \dots, 0)$  with probability  $1$ . Under the assumptions of exponential failure and repair distributions with parameters  $0 < \lambda_i < \infty$ ,  $0 < \mu_i < \infty$ ,  $i=1, \dots, n$ , the stochastic process  $X = \{X(t), t \geq 0\}$  forms a Markov process with state space  $S$ , consisting of  $2^n$  vectors of  $0$ 's and  $1$ 's. Let  $\{\tau_i, i \geq 0\}$  be the successive times of transition for the process  $X$  and  $Y = \{Y_i, i \geq 0\}$  be the successive states visited by  $X$ . If  $Y_i = \underline{x}$ ,  $\underline{x} = (x_1, \dots, x_n) \in S$ , then the sojourn interval in  $\underline{x}$ ,  $[\tau_i, \tau_{i+1})$ , is exponentially distributed with parameter  $\Sigma(x_i \mu_i + (1-x_i) \lambda_i)$  and further  $Y$  forms a Markov chain with state space  $S$  and transition matrix  $Q$ , of dimension  $2^n \times 2^n$  given by

$$\begin{aligned}
 Q(\underline{x}, \underline{y}) &= \frac{\lambda_i}{\Sigma(x_i \mu_i + (1-x_i) \lambda_i)} && \text{if } x_i=0, y_i=1, x_j=y_j \text{ for } j \neq i \\
 &= \frac{\mu_i}{\Sigma(x_i \mu_i + (1-x_i) \lambda_i)} && \text{if } x_i=1, y_i=0, x_j=y_j \text{ for } j \neq i \\
 &= 0 && \text{otherwise.}
 \end{aligned}$$

Let the function  $f: S \rightarrow N = \{0, 1, 2, \dots, n\}$  be defined by  $f(\underline{x}) = \Sigma x_i$ . The function  $f$  partitions the state space  $S$  into  $(n+1)$  parts  $A_0, \dots, A_n$ , where  $A_\ell = \{\underline{x} \in S: \Sigma x_i = \ell\}$ ,  $\ell \geq 0$ . Let

$$\delta(0) = \sum_{i=1}^n \lambda_i.$$



Under the assumption  $\lambda_i - \mu_i$  is constant over  $i$ , the quantity  $\delta(\ell) = \sum_{i=1}^{\ell} \mu_i + \sum_{i=\ell+1}^n \lambda_i$  is equal to  $\sum_{i=1}^n (x_i \mu_i + (1-x_i) \lambda_i)$  for all  $\underline{x} \in A_{\ell}$ . This means that the probability of transition from a state  $\underline{x} \in A_{\ell}$  to  $A_j$ ,  $j \neq \ell$ ,  $\ell \geq 0$ , is the same for all  $\underline{x} \in A_{\ell}$  in the embedded discrete time markov chain  $Y$ . Thus by Theorem 6.3.2, page 124, of Kemeny and Snell [8] the lumped process  $f(Y) = \{f(Y_i): i \geq 0\}$  is again a markov chain. Hence the process  $f(X) = \{f(X(t)): t \geq 0\}$  is a markov process with state space  $N$  and successive states visited given by  $f(Y)$  at times  $\{\tau_i, i \geq 0\}$ . Also  $[\tau_i, \tau_{i+1})$  is exponentially distributed with parameter  $\delta(\ell)$  if  $f(Y_i) = \ell$ . To compute the transition matrix of  $f(Y)$  we proceed as follows. Let  $M(\ell)$  be the cardinality of  $A_{\ell}$ . We can partition the matrix  $Q$  as

$$Q = \begin{bmatrix} Q_{00} & Q_{01} & \dots & Q_{0n} \\ Q_{10} & Q_{11} & \dots & Q_{1n} \\ \dots & \dots & \dots & \dots \\ Q_{n0} & Q_{n1} & \dots & Q_{nn} \end{bmatrix}, \quad (2.2)$$

where  $Q_{ij} = (Q(\underline{x}, \underline{y}))$ ,  $\underline{x} \in A_i$ ,  $\underline{y} \in A_j$ , is of order  $M(i) \times M(j)$ ,  $i, j \in N$ . Let  $i_{\ell} \in N$ ,  $1 \leq \ell \leq m$ ,  $m \geq 1$ . Then the joint probabilities of the lumped process  $f(Y)$  are given by

$$P[f(Y_{\ell}) = i_{\ell}, \ell = 1, \dots, m] = Q_{0i_1} Q_{i_1, i_2} \dots Q_{i_{m-1}, i_m} e_{i_m}, \quad (2.3)$$

where  $e_{i_m}$  is a column vector of  $M(i_m)$  one's. Note that  $Q_{ij} = 0$

if  $|i-j| \geq 2$ . Thus relation (2.3) shows that  $f(Y)$  is a birth and death process. The following lemma will be useful to compute the transition probabilities.

Lemma 2.1. Let  $Q_{ij}$  be defined as in (2.2). Then

$$Q_{01}Q_{12}\dots Q_{m-1,m}e_m = \frac{m! \sum_{1 \leq i_1 < i_2 < \dots < i_m \leq n} \lambda_{i_1} \lambda_{i_2} \dots \lambda_{i_m}}{\delta(0)\delta(1)\dots\delta(m-1)}, \quad (2.4)$$

for  $m \geq 1$ .

Proof: From (2.3) it follows that the L-H-S of (2.4) is the probability that  $m$  components fail in succession. These  $m$  components can be chosen in  $\binom{n}{m}$  ways. For each selection  $(i_1, \dots, i_m)$  there are  $m!$  arrangements of the components and each arrangement has probability  $(\lambda_{i_1} \dots \lambda_{i_m})/\delta(0)\dots\delta(m-1)$  and hence the R-H-S.

Let  $P = (p_{ij})$  be the transition matrix of the markov chain  $f(Y)$ . Using (2.3) and Lemma 2.1 we can easily verify that for  $1 \leq m \leq n-1$ ,

$$P_{mj} = \frac{\sum_{1 \leq i_1 < i_2 < \dots < i_{m+1} \leq n} \lambda_{i_1} \lambda_{i_2} \dots \lambda_{i_{m+1}}}{\delta(m) \sum_{1 \leq i_1 < \dots < i_m \leq n} \lambda_{i_1} \dots \lambda_{i_m}}, \quad \text{if } j=m+1 \quad (2.5)$$

$$= 1 - P_{m \ m+1}, \quad \text{if } j=m-1$$

$$= 0, \quad \text{otherwise,}$$

$$\text{and } P_{01} = P_{n \ n-1} = 1.$$

The preceding discussion also shows that the markov process  $f(X)$  is a birth and death process with birth rates  $\{p_{m m+1} \delta(m), 0 \leq m \leq n-1\}$  and death rates  $\{p_{m m-1} \delta(m), 1 \leq m \leq n\}$ . If the coherent system under consideration is a  $k$ -out-of- $n$  system then the life length of the system is just the first passage time to the set  $\{k+1, \dots, n\}$  for the process  $f(X)$ . Thus the life distribution of the  $k$ -out-of- $n$  system is IFR follows from the result in section 6.3 of Brown and Chaganty [4]. Summarizing the above discussion we have the following main result of this paper.

Theorem 2.2. Consider a  $k$ -out-of- $n$  system with independent repairable components. Assume that the life distributions are exponential with parameters  $\lambda_1, \dots, \lambda_n$  and repair distributions also exponential with parameters  $\mu_1, \dots, \mu_n$ . If  $\lambda_i - \mu_i = \Delta$  for all  $i$ , then the distribution of the system life is IFR.

Remark 2.3. If  $\lambda_1 = \lambda_2 = \dots = \lambda_n$  and  $\mu_1 = \mu_2 = \dots = \mu_n$  the above Theorem 2.2 shows that the life distribution of the  $k$ -out-of- $n$  system is IFR. This result was mentioned earlier in the introduction.

#### REFERENCES

- [1] Barlow, R. E. and Proschan, F. (1975). Statistical Theory of Reliability and Life Testing. Holt, Rhinehart and Winston, New York.
- [2] Birnbaum, Z. W., Esary, J. D., and Marshall, A. W. (1966). "Stochastic characterization of wearout for components and systems," Ann. Math. Statist., 37, 816-25.
- [3] Brown, M. (1975). "The first passage time distribution for a parallel exponential system with repair," Reliability and Fault Tree Analysis, edited by Barlow, R. E., Fussell, J., and Singpurwalla, N. Conference volume published by SIAM, Philadelphia.
- [4] Brown, M and Chaganty, N.R. (1980). "On the first passage time distribution for a class of markov chains," to appear in Ann. of Prob.
- [5] Esary, J. D. and Proschan, F. (1970). "A reliability bound for systems of maintained interdependent components," JASA, 65, 329-339.
- [6] Derman, C., Ross, S. M., and Schechner, Z. (1979). "A note on first passage times in birth and death and negative diffusion processes," unpublished manuscript.
- [7] Keilson, J. (1979). Markov Chain Models - Rarity and Exponentiality. Springer-Verlag, New York.
- [8] Kemeny, J. G. and Snell, J. L. (1960). Finite Markov Chains. D. Van Nostrand Co., Inc.. Princeton, N.J.
- [9] Ross, S. M. (1976). "On time to first failure in multicomponent exponential reliability systems," Journal of Stochastic Processes and their Appln., 4, 167-173.