



ADA 129622

(11)

NAVAL SUBMARINE MEDICAL RESEARCH LABORATORY

SUBMARINE BASE, GROTON, CONN.



MEMORANDUM REPORT 83-3

ANTHROPOMETRIC INDICES
AMONG U.S. NAVY DIVERS

by

M. L. Dembert and L. W. Mooney
with
J. F. Jekel

JUN 21 1983
H

Naval Medical Research and Development Command
Research Work Unit MR00001.01-5078

and partly supported

by a grant from United Technologies Corporation to Yale

Released by:

William C. Milroy, CAPT, MC, USN
Commanding Officer
Naval Submarine Medical Research Laboratory

8 June 1983

83 06 21 08 5

11

MEMORANDUM REPORT 83-3

ANTHROPOMETRIC INDICES AMONG U.S. NAVY DIVERS

by

LCDR Mark L. Dembert, MC, USN

and

Loren W. Mooney

Naval Submarine Medical Research Laboratory


with

James F. Jekel, M.D., M.P.H.

Department of Epidemiology and Public Health
Yale University School of Medicine, New Haven, Conn.

DTIC
ELECTE
JUN 21 1983
H D

Approved and Released by:



William C. Milroy, CAPT, MC, USN
Commanding Officer
Naval Submarine Medical Research Laboratory

Approved for public release; its distribution is unlimited

SUMMARY PAGE

THE PROBLEM

Only one large investigation of U.S. Navy diver anthropometry has been published to date. Additional data on anthropometric indices would provide a reference data bank which would assist in the resolution of operational, engineering and medical problems confronting Navy divers.

THE FINDINGS

Eighteen anthropometric indices obtained during the Longitudinal Health Survey have been analyzed from a cohort of 194 divers. These divers represent the spectrum of naval diving programs. Analysis was performed at two levels. First, statistics on the total cohort were computed. Second, the cohort was divided into two age strata which were felt to correspond to two levels of diving activity: very active and somewhat active. Statistics obtained for each index were fairly similar between strata.

APPLICATION

Knowledge of diver anthropometry enables this information to be applied to the design of naval diving suits and equipment, to research into diver performance and human factors research, to be incorporated into epidemiologic studies of diver health, and to contribute toward the periodic evaluation of physical standards for military diving personnel.

ADMINISTRATIVE INFORMATION

The initial data analysis of this work was supported by the Naval Medical Command, Naval Medical Research and Development Command Work Unit No. MR00001.01-5078. Subsequent analyses and the completion of the manuscript were supported by a grant from the United Technologies Corporation to Yale while the principal investigator (MLD) was a resident in preventive medicine at the Yale University School of Medicine.

The manuscript was submitted for review on 23 May 1983 and approved for publication on 8 June 1983. It has been designated as NSMRL Memo Rpt. 83-3.

Published by the Naval Submarine Medical Research Lab.

ABSTRACT

Anthropometric data obtained from 194 U.S. Navy divers who participated in the Longitudinal Health Survey were analyzed with descriptive statistics. Eighteen indices were examined, and the analysis was presented in two parts for each index. First, the mean value, its standard deviation, and percentiles were computed for the entire cohort. Second, the cohort was divided into two age strata, 20-30 years old and 31-42 years old. Within each stratum, the mean value, its standard deviation, and the range were calculated. When added to other published findings on U.S. Navy divers, these data have application as a reference source for personnel working in diving suit and equipment design, diving operations, human factors research, diving medicine, and epidemiology.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	



INTRODUCTION

Military anthropometry has been extensively investigated among U.S. Army and Marine Corps male and female personnel (1-2). It has also been incorporated into longitudinal studies of Naval aviators (3). However, military divers have been little characterized in terms of anthropometric indices.

From 1972-1978, the Naval Submarine Medical Research Laboratory conducted the initial phase of a Longitudinal Health Survey on a cohort of 197 U.S. Navy divers. The survey collected physical and historical information on fourteen multiphasic health profiles obtained during the initial examination of each diver. Anthropometry was one of these profiles, and relevant data were collected for several reasons. First, a review of the literature showed that there had been only one large study of U.S. Navy diver anthropometrics published to date (4). Other reports on associations between diver morbidity and physical characteristics considered few measures of body build or stature (5-6). Second, additional information on the characteristics of diver build could be incorporated into new designs for diving suits. Proper design is important for effective external temperature regulation of the body and it allows for better mobility and dexterity in performance. Third, information on diver build allows for the design of consoles in submersible vehicles as well as for the design of controls on diving equipment carried by the diver, e.g., regulator, gas mixture controls, straps and buckles. Finally, data collected on diver anthropometrics can be used in future epidemiologic studies of diver health that examine associations between anthropometry and risk for chronic diseases (e.g., coronary artery disease and diabetes mellitus), acute morbidity (e.g., musculoskeletal disorders such as low back pain), and diving-related problems (e.g., decompression sickness) (7).

METHODS

The purpose and design of the Longitudinal Health Survey have been described elsewhere (8-9). The initial findings and descriptive statistics on the cohort have been recently reported (10). This cohort was a nonrandom sample, as all who volunteered were accepted into the study. The cohort was nevertheless considered representative of the total Navy diving population, based on its distribution by rank, rate, and paygrade.

The eighteen anthropometric measures were taken on the advice of Mr. Robert White of the U.S. Army Natick Research and Development Command, Natick, Mass. All measurements were taken by the same examiner, who was trained by Mr. White. Weight was recorded on spring scales. Circumferences were measured with Keuffel and Esser Wytface tape. Linear parameters were measured with GPM anthropometers, except for the foot length, which was measured in a foot measuring box. Skinfold thickness determinations were made

with Lange skinfold calipers and were not obtained on all divers.

The individual method for each measurement was previously described (4,8-9), but this information is repeated here for the purpose of completeness of the report. Methods are combined with respective results and presented in Tables 1 through 18.

The original data for this study were collected during the Longitudinal Health Survey. The data for the present study were recoded and entered into the computer at Yale Computer Center, New Haven, Conn. All analyses were performed using programs from the Statistical Analysis System (11).

Upon the advice of Dr. Joseph Knapik of the Exercise Physiology Section of the U.S. Army Research Institute of Environmental Medicine, Natick, Mass., the mean, standard deviation and 5th, 10th, 25th, 50th, 75th, 90th, and 95th percentiles were computed for each measurement on the total cohort. This cohort was also broken into two age group strata for further analysis: 20-30, and 31-42. It was intuitively felt that more active divers would be in the former age interval, while less active divers would comprise the latter interval. This is supported by the findings of Berghage et al. (12), who found that approximately 69% of the active Navy divers in their survey were 30 years of age or younger. Also, the use of age intervals greater than five years (e.g., ten) is commonly used in descriptive epidemiologic studies and is predicated upon the available sample size. Stratification of the present cohort into smaller size age groups (e.g., five-year intervals) could obscure interpretation of the data because of the inherent large variability of values within such strata. Using two age strata, the mean, standard deviation, and range of values were computed for each index.

RESULTS

Tables 1-18 present the methodology and descriptive statistics for each anthropometric index.

DISCUSSION

The findings of this study are presented in a descriptive format and can serve as a reference for interested personnel in the fields of human factors, clothing and uniform design, diving medicine, environmental physiology, naval diving operations, and epidemiology.

A large data bank of anthropometric measures on U.S. Navy divers is important and should be enlarged in future investigations. This information can be used in the design of diving suits and equipment. It can also characterize populations of naval divers on the basis of body build and limb dimension, and on the amounts of subcutaneous

fat at various body sites. This latter information can be used in epidemiologic studies which try to relate risk for certain diseases and occupational injuries to multiphasic health variables. Results from these studies can also be incorporated into the redefinition of new standards of physical qualification for Navy divers.

REFERENCES

1. White RM. United States Army Anthropometry: 1946-1977. Technical Report Natick/TR-79/007. U.S. Army Natick Research and Development Command, Natick, Mass., July 1978.
2. White RM, Churchill E. United States Marine Corps Anthropometry. Technical Report Natick/TR-78/021. U.S. Army Research and Development Command, Natick, Mass., December 1977.
3. Oberman A, Mitchell RE, Graybiel A. Thousand Aviator Study Methodology. Monograph 11. U.S. Naval School of Aviation Medicine, Pensacola, Fla., July 1965.
4. Beatty HT, Berghage TE. Diver Anthropometrics. Navy Experimental Diving Unit Report No. 10-72. Panama City, Fla., June 1972.
5. Wise DA. Constitutional Factors in Decompression Sickness. Navy Experimental Diving Unit Report No. 2-63. Washington, D.C., April 1963.
6. Doll RE, Berghage TE. Interrelationships of Several Parameters of Decompression Sickness. Navy Experimental Diving Unit Report No. 7-65. Washington, D.C., March 1967.
7. Dembert ML, Jekel JF, Mooney LW. Health Risk Factors for the Development of Decompression Sickness among Navy Divers. Undersea Biomedical Research (submitted).
8. Sawyer RN, Baker JH. The Longitudinal Health Survey: I. Description. Naval Submarine Medical Research Laboratory Report No. 733. Groton, Conn., December 1972.
9. Tansey WA. The Longitudinal Health Study: A Multiphasic Medical Surveillance Program for U.S. Navy Submarine and Diving Personnel. Naval Submarine Medical Research Laboratory Report No. 786. Groton, Conn., May 1974.
10. Dembert ML, Mooney LW, Ostfeld AM, and Lacroix P. Multiphasic Health Profiles among Navy Divers. Undersea Biomedical Research 1983;10:45-61.
11. Helwig JT, Council KA, eds. SAS User's Guide. Carey, N.C.; Statistical Analysis System Institute, 1979, 303-306, 373-376.
12. Berghage TE, Rohrbaugh PA, Bachrach AJ, Armstrong FW. Navy Diving: Who's Doing It and Under What Conditions. Naval Medical Research Institute Report. Bethesda, Md., December 1975.

TABLE 1

Weight

Subject is weighed while wearing only undershorts. The unit of measure is kilograms.

Total Cohort		Age Strata	
		20-30	31-42
Number	194	Number	134
Mean value	81.2	Mean	80.0
Std. deviation	10.0	S.D.	9.5
Percentiles		Range	59.5-103.6
5th	65.9		60.9-110.0
10th	69.1		
25th	73.6		
50th	79.5		
75th	88.2		
90th	95.5		
95th	100.0		

Table 2

Height

Subject stands erect, with heels together and head level. Height is measured as the vertical distance from the top of the head (vertex). The unit of measure is centimeters.

Total Cohort		Age Strata	
		20-30	31-42
Number	194	Number	134
Mean	178.3	Mean	178.5
Std. Deviation	6.6	S.D.	6.7
Percentiles		Range	162.4-193.1
5th	167.8		159.1-190.0
10th	169.5		
25th	173.3		
50th	178.7		
75th	183.1		
90th	186.1		
95th	188.2		

TABLE 3

Functional Reach

Subject stands erect against a wall, with his scapulae touching the wall. His right arm is extended forward horizontally, with the tips of his thumb and index fingers pressed together. Functional reach is measured as the horizontal distance from the wall to the outer edge of the junction of the tips of the thumb and index fingers. The unit of measure is centimeters.

Total Cohort		Age Strata	
		20-30	31-42
Number	194	134	60
Mean	80.1	80.2	80.0
Std. Deviation	4.4	4.5	4.0
Percentiles		71.0-92.0	70.1-90.0
5th	73.2		
10th	74.4		
25th	76.8		
50th	80.2		
75th	83.1		
90th	86.0		
95th	87.2		

TABLE 4

Sitting Height

Subject sits erect with head level and with his feet resting on the floor. The seat is adjusted so that his knees are bent at right angles. Sitting height is measured as the vertical distance from the sitting surface to the top of the head (vertex). The unit of measure is centimeters.

Total Cohort		Age Strata	
		20-30	31-42
Number	194	134	60
Mean	92.6	92.6	92.5
Std. Deviation	3.7	3.9	3.3
Percentiles		80.4-102.8	82.2-97.9
5th	86.1		
10th	88.2		
25th	90.1		
50th	92.6		
75th	94.9		
90th	97.1		
95th	98.6		

TABLE 5

Chest Depth

Subject stands erect with his arms initially raised and then lowered after the anthropometer is in place under the right arm. The depth of the chest is measured at the level of the nipples during normal breathing. The unit of measure is centimeters.

Total Cohort		Age Strata	
		20-30	31-42
Number	194	134	60
Mean	24.8	24.3	25.9
Std. Deviation	2.1	1.9	2.3
Percentiles		18.0-29.7	21.4-31.5
5th	21.7		
10th	22.4		
25th	23.4		
50th	24.6		
75th	26.3		
90th	27.3		
95th	28.9		

TABLE 6

Chest Breadth

Subject stands erect with his arms initially raised and then lowered after the anthropometer is in place. The breadth of the chest is measured at the level of the nipples during normal breathing. The unit of measure is centimeters.

Total Cohort		Age Strata	
		20-30	31-42
Number	194	134	60
Mean	34.3	34.1	34.9
Std. Deviation	2.5	2.3	2.8
Percentiles		23.5-40.0	25.4-41.9
5th	30.9		
10th	31.6		
25th	32.8		
50th	34.3		
75th	35.8		
90th	37.5		
95th	38.5		

TABLE 7

Shoulders (Bideltoid) Breadth

Subject sits erect with his arms bent to form right angles at the elbows and with his elbows held against the body. The maximum breadth across the shoulders is measured at the level of the bulges of the deltoid muscles in the upper arms. The unit of measure is centimeters.

Total Cohort	Age Strata			
	20-30	31-42		
Number	194	Number 134	60	
Mean	48.7	Mean	48.5	49.2
Std. Deviation	2.4	S.D.	2.3	2.5
Percentiles		Range	43.2-54.7	44.2-55.6
5th	45.0			
10th	45.7			
25th	47.0			
50th	48.5			
75th	50.1			
90th	52.1			
95th	52.7			

TABLE 8

Hand Length

Subject sits with his right hand and fingers extended and palm up. The length of the right hand is measured from the wrist crease to the tip of the middle finger. The unit of measure is centimeters.

Total Cohort	Age Strata			
	20-30	31-42		
Number	194	Number 134	60	
Mean	19.3	Mean	19.3	19.1
Std. Deviation	0.9	S.D.	0.8	0.9
Percentiles		Range	17.2-21.5	16.9-21.0
5th	17.8			
10th	18.3			
25th	18.7			
50th	19.2			
75th	19.9			
90th	20.4			
95th	20.6			

TABLE 9

Hand Breadth

Subject sits with his right hand and fingers extended and palm up. The breadth of the hand is measured at the level of the knuckles (distal ends of the metacarpal bones). The unit of measure is centimeters.

Total Cohort		Age Strata	
		20-30	31-42
Number	194	134	60
Mean	9.3	9.3	9.2
Std. Deviation	0.4	0.4	0.4
Percentiles			
5th	8.6		
10th	8.8		
25th	9.0		
50th	9.2		
75th	9.6		
90th	9.8		
95th	10.0		
		8.5-10.7	8.0-10.2

TABLE 10

Foot Length

Subject stands erect with his weight evenly distributed on both feet. The maximum length of the right foot is measured from the back of the heel to the tip of the longest toe. The unit of measure is centimeters.

Total Cohort		Age Strata	
		20-30	31-42
Number	194	134	60
Mean	27.1	27.2	26.9
Std. Deviation	1.3	1.2	1.5
Percentiles			
5th	25.0		
10th	25.5		
25th	26.2		
50th	27.0		
75th	28.0		
90th	28.9		
95th	29.2		
		24.3-29.9	20.1-29.5

TABLE 11

Inspiratory Chest Circumference

Subject stands erect with his arms initially raised and then lowered after the tape is in place. The maximum horizontal circumference of the chest is measured at the level of the nipples at full inspiration. The unit of measure is centimeters.

Total Cohort		Age Strata	
		20-30	31-42
Number	194	Number	134
Mean	103.2	Mean	102.2
Std. Deviation	6.2	S.D.	6.0
Percentiles		Range	91.2-121.0
5th	94.0		94.3-120.1
10th	95.7		
25th	99.0		
50th	102.6		
75th	107.3		
90th	111.2		
95th	113.8		

TABLE 12

Expiratory Chest Circumference

Subject stands erect with his arms initially raised and then lowered after the tape is in place. The maximum horizontal circumference of the chest is measured at the level of the nipples at full expiration. The unit of measure is centimeters.

Total Cohort		Age Strata	
		20-30	31-42
Number	194	Number	134
Mean	96.7	Mean	95.4
Std. Deviation	6.2	S.D.	5.7
Percentiles		Range	82.2-110.7
5th	87.0		84.9-116.6
10th	89.6		
25th	92.4		
50th	96.6		
75th	99.5		
90th	105.5		
95th	108.4		

TABLE 13

Waist Circumference

Subject stands erect with abdomen relaxed. The maximum horizontal circumference of the waist is measured at the level of the umbilicus. The unit of measure is centimeters.

Total Cohort	Age Strata	
	20-30	31-42
Number	193	Number 134 59
Mean	88.5	Mean 86.4 93.4
Std. Deviation	8.5	S.D. 7.8 8.1
Percentiles		Range 71.2-106.8 75.5-110.0
5th	76.2	
10th	77.5	
25th	82.0	
50th	88.5	
75th	94.6	
90th	100.0	
95th	103.2	

TABLE 14

Vertical Trunk Circumference

Subject stands erect with his feet slightly apart. The vertical circumference of the trunk is measured with a steel tape passed through the crotch, to the right of the scrotum, and then over the midpoints of the right shoulder and right buttock. The unit of measure is centimeters.

Total Cohort	Age Strata	
	20-30	31-42
Number	193	Number 134 59
Mean	163.9	Mean 162.9 166.1
Std. Deviation	9.4	S.D. 9.8 7.9
Percentiles		Range 104.5-198.6 146.8-182.1
5th	151.0	
10th	153.0	
25th	159.2	
50th	163.5	
75th	169.2	
90th	174.5	
95th	178.2	

TABLE 15

Triceps Skinfold Thickness

The skinfold is picked up on the dorsum of the right arm, at a point midway between the acromial process of the scapula and the tip of the olecranon. Measurement is made with the forearm flexed to 90° and the arm pendant. The unit of measure is millimeters.

Total Cohort		Age Strata	
		20-30	31-42
Number	153	Number	104
Mean	17.7	Mean	17.2
Std. Deviation	7.8	S.D.	7.5
Percentiles		Range	3-35
5th	7		4-33
10th	8		
25th	12		
50th	17		
75th	23		
90th	29		
95th	30		

TABLE 16

Subscapular Skinfold Thickness

The skinfold is picked up under the angle of the right scapula. The fold is pointing slightly downward and outward. The unit of measure is millimeters.

Total Cohort		Age Strata	
		20-30	31-42
Number	153	Number	104
Mean	20.1	Mean	18.2
Std. Deviation	9.0	S.D.	9.2
Percentiles		Range	6-42
5th	9		8-46
10th	10		
25th	13		
50th	18		
75th	28		
90th	33		
95th	36		

TABLE 17

Suprailiac Skinfold Thickness

The skinfold is picked up one centimeter above and two centimeters medial to the right anterior superior iliac spine. The unit of measure is millimeters.

Total Cohort		Age Strata	
		20-30	31-42
Number	153	Number	104
Mean	24.3	Mean	23.1
Std. Deviation	11.5	S.D.	11.6
Percentiles		Range	5-50
5th	7		7-46
10th	9		
25th	14		
50th	23		
75th	34		
90th	41		
95th	45		

TABLE 18

Abdominal Skinfold Thickness

The skinfold is picked up at the level of the umbilicus and 2.5 centimeters to the right of it. The unit of measure is millimeters.

Total Cohort		Age Strata	
		20-30	31-42
Number	153	Number	104
Mean	24.8	Mean	22.0
Std. Deviation	13.2	S.D.	12.1
Percentiles		Range	4-58
5th	8		8-54
10th	9		
25th	14		
50th	22		
75th	34		
90th	45		
95th	48		

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NSMRL Memo Report 83-3	2. GOVT ACCESSION NO. AD-A129622	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Anthropometric Indices Among U. S. Navy Divers		5. TYPE OF REPORT & PERIOD COVERED Interim report
		6. PERFORMING ORG. REPORT NUMBER NSMRL Memo 83-3
7. AUTHOR(s) M. L. DEMBERT, LCDR, MSC, USN L. W. MOONEY and J. F. JEKEL, M.D., and M.P.H.		8. CONTRACT OR GRANT NUMBER(s) Partly supported by a grant from United Technologies to Yale
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Submarine Medical Research Laboratory Naval Submarine Base Nlon Groton, CT 06349		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Partly funded by MR00001.01-5078
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Medical Research and Development Command Naval Medical Command, National Capital Region Bethesda, MD 20814		12. REPORT DATE 8 June 1983
		13. NUMBER OF PAGES 13
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) anthropometric indices; naval diving suits and equipment design;		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Anthropometric data obtained from 194 U.S. Navy divers who participated in the Longitudinal Health Survey were analyzed with descriptive statistics. Eighteen indices were examined, and the analysis was presented in two parts for each index. First, the mean value, its standard deviation, and percentiles were computed for the entire cohort. Second, the cohort was divided into two age strata, 20-30 years old and 31-42 years old. Within each stratum, the mean value, its standard deviation, and the range were calculated. When added to other published findings on U. S. Navy		

DD FORM 1473
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6001

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

Item 20--continued

divers, these data have application as a reference source for personnel working in diving suit and equipment design, diving operations, human factors research, diving medicine and epidemiology.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)