

AD-A133 922


SOME NEW RESULTS ON GRUBBS' ESTIMATORS(U) FLORIDA STATE 1/1  
UNIV TALLAHASSEE DEPT OF STATISTICS  
D A BRINDLEY ET AL. JUN 83 FSU-STATISTICS-M648


UNCLASSIFIED

N00014-80-C-0093

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A133922

SOME NEW RESULTS ON GRUBBS ESTIMATORS  
Dennis A. Brinley  
Bristol-Myers Pharmaceutical Research and Development Divi  
and  
Ralph A. Bratley  
University of Georgia and Florida State University

The Florida State University  
Department  
of  
Statistics  
Tallahassee, Florida  
32306

OCT 24 1963

E

DIC FILE COPY

83 10 04 1965

SOME NEW RESULTS ON GRUBBS' ESTIMATORS \*

by

Dennis A. Brindley  
Bristol-Myers Pharmaceutical Research and Development Division  
and  
Ralph A. Bradley  
University of Georgia and Florida State University



FSU Statistics Report No. M648  
ONR Technical Report No. 159  
University of Georgia Statistics Report No. 154

Accession For		
NTIS ONLY		
DTIC		
UNCLASSIFIED		
CONFIDENTIAL		
SECRET		
Dist		
A		

June, 1983

The University of Georgia  
Department of Statistics and Computer Science  
Athens, Georgia 30602

The Florida State University  
Department of Statistics  
Tallahassee, Florida 32306

\* This work was supported by the Army, Navy and Air Force under Office of Naval Research Contract No. N00014-80-C-0093. Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document is for personal use only. Distribution is prohibited.

## SOME NEW RESULTS ON GRUBBS' ESTIMATORS

DENNIS A. BRINDLEY AND RALPH A. BRADLEY\*

→ Consider a two-way classification with  $n$  rows and  $r$  columns and the usual model of analysis of variance except that the error components of the model may have heterogeneous variances  $\sigma_j^2, j = 1, \dots, r$  by columns. Grubbs provided unbiased estimators  $Q_j$  of  $\sigma_j^2$  that depend on column sums of squared residuals  $S_j$ . → When  $r = 3$ , the joint distributions of the  $S_j$  and the  $Q_j$  are given for the first time in closed form.

Two tests proposed by Russell and Bradley are examined when  $r = 3$ , one for variance homogeneity and the second for one possible disparate variance. A very simple distribution is found for the test statistic of the first test and its non-null distribution is derived also. The distribution of the second test statistic was known to be the central variance-ratio distribution in the null case and now its ratio to a parameter of noncentrality is shown to have that same distribution in the non-null case.

When  $r = 4, n = 4$ , the joint distribution of the  $S_j$  is given also in closed form, but it is difficult to use. For  $r > 3$ , an approximate test of variance homogeneity is proposed, based on an extension of the Russell-Bradley statistic. → Extensive simulation studies show that the distribution of the test statistic may be approximated very well by a chi-square distribution. ←

---

\* Dennis A. Brindley is a research scientist with Bristol-Myers Pharmaceutical Research and Development Division, Evansville, Indiana 47630. Ralph A. Bradley is Research Professor, Department of Statistics and Computer Science, University of Georgia, Athens, Georgia 30602.

## 1. INTRODUCTION

Consider the two-way classification of observations  $y_{ij}$ ,  
 $i = 1, \dots, n$ ,  $j = 1, \dots, r$ , and the model,

$$y_{ij} = \mu_i + \beta_j + \varepsilon_{ij}, \quad (1)$$

when  $\mu_i$  represents the mean response of row  $i$ ,  $\beta_j$  represents the additional effect of column  $j$ ,  $\sum_j \beta_j = 0$ , and the  $\varepsilon_{ij}$  are independent, zero-mean, normal variates with expectation  $E(\varepsilon_{ij}^2) = \sigma_j^2$ . Model (1) differs from the usual model in that variances are column-related. Grubbs (1948) estimated  $\sigma_j^2$  using  $Q_j$ ,  $j = 1, \dots, r$ , where

$$Q_j = \frac{r}{(n-1)(r-2)} S_j - \frac{1}{(n-1)(r-1)(r-2)} \sum_{t=1}^r S_t \quad (2)$$

and

$$S_j = \sum_{i=1}^n (y_{ij} - y_{i.} - y_{.j} + y_{..})^2 \quad (3)$$

in which  $y_{i.}$ ,  $y_{.j}$  and  $y_{..}$  are respectively the means of the observations in the  $i$ -th row,  $j$ -th column, and complete table. Note that  $E(Q_j) = \sigma_j^2$ , but  $Q_j$  may be negative in certain situations.

Possible use of Grubbs' estimators, and inferences based upon them, arises frequently. Grubbs was concerned with the variabilities of observer-used electric clocks as timing instruments, Ehrenberg (1950) with the precisions of individuals in the sensory scoring of food samples, and Russell and Bradley (1958) with an application to three

fermentors in a distillery. More recently, Snee (1982) examined wheat-yield data and multiple-head machine data and suggested that apparent variance heterogeneity may be an indicator and manifestation of omitted interaction terms needed in the model. A substantial bibliography on Grubbs' estimators has developed and was summarized by Maloney (1973).

The use of Grubbs' estimators has been hampered by the difficulties encountered in developing appropriate inference procedures, although some results are available. Russell and Bradley gave two procedures. The first was based on the assumption,

$$A: \sigma_j^2 = \sigma^2, j = 1, \dots, (r - 1), \quad (4)$$

and provided a test of the hypothesis,

$$H_{01}: \sigma_r^2 = \sigma^2. \quad (5)$$

They showed that

$$F = [r(r - 2)Q_r + \sum_j Q_j] / r(\sum_j Q_j - Q_r) \quad (6)$$

has the central F-distribution with  $(n - 1)$  and  $(n - 1)(r - 2)$  degrees of freedom under A and  $H_{01}$ . In the special case with  $r = 3$ , they considered the hypothesis,

$$H_{02}: \sigma_1^2 = \sigma_2^2 = \sigma_3^2, \quad (7)$$

and showed that, under  $H_{02}$ , the statistic,

$$S = -(n - 1)\log T, \quad (8)$$

where

$$T = 3(Q_1Q_2 + Q_1Q_3 + Q_2Q_3)/(Q_1 + Q_2 + Q_3)^2, \quad (9)$$

has the central chi-square distribution with 2 degrees of freedom asymptotically with  $n$ . Shukla (1982) proposed a Bartlett-type statistic, which, in the notation of this paper, depends on  $\log T^*$  where  $T^* = \prod_j S_j / \bar{S}^r$  with  $\bar{S} = \sum_j S_j / r$ . He also approximated the distribution of  $\log T^*$ . Johnson (1962) specified  $j$  and  $j'$ ,  $j \neq j'$ , and used  $S_j / S_{j'}$  to test the hypothesis that  $\sigma_j^2 = \sigma_{j'}^2$ , obtaining a distribution that he approximated with a single F-distribution.

Ellenberg (1977), under the hypothesis that  $\sigma_j^2 = \sigma^2$ ,  $j = 1, \dots, r$ , derived the exact distribution of a subset,  $S_1, \dots, S_k$ ,  $k \leq r$ , of  $S_1, \dots, S_r$ . A complicated infinite series expression resulted that he regarded as intractable for further use.

In this article, a direct attempt is made to obtain the joint distribution of  $S_1, \dots, S_r$  and of  $Q_1, \dots, Q_r$  that is partially successful. Simple results are obtained when  $r = 3$ , the most important case. Then the exact distribution of  $T$  in (9) is derived both under  $H_{01}$  in (7) and under the general alternative hypothesis,  $H_{02}: \sigma_j^2 \neq \sigma^2$  for some  $j = 1, 2, 3$ . The non-null distribution of  $F$  in (6) is found also. When  $r = 4$ , the joint distribution of  $S_1, \dots, S_4$  is obtained under the extension of (7) for  $r = 4$ , but this distribution is complicated, although given in closed form. In general, when  $r > 3$ , distributional problems become very difficult; an approximate test procedure for equality of variances is then suggested.



## 2. DISTRIBUTION THEORY

We seek the joint distribution of  $S_1, \dots, S_r$ ,  $r \geq 3$ . Let  $\underline{y}$  be the column vector with typical element  $y_{ij}$ , the elements arranged in lexicographic order. Define  $\underline{A}_n$  to be the  $(n-1) \times n$  matrix with the  $(n-1)$  zero-sum rows of the  $n$ -dimensional Helmert matrix arranged so that row  $s$  has  $s$  elements  $1/[s(s+1)]^{1/2}$  followed by the element  $-s/[s(s+1)]^{1/2}$  and  $(n-s-1)$  zero elements. The usual  $(n-1)(r-1)$  error contrasts may be written as the elements of

$$\underline{e} = (\underline{A}_n \otimes \underline{A}_r) \underline{y}, \quad (10)$$

where the typical element of  $\underline{e}$  is  $e_{ij}$ ,  $i = 1, \dots, (n-1)$ ,  $j = 1, \dots, (r-1)$ , arranged in lexicographic order and  $\underline{A}_n \otimes \underline{A}_r$  is the direct product of  $\underline{A}_n$  and  $\underline{A}_r$ .

It is clear that  $E(\underline{e}) = \underline{0}$  and it may be shown that the covariance matrix of  $\underline{e}$  is  $\underline{\Sigma}_e = \underline{I}_{(n-1)} \otimes \underline{\Sigma}$ , where  $\underline{I}_m$  is the  $m$ -dimensional identity matrix. The matrix  $\underline{\Sigma}$  is  $(r-1)$ -square symmetric with diagonal elements,

$$\left( \sum_{j=1}^s \sigma_j^2 + s^2 \sigma_{s+1}^2 \right) / s(s+1), \text{ and } (s, t)\text{-elements,}$$

$$\left( \sum_{j=1}^t \sigma_j^2 - t \sigma_{t+1}^2 \right) / [t(t+1)s(s+1)]^{1/2}, \quad t < s, \quad s = 1, \dots, (r-1).$$

The form of  $\underline{\Sigma}_e$  indicates that  $\underline{e}$  may be partitioned into  $(n-1)$  independent  $(r-1)$ -dimensional vectors, each having covariance matrix  $\underline{\Sigma}$ ,

$\underline{e}' = (\underline{e}'_1, \dots, \underline{e}'_{n-1})$ . In effect,  $\underline{e}'_1, \dots, \underline{e}'_{n-1}$  may be regarded as  $(n-1)$  independent observation vectors from an  $(r-1)$ -dimensional multivariate normal population with mean vector  $\underline{0}$  and dispersion matrix  $\underline{\Sigma}$ .

Let  $\underline{V} = \sum_{i=1}^{n-1} \underline{e}'_i \underline{e}'_i$ . Then the  $r(r-1)/2$  distinct elements of  $\underline{V}$  have the Wishart density with  $(n-1)$  degrees of freedom and dispersion

matrix  $\underline{\Sigma}$  so that

$$f(\underline{V}; \underline{\Sigma}) = \frac{|\underline{V}|^{(n-r-1)/2} \exp(-\frac{1}{2} \text{tr} \underline{V} \underline{\Sigma}^{-1})}{2^{(n-1)(r-1)/2} \pi^{(r-1)(r-2)/4} |\underline{\Sigma}|^{(n-1)/2} \prod_{j=1}^{r-1} \Gamma[\frac{1}{2}(n-j)]} \quad (11)$$

for  $\underline{V}$  positive definite and zero otherwise,  $f$  being used generically to denote a density function. Furthermore, if  $v_{jj'}$  is the typical element of  $\underline{V}$ , algebraic reduction of (3) yields

$$S_j = [(j-1)/j] v_{j-1, j-1} + \sum_{t=j}^{r-1} [v_{tt}/t(t+1)] + \quad (12)$$

$$2 \left[ \sum_{j \leq t < t' < r} v_{tt'} / \{t(t+1)t'(t'+1)\}^{1/2} - \{(j-1)/j\}^{1/2} \sum_{t=j}^{r-1} v_{j-1, t} / \{t(t+1)\}^{1/2} \right],$$

$j = 1, \dots, r.$

When  $\sigma_j^2 = \sigma^2$ ,  $j = 1, \dots, r$ ,  $\underline{\Sigma} = \sigma^2 \underline{I}_{r-1}$  and  $\underline{\Sigma}^{-1} = \underline{I}_{r-1} / \sigma^2$ .

The statistics  $F$ ,  $S$ , and  $T$  in (6), (8) and (9) respectively, and any extensions to be considered, are scale invariant and we may take  $\sigma^2 = 1$  without loss of generality when considering the distributions of such statistics when  $\sigma_j^2 = \sigma^2$ ,  $j = 1, \dots, r$ . Then

$$f(\underline{V}; \underline{I}) = \frac{|\underline{V}|^{(n-r-1)/2} \exp(-\frac{1}{2} \text{tr} \underline{V})}{2^{(n-1)(r-1)/2} \pi^{(r-1)(r-2)/4} \prod_{j=1}^{r-1} \Gamma[\frac{1}{2}(n-j)]} \quad (13)$$

for  $\underline{V}$  positive definite and zero otherwise.

In this section, we have reduced the apparent dependence of the  $S_j$ , and the  $Q_j$ , on the  $nr$  original observations to dependence on  $\frac{1}{2}r(r-1)$  new variables with known joint distribution (11). It is clear from (12) that only a non-singular linear transformation is

needed to obtain the joint distribution of  $S_1$ ,  $S_2$  and  $S_3$  when  $r = 3$ . We turn to special cases.

### 3. SPECIAL CASES

**3.1 Joint Distributions,  $r = 3$ :** When  $r = 3$ , the joint distributions of  $S_1$ ,  $S_2$  and  $S_3$  and of  $Q_1$ ,  $Q_2$  and  $Q_3$  follow directly from (11), (12) and (2). Now

$$\begin{aligned} S_1 &= (v_{11}/2) + (v_{22}/6) + (v_{12}/\sqrt{3}), \\ S_2 &= (v_{11}/2) + (v_{22}/6) - (v_{12}/\sqrt{3}), \\ S_3 &= 2v_{22}/3, \end{aligned} \quad (14)$$

$$\underline{v} = \begin{bmatrix} v_{11} & v_{12} \\ v_{12} & v_{22} \end{bmatrix} \quad \text{and} \quad \underline{\Sigma} = \begin{bmatrix} (\sigma_1^2 + \sigma_2^2)/2 & (\sigma_1^2 - \sigma_2^2)/\sqrt{12} \\ (\sigma_1^2 - \sigma_2^2)/\sqrt{12} & (\sigma_1^2 + \sigma_2^2 + 4\sigma_3^2)/6 \end{bmatrix}.$$

The linear transformation (14) lets us write

$$\begin{aligned} f(S_1, S_2, S_3; \underline{\Sigma}) &= \frac{1}{\pi \Gamma(n-2)} \left( \frac{9}{4 \sum_{j < j'} \sigma_j^2 \sigma_{j'}^2} \right)^{\frac{1}{2}(n-1)} \left[ (\sum_j S_j)^2 - 2 \sum_j S_j^2 \right]^{\frac{1}{2}(n-4)} \\ &\quad \exp \left[ - \frac{1}{\sum_{j < j'} \sigma_j^2 \sigma_{j'}^2} \left( \sum_{j \neq j'} \sigma_j^2 S_j - \frac{1}{2} \sum_j \sigma_j^2 S_j \right) \right], \end{aligned} \quad (15)$$

where  $0 < S_j < 2 \sum_{j'} S_{j'}/3$ ,  $j = 1, 2, 3$ , and  $(\sum_j S_j)^2 - 2 \sum_j S_j^2 > 0$ . From (2), the density of  $Q_1$ ,  $Q_2$  and  $Q_3$  is

$$\begin{aligned} f(Q_1, Q_2, Q_3; \underline{\Sigma}) &= \frac{(n-1)^{n-1}}{4\pi \Gamma(n-2) \left( \sum_{j < j'} \sigma_j^2 \sigma_{j'}^2 \right)^{\frac{1}{2}(n-1)}} (Q_1 Q_2 + Q_1 Q_3 + Q_2 Q_3)^{\frac{1}{2}(n-4)} \\ &\quad \exp \left( - \frac{n-1}{2 \sum_{j < j'} \sigma_j^2 \sigma_{j'}^2} \sum_{j \neq j'} \sigma_j^2 Q_j \right), \end{aligned} \quad (16)$$

where  $Q_1 + Q_2 + Q_3 > 0$  and  $Q_1Q_2 + Q_1Q_3 + Q_2Q_3 > 0$ . Note that one  $Q_j$  may be negative but not two.

3.2 Russell-Bradley Test,  $r = 3$ : The non-null distribution of  $F$  in (6) may be found from (16) when  $\sigma_1^2 = \sigma_2^2 = \sigma^2$ ,  $\sigma_3^2 \neq \sigma^2$ . Now  $F = [1 + 4Q_3/(Q_1 + Q_2)]/3$  and its density is

$$f(F; \gamma) = \gamma^{\frac{1}{2}(n-1)} F^{\frac{1}{2}(n-3)} (\gamma + F)^{-(n-1)} / B[\frac{1}{2}(n-1), \frac{1}{2}(n-1)], \quad (17)$$

where  $F > 0$ ,  $\gamma = (\sigma^2 + 2\sigma_3^2)/3\sigma^2$ . Thus, for this test,  $F/\gamma$  has the central  $F$ -distribution with  $(n-1)$  and  $(n-1)$  degrees of freedom under both  $H_{01}$  in (5), for which  $\gamma = 1$ , and  $H_{a1}:\sigma_3^2 \neq \sigma^2$  when  $r = 3$ .

Details on this result are given by Brindley (1982). The transformation,  $x = Q_1 + Q_2$ ,  $y = Q_1/(Q_1 + Q_2)$ ,  $z = Q_3/(Q_1 + Q_2)$ , is used to obtain (17) from (16) since the marginal distribution of  $z$  may be obtained and  $F$  is directly dependent on  $z$ .

To illustrate the results of this subsection, suppose that a one-sided alternative hypothesis is considered, say  $H_{a1}:\sigma_3^2 > \sigma^2$ . Let  $F^*$  be a random variable with the central  $F$ -distribution with  $(n-1)$  and  $(n-1)$  degrees of freedom and let  $F_\alpha$  be such that  $P(F^* > F_\alpha) = \alpha$ . If  $F$  is computed from (6) with  $r = 3$ ,  $H_{01}$  in (5), given  $A$  in (4), is rejected at significance level  $\alpha$  if  $F > F_\alpha$ . The power of the test depends on  $\gamma$ , specified when  $\sigma_3^2/\sigma^2$  is specified. The power of the test for specified  $\gamma$  is evaluated simply as  $P(F^* > F_\alpha/\gamma)$ . The complementary one-sided alternative hypothesis or the two-sided alternative hypothesis,  $\sigma_3^2 \neq \sigma^2$ , are considered in similar ways.

3.3 Variance Homogeneity Test,  $r = 3$ : The distributions of  $T$  in (9) and of  $S$  in (8) may be obtained when  $r = 3$  under both  $H_{02}$  in (7) and the alternative hypothesis,  $H_{a2}: \sigma_j^2 \neq \sigma_{j'}^2$ , some  $j \neq j'$ ,  $j, j' = 1, 2, 3$ . Indeed, under  $H_{a2}$ , the distribution of  $T$  is

$$f(T; \lambda) = \frac{1}{2}(n-2)\lambda^{\frac{1}{2}(n-1)}T^{\frac{1}{2}(n-4)}G\left[\frac{1}{2}(n-1), n/2, 1, (1-T)(1-\lambda)\right], \quad (18)$$

where  $T > 0$ ,

$$\lambda = 3 \sum_{j < j'} \frac{\sigma_j^2 \sigma_{j'}^2}{(\sum_j \sigma_j^2)^2}, \quad (19)$$

and  $G$  is the usual hypergeometric function. When  $H_{02}$  is true,  $\lambda = 1$  and

$$f(T; 1) = \frac{1}{2}(n-2)T^{\frac{1}{2}(n-4)}, \quad T > 0, \quad (20)$$

a very simple density. Furthermore, it follows at once that the density of  $(n-2)S/(n-1)$  is the central chi-square distribution with 2 degrees of freedom, a result differing from the approximate one of Russell and Bradley (1958) only by the factor  $(n-2)/(n-1)$ .

The test of the hypothesis  $H_{02}$  may be based on  $(n-2)S/(n-1)$  and the chi-square distribution, with large values of the statistic significant. The test may also be done simply in terms of  $T$ , with small values of  $T$  significant. Indeed, under  $H_{02}$  with  $r = 3$ ,  $P(T < \alpha^{2/(n-2)}) = \alpha$ . The power of the  $\alpha$ -level test is

$$P(T < \alpha^{2/(n-2)} | \lambda) = \int_0^{\alpha^{2/(n-2)}} f(T, \lambda) dT$$

for specified  $\lambda$  with  $f(T, \lambda)$  given in (18).

3.4 Other Tests,  $r = 3$ : The joint densities (15) and (16) are relatively simple when  $\sigma_1^2 = \sigma_2^2 = \sigma_3^2 = \sigma^2$ ,  $\sigma^2 = 1$  without loss of generality.

Then we have

$$f(S_1, S_2, S_3) = \frac{3^{(n-1)/2}}{2^{n-1} \pi \Gamma(n-2)} [(\sum_j S_j)^2 - 2\sum_j S_j^2]^{(n-4)/2} e^{-\sum_j S_j/2}, \quad (21)$$

where  $0 < S_j < 2\sum_{j'} S_{j'}/3$ ,  $j = 1, 2, 3$ , and  $(\sum_j S_j)^2 - 2\sum_j S_j^2 > 0$ ,

and

$$f(Q_1, Q_2, Q_3) = \frac{(n-1)^{n-1}}{4\pi^3 (n-1)/2 \Gamma(n-2)} (Q_1 Q_2 + Q_1 Q_3 + Q_2 Q_3)^{(n-4)/2} e^{-(n-1)\sum_j Q_j/3}, \quad (22)$$

where  $\sum_j Q_j > 0$  and  $Q_1 Q_2 + Q_1 Q_3 + Q_2 Q_3 > 0$ . We have considered use of (21) to investigate the distribution of Shukla's statistic  $T^*$  for a test of variance homogeneity. The distribution of  $T^*$  is very intractable even when  $r = 3$  and its use cannot be recommended in view of the simple results obtained above for  $T$  when  $r = 3$ . Also, use of (21) to rederive the distribution of Johnson's statistic,  $S_j/S_{j'}$ ,  $j \neq j'$ , leads to no new insights or special simplifications.

3.5 A Joint Distribution,  $r = 4$ : The joint distribution of the elements of  $\underline{y}$  is given in (13), the number of variates on which  $S_1, \dots, S_r$  depend being  $\frac{1}{2} r(r-1)$ . We have been able to proceed to obtain the joint distribution of  $2r-3$  variates on which  $S_1, \dots, S_r$  depend, see Brindley (1982), but the joint distribution of  $S_1, \dots, S_r$  must be very complicated for general  $r$  and  $n$ .

The very special case with  $r = 4$  and  $n = 4$  has been completed.

Then

$$f(S_1, S_2, S_3, S_4) = 2^{-1/2} \pi^{-3/2} \min_{1 \leq j \leq 4} \{\sqrt{S_j}\} e^{-1/2 \sum_j S_j}$$

or

$$f(S_1, S_2, S_3, S_4) = (2\pi)^{-3/2} (\sum_j \sqrt{S_j} - 2 \max_{1 \leq j \leq 4} \{\sqrt{S_j}\}) e^{-1/2 \sum_j S_j}$$

as  $\sum_j S_j^2 - 2 \sum_{j < k} S_j S_k$  is less than  $-8(\prod_j S_j)^{1/2}$  or between  $-8(\prod_j S_j)^{1/2}$  and  $8(\prod_j S_j)^{1/2}$  respectively,  $S_j > 0$ ,  $j = 1, \dots, 4$ , and  $f(S_1, S_2, S_3, S_4)$  is zero otherwise.

It appears that the case with  $r = 3$ , an important case, may be the only situation for which simple distributions for test statistics may be found.

#### 4. THE USE OF CHARACTERISTIC FUNCTIONS

Ellenberg (1977) used the method of characteristic functions to obtain the joint distribution of a subset of  $k$  of  $S_1, \dots, S_r$ ,  $k \leq r$ , under the assumption that each  $\sigma_j^2 = 1$ . We summarize for  $k = r$  since coefficients in the series that he gives to represent the distribution are difficult to determine.

Let  $\phi(\underline{t})$  represent the characteristic function of  $S_1, \dots, S_r$ , where  $\underline{t} = (t_1, \dots, t_r)$ . From Ellenberg's formula (3.2) for  $\phi(\underline{t})$ , after algebraic reduction,

$$\phi(\underline{t}) = \prod_j (1 - 2it_j)^{-(n-1)/2} \left\{ \frac{1}{r} \sum_j (1 - 2it_j)^{-1} \right\}^{-(n-1)/2} \quad (23)$$

where  $i = \sqrt{-1}$  in (23). Since  $(1 - 2it)^{-k/2}$  is the characteristic function of a chi-square variate with  $k$  degrees of freedom, it is appropriate to expand  $\phi(t)$  in negative powers of the  $(1 - 2it_j)$ . The result is that

$$\phi(t) = \sum_{k=0}^{\infty} \frac{\Gamma(k + \frac{n-1}{2})}{k! \Gamma(\frac{n-1}{2})} \sum_{s=0}^k \binom{k}{s} \left(-\frac{1}{r}\right)^s \sum_{\alpha} \dots \sum_{\alpha} \frac{s!}{\prod_{\alpha} j_{\alpha}!} \prod_{\alpha} (1 - 2it_{\alpha})^{-(j_{\alpha} + \frac{n-1}{2})}$$

and

$$f(S_1, \dots, S_r) = \sum_{k=0}^{\infty} \frac{\Gamma(k + \frac{n-1}{2})}{k! \Gamma(\frac{n-1}{2})} \sum_{s=0}^k \binom{k}{s} \left(-\frac{1}{r}\right)^s \sum_{\alpha} \dots \sum_{\alpha} \frac{s!}{\prod_{\alpha} j_{\alpha}!} \prod_{\alpha} \frac{S_{\alpha}^{j_{\alpha} + \frac{n-1}{2} - 1}}{2^{j_{\alpha} + \frac{n-1}{2}} \Gamma(j_{\alpha} + \frac{n-1}{2})} \exp\left(-\frac{1}{2} \sum_j S_j\right), \quad (24)$$

where  $\alpha$  has the range  $1, \dots, r$  and  $S_j > 0$ ,  $j = 1, \dots, r$ , in (24).

It is far from apparent that (21) and (24) are identical when  $r = 3$ . Note that  $(\sum_j S_j)^2 - 2\sum_j S_j^2 > 0$  for (21) but this requirement does not accompany (24). Two verifications have been made.

The moment generating function,  $M_Q(\theta) = E\{\exp[(n-1)\sum_j \theta_j Q_j/3]\}$ , may be found from (22). This is done through use of the transformation,  $u = Q_1$ ,  $v = Q_1 + Q_2$ ,  $w = Q_3 + Q_1 Q_2 / (Q_1 + Q_2)$ , with integration with respect to  $w$ ,  $u$ , and  $v$  in turn. The result is that

$$M_Q(\theta) = \left\{1 - \frac{2}{3} \sum_j \theta_j + \frac{2}{3} (\theta_1 \theta_2 + \theta_1 \theta_3 + \theta_2 \theta_3) + \frac{1}{3} (\theta_1^2 + \theta_2^2 + \theta_3^2)\right\}^{-(n-1)/2}.$$

Since  $S_1 = (n-1)(4Q_1 + Q_2 + Q_3)/3$ ,  $S_2 = (n-1)(Q_1 + 4Q_2 + Q_3)/3$  and  $S_3 = (n-1)(Q_1 + Q_2 + 4Q_3)/3$ , the moment generating function of



$S_1, S_2$  and  $S_3$  is

$$M_{\underline{S}}(\underline{t}) = E(\exp \sum_j t_j S_j) = \left\{ 1 - \frac{4}{3} \sum_j t_j + \frac{4}{3}(t_1 t_2 + t_1 t_3 + t_2 t_3) \right\}^{-(n-1)/2},$$

corresponding to the characteristic function (23) for  $r = 3$  from which (24) was derived.

In the very special case with  $r = 3$  and  $n = 4$ , we have been able to sum the series in (24) to obtain the special form of (22). The method used seems awkward but an easier way has not been found. Details are given by Brindley (1982).

## 5. SIMULATIONS

Even though we have been unsuccessful in obtaining the joint distribution of  $Q_1, \dots, Q_r$  in usable form for  $r > 3$ , a test of variance homogeneity is needed. The statistic  $T$  in (9) yielded a simple distribution when  $r = 3$  and has intuitive appeal, particularly when written in the alternative form,

$$T = \left\{ 1 - \frac{(Q_1 - Q_2)^2 + (Q_1 - Q_3)^2 + (Q_2 - Q_3)^2}{2(Q_1 + Q_2 + Q_3)^2} \right\}.$$

We consider the statistic,

$$\chi_r = -(n - 1) \log T_r, \quad (25)$$

where

$$T_r = 2r \sum_{j < j'} Q_j Q_{j'} / (r - 1) (\sum_j Q_j)^2. \quad (26)$$

Note that  $T_3 = T$  and that  $T_r = 1$  in the event that all  $Q_j$  are equal. Choice of the multiplier  $(n - 1)$  in (25) was made after empirical investigation.

Let us suppose that, when  $\sigma_1^2 = \dots = \sigma_r^2$ ,  $x_r/a_r$  has the central chi-square distribution with  $\nu_r$  degrees of freedom. If the experiment is simulated a sufficient number of times,  $a_r$  and  $\nu_r$  may be evaluated.

Two series of simulations have been conducted, the first to determine  $a_r$  and  $\nu_r$  and the second to confirm and to extend the approximate procedure proposed to larger values of  $r$ . The method referenced in the Statistical Analysis Systems (SAS) user's guide was used to generate the required normal observations; properties of this generator are discussed by Lewis, Goodman and Miller (1969). In each simulation,  $nr$  independent observations on a standard normal variate were produced, grouped appropriately into  $n$  rows and  $r$  columns, and the value of  $X_r$  computed. For each  $r$  and  $n$ , 10,000 simulations were employed to compute the first and second moments of  $X_r$  from which  $a_r$  and  $\nu_r$  were estimated. The first series of simulations were done for  $r = 4, 5$  and  $6$ ,  $n = 10, 15, 20, 30$  and  $50$ . Results are given in Table I. Included also in Table I for comparison, are simulation results for  $r = 3$  and the statistic  $(n - 2)S/(n - 1) = (n - 2)X_3/(n - 1)$ ,  $S$  in (6), since this statistic is now known to have the chi-square distribution with 2 degrees of freedom.

The results shown in Table I are very interesting. It is apparent that  $\nu_r$  is very close to  $(r - 1)$ . Closer examination suggests that  $a_r$  is very close to  $2/(r - 1)(r - 2)$ . Accordingly, we suggest use of the statistic,

$$Y_r = - \frac{(n - 1)(r - 1)(r - 2)}{2} \log T_r, \quad (27)$$

TABLE I

Simulation Results to Determine  $a_r$  and  $v_r$ 

$r$	$n$	$a_r$	$v_r$
3	10	1.007	1.957
	15	1.039	1.922
	20	1.006	1.982
	30	0.954	2.075
	50	0.972	2.031
4	10	0.332	2.994
	15	0.330	3.027
	20	0.327	3.023
	30	0.341	2.921
	50	0.332	2.990
5	10	0.168	3.871
	15	0.164	3.924
	20	0.162	3.985
	30	0.163	3.993
	50	0.162	3.955
6	10	0.103	4.687
	15	0.099	4.903
	20	0.097	4.973
	30	0.098	4.946
	50	0.097	4.973

TABLE II

Simulated Significance Levels of  $Y_r$  Compared to Those of  $\chi^2_{r-1}$ 

r	n	Right-tail Significance Levels for $\chi^2_{r-1}$				
		0.01	0.05	0.10	0.20	0.50
3	10	0.010	0.048	0.098	0.199	0.502
	20	0.009	0.048	0.098	0.196	0.504
	30	0.011	0.049	0.098	0.200	0.491
	50	0.009	0.047	0.094	0.197	0.491
4	10	0.012	0.049	0.096	0.198	0.501
	20	0.011	0.048	0.098	0.200	0.502
	30	0.008	0.043	0.093	0.187	0.486
	50	0.007	0.046	0.098	0.197	0.489
5	10	0.012	0.048	0.093	0.184	0.486
	20	0.010	0.043	0.088	0.181	0.482
	30	0.011	0.047	0.095	0.191	0.487
	50	0.009	0.045	0.096	0.191	0.491
6	10	0.012	0.045	0.085	0.177	0.470
	20	0.009	0.045	0.088	0.180	0.474
	30	0.010	0.044	0.090	0.183	0.477
	50	0.008	0.042	0.085	0.177	0.477
8	20	0.009	0.045	0.086	0.175	0.459
	30	0.008	0.042	0.087	0.174	0.471
10	20	0.008	0.040	0.082	0.170	0.466
	30	0.009	0.040	0.084	0.177	0.470
12	20	0.009	0.042	0.082	0.164	0.450
	30	0.010	0.039	0.081	0.166	0.451

for the variance homogeneity test,  $Y_r$  to be taken to have the central chi-square distribution with  $(r - 1)$  degrees of freedom. Percentage points of the simulated distribution of  $Y_r$  are compared with those of chi square in Table II for  $r = 3, \dots, 6, 8, 10, \text{ and } 12$  for values of  $n$  shown.

The chi-square approximation to the distribution of  $Y_r$  seems remarkably good. In general, the approximation is very slightly conservative except for  $n = 10$  at the .01-level, but even here the agreement is very good. The approximate method proposed seems to effectively provide the desired test of variance homogeneity based on Grubbs' estimators for practical purposes.

## 6. BRIEF EXAMPLES

We examine the variance homogeneity test for two examples in the literature. The first example has  $r = 3$  and the exact test of Section 3.3 may be used. The second example has  $r = 4$  and the approximate method of Section 5 may be used. In both examples, the null hypothesis has

$$H_0: \sigma_j^2 = \sigma^2, j = 1, \dots, r, \text{ and the alternative hypothesis is}$$

$$H_a: \sigma_j^2 = \sigma_{j'}^2, \text{ for some } j \neq j'.$$

Russell and Bradley (1958) provided data on alcohol yields for three fermentors for  $n = 38$  days in a distillery. Day effects were judged to be important so Grubbs' estimators were used. They calculated  $Q_1 = 0.001537$ ,  $Q_2 = 0.001722$  and  $Q_3 = 0.000041$ . The test statistic  $T$  in (9) has the value 0.7659. From (20),  $P(T \leq T_0) = T_0^{(n-2)/2}$  and the P-value for the experiment is  $(0.7659)^{18} = 0.0082$ , indicating significantly different variances for the fermentors at the 0.01-level of significance.

Graybill (1954) gave the yields of  $r = 4$  varieties of wheat at  $n = 13$  locations in Iowa. It was suspected that error variances differed by varieties and that there were location effects. Calculation yields  $Q_1 = 875.40$ ,  $Q_2 = -84.92$ ,  $Q_3 = 451.47$  and  $Q_4 = 109.32$ . Now  $T_4 = 0.6109$  in (26) and  $Y_4$  in (27) has the value 17.74, highly significant when compared with significance levels of chi-square with 3 degrees of freedom. These data were considered also by Ellenberg (1977), Snee (1982), and others.

## 7. REMARKS

Small-sample distribution theory based on Grubbs' estimators appears to be very difficult in general, but surprisingly simple when  $r = 3$ . Statistical methods are most needed for smaller values of  $r$ . We have provided the necessary theory when  $r = 3$  and a good approximate test for variance homogeneity when  $r > 3$ . Some further investigation of the approximate test for small values of  $n$  may be desirable.

One warning should be issued. Tests on variances seem to be more sensitive to departures from the assumptions of normality than tests on means. This may be the case also for tests based on Grubbs' estimators and some investigation of the effects of nonnormality is suggested.

## ACKNOWLEDGMENTS

This work was supported by the Army, Navy and Air Force under Office of Naval Research Contract No. N00014-80-C-0093 with the Florida State University. Reproduction in whole or in part is permitted for any purpose of the United States Government. The major part of this research was done at the Florida State University, Tallahassee,

Florida 32306.

We are indebted also to Mead Johnson and Company, Evansville, Indiana for use of their IBM 3033 computer used in the simulation studies reported in Section 5.

#### REFERENCES

1. BRINDLEY, D. A. (1982), Some Results on the Distribution of Grubbs Estimators, Ph.D. Dissertation, Florida State University Library, Tallahassee, Florida.
2. EHRENBERG, A. S. C. (1950), "The Unbiased Estimation of Heterogeneous Error Variances," Biometrika, 37, 347-357.
3. ELLENBERG, J. H. (1977), "The Joint Distribution of Standardized Row Sums of Squares from a Balanced Two-way Layout," Journal of the American Statistical Association, 72, 407-411.
4. GRAYBILL, F. A. (1954), "Variance Heterogeneity in a Randomized Block Design," Biometrics, 10, 516-520.
5. GRUBBS, F. E. (1948), "On Estimating Precision of Measuring Instruments and Product Variability," Journal of the American Statistical Association, 72, 243-264.
6. JOHNSON, N. L. (1962), "Some Notes on the Investigation of Heterogeneity in Interactions," Revista Trabajos de Estadística, 13, 183-196.
7. LEWIS, P. A. W., GOODMAN, A. S. and MILLER, J. M. (1969), "A Pseudo-Random Number Generator for the System/360," I.B.M. Systems Journal, 8, 136-146.

8. MALONEY, C. J. (1973), "Grubbs' Estimators to Date," Proceedings of the Eighteenth Conference on the Design of Experiments in Army Research Development and Testing, Part 2, 523-547.
9. RUSSELL, T. S. and BRADLEY, R. A. (1958), "One-way Variances in a Two-way Classification," Biometrika, 45, 111-129.
10. SHUKLA, G. K. (1982). "Testing the Homogeneity of Variances in a Two-way Classification," Biometrika, 69, 411-416.
11. SNEE, R. D. (1982), "Nonadditivity in a Two-way Classification: Is It Interaction or Nonhomogeneous Variance?," Journal of the American Statistical Association, 77, 515-519.



Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1. REPORT NUMBER FSU Report No M648 ONR Report No 159	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and subtitle) Some New Results on Grubbs' Estimators	5. TYPE OF REPORT & PERIOD COVERED Technical Report	
7. AUTHOR(s) Dennis A. Brindley and Ralph A. Bradley	6. PERFORMING ORG. REPORT NUMBER FSU Report No. M648	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics Florida State University Tallahassee, FL 32306	8. CONTRACT OR GRANT NUMBER(s) ONR Contract No. N00014-80-C-0093	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Statistics and Probability Program Arlington, VA 22217	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	12. REPORT DATE June, 1983	
	13. NUMBER OF PAGES 19	
	15. SECURITY CLASS. (of this report) Unclassified	
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		

16. DISTRIBUTION STATEMENT (of this report)

Distribution unlimited.

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from report)

18. SUPPLEMENTARY NOTES

19. KEY WORDS

20. ABSTRACT (Continue on reverse side if necessary and identify by block number)

Consider a two-way classification with  $n$  rows and  $r$  columns and the usual model of analysis of variance except that the error components of the model may have heterogeneous variances  $\sigma_j^2$ ,  $j = 1, \dots, r$ , by columns. Grubbs provided unbiased estimators  $Q_j$  of  $\sigma_j^2$  that depend on column sums of squared residuals  $S_j$ . When  $r = 3$ , the joint distributions of the  $S_j$  and the  $Q_j$  are given for the first time in closed form.

Two tests proposed by Russell and Bradley are examined when  $r = 3$ , one for variance homogeneity and the second for one possible disparate variance. A very simple distribution is found for the test statistic of the first test and its non-null

## 20. Abstract (Continued)

distribution is derived also. The distribution of the second test statistic was known to be the central variance-ratio distribution in the null case and now its ratio to a parameter of noncentrality is shown to have the same distribution in the non-null case.

When  $r = 4$ ,  $n = 4$ , the joint distribution of the  $S_j$  is given also in closed form, but it is difficult to use. For  $r > 3$ , an approximate test of variance homogeneity is proposed, based on an extension of the Russell-Bradley statistic. Extensive simulation studies show that the distribution of the test statistic may be approximated very well by a chi-square distribution.

**END**

**FILMED**

**11-83**

**DTIC**