

AD-A134 121

REARRANGEMENT OF CLO50-
33-(PPH3)2-3-H-1-(R)-312-IRC2B9H10 TO CLO50-22-(P.. (U)
CALIFORNIA UNIV LOS ANGELES DEPT OF CHEMISTRY AND
BIOCHEMISTR .. J A DOI ET AL. 19 MAY 83 F/G 7/3


1/1

UNCLASSIFIED

NL



END
FILMED
1983



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

12

OFFICE OF NAVAL RESEARCH

Contract No. N00014-76-C-0390

Task No. NR 053-608

TECHNICAL REPORT NO. 123

Rearrangement of ~~Closo~~ 3,3-(PPh₃)₂-3-H-1-(R)-3,1,2-IrC₂B₉H₁₀ to
~~Closo~~-2,2-(PPh₃)₂-H-8-(R)-2,1,8-IrC₂B₉H₁₀. Synthesis and X-ray
Structure of ~~Closo~~-2,2-(PPh₃)₂-2-H-8-(C₆H₅)-2,1,8-IrC₂B₉H₁₀.

By

James A. Doi, Eugene A. Mizusawa, Carolyn B. Knobler, and M. Frederick Hawthorne*

AD-A134121

Prepared for Publication
in
Inorganic Chemistry

Department of Chemistry and Biochemistry
University of California
Los Angeles, California 90024

May 19, 1983

Reproduction in whole or part is permitted for
any purpose of the United States Government

Approved for Public Release; Distribution Unlimited

Accession For	
NTIS	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution _____	
Availability Codes	
_____ / or	
Dist _____	
A	



DTIC
ELECTE
OCT 25 1983

D

88 10 24 008

DTIC FILE COPY

Rearrangement of Closo-3,3-(PPh₃)₂-3-H-1-(R)-3,1,2-IrC₂B₉H₁₀
to Closo-2,2-(PPh₃)₂-2-H-8-(R)-2,1,8-IrC₂B₉H₁₀. Synthesis and X-ray Structure
of Closo-2,2-(PPh₃)₂-2-H-8-(C₆H₅)-2,1,8-IrC₂B₉H₁₀.

James A. Doi, Eugene A. Mizusawa, Carolyn B. Knobler, and M. Frederick
Hawthorne*

Department of Chemistry and Biochemistry
University of California, Los Angeles, California 90024

→ During ~~our~~ ^{the} continuing investigation of the chemistry of the known alkene
hydrogenation and isomerization catalyst,
closo-3,3-(PPh₃)₂-3-H-3,1,2-RhC₂B₉H₁₁¹ (Ia), the iridium congener and its
derivatives, closo-3,3-(PPh₃)₂-1-(R)-3-H-3,1,2-IrC₂B₉H₁₀, R = H (IIa)¹, R =
C₆H₅ (IIb)², R = CH₃ (IIc)² and R = 1'-(closo-1',2'-C₂B₁₀H₁₁) (IId)² were
prepared and notable differences between the reactivity of the Rh and Ir
systems were observed. Syntheses of IIb-d, the exo-nido complexes derived
from these compounds, and the dimeric complex derived from IIa will be
reported in a future paper. Although thermally induced migration of carbon
atoms over the surfaces of polyhedral cobaltacarboranes is a well-established
phenomenon,³ similar isomerization reactions are not so well known for other
metallacarboranes. The formation of 2,1,8 isomers of icosahedral
metalloacarboranes have been observed previously³ only during high temperature
vapor phase thermal isomerization of (η-C₅H₅)Co(C₂B₉H₁₁) isomers. The 2,1,8

isomer of this system was observed as the dominant kinetic product (74%) arising from the 3,1,2 isomer during such reactions at temperatures near 500°C; however, three other isomers were also detected in the reaction mixture. Here we report the ready polytopal rearrangement of **IIb** and **IIc** under mild thermolytic conditions in toluene solvent at the reflux temperature. In each of these examples the carbon atom bearing the substituent migrated in such a fashion as to produce the isomeric **closo-2,2-(PPh₃)₂-2-H-8-(C₆H₅)-2,1,8-IrC₂B₉H₁₀ (IIb)** and **closo-2,2-(PPh₃)₂-2-H-8-(CH₃)-2,1,8-IrC₂B₉H₁₀ (IIc)** species in 84% (30 h) and 33% (9 d) yields (heating time), respectively. None of the other possible isomers were produced in detectable quantities under the experimental conditions described. The rearranged species **IIId** was prepared by heating [(COD)Ir(PPh₃)₂]⁺ [nido-7-(1'-**closo-1',2'-C₂B₁₀H₁₁**)-7,8-C₂B₉H₁₁]⁻ in the presence of cyclohexane at the reflux temperature (41%). The unsubstituted **IIa** was not observed to rearrange after 5 d at reflux in toluene. The order of relative ease of rearrangement is apparently **IIId > IIb > IIc** in agreement with the idea that relief of steric strain provides some driving force for rearrangement by moving the bulky carbon substituent to the second belt (vertex 8) of the icosahedron, thus relieving interaction with the metal vertex and its triphenylphosphine ligands.

Below we describe the x-ray analysis of **IIb**, upon which the foregoing structural arguments are based.

The structure of **closo-2,2-(PPh₃)₂-H-8-(C₆H₅)-2,1,8-IrC₂B₉H₁₀** was determined by a single crystal diffraction study. The molecule is illustrated in the Figure⁴. Selected interatomic distances and angles are listed in Tables I and II. The crystal structure closely resembles that of the Rh

congener of Ia in its conformation about the metal. The most striking feature of the structure is the relative positions of the two carbon atoms of the C_2B_9 carborane ligand, which defines this complex as a member of the 2,1,8-isomer series⁴. The phenyl-bearing carbon vertex of the carborane has migrated to a position on the lower pentagonal belt, non-adjacent to the other carbon atom, resulting in a decrease in both electrostatic repulsions and steric interactions among the two carbon atoms and the Ir atom. The iridium atom, as expected, exhibits pseudo-octahedral coordination; the carborane ligand occupies three coordination sites and the two PPh_3 ligands and the hydride occupy the remaining sites. There is no fundamental distortion from the usual 12-vertex close icosahedral geometry.

Experimental Section

All manipulations were carried out in an inert-atmosphere with standard Schlenk techniques. Unless otherwise specified all solvents were purified by standard procedures and distilled under argon before use. Baker-analyzed silica gel (60-200 mesh) was used for column chromatography with reagent grade hexane and dichloromethane solvents. Infrared spectra were recorded as Nujol mineral oil mulls on a Perkin Elmer 521 dual grating spectrometer in the region 4000- 575 cm^{-1} . ^{31}P and 1H NMR spectra were obtained on a Bruker WP200 FT spectrometer. ^{11}B NMR spectra were recorded on a spectrometer built by Professor F. Anet of the Department of Chemistry, University of California, Los Angeles, CA.

Elemental analyses were by Schwartzkopf Microanalytical Laboratories, Woodside, NY. All melting points were determined on a Laboratory Devices Mel-Temp in open capillaries and are uncorrected.

Preparation of closo-2,2-(PPh₃)₂-2-H-8-(C₆H₅)-2,1,8-IrC₂B₉H₁₀ (IIIb). A 100 ml three neck flask was fitted with a reflux condenser and a gas inlet which in turn was connected to a vacuum/nitrogen manifold. The flask was stoppered, and the entire system was vacuum purged with nitrogen three times. To the flask was added 0.69 g (0.74 mmole) of IIb along with 35 ml benzene and a magnetic stir bar. The solution was heated to reflux and nitrogen gas was continually bubbled through the solution. After approximately 20 h of reflux, an additional 20 ml of toluene was added to the slurry of white solid and yellow solution. The slurry was refluxed an additional 10 h, cooled to room temperature and 0.58 g (0.62 mmole) of IIIb (84% yield) was collected. The complex was purified by column chromatography on silica gel and eluted with a 1:2 dichloromethane/hexane solvent mixture. The clear solution was reduced in volume on a rotary evaporator and yielded white microcrystals of IIIb.

Anal. Calcd for C₄₄H₄₆B₉P₂Ir: C, 57.07; H, 4.96; B, 10.51; P, 6.69; Ir, 20.75. Found: C, 56.90; H, 5.17; B, 10.24; P, 6.51; Ir, 20.95.

Infrared spectrum: 3060 (m), 2600 (s,sh) 2580 (vs), 2550 (vs), 2530 (s,sh), 2190 (m), 1490 (m,sh), 1480 (s), 1430 (s), 1180 (m), 1150 (m), 1085 (s), 1050 (m), 1010 (m), 990 (w), 925 (vw), 870 (vw), 815 (vw), 780 (w,sh), 760 (m), 740 (s,sh), 730 (s), 685 (vs) cm⁻¹.

NMR spectra (δ): ¹H (CDCl₃, room temperature) -10.6, triplet ²J_{P-H} = 26.6 Hz (terminal Ir-H), 0.0-3.0 (terminal B-H), 7.2, multiplet (phenyl); ¹¹B (127.048 MHz, ref. BF₃·Et₂O, CH₂Cl₂, room temperature) -0.1, 6.3, 9.4, 20.5; ³¹P{¹H} (10% C₆D₆ in THF, -60°C) 27.5, doublet, ²J_{P-P} = 19.5 Hz and 25.9, doublet, ²J_{P-P} = 20.7 Hz.

Preparation of $\text{closo-2,2-(PPh}_3)_2\text{-2-H-8-(CH}_3\text{)-2,1,8-IrC}_2\text{B}_9\text{H}_{10}$ (IIIc). The complex IIc (0.31 g, 0.36 mmole) in 40 ml of toluene was treated in the same manner as described for IIb for 9 days at reflux. After 2 additional days at room temperature, white microcrystals formed and were collected, yielding 0.067 g of IIIc. The filtrate was chromatographed, eluting with 1:2 dichloromethane-hexane. The clear second band gave an additional 0.0350 g (total 0.1016 g, 0.118 mmole, 33% yield) of product.

Anal. Calcd for $\text{C}_{39}\text{H}_{44}\text{B}_9\text{P}_2\text{Ir}$: C, 54.22; H, 5.09; B, 11.26; P, 7.17; Ir, 22.25. Found: C, 53.68; H, 5.18; B, 10.30; P, 7.06; Ir, 21.76.

Infrared spectrum: 3060 (m), 2620 (w), 2560 (vs,br), 2190 (w), 1480 (s), 1430 (s), 1310 (w), 1180 (m), 1150 (m), 1085 (s), 1023 (m), 1005 (m), 995 (m), 980 (m), 930 (vw), 910 (vw), 885 (vw), 760 (m), 750 (s), 735 (s), 690 (vs) cm^{-1} .

NMR spectra (δ): ^1H (CDCl_3 , -60°C) -10.5, triplet, $^2\text{J}_{\text{P-H}} = 25.5$ Hz (terminal Ir-H), 0.0-3.0 (terminal B-H), 1.2, singlet (methyl), 7.2, multiplet, (phenyl); ^{11}B (127.048 MHz, ref. $\text{BF}_3 \cdot \text{Et}_2\text{O}$, CH_2Cl_2 , room temperature) -1.8, 7.8, 9.9, 20.7; $^{31}\text{P}\{^1\text{H}\}$ (ref. D_3PO_4 , CDCl_3 , room temperature) 12.9, singlet.

Preparation of

$\text{closo-2,2-(PPh}_3)_2\text{-1-H-8-(1'-closo-1',2'-C}_2\text{B}_{10}\text{H}_{11}\text{)-2,1,8-IrC}_2\text{B}_9\text{H}_{10}$ (IIIId). The salt $[\text{CODIr(PPh}_3)_2]^+ [\text{nido-7-(1'-closo-1',2'-C}_2\text{B}_{10}\text{H}_{11}\text{)-7,8-C}_2\text{B}_9\text{H}_{11}]^-$ (0.30 g, 0.46 mmole) in 35 ml of cyclohexane was heated to reflux. After 24 h the slurry became a clear red solution. After 48 h a white solid was formed. Heating was continued for an additional 4 d. The slurry was cooled to room temperature and the white solid was filtered. The product IIIId (0.19 g, 41%) was collected.

Anal. Calcd for $\text{C}_{40}\text{H}_{52}\text{B}_{19}\text{P}_2\text{Ir}$: C, 48.43; H, 5.24; B, 20.70; P, 6.24; Ir,

19.37. Found: C, 48.43; H, 5.61; B, 20.14; P, 5.61; Ir, 20.00.

Infrared spectrum: 3060 (m), 2580 (vs, br), 2250 (w), 1480 (m, sh), 1475 (s, sh), 1430 (s), 1180 (m), 1150 (w), 1085 (s), 1065 (w), 1040 (w), 1020 (m), 1005 (m), 995 (m), 930 (vw), 845 (vw), 780 (w), 750 (s), 740 (s), 685 (vs) cm^{-1} .

NMR spectra (δ): ^1H (CDCl_3 , room temperature) -10.7, triplet, $^2J_{\text{P-H}} = 25.6$ Hz (terminal Ir-H), 0.0-3.0 (terminal B-H), 7.2, multiplet (phenyl); ^{11}B (111.9 MHz, ref. $\text{BF}_3 \cdot \text{Et}_2\text{O}$, CH_2Cl_2 , room temperature), 0.2, 4.5, 5.1, 11.7, 14.4, 21.4; $^{31}\text{P}\{^1\text{H}\}$ (ref. D_3PO_4 , CDCl_3 , room temperature) 12.0, doublet, $^2J_{\text{P-P}} = 21.9$ Hz and 10.7, doublet, $^2J_{\text{P-P}} = 20.8$ Hz.

Thermolysis of $\text{closo-3,3-(PPh}_3)_2\text{-3-H-3,1,2-IrC}_2\text{B}_9\text{H}_{11}$ (Ia). Ia (0.13 g, 0.15 mmole) was placed in a 25 ml 3 neck round bottom flask fitted with a water-cooled spiral condenser which was connected to a vacuum/nitrogen manifold through a gas inlet. The entire system was stoppered and vacuum purged with nitrogen three times. A magnetic stir bar and 15 ml of toluene were placed in the flask and the sample was heated to reflux. After 5 d the solution was cooled to room temperature. Only starting material was found. Constant monitoring by thin layer chromatography of the solution indicated no reaction.

$\text{Closo-2,2-(PPh}_3)_2\text{-2-H-8-(C}_6\text{H}_5\text{)-2,1,8-IrC}_2\text{B}_9\text{H}_{10}$ $1/2 \text{ CH}_2\text{Cl}_2$ was prepared as described above and crystals were grown by vapor diffusion using 1:1 chloroform/dichloromethane and hexane. A single crystal was mounted on a glass fiber. The automatic centering, autoindexing, and least-squares routines of a Syntex P2₁ diffractometer were used to determine the unit cell parameters of $a = 11.722(4)$, $b = 19.994(7)$, $c = 9.923(4)$ Å, $\alpha =$

100.17(3), $\beta = 107.59(3)$, $\gamma = 86.40(3)^\circ$ and $V = 2182(1) \text{ \AA}^3$ on the basis of 15 reflections. Some details of the data collection are presented in Table III. A graphite crystal was used to provide monochromatic $\text{MoK}_{\alpha 1}$ radiation (0.7107 \AA). The crystal density was found to be 1.48 g cm^{-3} (by flotation in $\text{CCl}_4/\text{heptane}$), while the calculated density was 1.47 g cm^{-3} based on $Z = 2$.

Intensity data ($h, \pm k, \pm l$) were collected with the theta-2theta scan technique to a limit of $2\theta = 45^\circ$. Reflections were scanned at a variable rate (between 2.5 and $14.65^\circ/\text{min}$) from 1° below the $\text{MoK}_{\alpha 1}$ to 1° above the $\text{MoK}_{\alpha 2}$ reflection. Intensities of three standard reflections were measured after every 50 reflections. No significant deviations were observed.

The data were corrected for Lorentz and polarization effects. The intensity of a reflection $I(hkl)$ and its estimated standard deviation $\sigma I(hkl)$ were calculated as described previously.⁵ Of 5778 unique reflections, 5025 with $I > 3\sigma(I)$ were considered to be observed and were included in subsequent calculations.

Solution and Refinement of the Structure. The coordinates of the Ir atom were determined by solution of a 3-dimensional Patterson map and the structure was solved in space group $P\bar{1}$ by heavy atom techniques. The 7 phenyl groups were described as rigid C_6 hexagons with $\text{C-C} = 1.391 \text{ \AA}$ and $\text{C-H} = 0.998 \text{ \AA}$. Positions for all hydrogen atoms not included as members of rigid groups were obtained from a difference map. In the final least-squares cycle 268 parameters were refined, including positional and anisotropic vibrational parameters for Ir, P, C_2B_9 , and CCl_2 (at half occupancy), positional and carbon isotropic vibrational parameters for the C_6H_5 groups and positional

parameters for the remaining nongroup H atoms, with the exception of the terminal Ir-H hydride H(2). For each H atom, B was fixed at 0.5 + B of the C or B atom to which that H is attached. Atomic positional parameters for atoms not included as members of rigid groups are given in Table IV. Scattering factors and anomalous dispersion terms were taken from International Tables for X-ray Crystallography⁶.

The final least-squares cycle converged at $R = 0.050$, $R_w = 0.064$ (refined by full-matrix least-squares procedure and based on F , $w = 1/\sigma^2(F_o)$; $R = \sum ||F_o| - |F_c|| / \sum |F_o|$; $R_w = [\sum w ||F_o| - |F_c||^2 / \sum w |F_o|^2]^{1/2}$; the "goodness of fit" was 2.30 as defined by $[\sum w (|F_o| - |F_c|)^2 / (N_o - N_v)]^{1/2}$ with $N_o = 5025$ and $N_v = 268$. In the final cycle of least-squares refinement no shift for a nonhydrogen nonsolvate atom was larger than 0.1 of its corresponding estimated standard deviation. All calculations were performed on a UCLA Departmental DEC VAX 11/780 using the UCLA Crystallographic Package (locally edited versions of CARESS, PROFILE, ORFLS, ORFFE, and ORTEP).

Acknowledgements

We thank the National Science Foundation and the Office of Naval Research for partial support of this research. We also thank Prof. R. Bau for use of the Syntex P2₁ diffractometer.

Supplementary Material Available: Listings of observed and calculated structure factor amplitudes, hydrogen positional parameters, group positional parameters and anisotropic thermal parameters of nonhydrogen atoms (pages). Ordering information is given on any current masthead page.

References

- 1a. Paxson, T.E.; Hawthorne, M.F. J. Amer. Chem. Soc., 1974, 96, 4674.
- 1b. Paxson, T.E. Ph.D. Thesis, University of California, Los Angeles, 1974.
2. Doi, J.A. Ph.D. Thesis, University of California, Los Angeles, 1980.
3. Kaloustian, M.K.; Wiersema, R.J.; Hawthorne, M. F. J. Amer. Chem. Soc., 1972, 94, 6679.
4. The cage numbering system in this paper is consistent with that favored by I.U.P.A.C. for close carboranes. Commission on Nomenclature of Inorganic Chemistry, Pure Appl. Chem., 1972, 30, 683.
5. Delaney, M.S.; Knobler, C.B; Hawthorne, M.F. Inorg. Chem., 1981, 20, 1341.
6. International Tables for X-ray Crystallography, Vol. IV. 1974, Birmingham, Kynoch Press.

Figure Caption

The molecular structure of $\text{closo-2,2-(PPh}_3)_2\text{-2-H-8-(C}_6\text{H}_5\text{)-2,1,8-Ir-C}_2\text{B}_9\text{H}_{10}$
 $1/2 \text{ CH}_2\text{Cl}_2$.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 123	2. GOVT ACCESSION NO. A134121	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) "Rearrangement of $\text{ClosO-3,3-(PPh}_3)_2\text{-3-H-1-(R)-3,1,2-IrC}_2\text{B}_9\text{H}_{10}$ to $\text{ClosO-2,2-(PPh}_3)_2\text{-2-H-8-(R)-2,1,8-IrC}_2\text{B}_9\text{H}_{10}$. Synthesis and X-ray Structure of $\text{ClosO-2,2-(PPh}_3)_2\text{-2-H-8-(C}_6\text{H}_5)_2\text{-2,1,8-IrC}_2\text{B}_9\text{H}_{10}$."		5. TYPE OF REPORT & PERIOD COVERED Interim
7. AUTHOR(s) James A. Doi, Eugene A. Mizusawa, Carolyn B. Knobler, and M. Frederick Hawthorne*		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS The University of California Department of Chemistry and Biochemistry Los Angeles, California 90024		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0390
11. CONTROLLING OFFICE NAME AND ADDRESS Chemistry Branch Office of Navy Research Washington, D.C. 20360		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-608
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE May 19, 1983
		13. NUMBER OF PAGES 11
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) <div style="border: 1px solid black; padding: 5px; text-align: center;">DISTRIBUTION STATEMENT A Approved for public release; Distribution Unlimited</div>		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Synthesis & structure Irida carborane Rearrangement of-		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		

APPENDIX

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
	1	Naval Weapons Center Attn: Dr. A.B. Amster, Chemistry Division China Lake, California 93555	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R.W. Drisko Port Hueneme, California 93401	1
ONR Branch Office Attn: Dr. L.H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A.L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (R,E&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division Naval Research Laboratory Washington, D.C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, GEN

No.
Copies

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco 96503

1

Mr. James Kelley
DTNSRDC Code 2803
Annapolis, Maryland 21402

1

TECHNICAL REPORT DISTRIBUTION LIST, 053

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. R.N. Grimes Department of Chemistry University of Virginia Charlottesville, Virginia 22901	1	Dr. M.H. Chisholm Department of Chemistry Indiana University Bloomington, Indiana 47401	1
Dr. D.B. Brown Department of Chemistry University of Vermont Burlington, Vermont 05401	1	Dr. B. Foxman Department of Chemistry Brandeis University Waltham, Massachusetts 02154	1
Dr. W.B. Fox Chemistry Division Naval Research Laboratory Code 6130 Washington, D.C. 20375	1	Dr. T. Marks Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. J. Adcock Department of Chemistry University of Tennessee Knoxville, Tennessee 37916	1	Dr. G. Geoffrey Department of Chemistry Pennsylvania State University University Park, Pennsylvania 16802	1
Dr. A. Cowley Department of Chemistry University of Texas Austin, Texas 78712	1	Dr. J. Zuckerman Department of Chemistry University of Oklahoma Norman, Oklahoma 73019	1
Dr. W. Hatfield Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514	1	Professor O.T. Beachley Department of Chemistry State University of New York Buffalo, New York 14214	1
Dr. D. Seyferth Department of Chemistry Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Professor P.S. Skell Department of Chemistry The Pennsylvania State University University Park, Pennsylvania 16802	1
Professor H. Abrahamson Department of Chemistry University of Oklahoma Norman, Oklahoma 73019	1	Professor K.M. Nicholas Department of Chemistry Boston College Chestnut Hill, Massachusetts 02167	1
Professor M. Newcomb Texas A&M University Department of Chemistry College Station, Texas 77843	1	Professor R. Neilson Department of Chemistry Texas Christian University Fort Worth, Texas 76129	1
		Professor Richard Eisenberg Department of Chemistry University of Rochester Rochester, New York 14627	1

END

FILMED

11-83

DTIC