

AD-A134 538


DIFFUSION ON VISCOUS FLUIDS EXISTENCE AND ASYMPTOTIC
PROPERTIES OF SOLUTIONS (U) WISCONSIN UNIV-MADISON
MATHEMATICS RESEARCH CENTER H BEIRAO-DA-VEIGA SEP 83
MRC-TSR-2565 DAAG29-80-C-0041


1/1

UNCLASSIFIED

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

A134538

(2)

MRC Technical Summary Report #2565

DIFFUSION ON VISCOUS FLUIDS, EXISTENCE
AND ASYMPTOTIC PROPERTIES OF SOLUTIONS

H. Beirão-da-Veiga

**Mathematics Research Center
University of Wisconsin—Madison
610 Walnut Street
Madison, Wisconsin 53705**

September 1983


(Received December 7, 1982)

**Approved for public release
Distribution unlimited**

DTIC FILE COPY

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709



39 11 00 150

UNIVERSITY OF WISCONSIN-MAIDISON
MATHEMATICS RESEARCH CENTER



DIFFUSION ON VISCOUS FLUIDS, EXISTENCE
AND ASYMPTOTIC PROPERTIES OF SOLUTIONS

H. Beirao-da-Veiga*

Technical Summary Report #2565

September 1983

ABSTRACT

We consider the motion of a mixture of two fluids, with a diffusion effect obeying Fick's law, for the derivation of the model (1.1) see Section 1 and references [2], [4], [5] and [6]. We consider the full non-linear problem (i.e., we don't omit the term λ^2 term in equation (1.1)). Moreover, we don't assume that λ/μ is small. We prove the existence of a (unique) local solution, the existence of a global solution for small data, and the exponential decay to the equilibrium solution; see Theorem A, Section 1.

AMS(MOS) Subject Classifications: 35K55, 76D99, 76R99.

Key Words: Non-homogeneous viscous fluids, non-linear parabolic equations, initial-boundary value problems

Work Unit Number 1 (Applied Analysis)

*Department of Mathematics, University of Trento, 38050 Povo (Trento) Italy.

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.

SIGNIFICANCE AND EXPLANATION

In this paper we consider the motion of a continuous medium consisting of two components, say water and a dissolved salt, with a diffusion effect obeying Fick's law. We denote by $v, w, \rho, \pi, \mu, \lambda$ the mean-volume velocity, mean-mass velocity, density, pressure, viscosity and diffusion constant, respectively. By using Fick's law we eliminate w from the equations and we obtain (1.1), where p is the modified pressure; see Section 1 and references [2], [4], [5] and [6]. The initial boundary conditions are given by equation (1.2).

Kazhikhov and Smagulov [5], [6] consider equation (1.1) for a small diffusion coefficient λ . More precisely, they assume that condition (1.3) holds; moreover, they omit the λ^2 term in equation (1.1)₁. Under these conditions they prove the existence of a unique local solution for the 3-dimensional motion (in the bi-dimensional case, solutions are global).

In our paper we consider the full equation (1.1), without assumption (1.3), and we prove: (i) the existence of a (unique) local solution; (ii) the existence of a global solution in time for small initial velocities and external forces, and for initial densities near-constant; (iii) the exponential decay (when $t \rightarrow +\infty$) of the solution (ρ, v) to the equilibrium solution $(\hat{\rho}, 0)$, if $f \equiv 0$. See Theorem A, Section 1.

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

DIFFUSION ON VISCOUS FLUIDS, EXISTENCE
AND ASYMPTOTIC PROPERTIES OF SOLUTIONS

H. Beirão-da-Veiga*

Main Notation

Ω : an open bounded set in \mathbb{R}^3 , locally situated on one side of its boundary Γ , a regular (say C^4) manifold.

$n = n(x)$: unit outward normal to Γ .

D_i, D_{ij}, D_t : $\partial/\partial x_i, \partial^2/\partial x_i \partial x_j, \partial/\partial t$.

$\|\cdot\|, (\cdot, \cdot)$: norm and scalar product in $L^2(\Omega)$.

H^k : Sobolev space $H^{k,2}(\Omega)$ with norm

$$\|\sigma\|_k^2 \equiv \sum_{|\alpha|=0}^k \|D^\alpha \sigma\|^2,$$

where

$$\|D^\alpha \sigma\|^2 \equiv \sum_{|\alpha|=1} \|D^\alpha \sigma\|^2,$$

Further,

$$\|D^\alpha \sigma\|_m^2 \equiv \sum_{|\alpha|=1} \|D^\alpha \sigma\|_m^2.$$

H_0^1 : Closure of $C_0^\infty(\Omega)$ in $H^1(\Omega)$.

$\|\cdot\|_\infty$: norm in $L^\infty(\Omega)$.

L^2, H^k, H_0^1 : Hilbert spaces of vectors $v = (v_1, v_2, v_3)$ such that

$v_i \in L^2, v_i \in H^k, v_i \in H_0^1$ ($i=1,2,3$), respectively. Corresponding

notation is used for other spaces of vector fields. Norms are defined

in the natural way, and denoted by the symbols used for the scalar

fields.

Let us introduce the following functional spaces (see, for instance, [7], [8] and [12] for their properties) =

*Department of Mathematics, University of Trento, 38050 Povo (Trento) Italy.

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.

$$H_N^k \equiv \{ \sigma \in H^k : \frac{\partial \sigma}{\partial n} = 0 \text{ on } \Gamma \text{ and } \int_{\Omega} \sigma(x) dx = 0 \}, k > 2.$$

$$V \equiv \{ v \in C_0^\infty(\Omega) : \operatorname{div} v = 0 \text{ in } \Omega \},$$

$$H = \{ v \in L^2 : \operatorname{div} v = 0 \text{ in } \Omega, v \cdot n = 0 \text{ on } \Gamma \},$$

$$V = \{ v \in H_0^1 : \operatorname{div} v = 0 \text{ on } \Omega \}.$$

H and V are the closure of v in $L^2(\Omega)$ and H_0^1 , respectively. Moreover $L^2 = H + G$, where $G \equiv \{ \nabla p : p \in H^1(\Omega) \}$. Denoting by P the orthogonal projection of L^2 onto H , we define the operator $A \equiv -PA$ on $D(A) \equiv H^2 \cap V$. One has

$$(Au, v) = ((u, v)) \equiv \sum_{i,j} (D_i u_j, D_i v_j), \forall u \in D(A), v \in V.$$

The norms $\| \cdot \|_2, \| \Delta \cdot \|$ are equivalent in $H_N^2, \| \cdot \|_3, \| \nabla \Delta \cdot \|$ are equivalent in H_N^3 and $\| v \|_2, \| \Delta v \|$ are equivalent in $D(A)$. We define $\| v \|_V^2 \equiv ((v, v))$; the norms $\| v \|_V, \| v \|_1$ are equivalent in V .

$L^2(0, T; X)$: Banach space of strongly measurable functions defined in $]0, T[$ with values in (a Banach space) X , for which

$$\| z \|_{L^2(0, T; X)}^2 \equiv \int_0^T \| z(t) \|_X^2 dt < +\infty.$$

$C(0, T; X)$: Banach space of X -vector valued continuous functions on $[0, T]$ endowed with the usual norm $\| z \|_{C(0, T; X)}$.

μ : viscosity (a positive constant).

λ : diffusion coefficient (a positive constant).

$v(t, x), v_0(x)$: mean-volume velocity. Initial mean-volume velocity.

$\rho(t, x), \rho_0(x)$: density of the mixture. Initial density.

Further,

$$m \equiv \inf_{x \in \Omega} \rho_0(x), \quad M \equiv \sup_{x \in \Omega} \rho_0(x),$$

$$\hat{\rho} \equiv \frac{1}{|\Omega|} \int_{\Omega} \rho_0(x) dx.$$

We assume that $m > 0$.

$\pi(t, x), p(t, x)$: pressure. Modified pressure

$$p = \pi + \lambda v \cdot \nabla p - \lambda^2 \Delta p + \lambda(2\mu + \mu') \Delta \log \rho.$$

$f(t, x)$: external mass-force.

We denote by $c, \bar{c}, c_0, c_1, c_2, \dots$ positive constant, depending at most on Ω and on the parameters μ, λ, m, M and $\hat{\rho}$. It is easy to derive at any stage of the proofs, the explicit dependence of the constants on the parameters.

For convenience we sometimes denote different constants by the same symbol c . Otherwise, we utilize the symbols $\bar{c}, c_k, k \in \mathbb{N}$.

1. Main results. In this paper we consider the motion of a viscous fluid consisting of two components, say, saturated salt water and water. The equations of the model are obtained, for example, in [2], [4], [5], and [6]. Let us give a brief sketch. Let ρ_1, ρ_2 be the characteristic densities (constants) of the two components, $v^{(1)}(t,x)$ and $v^{(2)}(t,x)$ their velocities, and $e(t,x), d(t,x)$ the mass and volume concentration of the first fluid. We define the density $\rho(t,x) \equiv d\rho_1 + (1-d)\rho_2$, and the mean-volume and mean-mass velocities $v \equiv d v^{(1)} + (1-d) v^{(2)}, w \equiv e v^{(1)} + (1-e)v^{(2)}$. Then the equations of motion are

$$\begin{cases} \rho[D_t w + (w \cdot \nabla)w - f] - \mu \Delta w - (\mu + \mu') \nabla \operatorname{div} w = -\nabla \pi, \\ \operatorname{div} v = 0, \\ D_t \rho + \operatorname{div}(\rho w) = 0. \end{cases}$$

On the other hand, Fick's diffusion law (see [2]) gives $w = v - \lambda \rho^{-1} \nabla \rho$. By elimination of w in the preceding equations one gets, after some calculations,

$$(1.1) \quad \begin{cases} \rho(D_t v + (v \cdot \nabla)v) - \mu \Delta v - \lambda [(v \cdot \nabla) \nabla \rho + (\nabla \rho \cdot \nabla)v] + \\ + \frac{\lambda^2}{\rho} [(\nabla \rho \cdot \nabla) \nabla \rho - \frac{1}{\rho} (\nabla \rho \cdot \nabla \rho) \nabla \rho + \Delta \rho \nabla \rho] = -\nabla p + \rho f, \\ D_t \rho + v \nabla \rho - \lambda \Delta \rho = 0, \\ \operatorname{div} v = 0. \end{cases}$$

We want to solve system (1.1) in $\Omega_T \equiv]0, T[\times \Omega$. Here p is the modified pressure. We add to system (1.1) the following initial boundary-value conditions.

$$(1.2) \quad \begin{cases} v = 0 & \text{on }]0, T[\times \Gamma, \\ \frac{\partial \rho}{\partial n} = 0 & \text{on }]0, T[\times \Gamma, \\ v|_{t=0} = v_0(x) & \text{in } \Omega, \\ \rho|_{t=0} = \rho_0(x) & \text{in } \Omega. \end{cases}$$

The first two conditions mean that there is no flux through the boundary.

In [5], [6] Kazhikhov and Smagulov consider the simplified system obtained from (1.1) by omitting the term containing λ^2 ; moreover, they assume that

$$(1.3) \quad \lambda < \frac{2\mu}{M-m}.$$

Under these conditions Kazhikhov and Smagulov state the existence of a local solution in time (global in the bidimensional case).

In our paper we take into account the full equation (1.1), and omit the condition (1.3). For this more general case we prove: (i) the existence of a (unique) local solution for arbitrary initial data and external force field; (ii) the existence of a unique global strong solution, for small initial data and external force field. Moreover, if $f \equiv 0$, the solution (ρ, v) decays exponentially to the equilibrium solution $(\hat{\rho}, 0)$. More precisely we prove the following result:

Theorem A. Let $v_0 \in \mathbb{V}, \rho_0 - \hat{\rho} \in H_N^2, f \in L^2(0, T; L^2)$. Then there exists $T_1 \in]0, T[$ such that problem (1.1), (1.2) is uniquely solveable in Q_{T_1} . Moreover $v \in L^2(0, T_1; \mathbb{H}^2) \cap C(0, T_1; \mathbb{V}), D_t v \in L^2(0, T_1; \mathbb{H}), \rho - \hat{\rho} \in L^2(0, T_1; H_N^3) \cap C(0, T_1; H_N^2), D_t \rho \in L^2(0, T_1; H^1)$ and $m < \rho(t, x) < M$.

Moreover, there exist positive constants k_1, k_2 , and k_3 depending at most on Ω, ν, λ and on the mean density $\hat{\rho}^{(1)}$ such that that if

$$(1.4) \quad \|v_0\|_1 + \|\rho_0 - \hat{\rho}\|_2 < k_1,$$

(1) Or, equivalently, depending on the total amount of mass $|\Omega| \hat{\rho} = \int_{\Omega} \rho_0(x) dx$.

and

(1.5) $\|f\|_{L^{\infty}(0,+\infty;L^2)} < k_2$,
 then the solution is global in time. If $f \equiv 0$, the solution (ρ, v) decays exponentially to the equilibrium solution $(\hat{\rho}, 0)$, i.e.

$$(1.6) \quad \|v(t)\|_1 + \|\rho(t) - \hat{\rho}\|_2 < (\|v_0\|_1 + \|\rho_0 - \hat{\rho}\|_2) e^{-k_3 t},$$

for every $t > 0$.

Theorem A also holds for coefficients μ, λ regularly dependent on ρ, v , provided they are strictly positive and bounded, in a neighborhood of the range of values of the initial data $\rho_0(x), v_0(x)$. This generalization can be done without any difficulty. Moreover, with standard techniques, one can prove that the solutions have more regularity (up to C^m) if the data are sufficiently regular and the usual compatibility conditions hold.

Local existence in the general case (i.e. with the λ^2 term and without (1.3)) was proved in the inviscid case by Beirão-da-Veiga, Serapioni and Valli in [1]. A similar result, in the viscous case and for $\Omega = \mathbb{R}^3$, was proved by Secchi [11]. For another approach (concerning Greffi's model) see [10].

2. The linearized equations. We start by proving the following theorem: Theorem 2.1 Let $\rho(t, x)$ be a measurable function verifying

$$(2.1) \quad 0 < m < \rho(t, x) < M, \quad \text{a.e., in } Q_T, \quad \text{let } F \in L^2(0, T; \mathbb{R}) \quad \text{and } v_0 \in V.$$

Then there exists a (unique) strong solution v of problem

$$(2.2) \quad \begin{cases} \rho D_t v - \mu \Delta v = -\nabla p + F & \text{in } Q_T, \\ \operatorname{div} v = 0 & \text{in } Q_T, \\ v = 0 & \text{on }]0, T[\times \Gamma, \\ v|_{t=0} = v_0(x) & \text{in } \Omega. \end{cases}$$

Moreover $v \in L^2(0, T; D(A)) \cap C(0, T; V)$, $D_t v \in L^2(0, T; \mathbb{R})$ and

$$(2.3) \quad \begin{aligned} & \mu \|v\|_{C(0, T; V)}^2 + m \|v'\|_{L^2(0, T; H)}^2 + \\ & + \frac{m\mu}{4M^2} \|\Delta v\|_{L^2(0, T; H)}^2 < \mu \|v_0\|_V^2 + \left(\frac{2}{m} + \frac{m}{2M^2}\right) \|F\|_{L^2(0, T; \mathbb{R})}^2 \end{aligned}$$

Proof. Let us write equation (2.2) in the equivalent form (2.4)

$$P(\rho D_t v) + \mu Av = F, \quad v|_{t=0} = v_0(X).$$

For brevity let us put

$$X = \{v: v \in L^2(0, T; D(A)), v' \in L^2(0, T; H)\}.$$

From well known results (see [9], Vol I, Chapter I : Theorem 3.1 with $y = H$, $X = D(A)$, $j = 0$; and (2.42) Proposition 2.1) it follows that $X \subset C(0, T; V)$.

We start by proving the a priori bound (2.3); an essential device is to introduce a parameter ϵ_0 in order to conveniently balance the estimates. In H take the inner product of (2.4) with $D_t v + \epsilon_0 A v$, $\epsilon_0 > 0$. Since $(v', Av) = 2^{-1} D_t \|v\|_V^2$ one gets

$$(2.5) \quad m \|D_t v\|^2 + \frac{\mu}{2} \frac{d}{dt} \|v\|_V^2 + \epsilon_0 \mu \|Av\|^2 < \\ < \|F\| \|D_t v\| + \epsilon_0 \|F\| \|Av\| + \epsilon_0 M \|D_t v\| \|Av\|.$$

By using the inequalities $\|F\| \|D_t v\| < 4^{-1} m \|D_t v\|^2 + m^{-1} \|F\|^2$, $\|F\| \|Av\| < 4^{-1} \mu \|Av\|^2 + \mu^{-1} \|F\|^2$ and $\|D_t v\| \|Av\| < (4M)^{-1} \mu \|Av\|^2 + \mu^{-1} M \|D_t v\|^2$ one gets

$$(2.6) \quad \frac{3}{4} m \|D_t v\|^2 + \frac{\mu}{2} \frac{d}{dt} \|v\|_V^2 + \epsilon_0 \frac{\mu}{2} \|Av\|^2 < \\ < \left(\frac{1}{m} + \frac{\epsilon_0}{\mu}\right) \|F\|^2 + \frac{\epsilon_0 M^2}{\mu} \|D_t v\|^2.$$

Now fix $\epsilon_0 = (4M^2)^{-1} m \mu$ and integrate equation (2.6) on $(0, T)$. This gives the a priori bound (2.3).

Define $\|v\|_X^2 \equiv$ "left hand side of equation (2.3)", $Y \equiv L^2(0, T; H) \times V$, and $\|(F, v_0)\|_Y^2 \equiv$ "right hand side of (2.3)".

We solve (2.4) by the continuity method. Define $\rho_\alpha \equiv (1-\alpha) \hat{\rho} + \alpha \rho$, $\alpha \in [0, 1]$. Clearly ρ_α verifies condition (2.1), for any α . Define $T_\alpha \equiv (1-\alpha)\bar{T} + \alpha T$, where

$$\begin{aligned} \bar{T}_v &\equiv (P(\rho D_t v) - \mu Av, v|_{t=0}) \in Y, \\ T_v &\equiv (P(\bar{\rho} D_t v) - \mu Av, v|_{t=0}) \in Y. \end{aligned}$$

Finally denote by γ the set of values $\alpha \in [0,1]$ for which problem (2.2) is solvable in X for every pair $(F, v_0) \in Y$. Clearly $0 \in \gamma$, because for this value of the parameter equation (2.6) because the linearized Navier-Stokes equation. Let us verify that γ is open and closed.

γ is open. Let $\alpha_0 \in \gamma$ and denote by $G(F, v_0) \equiv v$ the solution v of problem $T_{\alpha_0} v = (F, v_0)$. From (2.3) one gets $G \in L(Y; X)$ ⁽²⁾ with $\|G\|_{Y, X} < 1$. Equation $T_{\alpha_0 + \varepsilon} v = (F, v_0)$ can be written in the form

$$(2.7) \quad [1 - \varepsilon G(\bar{T} - T)]v = G(F, v_0).$$

Since $\|G(\bar{T} - T)\|_{X, Y} < \|\bar{T} - T\|_{X, Y}$, equation (2.7) is solvable for $|\varepsilon| < \|\bar{T} - T\|_{X, Y}^{-1}$ (by a Neuman expansion).

γ is closed. Let $\alpha_n \in \gamma$, $\alpha_n \rightarrow \alpha_0$, and let v_n be the solution of $T_{\alpha_n} v_n = (F, v_0)$. From (2.3) one has $\|v_n\|_X < \|(F, v_0)\|_Y$. Since X is an Hilbert space there exists a subsequence $v_{n_j} \rightarrow v \in X$, weakly in X . From $T, \bar{T} \in L(X; Y)$ one has $\bar{T}v_{n_j} \rightarrow \bar{T}v$, $Tv_{n_j} \rightarrow Tv$ weakly in Y . Hence $T_{\alpha_0} v = (F, v_0)$, i.e. $T_{\alpha_0} v = (F, v_0)$.

Let us now return to problem (1.1). Define

$$(2.8) \quad F(\rho, v) \equiv P \{-\rho(v \cdot \nabla)v + \lambda [(v \cdot \nabla)\nabla\rho + (\nabla\rho \cdot \nabla)v] + \frac{\lambda^2}{\rho} [(\nabla\rho \cdot \nabla)\nabla\rho - \frac{1}{\rho} (\nabla\rho \cdot \nabla\rho)\nabla\rho + \Delta\rho\nabla\rho] + \rho f\}.$$

For convenience we will use in the sequel the translation

$$(2.9) \quad \rho = \hat{\rho} + \sigma.$$

Recall that $\hat{\rho}$ is a given constant. To solve problem (1.1) and (1.2) in our functional framework is equivalent to finding $v \in L^2(0, T; D(A))$, $v_t \in L^2(0, T; H)$ and $\sigma \in L^2(0, T; H_N^3)$, $\sigma_t \in L^2(0, T; H^1)$ such that

$$(2.10) \quad \begin{cases} P(\rho D_t v) - \mu \Delta v = F(\rho, v), \\ v|_{t=0} = v_0(x), \\ D_t \sigma - \lambda \nabla \sigma = -v \cdot \nabla \sigma, \\ \sigma|_{t=0} = \sigma_0(x), \end{cases}$$

⁽²⁾ Banach space of linear continuous operator from Y into X , with norm $\| \cdot \|_{Y, X}$.

where $v_0 \in V$ and $\sigma_0(x) - \bar{\rho} \in H_N^2(\Omega)$, are given. Note that from the above conditions on σ it follows that $\sigma \in C(0, T; H_N^2)$.

We solve (2.10) by considering the linearized problem

$$(2.11) \quad \begin{cases} P(\rho D_t v) - \mu \Delta v = F(\bar{\rho}, \bar{v}) \equiv \bar{F}, \\ v|_{t=0} = v_0(x), \\ D_t \sigma - \lambda \Delta \sigma = -\bar{v} \cdot \nabla \bar{\sigma}, \\ \sigma|_{t=0} = \sigma_0(x), \end{cases}$$

and by proving the existence of a fixed point $(\bar{\rho}, \bar{v}) = (\rho, v)$ for the map $(\bar{\rho}, \bar{v}) \rightarrow (\rho, v)$ defined by (2.11).

In order to get a sufficiently strong estimate for the linearized equation (2.11)₃ we take in account the particular form of the data $\bar{v} \cdot \nabla \bar{\sigma}$. As for estimate (2.3) we will introduce a balance parameter $\varepsilon > 0$.

Theorem 2.2 Assume that $\bar{v} \in L^2(0, T; H^2) \cap C(0, T; H_0^1)$ and that
 $\bar{\sigma} \in L^2(0, T; H_N^3) \cap C(0, T; H_N^2)$. Then the solution $\sigma \in L^2(0, T; H_N^3)$, $\sigma' \in L^2(0, T; H^1)$ of
problem (2.11)₃, (2.11)₄ verifies the estimate

$$(2.12) \quad \begin{aligned} & \|\sigma\|_{C(0, T; H_N^2)}^2 + \|\sigma\|_{L^2(0, T; H_N^3)}^2 < \\ & < c_2 \|\sigma_0\|_2^2 + c_3 \varepsilon^{-3} T \left[\|\bar{v}\|_{C(0, T; H^2)}^6 + \|\bar{v}\bar{\sigma}\|_{C(0, T; H^1)}^6 \right] \\ & + c_3 \varepsilon \left[\|\bar{v}\|_{L^2(0, T; H^2)}^2 + \|\bar{v}\bar{\sigma}\|_{L^2(0, T; H^2)}^2 \right], \end{aligned}$$

for every positive ε verifying

$$(2.13) \quad \varepsilon < \frac{\lambda}{2c_1},$$

where c_1 is the constant in (2.16). Here c_1, c_2, c_3 are positive constants depending
only on Ω .

Proof. The existence of a solution σ in the required space follows with standard techniques from the a priori bound (2.12), or from [9], Volume II, Chapter 4, Theorem 5.2, with $H = H^1$. Let us prove (2.12):

By application of the operator Δ to both sides of (2.11)₃, by multiplication by $\Delta\sigma$, and by integration over Ω one gets $\frac{1}{2}D_t |\Delta\sigma|^2 + (\nabla(\lambda\Delta\sigma - \bar{v} \cdot \nabla\sigma), \nabla\Delta\sigma) = 0$. Hence

$$(2.14) \quad \frac{1}{2} \frac{d}{dt} |\Delta\sigma|^2 + \lambda |\nabla\Delta\sigma|^2 < \\ < (|\bar{Dv}\bar{D}\sigma| + |\bar{v}D^2\sigma|) |\nabla\Delta\sigma| .$$

By using Sobolev's embedding theorem $H^1 \hookrightarrow L^6$ and Hölder's inequality it follows that

$$(2.15) \quad \frac{1}{2} \frac{d}{dt} |\Delta\sigma|^2 + \lambda |\nabla\Delta\sigma|^2 < \\ < c(|\bar{Dv}|^{1/2} |\bar{Dv}|^{1/2} |\bar{v}\sigma|_1 + \\ + |\bar{v}|_1 |\bar{v}\sigma|_1^{1/2} |\nabla\Delta\sigma|^{1/2}) |\nabla\Delta\sigma| .$$

An utilization of abc $< (\beta\epsilon^{-3})a^4 + (\epsilon/2)b^2 + (\epsilon/2)c^4$, $\epsilon > 0$, leads to

$$(2.16) \quad \frac{1}{2} \frac{d}{dt} |\Delta\sigma|^2 + \lambda |\nabla\Delta\sigma|^2 < c \epsilon |\bar{Dv}|_1^2 + \\ + c_1 \epsilon |\nabla\Delta\sigma|^2 + \frac{c}{\epsilon^3} |\bar{v}|_1^4 |\bar{v}\sigma|_1^2 + \\ + c \epsilon |\nabla\Delta\sigma|^2 + \frac{c}{\epsilon^3} |\bar{v}|_1^2 |\bar{v}\sigma|_1^4 .$$

Hence for ϵ verifying (2.13) one has

$$(2.17) \quad \frac{d}{dt} |\Delta\sigma|^2 + \lambda |\nabla\Delta\sigma|^2 < \frac{c}{\epsilon^3} (|\bar{v}|_1^2 |\bar{v}\sigma|_1^4 + \\ + |\bar{v}|_1^4 |\bar{v}\sigma|_1^2) + c_9 \epsilon (|\bar{Dv}|_1^2 + |\nabla\Delta\sigma|^2),$$

where the constants c depend only on Ω . This proves inequality (2.12). Recall that

$$|\sigma|_2 < c|\Delta\sigma|, \quad |\sigma|_3 < c|\nabla\Delta\sigma| .$$

Remark. One could also consider the linearized equation $D_t \sigma + \bar{v} \cdot \nabla\sigma - \lambda\Delta\sigma = 0$ instead of (2.11)₃, then estimate (2.17) holds with $\bar{\sigma}$ replaced by σ and without the term $c_9 \epsilon |\nabla\Delta\sigma|^2$. In this case the solution σ of the linearized problem verifies the maximum principle (which doesn't hold for the solution of (2.11)₃). However, the linearization (2.11)₃ seems to be more in keeping with the linearization (2.11)₁. Besides, the maximum principle will be recovered for the solution of the full nonlinear problem (2.10)₃.

3. The nonlinear problem. Local existence. We will not take care of the explicit dependence on μ, λ, m, M ; some of the constants c, c_k , depend on these fixed quantities. In order to simplify the equations, we denote by K_0, K_1, K_2, \dots , constants depending on the norms of the initial data $|\bar{v}_0|_V$ and $|\sigma_0|_2$.

In this section we solve (2.10) by proving the existence of a fixed point $(\rho, v) = (\bar{\rho}, \bar{v})$ for system (2.11). Define

$$\begin{aligned} K_1 &\equiv \{ \bar{v} : \bar{v}|_{t=0} = v_0(x), \|\bar{v}\|_{L^2(0,T;H^2)}^2 + \\ &\quad + \|\bar{v}\|_{C(0,T;V)}^2 + \|\bar{v}'\|_{L^2(0,T;H)}^2 < 2c_4 \|v_0\|_V^2 \} \\ K_2 &\equiv \{ \bar{\sigma} : \bar{\sigma}|_{t=0} = \sigma_0(x), \|\bar{\sigma}\|_{L^2(0,T;H_N^3)}^2 + \|\bar{\sigma}\|_{C(0,T;H_N^2)}^2 \\ &\quad < 2c_2 \|\sigma_0\|_2^2, \|\bar{\sigma}'\|_{L^2(0,T;H^1)} < K_0, \\ &\quad \|\bar{\sigma} - \sigma_0\|_{C(\bar{Q}_T)} < \frac{m}{2} \} , \end{aligned}$$

where $c_4 \equiv \mu[\min\{\mu, M, (\mu M^2)/(4M^2)\}]^{-1}$ and $K_0 \equiv \lambda\sqrt{2c_2} \|\sigma_0\|_2 + \bar{c}\sqrt{4c_2c_4} \|v_0\|_V \|\sigma_0\|_2$.

Here $\bar{c} = \bar{c}(\Omega)$ is a positive constant such that

$$(3.1) \quad \|\bar{v} \cdot \bar{w}\|_1 < \bar{c} \|\bar{v}\|_V \|\bar{w}\|_2, \quad \forall \bar{v} \in V, \bar{w} \in H^2.$$

Note that for every $\bar{\sigma} \in K_2$ one has in Q_T

$$(3.2) \quad \frac{m}{2} < \bar{\rho}(t, x) < M + \frac{m}{2}.$$

We now evaluate the L^2 norm of $\bar{F} \equiv F(\bar{\rho}, \bar{v})$. By using Sobolev's embedding theorem $H^1 \hookrightarrow L^6$ and Hölder's inequality one easily gets

$$(3.3) \quad \begin{aligned} \|F(\bar{\rho}, \bar{v})\|_{L^2}^2 &< c \|\bar{v}\|_1^3 \|\bar{D}\bar{v}\|_1 + c \|\bar{v}\|_1^2 \|\bar{D}^2\bar{\sigma}\| \\ &\quad + \|\bar{D}^2\bar{\sigma}\|_1 + c \|\bar{v}\|_1 \|\bar{D}\bar{v}\|_1 \|\bar{D}\bar{\sigma}\|_1^2 + \\ &\quad + c \|\bar{D}\bar{\sigma}\|_1^3 \|\bar{D}^2\bar{\sigma}\|_1 + c \|\bar{D}\bar{\sigma}\|_1^6 + \\ &\quad + c \|f\|_{L^2}^2. \end{aligned}$$

Consequently,

$$(3.4) \quad \begin{aligned} \|F(\bar{\rho}, \bar{v})\|_{L^2(0,T;H)}^2 &< c (\|v_0\|_V + \|\sigma_0\|_2)^{7/2} T^{1/2} + \\ &\quad + c \|\sigma_0\|_2^6 T + c \|f\|_{L^2(0,T;H)}^2. \end{aligned}$$

Hence, by using (2.3), it follows that the solution v of (2.11)₁, (2.11)₂ verifies

$$(3.5) \quad \|v\|_{C(0,T;V)}^2 + \|v\|_{L^2(0,T;H^2)}^2 + \|v'\|_{L^2(0,T;L^2)}^2 < \\ < C_4 \|v_0\|_V^2 + K_1(\sqrt{T} + T) + C \|f\|_{L^2(0,T;H)}^2 .$$

On the other hand, from (2.12),

$$(3.6) \quad \| \sigma \|_{C(0,T;H_N^2)}^2 + \| \sigma \|_{L^2(0,T;H_N^3)}^2 < \\ < \{ C_2 \| \sigma_0 \|_2^2 + K_3 \epsilon^{-3} T + K_4 \epsilon \} .$$

Now we fix $\epsilon > 0$ such that $K_4 \epsilon < 2^{-1} C_2 \| \sigma_0 \|_2^2$. Finally, by fixing a sufficiently small $T > 0$, it follows that $v \in K_1$, $\sigma \in K_2$. The estimate for $D_t \sigma$ follows by using equation (2.11)₃ and (3.1). The estimate for the sup norm of $\sigma - \sigma_0$ in \bar{Q}_T is proved as follows:

Clearly, $\| \sigma(t) - \sigma_0 \|_1 < \int_0^t \| \sigma'(s) \|_1 ds < K_0 T^{1/2}$. On the other hand $\| \sigma(t) - \sigma_0 \|_{C(\bar{\Omega})} < C_5 \| \sigma(t) - \sigma_0 \|_1^{1/3} \| \sigma(t) - \sigma_0 \|_2^{2/3}$, where C_5 depends only on Ω ; recall that $H^{5/3}(\Omega) \hookrightarrow C(\bar{\Omega})$. Consequently

$$\| \sigma - \sigma_0 \|_{C(\bar{Q}_T)} < C_5 K_0^{1/3} T^{1/6} (C_6 \sqrt{2C_2} \| \sigma_0 \|_2 + \| \sigma_0 \|_2)^{2/3} ,$$

where $C_6 = C_6(\Omega)$ is a positive constant, such that $\| \sigma \|_{C(\bar{\Omega})} < C_6 \| \sigma \|_2$, $\forall \sigma \in H_N^2(\Omega)$.

Hence, by choosing (if necessary) a smaller value for T , one gets $\| \sigma - \sigma_0 \|_{C(\bar{Q}_T)} < m/2$.

Now we utilize Schauder's fixed point theorem. Clearly $K \equiv K_1 + K_2$ is a convex, compact set in $L^2(0,T;L^2) \times L^2(0,T;L^2)$. Let us denote by ϕ the map $\phi(\bar{\rho}, \bar{v}) = (\rho, v)$, defined by (2.11). Since $\phi(K) \subset K$, it is sufficient to prove that $\phi : K \rightarrow K$ is continuous in the L^2 topology. If $\bar{v}_n \rightarrow \bar{v}$ in $L^2(Q_T)$, $\bar{\rho}_n \rightarrow \bar{\rho}$ in $L^2(Q_T)$, it follows by compactness arguments that $\bar{v}_n \rightarrow \bar{v}$ weakly in $L^2(0,T;H^2)$ and in $H^1(0,T;L^2)$, and that $\bar{v}_n \rightarrow \bar{v}$ weakly in $L^2(0,T;H^2)$ and in $H^1(0,T;L^2)$. In particular $\bar{\rho}_n$ is a bounded sequence in $H^{1/2+\epsilon_1}(0,T;H^{2-\epsilon_2}) \hookrightarrow C^{0,\alpha}(\bar{Q}_T)$ ⁽³⁾, for suitable positive $\epsilon_1, \epsilon_2, \alpha$.

(3) α -Holder continuous functions in \bar{Q}_T .

Hence $\bar{\rho}_v \rightarrow \bar{\rho}$ uniformly in \bar{Q}_T . Moreover, \bar{v}_n and $\bar{\nabla} \bar{\rho}_n$ are bounded in $H^{1/2}(0, T; H^1)$ and in $H^{1/2}(0, T; H^1)$ respectively. Hence $\bar{v}_n \rightarrow \bar{v}$ and $\bar{\nabla} \bar{\rho}_n \rightarrow \bar{\nabla} \bar{\rho}$ strongly in the L^4 topology. It follows from (2.8) at $F(\bar{\rho}_n, \bar{v}_n) \rightarrow F(\bar{\rho}, \bar{v})$ as a distribution in Q_T ; consequently $F(\bar{\rho}_n, \bar{v}_n) \rightarrow F(\bar{\rho}, \bar{v})$ weakly in $L^2(Q_T)$, because $F(\bar{\rho}_n, \bar{v}_n)$ is a bounded sequence in this space. Analogously, $\bar{v}_n \cdot \bar{\nabla} \bar{\rho}_n \rightarrow \bar{v} \cdot \bar{\nabla} \bar{\rho}$ strongly in $L^2(Q_T)$. It follows from the linear equations (2.11) that $\bar{v}_n \rightarrow \bar{v}$ and $\bar{\rho}_n \rightarrow \bar{\rho}$ in $L^2(Q_T)$ and $L^2(Q_T)$, respectively. Hence Φ is continuous. This finishes the proof of the existence of a local solution. Uniqueness will be proved in Section 5.

4. Global solutions. Asymptotic behavior.

In this section the constants C_K depend at most on Ω and on the quantities μ, λ and $\hat{\rho}$, i.e. on the total amount of mass $|\hat{\Omega}| \hat{\rho}$. We assume that

$$(4.1) \quad \|\sigma_0\|_2 < (2C)^{-1} \hat{\rho},$$

hence $\hat{\rho}/2 < m < M < 3\hat{\rho}/2$. Let (ρ, v) be a solution of (1.1). From (2.6) for

$$\epsilon_0 = (4M^2)^{-1} m \mu, \text{ and from (3.3) one gets}$$

$$(4.2) \quad \frac{m}{2} |D_t v|^2 + \frac{\mu}{2} \frac{d}{dt} |v|_V^2 + \frac{m\mu^2}{8M^2} |\Delta v|^2 < \\ < C (|v|_1^2 + |\sigma|_2^3) (|v|_2 + |\sigma|_3) + C |\sigma|_2^6 + C |f|^2,$$

where C depends only on $\Omega, \mu, \hat{\rho}$. On the other hand, from (2.17) for $\epsilon = (2C_9)^{-1} \lambda$, one gets

$$(4.3) \quad \frac{d}{dt} |\Delta \sigma|^2 + \frac{\lambda}{2} |\nabla \Delta \sigma|^2 < C (|v|_1^6 + |\sigma|_2^6).$$

From (4.2) and (4.3) it easily follows that

$$\frac{d}{dt} \left(\frac{\mu}{2} |v|_V^2 + |\Delta \sigma|^2 \right) + \frac{m}{2} |D_t v|^2 + \\ + \frac{m\mu^2}{16M^2} |\Delta v|^2 + \frac{\lambda}{4} |\nabla \Delta \sigma|^2 < C (|v|_V^6 + |\Delta \sigma|^6 + |f|^2).$$

In particular, since $|Av| > C|v|_V$ and $|V\Delta\sigma| > C|\Delta\sigma|$, one has

$$(4.4) \quad \frac{d}{dt} (|v|_V^2 + |\Delta\sigma|^2) < -[C_{10} - C_{11} (|v|_V^2 + |\Delta\sigma|^2)^2] \cdot (|v|_V^2 + |\Delta\sigma|^2) + C_{12} |f|^2.$$

Hence (4.5)₁ below holds for every $t \in [0, +\infty[$ if

$$(4.5) \quad \begin{cases} C_{11} (|v_0|_V^2 + |\Delta\sigma_0|^2)^2 < \frac{C_{10}}{2} \\ C_{12} |f|_{L^\infty(0, +\infty; \mathbb{R})}^2 < \frac{C_{10}}{2} \sqrt{\frac{C_{10}}{2C_{11}}} \end{cases}$$

In fact, if $C_{11} (|v(t)|_V^2 + |\Delta\sigma(t)|^2)^2 = C_{10}/2$ it must be, from (4.4), that $\frac{d}{dt} (|v(t)|_V^2 + |\Delta\sigma(t)|^2) < 0$.

Let us now prove the last assertion in theorem A. Under the hypothesis (4.5)₁, it follows from (4.4) that

$$\frac{d}{dt} (|v|_V^2 + |\Delta\sigma|^2) < -\frac{C_{10}}{2} (|v|_V^2 + |\Delta\sigma|^2).$$

this proves (1.6). □

5. Uniqueness. We prove that the solution (ρ, v) of problem (1.1) is unique in the class in which existence was proved; see Theorem A. We remark that more careful calculations lead to uniqueness in a larger class.

Let $(\rho, v), (\bar{\rho}, \bar{v})$ be two solutions of problem (1.1), (1.2) and put $u = v - \bar{v}, \eta = \rho - \bar{\rho}$. By subtracting the equations (2.10)₁ written for (ρ, v) and $(\bar{\rho}, \bar{v})$ respectively, and by taking the inner product with u in \mathbb{H} one gets

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} (\rho u, u) + \mu |u|_V^2 &= -\frac{1}{2} (v \cdot \nabla \rho, u^2) + \\ &+ \frac{\lambda}{2} (\Delta \rho, u^2) - (u, D_t \bar{v} \cdot u) + (F - \bar{F}, u). \end{aligned}$$

By using $(\Delta \rho, u^2) = -(\nabla \rho, Du^2)$, we show that

$$(5.1) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} (\rho u, u) + \frac{\mu}{2} |u|_V^2 &< \frac{1}{2} |v|_\infty |\nabla \rho|_\infty |u|^2 + \\ &+ \frac{\lambda^2}{\mu} |\nabla \rho|_\infty^2 |u|^2 + C |D_t \bar{v}|^2 |u|^2 + \\ &+ \frac{\lambda}{4} |\Delta \eta|^2 + (F - \bar{F}, u). \end{aligned}$$

On the other hand, by subtracting equations (2.10)₃ written for (ρ, v) and for $(\bar{\rho}, \bar{v})$ respectively, and by taking the inner product with $\Delta \eta$ in $L^2(\Omega)$ one gets

$$(5.2) \quad \frac{1}{2} \frac{d}{dt} \|\nabla \eta\|^2 + \frac{\lambda}{2} \|\Delta \eta\|^2 < c \|\nabla \bar{\rho}\|_{\infty}^2 \|u\|^2 + c \|\nu\|_{\infty}^2 \|\nabla \eta\|^2.$$

By adding (5.1) and (5.2) it follows that

$$(5.3) \quad \frac{d}{dt} [(\rho u, u) + \|\nabla \eta\|^2] + \mu \|u\|_V^2 + \frac{\lambda}{2} \|\Delta \eta\|^2 < \theta_1(t) (\|u\|^2 + \|\nabla \eta\|^2) + (F - \bar{F}, u),$$

where $\theta_1(t)$ is a real integrable function on $[0, T]$.

On the other hand, by using Sobolev's embedding theorems and Hölder's inequality (and also $ab < \epsilon a^2 + \epsilon^{-1} b^2$) the reader easily verifies that given $\epsilon > 0$ there exists an integrable real function $\theta_2(t)$ (dependent on $\rho, \bar{\rho}, \nu, \bar{\nu}$ and on ϵ) such that

$$(5.4) \quad |(F - \bar{F}, u)| < \theta_2(t) \|u\|^2 + \epsilon (\|\eta\|_2^2 + \|u\|_1^2).$$

By using $\|u\|^2 < \mu^{-1} (\rho u, u)$, (5.3) and (5.4) it follows that

$$\frac{d}{dt} [(\rho u, u) + \|\nabla \eta\|^2] < (\theta_1(t) + \theta_2(t)) [(\rho u, u) + \|\nabla \eta\|^2].$$

Uniqueness follows now from Gronwall's lemma and from $u|_{t=0} = 0, \eta|_{t=0} = 0$. □

REFERENCES

1. H. Beirao-da-Veiga, R. Serapioni, A. Valli "On the Motion of Non-homogeneous fluids in the Presence of Diffusion", J. Math. Anal. Appl., 85 (1982), 179-191.
2. D.A. Frank-Kamenetskii "Diffusion and Heat Transfer in chemical Kinetics", Plenum ed., New York, London (1969).
3. E. Gagliardo "Caretterizzazioni delle tracce sulla Frontiera Relative ad alcune classi di Functioni in M Variabili", Rend. Sem. Mat. Univ. Padova, 27 (1957), 284-305.
4. V.N. Ignat'ev, B.G. Kuznetsov "A Model for the Diffusion of a Turbulent Boundary Layer in a Polymer", Cisl. Metody Meh. Splošn. Sredy, 4 (1973), 78-87 (in russian).
5. A.V. Kazhikhov, Sh. Smagulov, "The Correctness of Boundary-Value Problems in a Diffusion Model of an Inhomogeneous Liquid", Sov. Phys. Dokl., 22 (1977), 249-250.
6. A.V. Kazhikhov, Sh. Smagulov, "The Correctness of Boundary-Value Problems in a Certain Diffusion Model of an Inhomogeneous Fluid", Cisl. Metody Meh. Splošn. Sredy, 7 (1976), 75-92.
7. O.A. Ladyzhenskaya, "The Mathematical Theory of Viscous Incompressible Flow", Gordon and Breach, New York (1969).
8. J.L. Lions, "Quelques methodes de resolution des problemes aux limites non lineaires", Dunod, Gauthier-Villars, Paris (1969).
9. J.L. Lions, E. Magenes "Problemes aux limites nonhomogenes et applications", Vol. I,II, Dunod, Paris (1968).
10. G. Prouse, A. Zaretti "On the Inequalities Associated to a Model of Graffi for the Motion of a Mixture of two viscous Incompressible Fluids", preprint Dipartimento di Matematica - Politecuico di Milano (1982).

11. P. Secchi "On the Initial Value Problem for the Equations of Motion of Viscous Incompressible Fluids in Presence of Diffusion", Boll. U.M.I., 1-B(1982) 1117-1130.
12. R. Temam "Navier-Stokes Equations", North Holland Amsterdam (1977).

HB/jgb

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2565	2. GOVT ACCESSION NO. A 134538	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Diffusion on Viscous Fluids, Existence and Asymptotic Properties of Solutions	5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period	
	6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(s) H. Beirao-da-Veiga	8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Madison, Wisconsin 53706	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis	
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709	12. REPORT DATE September 1983	
	13. NUMBER OF PAGES 16	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Non-homogeneous viscous fluids, non-linear parabolic equations, initial-boundary value problems		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We consider the motion of a mixture of two fluids, with a diffusion effect obeying Fick's law; for the derivation of the model (1.1) see Section 1 and references [2], [4], [5] and [6]. We consider the full non-linear problem (i.e., we don't omit the term λ^2 term in equation (1.1)). Moreover, we don't assume that λ/μ is small. We prove the existence of a (unique) local		

ABSTRACT (Continued)

solution, the existence of a global solution for small data, and the exponential decay to the equilibrium solution; see Theorem A, Section 1.

60-A134 538


DIFFUSION ON VISCOUS FLUIDS. EXISTENCE AND ASYMPTOTIC
PROPERTIES OF SOLUTIONS. (U) WISCONSIN UNIV-MADISON
MATHEMATICS RESEARCH CENTER H BEIRAO-DA-VEIGA SEP 83
MRC-TSR-2565 DAAG29-80-C-0041


1/1

UNCLASSIFIED

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963-A

A134538

(2)

MRC Technical Summary Report #2565

DIFFUSION ON VISCOUS FLUIDS, EXISTENCE
AND ASYMPTOTIC PROPERTIES OF SOLUTIONS

H. Beirao-da-Veiga

**Mathematics Research Center
University of Wisconsin—Madison
610 Walnut Street
Madison, Wisconsin 53705**

September 1983


(Received December 7, 1982)

**Approved for public release
Distribution unlimited**

DTIC FILE COPY

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709



88 11 08 150

- a -

0010
COPY
MAILED

UNIVERSITY OF WISCONSIN-MADISON
MATHEMATICS RESEARCH CENTER

DIFFUSION ON VISCOUS FLUIDS, EXISTENCE
AND ASYMPTOTIC PROPERTIES OF SOLUTIONS

H. Beirão-da-Veiga*

Technical Summary Report #2565

September 1983

ABSTRACT

The document
We consider^s the motion of a mixture of two fluids, with a diffusion effect obeying Fick's law, for the derivation of the model (1.1) see Section 1 and references [2], [4], [5] and [6]. We consider^s the full non-linear problem (i.e., we don't omit the term λ^2 term in equation (1.1)). Moreover, we *don't assume* that λ/μ is small. We prove^s the existence of a (unique) local solution, the existence of a global solution for small data, and the exponential decay to the equilibrium solution; see Theorem A, Section 1.

↑

AMS(MOS) Subject Classifications: 35K55, 76D99, 76R99.

Key Words: Non-homogeneous viscous fluids, non-linear parabolic equations, initial-boundary value problems

Work Unit Number 1 (Applied Analysis)

*Department of Mathematics, University of Trento, 38050 Povo (Trento) Italy.

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.

SIGNIFICANCE AND EXPLANATION

In this paper we consider the motion of a continuous medium consisting of two components, say water and a dissolved salt, with a diffusion effect obeying Fick's law. We denote by $v, w, \rho, \pi, \mu, \lambda$ the mean-volume velocity, mean-mass velocity, density, pressure, viscosity and diffusion constant, respectively. By using Fick's law we eliminate w from the equations and we obtain (1.1), where p is the modified pressure; see Section 1 and references [2], [4],[5] and [6]. The initial boundary conditions are given by equation (1.2).

Kazhikhov and Smagulov [5],[6] consider equation (1.1) for a small diffusion coefficient λ . More precisely, they assume that condition (1.3) holds; moreover, they omit the λ^2 term in equation (1.1)₁. Under these conditions they prove the existence of a unique local solution for the 3-dimensional motion (in the bi-dimensional case, solutions are global).

In our paper we consider the full equation (1.1), without assumption (1.3), and we prove: (i) the existence of a (unique) local solution; (ii) the existence of a global solution in time for small initial velocities and external forces, and for initial densities near-constant; (iii) the exponential decay (when $t \rightarrow +\infty$) of the solution (ρ, v) to the equilibrium solution $(\hat{\rho}, 0)$, if $f \equiv 0$. See Theorem A, Section 1.

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

DIFFUSION ON VISCOUS FLUIDS, EXISTENCE
AND ASYMPTOTIC PROPERTIES OF SOLUTIONS

H. Beirão-da-Veiga*

Main Notation

Ω : an open bounded set in \mathbb{R}^3 , locally situated on one side of its boundary Γ , a regular (say C^4) manifold.

$n = n(x)$: unit outward normal to Γ .

D_i, D_{ij}, D_t : $\partial/\partial x_i, \partial^2/\partial x_i \partial x_j, \partial/\partial t$.

$\| \cdot \|, (\cdot, \cdot)$: norm and scalar product in $L^2(\Omega)$.

H^k : Sobolev space $H^{k,2}(\Omega)$ with norm

$$\| \sigma \|_k^2 \equiv \sum_{l=0}^k \| D^l \sigma \|^2,$$

where

$$\| D^l \sigma \|^2 \equiv \sum_{|\alpha|=l} \| D^\alpha \sigma \|^2,$$

Further,

$$\| D^l \sigma \|_m^2 \equiv \sum_{|\alpha|=l} \| D^\alpha \sigma \|_m^2.$$

H_0^1 : Closure of $C_0^\infty(\Omega)$ in $H^1(\Omega)$.

$\| \cdot \|_m$: norm in $L^m(\Omega)$.

L^2, H^k, H_0^1 : Hilbert spaces of vectors $v = (v_1, v_2, v_3)$ such that

$v_i \in L^2, v_i \in H^k, v_i \in H_0^1$ ($i=1,2,3$), respectively. Corresponding

notation is used for other spaces of vector fields. Norms are defined

in the natural way, and denoted by the symbols used for the scalar

fields.

Let us introduce the following functional spaces (see, for instance, [7], [8] and [12] for their properties) =

*Department of Mathematics, University of Trento, 38050 Povo (Trento) Italy.

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.

$$H_N^k \equiv \{ \sigma \in H^k : \frac{\partial \sigma}{\partial n} = 0 \text{ on } \Gamma \text{ and } \int_{\Omega} \sigma(x) dx = 0 \}, k > 2.$$

$$V \equiv \{ v \in C_0^\infty(\Omega) : \operatorname{div} v = 0 \text{ in } \Omega \},$$

$$H \equiv \{ v \in L^2 : \operatorname{div} v = 0 \text{ in } \Omega, v \cdot n = 0 \text{ on } \Gamma \},$$

$$V \equiv \{ v \in H_0^1 : \operatorname{div} v = 0 \text{ on } \Omega \}.$$

H and V are the closure of v in $L^2(\Omega)$ and H_0^1 , respectively. Moreover $L^2 = H + G$, where $G \equiv \{ \nabla p : p \in H^1(\Omega) \}$. Denoting by P the orthogonal projection of L^2 onto H , we define the operator $A \equiv -PA$ on $D(A) \equiv H^2 \cap V$. One has

$$(Au, v) = ((u, v)) \equiv \sum_{i,j} (D_i u_j, D_i v_j), \forall u \in D(A), v \in V.$$

The norms $\| \cdot \|_2, \| \Delta \cdot \|$ are equivalent in H_N^2 , $\| \cdot \|_3, \| \nabla \Delta \cdot \|$ are equivalent in H_N^3 and $\| v \|_2, \| \Delta v \|$ are equivalent in $D(A)$. We define $\| v \|_V^2 \equiv ((v, v))$; the norms $\| v \|_V, \| v \|_1$ are equivalent in V .

$L^2(0, T; X)$: Banach space of strongly measurable functions defined in $]0, T[$ with values in (a Banach space) X , for which

$$\| z \|_{L^2(0, T; X)}^2 \equiv \int_0^T \| z(t) \|_X^2 dt < +\infty.$$

$C(0, T; X)$: Banach space of X -vector valued continuous functions on $[0, T]$ endowed with the usual norm $\| z \|_{C(0, T; X)}$.

μ : viscosity (a positive constant).

λ : diffusion coefficient (a positive constant).

$v(t, x), v_0(x)$: mean-volume velocity. Initial mean-volume velocity.

$\rho(t, x), \rho_0(x)$: density of the mixture. Initial density.

Further,

$$m \equiv \inf_{x \in \Omega} \rho_0(x), \quad M \equiv \sup_{x \in \Omega} \rho_0(x),$$

$$\hat{\rho} \equiv \frac{1}{|\Omega|} \int_{\Omega} \rho_0(x) dx.$$

We assume that $m > 0$.

$\pi(t, x), p(t, x)$: pressure. Modified pressure

$$p = \pi + \lambda v \cdot \nabla \rho - \lambda^2 \Delta \rho + \lambda(2\mu + \mu') \Delta \log \rho.$$

$f(t, x)$: external mass-force.

We denote by $c, \bar{c}, c_0, c_1, c_2, \dots$ positive constant, depending at most on Ω and on the parameters μ, λ, m, M and $\hat{\rho}$. It is easy to derive at any stage of the proofs, the explicit dependence of the constants on the parameters.

For convenience we sometimes denote different constants by the same symbol c . Otherwise, we utilize the symbols $\bar{c}, c_k, k \in \mathbb{N}$.

1. Main results. In this paper we consider the motion of a viscous fluid consisting of two components, say, saturated salt water and water. The equations of the model are obtained, for example, in [2], [4], [5], and [6]. Let us give a brief sketch. Let ρ_1, ρ_2 be the characteristic densities (constants) of the two components, $v^{(1)}(t,x)$ and $v^{(2)}(t,x)$ their velocities, and $e(t,x), d(t,x)$ the mass and volume concentration of the first fluid. We define the density $\rho(t,x) \equiv d\rho_1 + (1-d)\rho_2$, and the mean-volume and mean-mass velocities $v \equiv d v^{(1)} + (1-d) v^{(2)}, w \equiv e v^{(1)} + (1-e)v^{(2)}$. Then the equations of motion are

$$\begin{cases} \rho[D_t w + (w \cdot \nabla)w - f] - \mu \Delta w - (\mu + \mu') \nabla \operatorname{div} w = -\nabla \pi, \\ \operatorname{div} v = 0, \\ D_t \rho + \operatorname{div}(\rho w) = 0. \end{cases}$$

On the other hand, Fick's diffusion law (see [2]) gives $w = v - \lambda \rho^{-1} \nabla \rho$. By elimination of w in the preceding equations one gets, after some calculations,

$$(1.1) \quad \begin{cases} \rho[D_t v + (v \cdot \nabla)v] - \mu \Delta v - \lambda [(\nabla \cdot \nabla) \nabla \rho + (\nabla \rho \cdot \nabla)v] + \\ + \frac{\lambda^2}{\rho} [(\nabla \rho \cdot \nabla) \nabla \rho - \frac{1}{\rho} (\nabla \rho \cdot \nabla \rho) \nabla \rho + \Delta \rho \nabla \rho] = -\nabla p + \rho f, \\ D_t \rho + v \nabla \rho - \lambda \Delta \rho = 0, \\ \operatorname{div} v = 0. \end{cases}$$

We want to solve system (1.1) in $Q_T \equiv]0, T[\times \Omega$. Here p is the modified pressure. We add to system (1.1) the following initial boundary-value conditions.

$$(1.2) \quad \begin{cases} v = 0 & \text{on }]0, T[\times \Gamma, \\ \frac{\partial \rho}{\partial n} = 0 & \text{on }]0, T[\times \Gamma, \\ v|_{t=0} = v_0(x) & \text{in } \Omega, \\ \rho|_{t=0} = \rho_0(x) & \text{in } \Omega. \end{cases}$$

The first two conditions mean that there is no flux through the boundary.

In [5], [6] Kazhikhov and Smagulov consider the simplified system obtained from (1.1) by omitting the term containing λ^2 ; moreover, they assume that

$$(1.3) \quad \lambda < \frac{2\mu}{M-m}.$$

Under these conditions Kazhikhov and Smagulov state the existence of a local solution in time (global in the bidimensional case).

In our paper we take into account the full equation (1.1), and omit the condition (1.3). For this more general case we prove: (i) the existence of a (unique) local solution for arbitrary initial data and external force field; (ii) the existence of a unique global strong solution, for small initial data and external force field. Moreover, if $f \equiv 0$, the solution (ρ, v) decays exponentially to the equilibrium solution $(\hat{\rho}, 0)$. More precisely we prove the following result:

Theorem A. Let $v_0 \in V, \rho_0 - \hat{\rho} \in H_N^2, f \in L^2(0, T; L^2)$. Then there exists $T_1 \in]0, T[$ such that problem (1.1), (1.2) is uniquely solvable in Q_{T_1} . Moreover $v \in L^2(0, T_1; H^2) \cap C(0, T_1; V), D_t v \in L^2(0, T_1; H), \rho - \hat{\rho} \in L^2(0, T_1; H_N^3) \cap C(0, T_1; H_N^2), D_t \rho \in L^2(0, T_1; H^1)$ and $m < \rho(t, x) < M$.

Moreover, there exist positive constants $k_1, k_2,$ and k_3 depending at most on Ω, μ, λ and on the mean density $\hat{\rho}$ ⁽¹⁾ such that that if

$$(1.4) \quad \|v_0\|_1 + \|\rho_0 - \hat{\rho}\|_2 < k_1,$$

(1) Or, equivalently, depending on the total amount of mass $|\Omega| \hat{\rho} = \int_{\Omega} \rho_0(x) dx$.

and

$$(1.5) \quad \|f\|_{L^2(0,+\infty;L^2)} < k_2,$$

then the solution is global in time. If $f \equiv 0$, the solution (ρ, v) decays exponentially to the equilibrium solution $(\hat{\rho}, 0)$, i.e.

$$(1.6) \quad \|v(t)\|_1 + \|\rho(t) - \hat{\rho}\|_2 < (\|v_0\|_1 + \|\rho_0 - \hat{\rho}\|_2) e^{-k_3 t},$$

for every $t > 0$.

Theorem A also holds for coefficients μ, λ regularly dependent on ρ, v , provided they are strictly positive and bounded, in a neighborhood of the range of values of the initial data $\rho_0(x), v_0(x)$. This generalization can be done without any difficulty. Moreover, with standard techniques, one can prove that the solutions have more regularity (up to C^∞) if the data are sufficiently regular and the usual compatibility conditions hold.

Local existence in the general case (i.e. with the λ^2 term and without (1.3)) was proved in the inviscid case by Beirão-da-Veiga, Serapioni and Valli in [1]. A similar result, in the viscous case and for $\Omega = \mathbb{R}^3$, was proved by Secchi [11]. For another approach (concerning Greffi's model) see [10].

2. The linearized equations. We start by proving the following theorem: Theorem 2.1 Let $\rho(t, x)$ be a measurable function verifying

$$(2.1) \quad 0 < m < \rho(t, x) < M, \quad \text{a.e., in } Q_T, \quad \text{let } F \in L^2(0, T; H) \quad \text{and } v_0 \in V.$$

Then there exists a (unique) strong solution v of problem

$$(2.2) \quad \begin{cases} \rho D_t v - \mu \Delta v = -\nabla p + F & \text{in } Q_T, \\ \operatorname{div} v = 0 & \text{in } Q_T, \\ v = 0 & \text{on }]0, T[\times \Gamma, \\ v|_{t=0} = v_0(x) & \text{in } \Omega. \end{cases}$$

Moreover $v \in L^2(0, T; D(A)) \cap C(0, T; V)$, $D_t v \in L^2(0, T; H)$ and

$$(2.3) \quad \begin{aligned} \mu \|v\|_{C(0, T; V)}^2 + m \|v\|_{L^2(0, T; H)}^2 + \\ + \frac{m\mu}{4m^2} \|Av\|_{L^2(0, T; H)}^2 < \mu \|v_0\|_V^2 + \left(\frac{2}{m} + \frac{m}{2M^2}\right) \|F\|_{L^2(0, T; H)}^2 \end{aligned}$$

Proof. Let us write equation (2.2) in the equivalent form (2.4)

$$P(\rho D_t v) + \mu \Delta v = F, \quad v|_{t=0} = v_0(X).$$

For brevity let us put

$$X = \{v; v \in L^2(0, T; D(\Lambda)), v' \in L^2(0, T; H)\}.$$

From well known results (see [9], Vol I, Chapter I : Theorem 3.1 with $y = H$, $X = D(\Lambda)$, $j = 0$; and (2.42) Proposition 2.1) it follows that $X \subset C(0, T; V)$.

We start by proving the a priori bound (2.3); an essential device is to introduce a parameter ε_0 in order to conveniently balance the estimates. In H take the inner product of (2.4) with $D_t v + \varepsilon_0 \Delta v$, $\varepsilon_0 > 0$. Since $(v', \Delta v) = 2^{-1} D_t |v|_V^2$ one gets

$$(2.5) \quad \begin{aligned} m |D_t v|^2 + \frac{\mu}{2} \frac{d}{dt} |v|_V^2 + \varepsilon_0 \mu |\Delta v|^2 < \\ < |F| |D_t v| + \varepsilon_0 |F| |\Delta v| + \varepsilon_0 M |D_t v| |\Delta v|. \end{aligned}$$

By using the inequalities $|F| |D_t v| < 4^{-1} m |D_t v|^2 + m^{-1} |F|^2$, $|F| |\Delta v| < 4^{-1} \mu |\Delta v|^2 + \mu^{-1} |F|^2$ and $|D_t v| |\Delta v| < (4M)^{-1} \mu |\Delta v|^2 + \mu^{-1} M |D_t v|^2$ one gets

$$(2.6) \quad \begin{aligned} \frac{3}{4} m |D_t v|^2 + \frac{\mu}{2} \frac{d}{dt} |v|_V^2 + \varepsilon_0 \frac{\mu}{2} |\Delta v|^2 < \\ < \left(\frac{1}{m} + \frac{\varepsilon_0}{\mu}\right) |F|^2 + \frac{\varepsilon_0 M^2}{\mu} |D_t v|^2. \end{aligned}$$

Now fix $\varepsilon_0 = (4M^2)^{-1} \mu$ and integrate equation (2.6) on $(0, T)$. This gives the a priori bound (2.3).

Define $|v|_X^2 \equiv$ "left hand side of equation (2.3)", $Y \equiv L^2(0, T; H) \times V$, and $|(F, v_0)|_Y^2 \equiv$ "right hand side of (2.3)".

We solve (2.4) by the continuity method. Define $\rho \alpha \equiv (1-\alpha) \hat{\rho} + \alpha \rho$, $\alpha \in [0, 1]$. Clearly ρ_α verifies condition (2.1), for any α . Define $T_\alpha \equiv (1-\alpha) \bar{T} + \alpha T$, where

$$\begin{aligned} T_v &\equiv (P(\rho D_t v) - \mu \Delta v, v|_{t=0}) \in Y, \\ \bar{T}_v &\equiv (P(\bar{\rho} D_t v) - \mu \Delta v, v|_{t=0}) \in Y. \end{aligned}$$

Finally denote by γ the set of values $\alpha \in [0,1]$ for which problem (2.2) is solvable in X for every pair $(F, v_0) \in Y$. Clearly $0 \in \gamma$, because for this value of the parameter equation (2.6) because the linearized Navier-Stokes equation. Let us verify that γ is open and closed.

γ is open. Let $\alpha_0 \in \gamma$ and denote by $G(F, v_0) \equiv v$ the solution v of problem $T_{\alpha_0} v = (F, v_0)$. From (2.3) one gets $G \in L(Y; X)$ ⁽²⁾ with $\|G\|_{Y, X} < 1$. Equation $T_{\alpha_0 + \varepsilon} v = (F, v_0)$ can be written in the form

$$(2.7) \quad [1 - \varepsilon G(\bar{T} - T)]v = G(F, v_0).$$

Since $\|G(\bar{T} - T)\|_{X, Y} < \|(\bar{T} - T)\|_{X, Y}^{-1}$, equation (2.7) is solvable for $|\varepsilon| < \|(\bar{T} - T)\|_{X, Y}^{-1}$ (by a Neuman expansion).

γ is closed. Let $\alpha_n \in \gamma$, $\alpha_n \rightarrow \alpha_0$, and let v_n be the solution of $T_{\alpha_n} v_n = (F, v_0)$. From (2.3) one has $\|v_n\|_X < \|(F, v_0)\|_Y$. Since X is an Hilbert space there exists a subsequence $v_{n'} \rightarrow v \in X$, weakly in X . From $T, \bar{T} \in L(X; Y)$ one has $\bar{T}v_{n'} \rightarrow \bar{T}v$, $Tv_{n'} \rightarrow Tv$ weakly in Y . Hence $T_{\alpha_0} v = (F, v_0)$, i.e. $T_{\alpha_0} v = (F, v_0)$.

Let us now return to problem (1.1). Define

$$(2.8) \quad F(\rho, v) \equiv P \{-\rho(v \cdot \nabla)v + \lambda[(v \cdot \nabla)\nabla\rho + (\nabla\rho \cdot \nabla)v] + \frac{\lambda^2}{\rho} [(\nabla\rho \cdot \nabla)\nabla\rho - \frac{1}{\rho}(\nabla\rho \cdot \nabla\rho)\nabla\rho + \Delta\rho\nabla\rho] + \rho f\}.$$

For convenience we will use in the sequel the translation

$$(2.9) \quad \rho = \hat{\rho} + \sigma.$$

Recall that $\hat{\rho}$ is a given constant. To solve problem (1.1) and (1.2) in our functional framework is equivalent to finding $v \in L^2(0, T; D(A))$, $v_t \in L^2(0, T; H)$ and $\sigma \in L^2(0, T; H_N^3)$, $\sigma_t \in L^2(0, T; H^1)$ such that

$$(2.10) \quad \begin{cases} P(\rho D_t v) - \mu \Delta v = F(\rho, v), \\ v|_{t=0} = v_0(x), \\ D_t \sigma - \lambda \nabla \sigma = -v \cdot \nabla \sigma, \\ \sigma|_{t=0} = \sigma_0(x), \end{cases}$$

⁽²⁾ Banach space of linear continuous operator from Y into X , with norm $\|\cdot\|_{Y, X}$.

where $v_0 \in V$ and $\sigma_0(x) - \hat{p} \in H_N^2(\Omega)$, are given. Note that from the above conditions on σ it follows that $\sigma \in C(0, T; H_N^2)$.

We solve (2.10) by considering the linearized problem

$$(2.11) \quad \begin{cases} P(\rho D_t v) - \mu \Delta v = F(\bar{\rho}, \bar{v}) \equiv \bar{F}, \\ v|_{t=0} = v_0(x), \\ D_t \sigma - \lambda \Delta \sigma = -\bar{v} \cdot \nabla \bar{\sigma}, \\ \sigma|_{t=0} = \sigma_0(x), \end{cases}$$

and by proving the existence of a fixed point $(\bar{\rho}, \bar{v}) = (\rho, v)$ for the map $(\bar{\rho}, \bar{v}) \rightarrow (\rho, v)$ defined by (2.11).

In order to get a sufficiently strong estimate for the linearized equation (2.11) we take in account the particular form of the data $\bar{v} \cdot \nabla \bar{\sigma}$. As for estimate (2.3) we will introduce a balance parameter $\varepsilon > 0$.

Theorem 2.2 Assume that $\bar{v} \in L^2(0, T; H^2) \cap C(0, T; H_0^1)$ and that $\bar{\sigma} \in L^2(0, T; H_N^3) \cap C(0, T; H_N^2)$. Then the solution $\sigma \in L^2(0, T; H_N^3)$, $\sigma' \in L^2(0, T; H^1)$ of problem (2.11)₃, (2.11)₄ verifies the estimate

$$(2.12) \quad \begin{aligned} & \|\sigma\|_{C(0, T; H_N^2)}^2 + \|\sigma\|_{L^2(0, T; H_N^3)}^2 < \\ & < c_2 \|\sigma_0\|_2^2 + c_3 \varepsilon^{-3} T \left[\|\bar{v}\|_{C(0, T; H^2)}^6 + \|\bar{v}\bar{\sigma}\|_{C(0, T; H^1)}^6 \right] \\ & + c_3 \varepsilon \left[\|\bar{v}\|_{L^2(0, T; H^2)}^2 + \|\bar{v}\bar{\sigma}\|_{L^2(0, T; H^2)}^2 \right], \end{aligned}$$

for every positive ε verifying

$$(2.13) \quad \varepsilon < \frac{\lambda}{2c_1},$$

where c_1 is the constant in (2.16). Here c_1, c_2, c_3 are positive constants depending only on Ω .

Proof. The existence of a solution σ in the required space follows with standard techniques from the a priori bound (2.12), or from [9], Volume II, Chapter 4, Theorem 5.2, with $H = H^1$. Let us prove (2.12):

By application of the operator Δ to both sides of (2.11)₃, by multiplication by $\Delta\sigma$, and by integration over Ω one gets $\frac{1}{2}D_t |\Delta\sigma|^2 + (\nabla(\lambda\Delta\sigma - \bar{v} \cdot \nabla\bar{\sigma}), \nabla\Delta\sigma) = 0$. Hence

$$(2.14) \quad \frac{1}{2} \frac{d}{dt} |\Delta\sigma|^2 + \lambda |\nabla\Delta\sigma|^2 < \\ < (|\text{DvD}\bar{\sigma}| + |\bar{v} \cdot \nabla^2 \bar{\sigma}|) |\nabla\Delta\sigma|.$$

By using Sobolev's embedding theorem $H^1 \hookrightarrow L^6$ and Hölder's inequality it follows that

$$(2.15) \quad \frac{1}{2} \frac{d}{dt} |\Delta\sigma|^2 + \lambda |\nabla\Delta\sigma|^2 < \\ < c (|\text{Dv}\bar{v}|^{1/2} |\text{Dv}\bar{v}|^{1/2} |\nabla\bar{\sigma}|_1 + \\ + |\bar{v}|_1 |\nabla\bar{\sigma}|_1^{1/2} |\nabla\Delta\sigma|^{1/2}) |\nabla\Delta\sigma|.$$

An utilization of abc $< (8\epsilon^{-3})a^4 + (\epsilon/2)b^2 + (\epsilon/2)c^4$, $\epsilon > 0$, leads to

$$(2.16) \quad \frac{1}{2} \frac{d}{dt} |\Delta\sigma|^2 + \lambda |\nabla\Delta\sigma|^2 < c \epsilon |\text{Dv}\bar{v}|_1^2 + \\ + c_1 \epsilon |\nabla\Delta\sigma|^2 + \frac{c}{\epsilon} |\bar{v}|_1^4 |\nabla\bar{\sigma}|_1^2 + \\ + c \epsilon |\nabla\Delta\sigma|^2 + \frac{c}{\epsilon} |\bar{v}|_1^2 |\nabla\bar{\sigma}|_1^4.$$

Hence for ϵ verifying (2.13) one has

$$(2.17) \quad \frac{d}{dt} |\Delta\sigma|^2 + \lambda |\nabla\Delta\sigma|^2 < \frac{c}{\epsilon} (|\bar{v}|_1^2 |\nabla\bar{\sigma}|_1^4 + \\ + |\bar{v}|_1^4 |\nabla\bar{\sigma}|_1^2) + c_9 \epsilon (|\text{Dv}\bar{v}|_1^2 + |\nabla\Delta\sigma|^2),$$

where the constants c depend only on Ω . This proves inequality (2.12). Recall that $|\sigma|_2 < c|\Delta\sigma|$, $|\sigma|_3 < c|\nabla\Delta\sigma|$.

Remark. One could also consider the linearized equation $D_t \sigma + \bar{v} \cdot \nabla\sigma - \lambda\Delta\sigma = 0$ instead of (2.11)₃; then estimate (2.17) holds with $\bar{\sigma}$ replaced by σ and without the term $c_9 \epsilon |\nabla\Delta\sigma|^2$. In this case the solution σ of the linearized problem verifies the maximum principle (which doesn't hold for the solution of (2.11)₃). However, the linearization (2.11)₃ seems to be more in keeping with the linearization (2.11)₁. Besides, the maximum principle will be recovered for the solution of the full nonlinear problem (2.10)₃.

3. The nonlinear problem. Local existence. We will not take care of the explicit dependence on μ, λ, m, M ; some of the constants c, c_k , depend on these fixed quantities. In order to simplify the equations, we denote by K_0, K_1, K_2, \dots , constants depending on the norms of the initial data $|\bar{v}_0|_V$ and $|\sigma_0|_2$.

In this section we solve (2.10) by proving the existence of a fixed point $(\rho, v) = (\bar{\rho}, \bar{v})$ for system (2.11). Define

$$\begin{aligned} K_1 &\equiv \{ \bar{v} : \bar{v}|_{t=0} = v_0(x), \|\bar{v}\|_{L^2(0,T;H^2)}^2 + \\ &\quad + \|\bar{v}\|_{C(0,T;V)}^2 + \|\bar{v}'\|_{L^2(0,T;H)}^2 < 2 C_4 \|v_0\|_V^2 \} \\ K_2 &\equiv \{ \bar{\sigma} : \bar{\sigma}|_{t=0} = \sigma_0(x), \|\bar{\sigma}\|_{L^2(0,T;H_N^3)}^2 + \|\bar{\sigma}\|_{C(0,T;H_N^2)}^2 \\ &\quad < 2 c_2 \|\sigma_0\|_2^2, \|\bar{\sigma}'\|_{L^2(0,T;H^1)} < K_0, \\ &\quad \|\bar{\sigma} - \sigma_0\|_{C(\bar{Q}_T)} < \frac{m}{2} \} , \end{aligned}$$

where $C_4 \equiv \mu [\min \{ \mu, M, (m\mu^2)/(4M^2) \}]^{-1}$ and $K_0 \equiv \lambda \sqrt{2c_2} \|\sigma_0\|_2 + \bar{c} \sqrt{4c_2 c_4} \|v_0\|_V \|\sigma_0\|_2$.

Here $\bar{c} = \bar{c}(\Omega)$ is a positive constant such that

$$(3.1) \quad \|\bar{v} \cdot \bar{w}\|_1 < \bar{c} \|\bar{v}\|_V \|\bar{w}\|_2, \quad \forall \bar{v} \in V, \bar{w} \in H^2.$$

Note that for every $\bar{\sigma} \in K_2$ one has in Q_T

$$(3.2) \quad \frac{m}{2} < \bar{\rho}(t, x) < M + \frac{m}{2}.$$

We now evaluate the L^2 norm of $\bar{F} \equiv F(\bar{\rho}, \bar{v})$. By using Sobolev's embedding theorem $H^1 \hookrightarrow L^6$ and Hölder's inequality one easily gets

$$(3.3) \quad \begin{aligned} \|F(\bar{\rho}, \bar{v})\|^2 &< c \|\bar{v}\|_1^3 \|D\bar{v}\|_1 + c \|\bar{v}\|_1^2 \|D^2\bar{\sigma}\| \\ &\quad + \|D^2\bar{\sigma}\|_1 + c \|\bar{v}\|_1 \|D\bar{v}\|_1 \|D\bar{\sigma}\|_1^2 + \\ &\quad + c \|D\bar{\sigma}\|_1^3 \|D^2\bar{\sigma}\|_1 + c \|D\bar{\sigma}\|_1^6 \\ &\quad + c \|f\|^2. \end{aligned}$$

Consequently,

$$(3.4) \quad \begin{aligned} \|F(\bar{\rho}, \bar{v})\|_{L^2(0,T;H)}^2 &< c (\|v_0\|_V + \|\sigma_0\|_2)^{7/2} T^{1/2} + \\ &\quad + c \|\sigma_0\|_2^6 T + c \|f\|_{L^2(0,T;H)}^2. \end{aligned}$$

Hence, by using (2.3), it follows that the solution v of (2.11)₁, (2.11)₂ verifies

$$(3.5) \quad |v|^2_{C(0,T;V)} + |v|^2_{L^2(0,T;H^2)} + |v'|^2_{L^2(0,T;L^2)} < \\ < C_4 |v_0|^2_V + K_1(\sqrt{T} + T) + C |f|^2_{L^2(0,T;H)} .$$

On the other hand, from (2.12),

$$(3.6) \quad |\sigma|^2_{C(0,T;H^2_N)} + |\sigma|^2_{L^2(0,T;H^3_N)} < \\ < \{ C_2 |\sigma_0|^2_2 + K_3 \varepsilon^{-3} T + K_4 \varepsilon \} .$$

Now we fix $\varepsilon > 0$ such that $K_4 \varepsilon < 2^{-1} C_2 |\sigma_0|^2_2$. Finally, by fixing a sufficiently small $T > 0$, it follows that $v \in K_1, \sigma \in K_2$. The estimate for $D_t \sigma$ follows by using equation (2.11)₃ and (3.1). The estimate for the sup norm of $\sigma - \sigma_0$ in \bar{Q}_T is proved as follows:

Clearly, $|\sigma(t) - \sigma_0|_1 < \int_0^t |\sigma'(s)|_1 ds < K_0 T^{1/2}$. On the other hand $|\sigma(t) - \sigma_0|_{C(\bar{\Omega})} < C_5 |\sigma(t) - \sigma_0|_1^{1/3} |\sigma(t) - \sigma_0|_2^{2/3}$, where C_5 depends only on Ω ; recall that $H^{5/3}(\Omega) \hookrightarrow C(\bar{\Omega})$. Consequently

$$|\sigma - \sigma_0|_{C(\bar{Q}_T)} < C_5 K_0^{1/3} T^{1/6} (C_6 \sqrt{2C_2} |\sigma_0|_2 + |\sigma_0|_2)^{2/3} ,$$

where $C_6 = C_6(\Omega)$ is a positive constant, such that $|\sigma|_{C(\bar{\Omega})} < C_6 |\sigma|_2, \forall \sigma \in H^2_N(\Omega)$.

Hence, by choosing (if necessary) a smaller value for T , one gets $|\sigma - \sigma_0|_{C(\bar{Q}_T)} < \varepsilon/2$.

Now we utilize Schauder's fixed point theorem. Clearly $K \equiv K_1 + K_2$ is a convex, compact set in $L^2(0,T;L^2) \times L^2(0,T;L^2)$. Let us denote by Φ the map $\Phi(\bar{\rho}, \bar{v}) = (\rho, v)$, defined by (2.11). Since $\Phi(K) \subset K$, it is sufficient to prove that $\Phi : K \rightarrow K$ is continuous in the L^2 topology. If $\bar{v}_n + \bar{v}$ in $L^2(Q_T), \bar{\rho}_n + \bar{\rho}$ in $L^2(Q_T)$, it follows by compactness arguments that $\bar{v}_n + \bar{v}$ weakly in $L^2(0,T;H^2)$ and in $H^1(0,T;L^2)$, and that $\bar{v}_n + \bar{v}$ weakly in $L^2(0,T;H^2)$ and in $H^1(0,T;L^2)$. In particular $\bar{\rho}_n$ is a bounded sequence in $H^{1/2+\varepsilon_1}(0,T;H^{2-\varepsilon_2}) \hookrightarrow C^{0,\alpha}(\bar{Q}_T)$ ⁽³⁾, for suitable positive $\varepsilon_1, \varepsilon_2, \alpha$.

(3) α -Holder continuous functions in \bar{Q}_T .

Hence $\bar{\rho}_v \rightarrow \bar{\rho}$ uniformly in \bar{Q}_T . Moreover, \bar{v}_n and $\bar{\nabla} \bar{\rho}_n$ are bounded in $H^{1/2}(0, T; H^1)$ and in $H^{1/2}(0, T; H^1)$ respectively. Hence $\bar{v}_n \rightarrow \bar{v}$ and $\bar{\nabla} \bar{\rho}_n \rightarrow \bar{\nabla} \bar{\rho}$ strongly in the L^4 topology. It follows from (2.8) that $F(\bar{\rho}_n, \bar{v}_n) \rightarrow F(\bar{\rho}, \bar{v})$ as a distribution in Q_T ; consequently $F(\bar{\rho}_n, \bar{v}_n) \rightarrow F(\bar{\rho}, \bar{v})$ weakly in $L^2(Q_T)$, because $F(\bar{\rho}_n, \bar{v}_n)$ is a bounded sequence in this space. Analogously, $\bar{v}_n \cdot \bar{\nabla} \bar{\rho}_n \rightarrow \bar{v} \cdot \bar{\nabla} \bar{\rho}$ strongly in $L^2(Q_T)$. It follows from the linear equations (2.11) that $\bar{v}_n \rightarrow \bar{v}$ and $\bar{\rho}_n \rightarrow \bar{\rho}$ in $L^2(Q_m)$ and $L^2(Q_m)$, respectively. Hence Φ is continuous. This finishes the proof of the existence of a local solution. Uniqueness will be proved in Section 5.

4. Global solutions. Asymptotic behavior.

In this section the constants C_K depend at most on Ω and on the quantities μ, λ and $\hat{\rho}$, i.e. on the total amount of mass $|\Omega| \hat{\rho}$. We assume that

$$(4.1) \quad \|\sigma_0\|_2 < (2C)^{-1} \hat{\rho},$$

hence $\hat{\rho}/2 < m < M < 3\hat{\rho}/2$. Let (ρ, v) be a solution of (1.1). From (2.6) for $\epsilon_0 = (4M^2)^{-1} m \mu$, and from (3.3) one gets

$$(4.2) \quad \frac{m}{2} \|D_t v\|^2 + \frac{m}{2} \frac{d}{dt} \|v\|_V^2 + \frac{m\mu}{8M^2} \|\Delta v\|^2 < \\ < C (\|v\|_1^2 + \|\sigma\|_2^3) (\|v\|_2 + \|\sigma\|_3) + C \|\sigma\|_2^6 + C \|f\|^2,$$

where C depends only on $\Omega, \mu, \hat{\rho}$. On the other hand, from (2.17) for $\epsilon = (2C_9)^{-1} \lambda$, one gets

$$(4.3) \quad \frac{d}{dt} \|\Delta \sigma\|^2 + \frac{\lambda}{2} \|\nabla \Delta \sigma\|^2 < C (\|v\|_1^6 + \|\sigma\|_2^6).$$

From (4.2) and (4.3) it easily follows that

$$\frac{d}{dt} \left(\frac{m}{2} \|v\|_V^2 + \|\Delta \sigma\|^2 \right) + \frac{m}{2} \|D_t v\|^2 + \\ + \frac{m\mu}{16M^2} \|\Delta v\|^2 + \frac{\lambda}{4} \|\nabla \Delta \sigma\|^2 < C (\|v\|_V^6 + \|\Delta \sigma\|^6 + \|f\|^2).$$

In particular, since $|Av| > C|v|_V$ and $|V\Delta\sigma| > C|\Delta\sigma|$, one has

$$(4.4) \quad \frac{d}{dt} (|v|_V^2 + |\Delta\sigma|^2) < -[C_{10} - C_{11} (|v|_V^2 + |\Delta\sigma|^2)^2] \cdot (|v|_V^2 + |\Delta\sigma|^2) + C_{12} |f|^2.$$

Hence (4.5)₁ below holds for every $t \in [0, +\infty[$ if

$$(4.5) \quad \begin{cases} C_{11} (|v_0|_V^2 + |\Delta\sigma_0|^2)^2 < \frac{C_{10}}{2} \\ C_{12} |f|_{L^\infty(0, +\infty; \mathbb{R})}^2 < \frac{C_{10}}{2} \sqrt{\frac{C_{10}}{2C_{11}}} \end{cases}$$

In fact, if $C_{11} (|v(t)|_V^2 + |\Delta\sigma(t)|^2)^2 = C_{10}/2$ it must be, from (4.4), that $\frac{d}{dt} (|v(t)|_V^2 + |\Delta\sigma(t)|^2) < 0$.

Let us now prove the last assertion in theorem A. Under the hypothesis (4.5)₁, it follows from (4.4) that

$$\frac{d}{dt} (|v|_V^2 + |\Delta\sigma|^2) < -\frac{C_{10}}{2} (|v|_V^2 + |\Delta\sigma|^2).$$

this proves (1.6). □

5. Uniqueness. We prove that the solution (ρ, v) of problem (1.1) is unique in the class in which existence was proved; see Theorem A. We remark that more careful calculations lead to uniqueness in a larger class.

Let $(\rho, v), (\bar{\rho}, \bar{v})$ be two solutions of problem (1.1), (1.2) and put $u = v - \bar{v}, \eta = \rho - \bar{\rho}$. By subtracting the equations (2.10)₁ written for (ρ, v) and $(\bar{\rho}, \bar{v})$ respectively, and by taking the inner product with u in \mathbb{H} one gets

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} (\rho u, u) + \mu |u|_V^2 &= -\frac{1}{2} (v \cdot \nabla \rho, u^2) + \\ &+ \frac{\lambda}{2} (\Delta \rho, u^2) - (u, D_t \bar{v} \cdot u) + (F - \bar{F}, u). \end{aligned}$$

By using $(\Delta \rho, u^2) = -(\nabla \rho, Du^2)$, we show that

$$(5.1) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} (\rho u, u) + \frac{\mu}{2} |u|_V^2 &< \frac{1}{2} |v|_\infty |\nabla \rho|_\infty |u|^2 + \\ &+ \frac{\lambda^2}{\mu} |\nabla \rho|_\infty^2 |u|^2 + C |D_t \bar{v}|^2 |u|^2 + \\ &+ \frac{\lambda}{4} |\Delta \eta|^2 + (F - \bar{F}, u). \end{aligned}$$

On the other hand, by subtracting equations (2.10)₃ written for (ρ, v) and for $(\bar{\rho}, \bar{v})$ respectively, and by taking the inner product with $\Delta \eta$ in $L^2(\Omega)$ one gets

$$(5.2) \quad \frac{1}{2} \frac{d}{dt} \|\nabla \eta\|^2 + \frac{\lambda}{2} \|\Delta \eta\|^2 < c \|\nabla \bar{\rho}\|_{\infty}^2 \|u\|^2 + c \|\nu\|_{\infty}^2 \|\nabla \eta\|^2.$$

By adding (5.1) and (5.2) it follows that

$$(5.3) \quad \frac{d}{dt} [(pu, u) + \|\nabla \eta\|^2] + \mu \|u\|_V^2 + \frac{\lambda}{2} \|\Delta \eta\|^2 < \theta_1(t) (\|u\|^2 + \|\nabla \eta\|^2) + (F - \bar{F}, u),$$

where $\theta_1(t)$ is a real integrable function on $[0, T]$.

On the other hand, by using Sobolev's embedding theorems and Hölder's inequality (and also $ab < \epsilon a^2 + \epsilon^{-1} b^2$) the reader easily verifies that given $\epsilon > 0$ there exists an integrable real function $\theta_2(t)$ (dependent on $\rho, \bar{\rho}, v, \bar{v}$ and on ϵ) such that

$$(5.4) \quad |(F - \bar{F}, u)| < \theta_2(t) \|u\|^2 + \epsilon (\|\eta\|_2^2 + \|\eta\|_1^2).$$

By using $\|u\|^2 < m^{-1}(pu, u)$, (5.3) and (5.4) it follows that

$$\frac{d}{dt} [(pu, u) + \|\nabla \eta\|^2] < (\theta_1(t) + \theta_2(t)) [(pu, u) + \|\nabla \eta\|^2].$$

Uniqueness follows now from Gronwall's lemma and from $u|_{t=0} = 0, \eta|_{t=0} = 0$. □

REFERENCES

1. H. Beirao-da-Veiga, R. Serapioni, A. Valli "On the Motion of Non-homogeneous fluids in the Presence of Diffusion", J. Math. Anal. Appl., 85 (1982), 179-191.
2. D.A. Frank-Kamenetskii "Diffusion and Heat Transfer in chemical Kinetics", Plenum ed., New York, London (1969).
3. E. Gagliardo "Caratterizzazioni delle tracce sulla Frontiera Relative ad alcune classi di Funzioni in N Variabili", Rend. Sem. Mat. Univ. Padova, 27 (1957), 284-305.
4. V.N. Ignat'ev, B.G. Kuznetsov "A Model for the Diffusion of a Turbulent Boundary Layer in a Polymer", Cisl. Metody Meh. Splosn. Sredy, 4 (1973), 78-87 (in russian).
5. A.V. Kazhikhov, Sh. Smagulov, "The Correctness of Boundary-Value Problems in a Diffusion Model of an Inhomogeneous Liquid", Sov. Phys. Dokl., 22 (1977), 249-250.
6. A.V. Kazhikhov, Sh. Smagulov, "The Correctness of Boundary-Value Problems in a Certain Diffusion Model of an Inhomogeneous Fluid", Cisl. Metody Meh. Splosn. Sredy, 7 (1976), 75-92.
7. O.A. Ladyzhenskaya, "The Mathematical Theory of Viscous Incompressible Flow", Gordon and Breach, New York (1969).
8. J.L. Lions, "Quelques methodes de resolution des problemes aux limites non lineaires", Dunod, Gauthier-Villars, Paris (1969).
9. J.L. Lions, E. Magenes "Problemes aux limites nonhomogenes et applications", Vol. I,II, Dunod, Paris (1968).
10. G. Prouse, A. Zaretti "On the Inequalities Associated to a Model of Graffi for the Motion of a Mixture of two viscous Incompressible Fluids", preprint Dipartimento di Matematica - Politecnico di Milano (1982).

11. P. Secchi "On the Initial Value Problem for the Equations of Motion of Viscous Incompressible Fluids in Presence of Diffusion", Boll. U.M.I., 1-3(1982) 1117-1130.
12. R. Temam "Navier-Stokes Equations", North Holland Amsterdam (1977).

HB/jgb

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER 2565	2. GOVT ACCESSION NO. A 134538	3. RECIPIENT'S CATALOG NUMBER	
4. TITLE (and Subtitle) Diffusion on Viscous Fluids, Existence and Asymptotic Properties of Solutions		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period	
		6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(s) H. Beirao-da-Veiga		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of Wisconsin 610 Walnut Street Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis	
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE September 1983	
		13. NUMBER OF PAGES 16	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Non-homogeneous viscous fluids, non-linear parabolic equations, initial-boundary value problems			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We consider the motion of a mixture of two fluids, with a diffusion effect obeying Fick's law; for the derivation of the model (1.1) see Section 1 and references [2], [4],[5] and [6]. We consider the full non-linear problem (i.e., we don't omit the term λ^2 term in equation (1.1 ₁)). Moreover, we don't assume that λ/μ is small. We prove the existence of a (unique) local			

ABSTRACT (Continued)

solution, the existence of a global solution for small data, and the exponential decay to the equilibrium solution; see Theorem A, Section 1.