

AD-A135 143

OPTICAL RADIO AND COMPUTER SIMULATION STUDIES OF THE
BRAZIL IONOSPHERIC M..(U) BOSTON UNIV MA DEPT. OF
ASTRONOMY J BAUMGARDNER ET AL. 31 JUL 83 SCIENTIFIC-1


1/1


UNCLASSIFIED

AFGL-TR-83-0196 F19628-81-K-0051

F/G 4/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

12

OPTICAL, RADIO AND COMPUTER SIMULATION STUDIES OF THE BRAZIL IONOSPHERIC MODIFICATION EXPERIMENT OF 8 SEPTEMBER 1982

AD-A135143

Jeffrey Baumgardner, Michael Mendillo, Bruce Herniter, and Harlan Spence

Department of Astronomy
Boston University
725 Commonwealth Avenue
Boston, Massachusetts 02215

Scientific Report No. 1
1 September 1982 - 30 June 1983

31 July 1983

Approved for public release; distribution unlimited.

AIR FORCE GEOPHYSICS LABORATORY
AIR FORCE SYSTEMS COMMAND
UNITED STATES AIR FORCE
HANSCOM AFB, MASSACHUSETTS 01731

SEARCHED
SERIALIZED
NOV 30 1983
S A


DTIC FILE COPY

83 11 28 242

This report has been reviewed by the ESD Public Affairs Office (PA) and is releasable to the National Technical Information Service (NTIS).

This technical report has been reviewed and is approved for publication.


JOHN A. KLOBUCHAR
Contract Manager


HERBERT C. CARLSON
Branch Chief

FOR THE COMMANDER


RITA C. SAGALYN
Division Director

Qualified requestors may obtain additional copies from the Defense Technical Information Center. All others should apply to the National Technical Information Service.

If your address has changed, or if you wish to be removed from the mailing list, or if the addressee is no longer employed by your organization, please notify AFGL/DAA/ Hanscom AFB, MA 01731. This will assist us in maintaining a current mailing list.

Do not return copies of this report unless contractual obligations or notices on a specific document requires that it be returned.

Unclassified

1.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFGL - TR-83-0196	2. GOVT ACCESSION NO. AD-A135143	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Optical, Radio and Computer Simulation Studies of the Brazil Ionospheric Modification Experiment of 8 September 1982		5. TYPE OF REPORT & PERIOD COVERED Scientific Report No. 1 1 Sept. 1982 - 30 June 1983
7. AUTHOR(s) Jeffrey Baumgardner, Michael Mendillo, Bruce Herniter, and Harlan Spence		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Astronomy Department 725 Commonwealth Ave. Boston, MA 02215		8. CONTRACT OR GRANT NUMBER(s) F-19628-81-K-0051
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Geophysics Laboratory Hanscom AFB, MA 01731 Contract Monitor: J.A. Klobuchar / PHY		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 62101F 464309AA ILIR2G
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE 31 July 1983
		13. NUMBER OF PAGES 32
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES This research was partially supported by the Air Force In-House Laboratory Independent Research Fund		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Ionospheric Modifications, Ionospheric Holes, Airglow Imaging, Total Electron Content, Computer Simulation		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) -The Brazil Ionospheric Modification Experiments (BIME) provided an opportunity to conduct simultaneous airglow and electron content observations during a space plasma "active experiments" campaign in Natal, Brazil, in September 1982. A period of background data was also obtained, and computer simulation studies were conducted for the background and artificially dis- turbed conditions. The BIME - 1 release produced a depletion in TEC of 1-1.5 TEC units that was accompanied by an airglow burst of approximately		

DD FORM 1473 1 JAN 73 EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

500 R at 6300 A. Wide-angle imaging at 6300 A revealed an airglow cloud that extended to 150 km in radius. Post mission simulation studies carried out using the Boston University Finite Element Simulation (FES) Code were in good agreement with the observations.



Accession For	
NTIS GRA&I	
DTIC TAB	
Unannounced	
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

1. OBSERVATIONS

The Boston University observing site at Jundiai (5.88°S , 35.3°W) was chosen using a nominal rocket trajectory and an average position for SIRIO of 0° , 15°W . With a release placed at 320 km, a ray path from Jundiai to the average position of SIRIO would have passed within 5 km of the center of the release. Using post flight trajectory data and an updated position for SIRIO, the actual miss distance for BIME-1 was ~ 40 km. This was mostly due to an error in the original position for SIRIO. Figure 1 illustrates the viewing geometry for Jundiai.

The instrumentation at Jundiai included: 1) a VHF Polarimeter, 2) a 1° field of view photon counting system using 7774, 6300, 6200, and 5577 \AA filters, 3) two all-sky intensified cameras operating at the same wavelengths as the photon counter.

The BIME-1 release occurred on 8 September at 21:45:20 UT (19:23 LMT). A 40 minute section of the Polarimeter record is shown in Figure 2. The top trace is the amplitude record and the bottom trace records Faraday rotation angle. The release is not evident on the amplitude record; however, it can be seen on the phase record as a change in slope of the normal decay in TEC. The release caused a 1 to 1.5 TEC unit drop to occur in about 5 minutes with some evidence for recovery at later times.

Figure 3 shows the recorded 6300 \AA airglow intensity (uncorrected for atmosphere transmission) obtained with the photon counter that was coaligned with the ray path to SIRIO. The pre-event level of ~ 40 Rayleighs was enhanced by a factor of about 15 to ~ 550 R.

The peak brightness occurred 3 minutes after the release and decayed to the pre-event background in about 20 minutes. The 6200 \AA channel recorded no change from pre-event levels as did the 7774 \AA channel. There is some weak evidence for a ~ 50 R enhancement in 5577. The background at 5577 may have been as high as




Figure 1. Observing Geometry




Figure 2. TEC and Amplitude Observations Using the SIRIO - Jundiai Ray Path During BIME-1, 8 September 1982.




Figure 3. 6300A Photometer Curve

~400 R masking this increase.

The intensified cameras were operated in 60° field of view modes for the first 4 minutes after the release. System I had 6300 Å and 6200 Å filters. System II used a 5577 filter for the first 4 minutes, then started sequencing through 6300, 6200, 5577 and 7774 Å. No effect was seen in this camera other than at 6300 Å. After ~4 minutes the airglow cloud was observed to be filling the 60° field of view of the cameras and the lenses were switched to 180° field of view. Two examples of 6300 Å photographs are shown in Figures 4 and 5 (Figure 5 also shows the Sonda Rocket). The brightness of the photographed cloud is consistent with the photometer measurements; the photographs show the cloud to be ~150 km in radius. Photographs taken about 1 hour after the release show weak evidence for a field aligned depletion in the 6300 Å background to the east of the release point.

The background ionosphere at Natal during the September 1982 BIME/Colored Bubbles campaign proved to be more stable than we had hoped, with episodes of natural spread F and scintillations increasing as the month progressed. On September 12, a well defined period of scintillation occurred with corresponding bite-outs in the TEC record.


Also on this night airglow structure was seen by the photometer as well as the more sensitive of the two all-sky cameras. Figure 6 shows the diurnal TEC curve for this night (the average behavior of TEC during the campaign is marked by X's). The two bite-outs in TEC are seen around 21:30 and 22:30. Figure 7 is an expanded view showing the 2 hours around the time when the depletions were passing through the polarimeter and photometer beams. Notice that the 7774 Å record very closely tracks the TEC record while the two 6300 Å channels appear to have little or no correlation to the TEC record.

All sky photographs were taken during this period and are shown in Figures 8, 9 and 10. Figure 8 is a photograph at 7774 that clearly shows 2 depletions--

W

N

S



E

6300A BIME 1
5 MINUTES AFTER RELEASE

Figure 4. All Sky Image At T=5min (from 35mm slide)



6300A BIME 1
7 MINUTES AFTER RELEASE

Figure 5. All Sky Image Showing SONDA (from a 35mm slide)






Figure 6. TEC Curve for September 12




7774A SEPTEMBER 12
21:54 U.T.

Figure 8. All Sky Image With 7774 Depletions (from a 35mm slide)



6300A SEPTEMBER 12
21:55 U.T.


Figure 9. All Sky Image With 6300 Structures (from a 35mm slide)



7774A SEPTEMBER 12

22:29 U.T.

Figure 10. All Sky Image K111-774 Reflections (From a 35mm slide)



6300A SEPTEMBER 14
22:40 U.T.

Figure 11. All Sky Image With 6300 Depletions (From 0.3deg. to 1.5 deg.)



6300A SEPTEMBER 18

01:07 U.T.

Figure 12. All Sky Image With 6300 Depletions (from a 35mm slide)

the one nearest the center of the field of view is the one that causes the bite-out in TEC at $\sim 21:50$. The depletions near the edge of the field of view must have formed to the east of the ray path to SIRIO and drifted farther to the east, as its signature does not appear on the polarimeter records.


Figure 9 is a 6300 \AA photograph taken just 1 minute after the 7774 exposure. Rather than a depletion there appears to be a ridge of 6300 \AA airglow between the two 7774 depletions. Figure 10 is another 7774 exposure taken about 40 minutes after Figure 8 (this is about the time that the second bite-out occurred in the TEC record). Other instances of optical depletions in 6300 \AA are shown in Figures 11 and 12. Out of the 13 nights that optical data were taken at Jundiai 5 nights showed evidence of depletions.

II. MODEL RESULTS

The geometry of the Boston University Finite Element Simulation (FES) Code is shown in Figure 13. The model uses cylindrical symmetry with 20 height bins and 10 radius bins. The allowed chemistry (reactants and products) for the BIME series of releases is shown on the lower left-hand side of Figure 13, while species that are allowed to diffuse are shown on the right. A complete description of the model is given elsewhere (Vance and Mendillo, 1981). The initial conditions for the BIME-1 simulation are outlined in Figure 14. The geomagnetic index $A_p=100$ was used for the input to the MSIS neutral atmosphere model which is the three-day average for 6-7-8 September. We feel that this high A_p index value was warranted due to the major magnetic storm that preceded the BIME-1 experiment. The baseline electron density profile used as input to the model was a composite profile from in-situ rocket, ionosonde and TEC data. The amount of released material available for chemistry in the simulation was 22 kg of H_2O and 19 kg of CO_2 . This represents 40% of the H_2O released and 100% of the CO_2 . This is probably a lower limit of the amount of H_2O available for chemistry -- the actual amount

Boston University

FINITE ELEMENT SIMULATION CODE



Chemistry

O_2 O^+
 N_2 O_2^+
 CO_2 NO^+
 H_2O H_2O^+
 E^-
 $O(^1D)$

Transport

H_2O O^+
 CO_2 O_2^+
 $O(^1D)$ NO^+
 H_2O^+
 E^-

Figure 13. Simulation Geometry

INITIAL CONDITIONS

- MSIS NEUTRAL ATMOSPHERE: EVALUATED FOR BIME 1 CONDITIONS
 $A_p=100$
- $N_e(h)$ PROFILE: OBSERVED BOTTOM SIDE PROFILE (BUCHAU)
 TOPSIDE DERIVED TO FIT TEC (KLOBUCHAR)
- RELEASE: 60% H_2O FROZEN, ALL CO_2 AVAILABLE (BEST FIT TO
 LAGOPEDO RESULTS)
- REACTION RATES: ALL FROM RECENT REVIEW ARTICLES
- AIRGLOW PRODUCTION: $O_2^+ + e^- \rightarrow kO^* + (2-k)O$
 (6300A FROM CO_2 CHAIN) $k=1.0$ (REF. $0.9 \leq k \leq 1.33$)
 (NO 6300A FROM H_2O CHAIN)
 QUENCHING OF O^* FROM $N_2, O_2, CO_2, H_2O, e^-$

Figure 14. Initial Conditions

may be as much as twice this value. The effect of increasing the amount of H_2O available for chemistry would be to increase the size and depth of the TEC "hole" but have little or no effect on the airglow since all of the modification-induced 6300 \AA airglow in the model comes from reactions initiated by the CO_2 .

The results of a 10 minute simulation of the BIME-1 release are summarized in Figures 15 through 24. It can be seen from Figures 15 and 16 that the release steepened the bottom side gradient and that the effect of the release did not extend much above 360 km. In characterizing the maximum effect of the release on electron densities, two plots were prepared: Figure 17 shows the electron density as a function of radius from the release point at the release height of 320 km. Figure 18 shows the same data but in terms of percent change from baseline conditions. From the second figure it can be seen that the maximum effect was a $\sim 70\%$ depletion in electron density at the release point and the radius at which there was only a 1% effect was ~ 100 km.

The effect of the release on the TEC is shown in Figures 19 and 20. The maximum change in TEC from Figure 19 is ~ 1.5 TEC units. Figure 20 shows that at 80 km radius the effect on TEC is about 1%--similar to the size of the "hole" in electron density at the release height.

The effect of the release on 6300 \AA airglow is summarized in the remaining figures. The baseline volume emission profile (O^1D concentration) and the effect of the release is shown in Figure 21. Figure 22 shows the time history of 6300 \AA airglow at different radii from the release point. It can be seen that the model predicts a fairly strong gradient in brightness at early times--a factor of four within 60 km of the cloud center. This is somewhat steeper than was observed by the all-sky cameras. The size with the airglow cloud predicted by the release is shown in Figures 23 and 24. Figure 24 shows that the radius at which the percent change in airglow is $\sim 1\%$ is an excess of 150 km, and that at 80 km where the per-

BOSTON UNIVERSITY FES CODE RESULTS: BIME-1




Figure 15. Electron Profile at T=5 min, R=0

BOSTON UNIVERSITY FES CODE RESULTS: BIME-1




Figure 16. Electron Profile at T=10 min, R=0

BOSTON UNIVERSITY FES CODE RESULTS: BIME-1




Figure 17. Electron Density vs. R at H= 326 km.

BOSTON UNIVERSITY FES CODE RESULTS: BIME-1





Figure 18. Percent Change in Electron Density vs. Radius

BOSTON UNIVERSITY FES CODE RESULTS: BIME-1



Total Electron Content

Figure 19, TEC vs. Time

BOSTON UNIVERSITY FES CODE RESULTS: BIME-1




Figure 20. Percent Change in TEC vs. Radius

BOSTON UNIVERSITY FES CODE RESULTS: BIME--1




Figure 21. O(¹D) Concentration vs. H

BOSTON UNIVERSITY FES CODE RESULTS: BIME-1




Figure 22. 6300 Airglow vs. Time

BOSTON UNIVERSITY FES CODE RESULTS: BIME-1




Figure 23. 6300 Airglow vs.

BOSTON UNIVERSITY FES CODE RESULTS: BIME-1




Figure 24. Percent Change in Airglow Vs. Radius

cent change in TEC was $\sim 1\%$ the percent change in airglow is $\sim 50\%$.

A more detailed reduction of the BIME-1 airglow images and a refined set of simulation runs are currently in progress.

Reference

Vance, B., and Mendillo, M., Computer Simulation Techniques for Artificial Modification of the Ionosphere, AFGL-TR-81-0119, Air Force Geophysics Laboratory, Hanscom AFB, MA, April 1981. ADA102302

Acknowledgements

We wish to express our appreciation to a great many individuals who contributed to the success of our participation in the BIME campaign. At the Air Force Geophysics Laboratory, J.A. Klobuchar (Contract Monitor) provided valuable assistance in site selection, polarimeter operations, radio communications and overall guidance throughout the course of the project. Dr. E. Weber provided the critical final support via the LDF program that allowed our participation in the program. In addition, he provided valuable technical advice and equipment transportation support. Mr John Lloyd and Dr. J. Bucheau offered help and encouragement in many ways. In Brazil, Prof. Liacir Lucena of the Federal University of Rio Grande of the North was a gracious host and provided for all of our local needs at the University's Field Station in Jundiai. The hospitality of the Brazilian people and government officials were a most valuable resource throughout the campaign. At Boston University, Prof. K. Janes (Chairman of the Astronomy Department) kindly allowed us to take the Observatory's photometer and computer control system for use in the experiments.

**ATE
LME**