

HD-A136 308

PRODUCTS OF PHOTODEGRADATION FOR COUMARIN LASER DYES
(U) BOSTON UNIV MA DEPT OF CHEMISTRY G JONES ET AL.
31 OCT 83 TR-7 N00014-79-C-0054

1/1


UNCLASSIFIED

F/G 11/3

NL

END

1/1
1/1
1/1



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

12

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 7	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Products of Photodegradation for Coumarin Laser Dyes		5. TYPE OF REPORT & PERIOD COVERED Technical, 1/1/82-10/31/82
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) G. Jones, II, W.R. Jackson, S. Kanoktanaporn, and W.R. Bergmark		8. CONTRACT OR GRANT NUMBER(s) N00014-79-C-0054
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Boston University Boston, Massachusetts 02215		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 395-609
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Eastern/Central Regional Office 666 Summer Street, Boston MA 02210		12. REPORT DATE October 31, 1983
		13. NUMBER OF PAGES 14
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Reproduction in whole or in part is permitted for any purpose of the United States Government. Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Prepared for publication in Optics Communications.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) coumarin laser dyes, laser dye photodegradation, dye photolysis mechanisms, radical intermediates, laser dye triplet yields		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Photolysis products of the coumarin dyes have been investigated. Amine group dealkylation is observed for fluorinated coumarins. A reduction product, consistent with a photolysis mechanism involving free radicals, is found for coumarin 1. A new type of photooxidation product is identified for the fluorinated rigid dye, C153, in aerated media.		

DTIC
ELECTE
DEC 27 1983
S H D

A136308

DMC FILE COPY

DD FORM 1473 EDITION OF 1 NOV 65 IS OBSOLETE
1 JAN 73 S/N 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

83

50

PRODUCTS OF PHOTODEGRADATION FOR COUMARIN LASER DYES

Guilford Jones, II,* W. R. Bergmark, and W. R. Jackson

Department of Chemistry, Boston University, Boston MA 02215 USA

Abstract

↓
The products of conventional photolysis of the coumarin laser dyes, C1, C35, C153, and C152 have been investigated. The previously reported dealkylation of C1 is documented for the fluorinated dyes, C35, and C152 in deaerated solvents. In addition, a reduction product is identified for C1, consistent with a radical mechanism for decomposition. Evidence is provided that the concentration quenching (self quenching) of singlet dye is important to the degradation mechanism. For the rigid dye, C153, a photooxidation product involving the amine functionality results from decomposition in aerated media. For several dyes, very low triplet yields have been measured.
↑



Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

The stability of coumarin dyes during dye laser operation has been of interest for some years.^{1,2} The effects of medium, cover gas, dye structure, and optical filters have been investigated extensively.³ Quantum efficiencies for photodecomposition have been reported in rare cases,^{2a,4} and the effects of additives which improve performance under lasing conditions have been documented.⁵ The products of photodegradation for a single coumarin dye have been identified in the early report of Winters, Mandelberg, and Mohr.⁶ We wish to establish that the dealkylation process noted by the latter authors for C1 is also important for other related dyes. Furthermore, another type of product accompanying dealkylation and characteristic of radical disproportionation has been identified for C1. In addition, the decomposition of a dye having a "rigid" structure (C153), which is known for its stability,¹ has also been investigated.

Product studies. Dyes chosen for study are the 7-aminocoumarins shown below which are commercially available ("laser grade") from Eastman (Cxx numbers as shown) or from Exciton Chemical Co. (1a = C460, 1c = C485, 2 = C540A). Solutions of the dyes were photolyzed using either a 450 W Hanovia medium pressure Hg lamp, immersion well apparatus, and a Pyrex glass filter (>300 nm) or, for comparative runs, a Rayonet chamber reactor in which 13 x 150 mm Pyrex test tubes were irradiated.⁷ The samples were purged with nitrogen or argon except where noted. Since higher concentrations of dye led to more rapid disappearance (vide infra), relatively concentrated solutions were generally employed for product studies (10^{-2} - 10^{-3} M). Lengthy irradiation resulted in loss of dye which could be followed by uv-vis absorption or nuclear magnetic resonance (nmr) spectrometry. A general result was the deposition of a film which was difficult to characterize

(i.e., nmr analysis revealed mainly dye disappearance). However, small quantities of identifiable products were apparent from glc analysis (2m 10% OV-101 column, 200°) on photolysis of 1 in acetonitrile. Product 2a was identified by comparison of reported spectral data;⁶ the analogous 2b was obtained after column chromatography of the photoproduct (silica gel, 5% ethyl acetate - hexane eluent) followed by recrystallization from toluene: fine yellow crystals, mp 158-159° (10% isolated yield). Elemental analysis for $C_{11}F_3H_4NO_2$ was: 56.21 %C, 3.98 %H, and 5.40 %N (found); 56.04 %C, 3.92 %H, and 5.45 %N (theory). The spectral data consistent with structure 1b were as follows:

uv (methanol): $\lambda_{max} = 390 \text{ nm}$

ir (film): 3370, 2990, 1725, 1645, 1620, 1135, 1170, 1230
and 1300 cm^{-1}

nmr ($CDCl_3$): δ 7.40 (m, arom., 1H), 6.45 (m, arom. and
-CH), 4.40 (broad m, N-H), 3.28 (q, $-CH_2$, 2H),
and 1.20 (t, $-CH_3$, 3H)

Another product originating from 1a could not be isolated but was identified as 3 by glc - mass spectral analysis. The major ms peaks and assignments were as follows:

ms: m/e 233 (mol. ion), 218 ($-CH_3$), 190 ($-CH_3$, $-CO$ or
 C_8H_4), and 148 ($190 - CH_3CO$)

Although 2a and the previously unidentified 3 failed to accumulate appreciably with photolysis time (ca. 10% maximum), their appearance proved interesting in that 3 was favored at the shortest times possible for glc detection whereas 2a was the more important volatile component after lengthy irradiation periods. Confirming the rather erratic nature of photodegradation, the disappearance of native dye appeared to be accelerated following an induction period in which about 10% of the starting dye was decomposed. Glc analysis also revealed small quantities of the fully dealkylated dyes (identified by direct comparison with the commercially available C120 and C151 (4) from 2a⁶ and 2b, respectively).

Coumarin 1c was also investigated to establish the dependence of photodealkylation on substitution at the amino group. The rate of decomposition in deaerated acetonitrile was in fact comparable to that of 1b and products of dealkylation were apparent from nmr and glc analysis (only the common product 4 was thoroughly identified). Photodecomposition of coumarins 1 as a function of solvent was followed by glc analysis of dealkylation product at relatively low conversion (anthracene used as an internal standard). Parallel irradiation showed similar results for acetonitrile, toluene, and cyclohexane with some acceleration in chloroform (ca. 3-fold) and a retardation in alcohols (>10-fold).

Photolysis of deaerated acetonitrile solutions of coumarin 153 (7) led to decomposition (significantly slower than 1) to intractable material with no appearance of an identifiable monomeric product. However, on photolysis of air-saturated solutions, a photoproduct was obtained after preparative thin-layer chromatography (Analtech 2000 μ silica gel (GF) plates, 3:2 hex-

ane - ethyl acetate eluent). The product ($R_f = 0.5$), a solid with reddish coloration recrystallized from methanol/water, mp 193-207°. Elemental analysis provided the following data: for $C_{16}F_3H_{11}NO_3$, %C, 59.38, %H, 3.75, %N, 4.29 (found); %C, 59.44, %H, 3.72, %N, 4.33 (theory). The spectral data were consistent with formation of a pair of photooxidation products, the isomeric amides, 3 and 2, which crystallized as a mixture.

uv (ethanol): λ_{max} 353 nm

ir (film): 1735, 1690, 1615, 1415, 1370, 1295, 1135 cm^{-1}

nmr ($CDCl_3$): δ 7.40 (s, arom., 1H), 6.70 (s, vinyl, 1H),
3.80 (m, N-CH₂, 2H), 2.80 (m, -CH₂, 2H),
1.1-2.2 (m, CH₂, 6H)

Mechanistic studies. For the purposes of establishing a mechanism of photodegradation, the effect of adding a triplet quencher was investigated. With 0.1 M 1,3-cyclohexadiene (CHD) or with 0.06 M trans-stilbene, neither formation of products, 2a and 3, nor the rate of net destruction of 1a was altered. As a control the quenching of dye triplets by these additives was investigated by flash photolysis. The triplet of 1a was observed (λ_{max} 625 nm) as previously reported⁸ on flash lamp irradiation in argon-purged acetonitrile (Xe source,⁹ pulse duration = ca. 30 μs FWHM). This transient was absent on the addition of triplet quenchers.

Although the quenching results for 1a did not favor a photolysis mechanism involving dye triplets, the appearance of triplets for other dyes

remained of interest. Intersystem crossing yields were measured for three dyes in two solvents using the triplet counting technique in which the photodimerization of CHD⁹ via energy transfer from dye was followed with reference to dimerization sensitized by benzophenone (whose triplet yield (1.0) is known)(Table).¹¹

With a singlet mechanism for photodegradation in mind, dependences on dye concentration were investigated. In fact the appearance of 3 at low conversion (ca. 0.1%) was noticeably accelerated at high dye concentrations (0.01-0.25 M). The profile of concentration dependence is consistent with a singlet self-quenching mechanism according to Scheme 1. Along with values for singlet lifetime for 1a in very dilute solution ($\tau = 1 / (k_f + k_d)$),¹¹ the data provide a source for the rate constant sum, $k_{oq} + k_r = 7.8 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ referring to the sum of rates for concentration quenching and bimolecular reaction. A similar value for these rates ($2.9 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$) has been independently obtained by measuring the concentration quenching of fluorescence as reported separately for several coumarin dyes.¹² The plot of quantum yield data for formation of 3 (eqn 1) provided an estimate of the maximum (extrapolated) yield of product 3, $(k_r + k_{oq})/k_r = 3 \times 10^{-4}$, which showed that the photoreduction of 1a constitutes a small part of total (unproductive) concentration quenching.

Summary. The photodealkylation reaction for 7-dialkylaminocoumarins in deaerated media has been extended to fluorinated analogs, 1b and 1c. Another type of product, whose structure has been tentatively assigned to photoreduction product 3, can be ascribed to formation of radicals 5 and 6 as previously proposed by von Trebra and Koch.¹³ This type of intermediate

may arise from bimolecular quenching by ground state dye (self reaction of dye or concentration quenching) via rather inefficient hydrogen atom abstraction (or the equivalent steps of electron transfer and proton transfer). Radical 6 is capable of disproportionation to reduction product 3 and starting dye and the initiation of dye polymerization, whereas 5 is a likely intermediate for dealkylation either via coupling reaction with 6¹³ or by way of further electron/proton transfer.

The rigid coumarin, 7, is less photoactive in oxygen-free solutions than the analogs, 1, presumably due to low productivity in radical formation from its less energetic excited singlet (concentration quenching is nonetheless observed¹³). On the other hand, photooxidation in aerated solution is important. Formation of products 8 and 9 is consistent with the quenching of dye singlets by oxygen, a step which has been documented,^{11b,14} and formation of radicals (via hydrogen or electron/proton transfer). Photodecomposition proceeds as shown in Scheme 2 for reaction at a position adjacent to nitrogen, reminiscent of the path which has been proposed for oxidation of tertiary amines.¹⁵ The reaction observed for the amine functional group contrasts with the reported photooxidation of 1a⁶ (reaction at the 4-methyl group) and portends a more general destructive role for oxygen in the aminocoumarin series (a similar rate of degradation for a variety of coumarins under lasing conditions for air-saturated solutions has been reported.⁵

The results overall point to the importance of dye self quenching in photodegradation, a phenomenon which is alleviated but not entirely eliminated at low but practical dye concentrations. Alcohol (protic) solvents

reduce photodecomposition (but not concentration quenching ¹³) presumably by inhibition of radical formation through hydrogen-bonding to the coumarin carbonyl group, a particularly effective interaction for the excited singlet dye.¹¹ The potential for destruction by oxygen in aerated samples is also reiterated.^{3, 8b, 6, 16} However, it should be noted that for the general case of laser dye stability the presence of oxygen is not necessarily defeating if there is a significant role for dye triplets in photodecomposition or in competitive absorption at lasing wavelengths.

Acknowledgements. Support of this research by the U. S. Office of Naval Research is gratefully acknowledged.

References


1. Reynolds, G. A.; Drexhage, K. H. Optics Commun. 1975, 13, 222.
2. a. Schimitschek, E. J.; Trias, J. A.; Hammond, P. R.; Atkins, R. L. Optics Commun. 1974, 11, 352; b. Schimitschek, E. J.; Trias, J. A.; Hammond, P. R.; Henry, R. A.; Atkins, R. L. ibid. 1976, 16, 313
3. a. Fletcher, A. N. Appl. Phys. B 1983, 31, 19; b. Fletcher, A. N.; Bliss, D. E. ibid. 1978, 16, 289.
4. Gauglitz, G. Z. Phys. Chem. (Wiesbaden) 1978, 113, 217.
5. a. von Trebra, R. J.; Koch, T. H. Appl. Phys. Lett. 1983, 42, 129; b. von Trebra, R. J.; Koch, T. H. Chem. Phys. Lett. 1982, 93, 315.
6. Winters, B. H.; Mandelberg, H. I.; Mohr, W. B. Appl. Phys. Lett. 1974, 25, 723.
7. Jones, II, G.; Santhanam, M.; Chiang, S. H. J. Am. Chem. Soc. 1980, 102, 6088.
8. Dempster, D. N.; Morrow, T.; Quinn, M. F. J. Photochem. 1973/74, 2, 29.
9. Caspari, G.; Hughes, R. G.; Endicott, J. F.; Hoffman, M. Z. J. Am. Chem. Soc. 1970, 92, 6801.

10. Vesley, G. F.; Hammond, G. S. Mol. Photochem. 1973, 5, 367.
11. For discussion of coumarin singlet decay in lieu of intersystem crossing and medium effects thereon, see (a) Jones, G. II; Jackson, W. R.; Halpern, A. M. Chem. Phys. Lett. 1980, 72, 391; (b) Jones, G. II; Jackson, W. R.; Choi, C.; Bergmark, W. R. J. Am. Chem. Soc. submitted.
12. Jones, G. II; Bergmark, W. R. J. Photochem. submitted.
13. Wamser, C. C.; Hammond, G. S.; Chang, C. T.; Baylor, C. J. Am. Chem. Soc. 1970, 92, 6362.
14. Rubin, R. F.; Fletcher, A. N. Chem. Phys. Lett. 1983, 99, 49.
15. Bartholomew, R. F.; Davidson, R. S.; Howell, M. J. J. Chem. Soc. (C) 1971, 2804.
16. Hammond, P. R.; Fletcher, A. N.; Bliss, D. E.; Henry, R. A.; Atkins, R. L.; Moore, D. W. Appl. Phys. 1976, 2, 67.


Table. Intersystem crossing quantum yields for coumarin dyes.


	$\Phi_t(\text{C}_6\text{H}_{12})$	$\Phi_t(\text{C}_2\text{H}_5\text{OH})$
<u>1a</u>	0.30	0.006
<u>1b</u>	0.043	<0.001
<u>1</u>	0.053	<0.001

Scheme 1



Scheme 2






1a, $R = C_2H_5$, $R' = CH_3$ [C1]


1b, $R = C_2H_5$, $R' = CF_3$ [C35]

1c, $R = CH_3$, $R' = CF_3$ [C152]


2




3




4




5




6



7 (C153)



8



9

FILM

2-8