

AD-A136 405


LINEAR COMPLEMENTARITY PROBLEMS SOLVABLE BY AN  
EFFICIENT POLYNOMIALLY BOUND (U) WISCONSIN UNIV-MADISON  
MATHEMATICS RESEARCH CENTER J PANG NOV 83 MRC-TSR-2597  
DAAG29-80-C-0041


1/1

UNCLASSIFIED

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

4

A136405

MRC Technical Summary Report #2597

LINEAR COMPLEMENTARITY PROBLEMS  
SOLVABLE BY AN EFFICIENT POLYNOMIALLY  
BOUNDED PIVOTING ALGORITHM

Jong-Shi Pang

**Mathematics Research Center  
University of Wisconsin—Madison  
610 Walnut Street  
Madison, Wisconsin 53705**

November 1983

(Received September 29, 1983)

DTIC  
SELECTED  
NOV 23 1983  
A

DTIC FILE COPY

Approved for public release  
Distribution unlimited

Sponsored by  
U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

83 12 27 066

UNIVERSITY OF WISCONSIN-MADISON  
MATHEMATICS RESEARCH CENTER

LINEAR COMPLEMENTARITY PROBLEMS SOLVABLE BY AN EFFICIENT  
POLYNOMIALLY BOUNDED PIVOTING ALGORITHM

Jong-Shi Pang\*

Technical Summary Report #2597  
November 1983

ABSTRACT

Applied to two important classes of linear complementarity problems defined by an  $n \times n$  matrix, the parametric principal pivoting algorithm, using a suitably chosen (and easily computable) parametric vector, terminates with a desired solution after at most  $n$  pivot operations. Since each pivot can be performed using at most  $O(n^2)$  arithmetic operations, the total computational complexity of the algorithm for solving these linear complementarity problems is no more than  $O(n^3)$ . In one of the two classes of problems being studied, the complexity is  $O(n^2)$  because the matrix involved is 5-diagonal which allows each pivot to be performed in linear time. Some discussion in connection with Lemke's well-known almost complementary pivoting algorithm is also addressed.

AMS (MOS) Subject Classification: 90C20, 90C33

Key Words: Linear complementarity, parametric principal pivoting, efficient algorithm, polynomially bounded, concave regression, hidden Z, diagonally dominant

Work Unit Number 5 - Optimization and Large Scale Systems

\*School of Management and Administration, The University of Texas at Dallas, Box 688, Richardson, TX 75080, U.S.A.

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.

**SIGNIFICANCE AND EXPLANATION**

Many well-known pivoting methods for solving the linear complementarity problem have been shown to exhibit an exponential computational complexity. In this paper, we identify two classes of linear complementarity problems which can be solved by a numerically efficient as well as polynomially bounded pivoting algorithm. One of these classes of problems arises from the concave regression problem which has practical applications in statistics.



*[Faint, mostly illegible text and markings, possibly a stamp or administrative notes, with a large handwritten mark resembling a stylized 'A' or '7' at the top right.]*

---

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

LINEAR COMPLEMENTARITY PROBLEMS SOLVABLE BY AN EFFICIENT  
POLYNOMIALLY BOUNDED PIVOTING ALGORITHM

Jong-Shi Pang\*

1. INTRODUCTION.

It is known that the general linear complementarity problem defined by an arbitrary matrix is NP-complete [5]. Hence, it is unlikely that there will be a polynomially bounded algorithm for solving an arbitrary linear complementarity problem. With respect to several most notable pivoting methods, examples of problems have been constructed which show that these methods can require an exponential number of pivots [2, 12, 22]. These cited studies are all theoretical in nature and provide the worst-case analysis of the linear complementarity problem. From a practical point of view, it is more desirable to be able to identify classes of problems (with applications) which are solvable by an algorithm that is both numerically efficient as well as polynomially bounded. Except for the trivial ones (e.g. those defined by triangular matrices) the class of linear complementarity problems with a Z-matrix is perhaps the best known member belonging to such a category [4, 30]. (Another related class can be found in [25].) These latter complementarity problems have applications in the numerical solution of free boundary value problems, optimal stopping, isotonic regression and others.

In this paper, we identify two classes of linear complementarity problems and show that with a suitably chosen (and easily computable) parametric vector, the parametric principal pivoting algorithm [6, 27] will compute a solution after at most  $n$  pivot iterations where  $n$  is the order of the matrix defining the complementarity problems. One such class arises from the concave regression problem which has practical applications in statistical regression analysis. The latter problem was first formulated by Hildreth [13] for the estimation of marginal productivity curves and is concerned, in general, with finding a least-square estimate of a certain functional relationship between some dependent

---

\*School of Management and Administration, The University of Texas at Dallas, Box 688, Richardson, TX 75080, U.S.A.

---

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.

and independent variables which is known to be concave. Recent references on the concave regression problem include [11, 14, 31]. The resulting class of linear complementarity problems is defined by a 5-diagonal symmetric positive definite but non-Z matrix. The 5-diagonal structure greatly simplifies the practical implementation of the parametric principal pivoting algorithm which results in an efficient  $O(n^2)$  method for solving the concave regression problem.

The primary reason why a great deal of emphasis is placed on the concave regression problem is because it is this practical problem which has challenged us to develop an algorithm that is both numerically efficient and polynomially bounded and has subsequently led us into this entire study. The desire to derive such a fast algorithm is in turn motivated by a close relative of the concave regression problem, namely, the isotonic regression problem. This latter problem has been well studied in statistics [1] and a linear-time algorithm has been developed [17, 31]. From a complementarity point of view, the isotonic regression problem can be formulated as a linear complementarity problem with a tridiagonal Stieltjes matrix and thus its solution by a linear-time algorithm is to be expected [4, 9, 10]. On the other hand, as we shall see, the matrix defining the linear complementarity problem arising from the concave regression problem is not Z.

The other class of linear complementarity problems identified in this paper consists of those defined by a matrix whose transpose is hidden Minkowski. A hidden Minkowski matrix is a P-matrix which is hidden Z. The class of hidden Z-matrices was introduced by Mangasarian [20] in connection with the study of solving linear complementarity problems as linear programs (see also [8]). The name "hidden Z" was coined because of the relation to a hidden Leontief matrix [23]. Specifically, a matrix M is hidden Z if there exist Z-matrices X and Y such that the two conditions below are satisfied

$$MX = Y \quad (1)$$

$$r^T X + s^T Y > 0 \text{ for some } r, s \geq 0. \quad (2)$$

Obviously, a Z-matrix is hidden Z. Many basic properties of a hidden Minkowski matrix have been obtained in [24]. In particular, it is known that an H-matrix with positive diagonals is hidden Minkowski. (A matrix M is H if its comparison matrix  $\bar{M}$  defined

by

$$\bar{M}_{ij} = \begin{cases} -|M_{ij}| & \text{if } i \neq j \\ |M_{ii}| & \text{if } i = j \end{cases}$$

is P.) Examples of H-matrices include the column (or row) strictly diagonally dominant matrices and of course the Minkowski matrices. Other examples of hidden Minkowski matrices can be found in [24]. Since the transpose of an H-matrix is obviously an H-matrix, it follows from our analysis that a linear complementarity problem defined by an H-matrix with positive diagonals can be solved by an effective  $O(n^3)$  pivoting algorithm. In particular, so can a problem with a strictly diagonally dominant matrix having positive diagonal entries.

Although Mangasarian [20] has shown that a linear complementarity problem with a hidden Z-matrix can be solved by a linear program (using an easily computable objective function), the only polynomially bounded algorithm for solving a general linear program [16] is known to be numerically inefficient [3]. The parametric principal pivoting algorithm, on the other hand, has been shown to exhibit good numerical performance even on problems of fairly large size [28, 29].

The question of efficiently identifying whether an arbitrary matrix is hidden Z remains unsolved. (Basically, the difficulty has to do with the nonlinearity of the second defining condition (2).) However, that of checking if a matrix is hidden Minkowski can be effectively answered by solving two linear programs [26]. The procedure described in the cited reference will find the two Z-matrices X and Y satisfying (1) and (2) if M is indeed hidden Minkowski. For certain subclasses (like the H-matrices with positive diagonals), the matrices X and Y can be obtained easily without solving any linear program (see [8, 20, 24] for more such subclasses). Related to this discussion is the open question of whether the transpose of a hidden Z-matrix is hidden Z.

The organization of the remainder of this paper is as follows. In the next section, we give a quick review of the parametric principal pivoting algorithm for solving a linear complementarity problem with an  $n \times n$  P-matrix and state a general (sufficient) condition under which the algorithm will terminate with the desired solution after at most  $n$

pivots. We also discuss an efficient implementation of the algorithm when the condition is satisfied and establish its  $O(n^3)$  computational complexity. Section 3 deals with the concave regression problem. We show how this practical problem can be formulated as a special linear complementarity problem and demonstrate that the condition guaranteeing the linear termination of the parametric principal pivoting algorithm is indeed satisfied by this special problem. In Section 4, we study a general linear complementarity problem defined by a P-matrix whose transpose is hidden  $Z$  and show that the same sufficient condition on the parametric principal pivoting algorithm is also satisfied by this class of problems. Finally, in the fifth and last section, we extend our discussion to Lemke's almost complementary pivoting algorithm [18, 19] and show that the same condition given in Section 1 is also sufficient for Lemke's method to terminate in at most  $n + 1$  pivots with a desired solution when applied to a general linear complementarity problem with an  $n \times n$  nondegenerate matrix.

## 2. THE PARAMETRIC PRINCIPAL PIVOTING ALGORITHM

We find it useful to quickly review the parametric principal pivoting algorithm in terms of its practical implementation [6, 27]. Given the linear complementarity problem (LCP)

$$y = q + Mx \geq 0, \quad x \geq 0, \quad y^T x = 0 \quad (3)$$

where the matrix  $M$  is  $P$ , we augment it by a parametric vector  $p$  and consider the parametric LCP

$$y = q + \theta p + Mx \geq 0, \quad x \geq 0, \quad y^T x = 0$$

where  $\theta$  is a parameter to be driven to zero. The vector  $p$  is chosen positive. Assume that several (principal) pivots have been performed. Let  $L(K)$  denote the currently basic (nonbasic)  $x$ -variables. (Initially,  $L = \phi$ .) With respect to these index sets, the canonical tableau may be written in the form

$$\begin{array}{c} \theta \quad y_L \quad x_K \\ x_L = \begin{array}{|c|c|c|c|} \hline \bar{q}_L & \bar{p}_L & M_{LL}^{-1} & -M_{LL}^{-1}M_{LK} \\ \hline \end{array} \\ y_K = \begin{array}{|c|c|c|c|} \hline \bar{q}_K & \bar{p}_K & M_{KL}M_{LL}^{-1} & (M/M_{LL}) \\ \hline \end{array} \end{array}$$

where  $(\bar{q}_L, \bar{p}_L)$  is the (unique) solution to the system of linear equations

$$M_{LL}(\bar{q}_L, \bar{p}_L) = -(q_L, p_L)$$

and

$$(\bar{q}_K, \bar{p}_K) = (q_K, p_K) + M_{KL}(\bar{q}_L, \bar{p}_L),$$

and where  $(M/M_{LL})$  denotes the Schur complement of  $M_{LL}$  in  $M$ :

$$(M/M_{LL}) = M_{KK} - M_{KL}M_{LL}^{-1}M_{LK}.$$

(See [7] for various properties of the Schur complement.)

To determine the next pivot, the ratio test is performed:

$$\bar{\theta} = \max\{-\bar{q}_i/\bar{p}_i : \bar{p}_i > 0\}. \quad (4)$$

If  $\bar{p} \leq 0$  or  $\bar{\theta} \leq 0$ , then the desired solution to the original LCP(3) is obtained as

$$x^* = (\bar{q}_L, 0).$$

Otherwise, let  $k$  be a maximizing index in (4). Update the index set  $L$  (and its

complement  $K$ ) by the rule:

$$L_{\text{new}} = \begin{cases} L_{\text{old}} \setminus \{k\} & \text{if } k \in L_{\text{old}} \\ L_{\text{old}} \cup \{k\} & \text{if } k \notin L_{\text{old}}. \end{cases}$$

This completes one (pivot) iteration of the algorithm.

Now suppose that the vector  $p$  is chosen so that

$$M_{LL}^{-1} p_L \geq 0 \text{ for all } L. \quad (5)$$

Then the maximizing index  $k$  can never occur in  $L_{\text{old}}$ . Thus the cardinality of the set  $L$  is increasing by one at each pivot. Consequently, unless the algorithm has already terminated (with a desired solution), it will continue until the complement  $K$  reaches empty, at which point, we have  $\bar{p} = -M^{-1}p \leq 0$  and the algorithm terminates. We have therefore proved

**Theorem 1.** Let  $M$  be a  $P$ -matrix of order  $n$ . If there exists a vector  $p > 0$  such that condition (5) holds, then the parametric principal pivoting algorithm, using  $p$  as the parametric vector, will compute a solution to the LCP(3) in at most  $n$  pivots.

**Remarks.** (i) No nondegeneracy assumption is needed in Theorem 1.

(ii) Underlying the proof of Theorem 1 is the key idea that once an  $x$ -variable becomes basic, it will stay basic until termination. Loosely stated, Theorem 1 asserts that all the basic  $x$ -variables can be identified in no more than  $n$  pivot steps. Of course, once those variables are determined, the desired solution to (3) is readily obtained.

Based on the idea pointed out in Remark (ii) above, it is possible to simplify the implementation of the parametric principal pivoting algorithm. Indeed, since a basic  $x$ -variable can not become nonbasic again, we need to keep track of the nonbasic components  $(\bar{q}_K, \bar{p}_K)$  only and restrict the ratio comparisons (4) to such components. Moreover, exploiting the fact that the index set  $K$  decreases by one element  $k$  (the maximizing index) at each pivot, we may update these nonbasic components by the following recursive formula which does not require the knowledge of the basic components:

$$(\bar{q}_{K'}, \bar{p}_{K'})_{\text{new}} = (\bar{q}_{K'}, \bar{p}_{K'})_{\text{old}} + (M_{K'L} - M_{K'L} \bar{M}_{Lk}) (\bar{q}_k, \bar{p}_k) \quad (61)$$

where  $K' = K \setminus \{k\}$  and  $\bar{M}_{Lk}$  is the solution to the system of linear equations

$$M_{LL} \bar{M}_{Lk} = M_{Lk} \quad (6ii)$$

(The formulas (6i) and (6ii) are easy to verify.)

Summarizing the discussion, we give below a step-by-step implementation of the parametric principal pivoting algorithm for solving the LCP(3), assuming that a vector  $p$  has been chosen as specified in Theorem 1.

Step 0. (Initialization) Set  $L = \emptyset$  and  $K = \{1, \dots, n\}$ . Set  $\bar{p} = p$  and  $\bar{q} = q$ .

Step 1. (Termination Test) Determine the critical value

$$\bar{\theta} = \max\{-\bar{q}_i / \bar{p}_i : i \in K \text{ and } \bar{p}_i > 0\}. \quad (4)'$$

If  $\bar{p}_k \leq 0$  or  $\bar{\theta} \leq 0$ , go to Step 3. Otherwise, let  $k \in K$  be a maximizing index.

Step 2. (Update of the Nonbasic Components) If  $K = \{k\}$ , set  $K = \emptyset$  and  $L = \{1, \dots, n\}$ , go to Step 3. Otherwise, solve the system of linear equations (6ii) and compute (6i).

Set  $K = K'$  and  $L = L \cup \{k\}$ . Go to Step 1.

Step 3. (Output of Solution) Solve the system of linear equations for  $x_L^*$ :

$$M_{LL} x_L^* = -q_L.$$

The vector  $x^* = (x_L^*, 0)$  is the unique solution to the LCP (3). Stop

The success of the above algorithm clearly hinges on the ability to find the crucial vector  $p$ . Continuing to assume that it is available, we analyze the complexity of the algorithm. Step 1 requires  $O(n)$  comparisons. By using an adaptive matrix factorization (such as LU) scheme, the system (6ii) can be solved in  $O(n^2)$  time, so can the one in Step 3. The computations in (6i) can be achieved in the same amount of time. Since the algorithm will terminate after at most  $n$  passes through Steps 1 and 2 and since the storage requirement is obviously no more than  $O(n^2)$ , the overall complexity becomes  $O(n^3)$ . If the matrix  $M$  is banded with small width (such as the one to be analyzed in the next section), both Steps 2 and 3 can be carried out in linear time (even without any adaptive procedure). In this case, the complexity reduces to  $O(n^2)$ .

Remarks. (i) The above complexity analysis assumes that the systems of linear equations (6ii) are solved by a direct method (such as Gaussian elimination). For problems of very large size (say when  $n$  is a few thousands), it may be more advantageous to solve such

systems by iterative methods (like SOR). In this case, the analysis would not be applicable because of the infinite nature of the iterative schemes.

(ii) The implementation given above ignores the basic components  $(\bar{q}_L, \bar{p}_L)$  until termination actually occurs. If for some reason they are needed, they can be obtained recursively from the following formulas:

$$(\bar{q}_L, \bar{p}_L)_{\text{new}} = (\bar{q}_L, \bar{p}_L)_{\text{old}} - M_{Lk}(\bar{q}_k, \bar{p}_k)_{\text{new}}$$

$$(\bar{q}_k, \bar{p}_k)_{\text{new}} = -(\bar{q}_k, \bar{p}_k)_{\text{old}} / (M_{kk} - M_{kL} \bar{M}_{Lk}) .$$





of consecutive rows and columns, then  $M_{LL}$  is of the same type as  $M$ . Indeed, if

$$L = \{j, j+1, \dots, j+k\},$$

then  $M_{LL}$  is of type  $(\{\beta_i\}_{i=j}^{j+k+1}, \{w_i\}_{i=j}^{j+k+2})$ .

By the  $k$ -th forward (backward) leading principal submatrix of  $M$ , we mean the submatrix consisting of the first (last)  $k$  rows and  $k$  columns. The following few lemmas establish certain invariance properties of the Schur complements of leading principal submatrices in  $M$ . They are all stated in terms of forward leading principal submatrices. Similar results hold for backward leading principal submatrices. The first lemma says that the Schur complement of  $M_{11}$  in  $M$  is of the same type as  $M$ . Moreover, the scalars defining this Schur complement are not much different from those defining  $M$  and can be obtained easily from some simple expressions.

Lemma 1. The Schur complement  $(M/M_{11})$  is of type

$$(\{\tilde{\beta}_2, \{\beta_i\}_{i=3}^{n+1}\}, \{\tilde{w}_2, \tilde{w}_3, \{w_i\}_{i=4}^{n+2}\})$$

where

$$\tilde{\beta}_2 = \beta_1 \beta_2 (w_1^{-1} \beta_1 + w_2^{-1} (\beta_1 + \beta_2)) / (w_1^{-1} \beta_1^2 + w_2^{-1} (\beta_1 + \beta_2)^2)$$

$$\tilde{w}_2 = M_{11} / (w_2^{-1} (w_1^{-1} \beta_1^2 + w_3^{-1} \tilde{\beta}_2 (\beta_1 + \beta_2)))$$

$$\tilde{w}_3 = M_{11} / (w_3^{-1} (w_1^{-1} \beta_1^2 + w_2^{-1} (\beta_1 + \beta_2)^2)).$$

Proof. This follows from a straightforward computation of  $(M/M_{11})$ .

Generalizing Lemma 1, we have

Lemma 2. Let  $N_k$  denote the  $k$ -th forward leading principal submatrix of  $M$ . Then the Schur complement  $(M/N_k)$  is of type

$$(\{\tilde{\beta}_{k+1}, \{\beta_i\}_{i=k+2}^{n+1}\}, \{\tilde{w}_{k+1}, \tilde{w}_{k+2}, \{w_i\}_{i=k+3}^{n+2}\})$$

for some suitable scalars  $\tilde{\beta}_{k+1}$ ,  $\tilde{w}_{k+1}$  and  $\tilde{w}_{k+2}$ .

Proof. This can be proved by induction, utilizing Lemma 1 and the fact that

$$(M/N_k) = (\bar{M}/\bar{N}_{k-1})$$

where  $\bar{M} = (M/M_{11})$  and  $\bar{N}_{k-1}$  denotes the  $(k-1)$ -st leading principal submatrix of  $\bar{M}$ .

Lemma 3. Let  $M_{LL}$  be a principal submatrix of  $M$ . Let  $N_k$  be the  $k$ -th forward leading principal submatrix of  $M_{LL}$ . Then the Schur complement  $(M_{LL}/N_k)$  is a principal submatrix of some matrix  $M'$  of the same type as  $M$ .

Proof. By means of an inductive argument, like the one used in Lemma 2, it suffices to prove for  $k = 1$ . If  $M_{LL}$  consists of consecutive rows and columns from  $M$ , then  $M_{LL}$  is itself a matrix of the same type as  $M$  and thus the conclusion follows from Lemma 1. On the other hand, if some rows (and columns) in  $M_{LL}$  are not consecutive, we may fill in those missing rows and columns to get a larger principal submatrix  $M_{L'L'}$  with  $L \subseteq L'$ . Then  $(M_{LL}/(M_{LL})_{11})$  is a principal submatrix of  $(M_{L'L'}/(M_{LL})_{11})$  which by Lemma 1 again, is a matrix of the same type as  $M$ .

The following lemma is easy to see. It allows us to perform some scaling operation in the main inductive proof of Theorem 2.

Lemma 4. Let  $N$  be obtained from the matrix  $M$  by deleting its second row and second column and then scaling the first row and the first column by a positive scalar  $\delta$  (the  $(1,1)$ -entry is thus scaled by  $\delta^2$ ). Then  $N$  is equal to the principal submatrix obtained from the matrix  $M'$  by deleting its second row and second column where  $M'$  is of type

$$(\{\delta\beta_1, \delta\beta_2, \{\beta_i\}_{i=3}^{n+1}\}, \{w_i\}_{i=1}^{n+2}).$$

Remark. We should point out that if we perform the same scaling operation to the matrix  $M$  itself, the resulting matrix will not be of the same type as  $M$ .

In [31], it was proved that if the set of data points  $\{a_i\}_{i=1}^{n+2}$  is convex, then the least-square concave fit must be a straight line. This interesting geometric fact has an important algebraic consequence, namely, that the inverse of the matrix  $M$  is nonnegative. To see this, we translate the geometric statement in the context of the LCP( $\delta$ ), obtaining the implication:

$$Aa \leq 0 \implies y^* = Au^* = 0$$

which is equivalent to

$$Aa \leq 0 \implies x^* = -(AW^{-1}A^T)^{-1}Aa \geq 0.$$

Since the matrix  $A$  has full row rank, the above implication is in turn equivalent to



Remark. Lemma 6 is believed to have its own interest. For example, it provides an explicit inverse for the matrix

$$\begin{bmatrix} 2 & -1 & & & \\ -1 & 2 & -1 & & \\ & \ddots & \ddots & \ddots & \\ & & -1 & 2 & -1 \\ & & & -1 & 2 \end{bmatrix}$$

which occurs very often in the discretization of elliptic partial differential equations.

In order to establish Lemma 5, we first show that a certain entry of  $M^{-1}$  is positive.

Lemma 7.  $(M^{-1})_{1,n} > 0$ .

Proof. Applying the Sherman-Morrison-Woodbury formula [15, p. 124] to (9), we obtain

$$M^{-1} = (B\tilde{W}^{-1}B^T)^{-1} - (w_1^{-1/2}\beta_1\bar{e}_1^T, w_m^{-1/2}\beta_m\bar{e}_n^T) \left[ I_{2 \times 2} + \begin{pmatrix} -1/2 & \\ w_1 & \beta_1\bar{e}_1^T \\ w_m & 1/2\beta_m\bar{e}_n^T \end{pmatrix} (B\tilde{W}^{-1}B^T)^{-1} (w_1^{-1/2}\beta_1\bar{e}_1, w_m^{-1/2}\beta_m\bar{e}_n) \right] \begin{pmatrix} -1/2\beta_1\bar{e}_1 \\ w_m^{-1/2}\beta_m\bar{e}_n \end{pmatrix}$$

where

$$(\bar{e}_1, \bar{e}_n) = (B\tilde{W}^{-1}B^T)^{-1}(e_1, e_n).$$

Define three scalars

$$\gamma = \sum_{k=2}^{m-1} w_k(\alpha_k - \alpha_1)(\alpha_m - \alpha_k), \quad \delta = \sum_{k=2}^{m-1} w_k(\alpha_m - \alpha_k)^2 \quad \text{and} \quad \sigma = \sum_{k=2}^{m-1} w_k(\alpha_k - \alpha_1)^2.$$

Then, by Lemma 6 and an easy manipulation, we obtain

$$(M^{-1})_{1,n} = e_1^T M^{-1} e_n = \frac{(\alpha_2 - \alpha_1)(\alpha_m - \alpha_{m-1})}{(\alpha_m - \alpha_1)^2} (\gamma - (\delta, \gamma)G^{-1}(\gamma))$$

where  $G$  is the  $2 \times 2$  positive definite matrix

$$G = \begin{bmatrix} \delta + w_1(\alpha_m - \alpha_1)^2 & \gamma \\ \gamma & \sigma + w_m(\alpha_m - \alpha_1)^2 \end{bmatrix}.$$

By the Schur determinantal formula (see [7]), we deduce

$$(M^{-1})_{1,n} = \frac{(\alpha_2 - \alpha_1)(\alpha_m - \alpha_{m-1})}{(\alpha_m - \alpha_1)^2} \frac{\det R}{\det G}$$

where  $R$  is the  $3 \times 3$  matrix

$$R = \begin{bmatrix} \gamma & \delta & \gamma \\ \gamma & & G \\ \sigma & & \end{bmatrix} .$$

Obviously,

$$\det R = w_1 w_m (\alpha_m - \alpha_1)^4 \gamma > 0 .$$

Thus,  $(M^{-1})_{1,n} > 0$  as desired.

Proof of Lemma 5. We use induction on  $n$ , which is the order of the matrix  $M$ . It is obvious that the assertion is true for  $n = 1$ . Suppose that it is true for  $k < n$ . Let  $M$  be a matrix of type  $(\{\beta_i\}_{i=1}^{n+1}, \{w_i\}_{i=1}^{n+2})$ . We may write

$$M = \begin{pmatrix} M_{11} & a^T \\ a & N \end{pmatrix}$$

where  $N$  is the  $(n-1)$ -st backward leading principal submatrix of  $M$ . According to Lemma 1, the Schur complement  $(M/M_{11})$  is of the same type as  $M$ . Thus, the induction hypothesis implies that  $\bar{N} = (M/M_{11})^{-1}$  is positive. We have (see [7] e.g.)

$$M^{-1} = \frac{1}{\delta} \begin{pmatrix} 1 & -a^T \bar{N}^{-1} \\ -\bar{N}^{-1} a & \delta \bar{N} \end{pmatrix}$$

where  $\delta = (M/N)$ . Since  $M$  is positive definite,  $\delta > 0$ . Consequently, with the possible exception of the second to last entries in the first row and first column,  $M^{-1}$  is positive. Similarly, applying the same argument to the partitioning

$$M = \begin{pmatrix} N' & b \\ b^T & M_{nn} \end{pmatrix}$$

where  $N'$  is the  $(n-1)$ -st forward leading principal submatrix of  $M$ , we can deduce that with the possible exception of  $(M^{-1})_{1,n}$  and  $(M^{-1})_{n,1}$ , (these latter two entries







$$\begin{bmatrix} \frac{1}{p_2} \bar{N}_2 & \frac{1}{p_2} Q_2^T & & & \\ & \frac{1}{p_2} Q_2 & N_3 & & \\ & & & Q_3^T & \\ & & & & \ddots \\ & & & & & Q_{\ell-2}^T \\ & & & & & & N_{\ell-1} & & \\ & & & & & & & Q_{\ell-1}^T & \\ & & & & & & & & N_{\ell} \end{bmatrix} \begin{bmatrix} \tilde{p}_2 \\ \tilde{p}_3 \\ \vdots \\ \tilde{p}_{\ell-1} \\ \tilde{p}_{\ell} \end{bmatrix} = \begin{bmatrix} 1 \\ p_3 \\ \vdots \\ p_{\ell-1} \\ p_{\ell} \end{bmatrix} \quad (14)$$

By Lemma 4 again, the matrix defining the above system is a principal submatrix of some matrix of the same type as  $M$ . Thus, the inductive hypothesis implies  $\tilde{p}_i > 0$  ( $i = 2, \dots, \ell$ ). To show  $\tilde{p}_1 > 0$ , perform a principal pivot on the third to last block  $N_i$  ( $i = 3, \dots, \ell$ ) in the system (10) (since both  $N_1$  and  $N_2$  are single-entry blocks and  $L$  has  $k \geq 3$  entries, there must be at least one block  $N_i$  ( $i \geq 3$ )), obtaining a  $2 \times 2$  system from which the positivity of  $\tilde{p}_1$  follows easily.

Finally, suppose that the second block  $N_2$  is of order larger than 1. Then writing

$$N_2 = \begin{pmatrix} \bar{N}_2 & b \\ b^T & \delta' \end{pmatrix}$$

and applying an argument similar to the one used when  $N_1$  is of order larger than 1, we can easily deduce that  $\tilde{p}_L > 0$  as desired.

Summarizing, we always have  $\tilde{p}_L > 0$  for any index set  $L$ . This completes the proof.

Remark. The reason why the scaling in the systems (12) and (14) is needed is because the induction hypothesis is applicable only when the right-hand vector in (12) and (14) consists of all ones.

4. THE CASE OF A HIDDEN MINKOWSKI MATRIX.

Returning to the general LCP(3), we prove

Theorem 3. Suppose that  $M^T$  is hidden Minkowski. Let  $X$  and  $Y$  be two  $Z$ -matrices satisfying

$$M^T X = Y \quad (15i)$$

$$r^T X + s^T Y > 0 \text{ for some vectors } r, s \geq 0. \quad (15ii)$$

Then condition (5) holds for any vector  $p > 0$  satisfying  $X^T p > 0$ . Consequently, the conclusion of Theorem 1 applies to the LCP(3).

Before proving Theorem 3, we compare it with the results obtained in [8,20] concerning the solvability of an LCP with a hidden  $Z$ -matrix as a linear program. Let  $M, X, Y$  and  $p$  be as given in Theorem 3. Then the (unique) solution to the LCP

$$y = q + M^T x \geq 0, \quad x \geq 0 \text{ and } y^T x = 0 \quad (16)$$

can be obtained by solving the LP

$$\begin{aligned} &\text{minimize } p^T x \\ &\text{subject to } q + M^T x \geq 0 \text{ and } x \geq 0. \end{aligned}$$

Theorem 3 implies that the same vector  $p$  can be used to start the parametric principal pivoting algorithm for solving the LCP

$$y = q + Mx \geq 0, \quad x \geq 0 \text{ and } y^T x = 0 \quad (3)$$

and the algorithm will terminate after at most  $n$  pivots. Notice that (3) is defined by the matrix  $M$  whereas (16) is by its transpose  $M^T$ .

An example of a vector  $p > 0$  satisfying  $X^T p > 0$  is given by

$$p = r + Ms \quad (17)$$

where  $r$  and  $s$  satisfy (15ii), see [8]. More generally, the vector  $p$  computed by solving the system of linear equations

$$X^T p = e \quad (17)'$$

for any positive right-hand vector  $e$  will have the same required properties. Indeed, the fact that  $M^T$  is  $P$  implies that  $X^T$  is itself Minkowski (see Lemma 8 below). Hence the solution to (17)' is  $p = X^{-T} e > 0$ . In any case,  $p$  can be obtained in at most  $O(n^3)$

time. Consequently, by the analysis of Section 2, the parametric principal pivoting algorithm will solve the LCP(3) in  $O(n^3)$  time.

To prove Theorem 3, we quote the lemma below which summarizes two useful properties of a hidden Minkowski matrix. A proof of the lemma can be found in [24].

Lemma 8. Let  $M$ ,  $X$  and  $Y$  be as given in Theorem 3. Then for any index set  $L$ ,

$$(i) \quad M_{LL}^T(X/X_{KK}) = (W/X_{KK})$$

where  $K$  is the complement of  $L$  and

$$W = \begin{pmatrix} Y_{LL} & Y_{LK} \\ X_{KL} & X_{KK} \end{pmatrix}.$$

(ii) The matrix  $W$  is Minkowski. (In particular, so are  $X$  and  $Y$ .)

Proof of Theorem 3. We first remark that if  $X^T p > 0$ , then for any complementary index sets  $L$  and  $K$ ,

$$(X^T/(X^T)_{KK})_{PL} > 0.$$

From Lemma 8(i), we obtain

$$(M^T)_{LL} = (W/X_{KK})(X/X_{KK})^{-1}$$

which implies

$$(M_{LL})^{-1} = (W/X_{KK})^{-T}(X/X_{KK})^T$$

Thus, it follows that

$$(M_{LL})^{-1}_{PL} = (W/X_{KK})^{-T}(X/X_{KK})^T_{PL} \geq 0$$

because  $(W/X_{KK})$  is Minkowski (implied by Lemma 8(ii)) and thus has a nonnegative inverse, and  $(X/X_{KK})^T = (X^T/(X^T)_{KK})$ . This completes the proof.

The preceding discussion shows that in general the parametric vector  $p$  can be calculated from either (17) or (17)'. In the Corollary below, we identify two classes of matrices  $M$  for which the vector  $p$  is available trivially.

Corollary 1. (i) If  $M = Y^T + ab^T$  where  $Y$  is a Minkowski matrix and  $a$  and  $b$  are positive vectors, then  $p = a$  satisfies condition (5).

(ii) If  $M$  has positive diagonal entries and is strictly row diagonally dominant, i.e.,

$$M_{ii} > \sum_{j \neq i} |M_{ij}| \quad \text{all } i,$$

then the vector  $p = (p_i)$  defined by

$$p_i = M_{ii} + \sum_{j: M_{ij} < 0} M_{ij} \quad \text{all } i$$

satisfies condition (5).

Proof. It is known that a matrix  $M$  satisfying either (i) or (ii) is hidden Minkowski (see [8] e.g.). It is also obvious that the vector  $p$  in either case is positive. Thus it remains to verify that  $X^T p > 0$  where  $X$  is as given in Theorem 3. In case (i), we have

$$M^T = Y(I + Y^{-1}ba^T).$$

Thus

$$X^T = (I + Y^{-1}ba^T)^{-T} = I - \frac{ab^T Y^{-T}}{1 + b^T Y^{-T} a}.$$

Hence, it follows that

$$X^T a = a / (1 + b^T Y^{-T} a) > 0.$$

In case (ii), we may write (see [8])

$$M = 2A - B$$

where  $B = \bar{M}$  is the comparison matrix of  $M$  and  $A = (M + \bar{M})/2$ . It is easy to show that the required  $X^T$  is given by (see [8])

$$X^T = BA^{-1}.$$

With the vector  $p$  defined as specified, we easily deduce that  $A^{-1}p$  is the vector of all ones. Thus  $X^T p > 0$  because  $M$  is strictly row diagonally dominant.

Remark. The matrix  $M$  arising from the concave regression problem (cf. (8)') provides another example for which the parametric vector  $p$  is available trivially.

If  $M$  is an  $H$ -matrix with positive diagonals and if  $d$  is a positive vector such that  $\bar{M}d > 0$ , then the same proof of case (ii) in the above Corollary shows that the vector  $p = (M + \bar{M})d/2$  satisfies condition (5).

5. ON LEMKE'S METHOD.

It is rather evident that the parametric principal pivoting algorithm presented in Section 2 is intimately related to Lemke's well-known almost complementary pivoting algorithm [18, 19]. Indeed, in [21], McCammon pointed out that Lemke's method can be implemented as a parametric pivot scheme in which the artificial variable is treated as a parameter (see also [19]). McCammon also discussed the relationship between this parametric version of Lemke's method and principal pivoting. As a result of such a connection, it is natural to ask whether the existence of a positive vector  $p$  satisfying condition (5) will imply a linear termination for Lemke's method applied to a LCP with a matrix which is not necessarily  $P$ . The result below gives an affirmative answer to this question. Recall that a matrix  $M$  is nondegenerate if all its principal submatrices are nonsingular.

Theorem 4. Let  $M$  be an  $n \times n$  nondegenerate matrix. Suppose that there is a positive vector  $p$  satisfying condition (5). Then Lemke's almost complementary pivoting algorithm using  $p$  as the artificial vector, will terminate with a desired solution to the LCP(3) after at most  $n + 1$  pivots.

Proof. Consider a current iteration (after the first) of Lemke's method. We have available an index set  $L$  corresponding to the basic  $x$ -variables and an index  $t \notin L$  such that  $(y_t, x_t)$  is the nonbasic pair. The canonical tableau may be written as (only the key entries are displayed)

$$\begin{array}{c} x_L \\ \theta \\ y_J \end{array} = \begin{array}{c} y_L \quad y_t \quad x_t \quad x_J \\ \left[ \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & - \begin{pmatrix} M_{LL} & P_L \\ M_{tL} & P_t \end{pmatrix}^{-1} \begin{pmatrix} M_{Lt} \\ M_{tt} \end{pmatrix} & \\ \hline & & & \\ \hline \end{array} \right] \end{array} \quad (17)$$

where  $J$  is the complement of  $\{t\} \cup L$ .

Assume that  $y_t$  has just become nonbasic so that  $x_t$  is the incoming variable. We claim that condition (5) implies that the next pivot will not occur in an  $x_L$ -row. Indeed, after an easy computation, it is easy to verify that the  $(x_L, x_t)$ -entry in the above

tableau is given more explicitly by

$$\begin{pmatrix} M_{tt} - M_{tL} M_{LL}^{-1} M_{Lt} \\ P_t - M_{tL} M_{LL}^{-1} P_L \end{pmatrix} M_{LL}^{-1} P_L - M_{LL}^{-1} M_{Lt} . \quad (18)$$

(The fact that  $M$  is nondegenerate (implying  $M_{LL}^{-1}$  exists) and the nonsingularity of the basis matrix

$$B = \begin{pmatrix} M_{LL} & P_L \\ M_{tL} & P_t \end{pmatrix}$$

guarantee the nonvanishing of the denominator. The nonsingularity of the basis  $B$  is in turn guaranteed by the pivot operations.) By (5), the vector

$$\begin{pmatrix} \bar{P}_L \\ \bar{P}_t \end{pmatrix} = \begin{pmatrix} M_{LL} & M_{Lt} \\ M_{tL} & M_{tt} \end{pmatrix}^{-1} \begin{pmatrix} P_L \\ P_t \end{pmatrix} > 0 .$$

It is easy to show that

$$\begin{aligned} \bar{P}_L &= M_{LL}^{-1} P_L - \left( \frac{P_t - M_{tL} M_{LL}^{-1} P_L}{M_{tt} - M_{tL} M_{LL}^{-1} M_{Lt}} \right) M_{LL}^{-1} M_{Lt} \\ \bar{P}_t &= (P_t - M_{tL} M_{LL}^{-1} P_L) / (M_{tt} - M_{tL} M_{LL}^{-1} M_{Lt}) . \end{aligned}$$

(Again, the nondegeneracy of  $M$  implies that the denominator in the above two expressions is nonzero.) Thus, it follows from (18) that the  $(x_L, x_t)$ -entry in tableau (17) is given by the vector  $\bar{P}_L / \bar{P}_t$  which is nonnegative. Consequently, the increase of  $x_t$  will not decrease the values of the already-basic  $x_L$ -variables. Therefore, the next pivot will occur either in the  $\theta$ -row (in which case a solution to the LCP(3) is obtained) or in a  $y_j$ -row (in which case the index set  $L$  increases by one element and the argument just given repeats itself).

Next, we show that the  $(\theta, x_t)$ -entry in tableau (17) is negative. Indeed, this entry is  $-1/\bar{P}_t$ . Thus it is negative. Consequently, the increase of  $x_t$  is bounded above by  $\theta$ . Hence, termination on a secondary ray can not occur.

Summarizing, we have proven that once an  $x$ -variable has become basic, it will stay basic until termination occurs. Since there are only  $n$   $x$ -variables, the algorithm terminates after at most  $n + 1$  pivots. Since termination on a ray is ruled out, a desired solution to the LCP(3) will be obtained. This establishes the theorem.

Remark. i) As in Theorem 1, no nondegeneracy assumption is needed to prove Theorem 4.

ii) The matrix

$$\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$$

satisfies the assumptions of Theorem 4 (with  $p = (1,1)^T$ ). But it is not  $P$ .

Theorem 4 may be considered an extension of Theorem 1 to the case of a non- $P$  matrix. Proofs to both theorems are based on the same key idea that an  $x$ -variable once becomes basic, stays basic. At present, it is not clear to us what class of matrices  $M$  (besides those considered in the last two sections) will produce an easily calculable vector  $p$  satisfying the required properties. We leave this as an open question for further investigation.

It is interesting to contrast Theorem 4 with the worst-case studies of [12, 22]. In these earlier studies, examples have been presented which demonstrate that Lemke's method can require an exponential number of pivots. On the other hand, Theorem 4 guarantees that for certain class of problems, the polynomial-time complexity of Lemke's method can be established. Although a complete knowledge of such problems is not yet available, the examples in Sections 3 and 4, together with those that are previously known should have amply demonstrated that the class is non-void and indeed contains some very interesting applications of the LCP.

#### REFERENCES

1. R. E. Barlow, D. J. Bartholomew, J. M. Bremner and H. D. Brunk, Statistical inference under order restrictions, John Wiley & Sons (New York 1972).
2. J. R. Birge and A. Gana, "Computational complexity of van der Heyden's variable dimension algorithm and Dantzig-Cottle's principal pivoting method for solving LCP's", Mathematical Programming 26 (1983) 316-325.
3. R. G. Bland, D. Goldfarb and M. J. Todd, "The ellipsoid method: A survey", Operations Research 29 (1981) 1039-1091.
4. R. Chandrasekaran, "A special case of the linear complementarity problem", Opsearch 7 (1970) 263-268.
5. S. J. Chung, "A note on the complexity of LCP: The LCP is strongly NP-complete", Technical Report 792, Department of Industrial and Operations Engineering, The University of Michigan-Ann Arbor (1979).
6. R. W. Cottle, "Monotone solutions of the parametric linear complementarity problem", Mathematical Programming 3 (1972) 210-214.
7. R. W. Cottle, "Manifestations of the Schur complement", Linear Algebra and its Applications 8 (1974) 189-211.
8. R. W. Cottle and J. S. Pang, "On solving linear complementarity problems as linear programs", Mathematical Programming Study 7 (1978) 88-107.
9. R. W. Cottle and R. S. Sacher, "On the solution of large structured linear complementarity problems: The tridiagonal case", Journal of Applied Mathematics and Optimization 3 (1977) 321-340.
10. C. W. Cryer, "The efficient solution of linear complementarity problems for tridiagonal Minkowski matrices", ACM Transactions of Mathematical Software 9 (1983) 199-214.
11. W. Dent, T. Robertson and W. Johnson, "Concave regression", unpublished manuscript, University of Iowa (1977).
12. Y. Fathi, "Computational complexity of LCPs associated with positive definite symmetric matrices", Mathematical Programming 17 (1979) 335-344.

13. C. Hildreth, "Point estimates of ordinates of convex functions", Journal of the American Statistical Association 49 (1954) 598-619.
14. C. A. Holloway, "On the estimation of convex functions", Operations Research 27 (1979) 401-407.
15. A. S. Householder, The theory of matrices in numerical analysis, Dover Publication Inc. (New York 1964).
16. L. G. Khachiyan, "A polynomial algorithm in linear programming", Soviet Mathematics Doklady 20 (1979) 191-194.
17. J. B. Kruskal, "Nonmetric multidimensional scaling: A numerical method", Psychometrika 29 (1964) 115-129.
18. C. E. Lemke, "Bimatrix equilibrium points and mathematical programming", Management Science 4 (1965) 681-689.
19. C. E. Lemke, "Recent results on complementarity problems", in Nonlinear Programming, J. B. Rosen, O. L. Mangasarian and K. Ritter, eds., Academic Press (New York 1970) 349-384.
20. O. L. Mangasarian, "Linear complementarity problems solvable by a single linear program", Mathematical Programming 10 (1976) 263-270.
21. S. R. McCammon, "Complementary pivoting", Ph.D. dissertation, Department of Mathematical Sciences, Rensselaer Polytechnique Institute (1970).
22. K. G. Murty, "Computational complexity of complementary pivot methods", Mathematical Programming Study 7 (1978) 61-73.
23. J. S. Pang, "On cone orderings and the linear complementarity problem", Linear Algebra and its Applications 22 (1978) 267-281.
24. J. S. Pang, "Hidden Z-matrices with positive principal minors", Linear Algebra and its Applications 23 (1979) 201-215.
25. J. S. Pang, "On a class of least-element complementarity problem", Mathematical Programming 16 (1979) 111-126.
26. J. S. Pang, "On discovering hidden Z-matrices", in Constructive approaches to mathematical models, Coffman and G. Fix, eds., (Academic Press 1979) 231-241.

27. J. S. Pang, "A new and efficient algorithm for a class of portfolio selection problems", Operations Research 28 (1980) 754-767.
28. J. S. Pang, I. Kaneko and W. P. Hallman, "On the solution of some (parametric) linear complementarity problems with applications to portfolio selection, structural engineering and actuarial graduation", Mathematical Programming 16 (1979) 325-347.
29. J. S. Pang and P. S. C. Lee, "A parametric linear complementarity technique for the computation of equilibrium prices in a single commodity spatial model", Mathematical Programming 20 (1981) 81-102.
30. R. Saigal, "A note on a special linear complementarity problem", Opsearch 7 (1970) 175-183.
31. C. F. Wu, "Some algorithms for concave and isotonic regression", TIMS Studies in Management Sciences 19 (1982) 105-116.

JSP/ed

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2597	2. GOVT ACCESSION NO. <b>AD-A136405</b>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) LINEAR COMPLEMENTARITY PROBLEMS SOLVABLE BY AN EFFICIENT POLYNOMIALLY BOUNDED PIVOTING ALGORITHM	5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period	
	6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(s) Jong-Shi Pang	8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Madison, Wisconsin 53706	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 5 - Optimization and Large Scale Systems	
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709	12. REPORT DATE November 1983	
	13. NUMBER OF PAGES 28	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Linear complementarity, parametric principal pivoting, efficient algorithm, polynomially bounded, concave regression, hidden Z, diagonally dominant		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Applied to two important classes of linear complementarity problems defined by an $n \times n$ matrix, the parametric principal pivoting algorithm, using a suitably chosen (and easily computable) parametric vector, terminates with a desired solution after at most $n$ pivot operations. Since each pivot can be performed using at most $O(n^2)$ arithmetic operations, the total (cont.)		

ABSTRACT (cont.)

computational complexity of the algorithm for solving these linear complementarity problems is no more than  $O(n^3)$ . In one of the two classes of problems being studied, the complexity is  $O(n^2)$  because the matrix involved is 5-diagonal which allows each pivot to be performed in linear time. Some discussion in connection with Lemke's well-known almost complementary pivoting algorithm is also addressed.

DA  
FIL