

AD-A141 185

IONIC REACTIONS IN THE EARTH'S UPPER ATMOSPHERE(U)  
OKLAHOMA UNIV NORMAN DEPT OF PHYSICS AND ASTRONOMY  
T M MILLER MAR 84 AFOSR-TR-84-0345 AFOSR-83-0177

1/1

UNCLASSIFIED


F/G 4/1

NL



100

[The rest of the page is heavily obscured by a large black redaction box.]



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

**AFOSR-TR- 84 - 0345**

APR 3 1984

11

**FINAL REPORT**

**GRANT NO. AFOSR-83-0177**

submitted to

**Air Force Office of Scientific Research  
"Minigrant" Program**

**Lt. Col. Ted Cress, Atmospheric Sciences**

project entitled

**Ionic Reactions in the Earth's Upper Atmosphere**

Prepared by

**Dr. Thomas M. Miller  
Associate Professor of Physics  
Department of Physics and Astronomy  
(405) 325-3961**

**DTIC  
SELECTED  
MAY 17 1984**

Submitted by

**Office of Research Administration  
University of Oklahoma  
1000 Asp, Room 314  
Norman, Oklahoma 73019**

March, 1984

Approved for public release,  
distribution unlimited

**84 05 15 237**

**AD-A141 185**

**DTIC FILE COPY**

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION <b>Unclassified</b>		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; Distribution unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		5. MONITORING ORGANIZATION REPORT NUMBER(S) <b>AFOSR-TR- 84 - 0345</b>	
6a. NAME OF PERFORMING ORGANIZATION Department of Physics and Atmospheric Sciences	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION <b>AFOSR/NC</b>	
6c. ADDRESS (City, State and ZIP Code) University of Oklahoma Norman, OK 73019		7b. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling, AFB, D.C. 20332	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION <b>AFOSR</b>	8b. OFFICE SYMBOL (If applicable) <b>NC</b>	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER <b>AFOSR-83-0177</b>	
8c. ADDRESS (City, State and ZIP Code) Bldg 410 Bolling AFB, D.C. 20332		10. SOURCE OF FUNDING NOS.	
		PROGRAM ELEMENT NO. <b>61102F</b>	TASK NO. <b>D9</b>
		PROJECT NO. <b>2310</b>	WORK UNIT NO.
11. TITLE (Include Security Classification) <b>Ionic Reactions in the Earth's Atmosphere</b>			
12. PERSONAL AUTHOR(S) <b>Dr. Thomas M. Miller</b>			
13a. TYPE OF REPORT <b>Final</b>	13b. TIME COVERED FROM <b>1 May 83</b> TO <b>30 Apr 84</b>	14. DATE OF REPORT (Yr., Mo., Day) <b>March 1984</b>	15. PAGE COUNT <b>8</b>
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB. GR.	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
<p>Measurements made on the selected-ion flow-tube at the Air Force Geophysics Laboratory on the temperature dependence of several ion-molecule reaction rate coefficients. The <math>N^+ + CO</math> and <math>O^+ + NO</math> systems were studied over a range of 9 - 450° K. Measurements were also made on <math>O^+ + O_2</math>, <math>CO^+ + O_2</math>, and <math>CO_2^+ + O_2</math> over the range 90 - 450° K. Comparison with available earlier results showed agreement within experimental uncertainty. A flowing afterglow apparatus was developed at Oklahoma for the study of ion interactions. Positive-ion/negative ion recombination was studied in this apparatus. Preliminary work was carried out for the <math>Cl_2^+ + Cl</math> system and photographic documentation is used to determine the strength of the recombination process.</p>			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input checked="" type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION <b>Unclassified</b>	
22a. NAME OF RESPONSIBLE INDIVIDUAL <b>Ted S. Cress, Lt Col, USAF</b>		22b. TELEPHONE NUMBER (Include Area Code) <b>(202) 767-4960</b>	22c. OFFICE SYMBOL <b>NC</b>

## ABSTRACT OF OBJECTIVES AND ACCOMPLISHMENTS

Our primary objective has been to complete data analysis and a journal publication on the temperature dependence of several ion-molecule reaction rate coefficients. The experimental work was carried out at the Air Force Geophysics Laboratory in collaboration with Dr. John F. Paulson.

The AFGL Selected-Ion Flow-Tube had not been used at temperatures other than 300°K previously. Relatively simple ion-molecule systems were chosen for this initial work.  $N^+ + CO$  and  $C^+ + NO$  were studied over the range 9-450°K in order to compare with recent beam-gas experiments at Universitat Freiburg. In the case of  $C^+ + NO$ , the partial rate coefficients for the product channels had not been previously measured, even at room temperature.

We also made measurements on  $C^+ + O_2$ ,  $CO^+ + O_2$ , and  $CO_2^+ + O_2$  over the range 90-450°K. Previous work was at 300°K except for one  $C^+ + O_2$  datum at 100°K, and high temperature data on  $CO_2^+ + O_2$ .

Where comparisons with earlier work were possible the agreement was within experimental uncertainties except for the product branching ratios measured in an ion cyclotron resonance apparatus by Anicich, Huntress, and Futrell for  $C^+ + O_2$ .

An article on these experiments and a detailed discussion of the results has been accepted by the Journal of Chemical Physics. Page proofs are attached to this report.

We have also continued our study of the perturbation of the gas flow in flow-tube reactors by high-speed helium injectors and have decided on a redesign of the injector which will eliminate a shock wave in the flow tube.

We have also completed the construction of a related apparatus at the University of Oklahoma for the study of ion interactions. This flowing afterglow apparatus is being used initially for research on positive-ion/negative-ion recombination. If the ion density is high enough ( $> 10^9 \text{ cm}^{-3}$ ) the loss of plasma is dominated by ion-ion recombination, and a measurement of ion density vs flow time from a

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFOSR)  
NOTICE OF TRANSFER TO DTIC


This technical report has been reviewed and approved for distribution under AFOSR-12.  
Distribution is unlimited.

MATTHEW J. KENNER  
Chief, Technical Information Division


reference point upstream in the apparatus yields the recombination rate. A mass spectrometer is used to identify the ion species present in the plasma at the downstream end.

Preliminary work has been carried out for the simple  $\text{Cl}_2^+ + \text{Cl}^-$  system. The attached photograph shows the apparatus in action. The bright white glow at the left is due to microwave excitation of the flowing helium gas. The blue glow just to the right follows injection of chlorine gas into the electron-ion plasma. Electrons rapidly attach to  $\text{Cl}_2$  to form  $\text{Cl}_2^-$ . The blue glow is a result of electron transfer into excited states of  $\text{Cl}_2$  via  $\text{Cl}_2^+ + \text{Cl}^- \rightarrow \text{Cl}_2^* + \text{Cl} \rightarrow \text{Cl}_2 + \text{Cl} + \text{light}$ .

Thus, the rate of decay of the plasma which is evident in the photograph is indicative of the strength of the ion-ion recombination process. A langmuir probe is used to determine the ion density vs flow time (or distance). However, we have been using a monochromator to observe the light given off as an alternative measurement scheme. A number of advantages may ensue as outlined in our recent AFOSR proposal, "Experimental Studies of Atmospheric Ions: Ion-Ion and Photon-Ion Interactions", submitted in January, 1984.



A-1



Running Title: Temperature dependence of ion-molecule reactions

# Temperature dependence of the ion-molecule reactions $N^+ + CO_2^+ + NO$ , and $C^+, CO^+, CO_2^+ + O_2$ from 90–450 K

Thomas M. Miller<sup>a</sup>, Rodney E. Wetterskog<sup>a</sup>, and John F. Paulson*Air Force Geophysics Laboratory, Hanscom Air Force Base, Bedford, Massachusetts 01731*

(Received 28 November 1983; accepted 23 January 1984)

A variable-temperature, selected-ion flow tube was used to study the ion-molecule reactions  $N^+ + CO_2^+ + NO$ ,  $C^+ + NO$ ,  $C^+ + O_2$ ,  $CO^+ + O_2$ , and  $CO_2^+ + O_2$  over the temperature range of 90–450 K. We have measured reaction rate coefficients and have determined branching ratios for different reaction channels. The two  $C^+$  reactions are found to be essentially gas kinetic in rate for 90–450 K. The other three reaction rate coefficients are found to decrease with temperature.

PACS numbers: 34.70

## I. INTRODUCTION

We have used a selected-ion flow tube to study several ion-molecule reactions as a function of temperature, over the range 90–450 K. We have studied the temperature dependence of reactions between  $N^+ + CO$  and between  $C^+ + NO$  in order to tie in with the remarkable supra-thermal data from Universität Freiburg.<sup>1,2</sup> Previous research on these two reactions has been carried out only at room temperature. Further, measurements of the product branching ratios were needed for comparison with the beam data. We have also studied reactions with  $O_2$  for positive ions formed by electron impact on  $CO_2$ , namely  $C^+$ ,  $CO^+$ , and  $CO_2^+$ ,<sup>3</sup> in order to compare with previous data at one or more temperatures.<sup>4–11</sup>

## II. EXPERIMENTAL APPARATUS, METHOD, AND ANALYSIS

The selected-ion flow tube (SIFT) apparatus was developed by Adams and Smith<sup>12</sup> at the University of Birmingham (UK) around 1975, an outgrowth of the well-established flowing afterglow technique<sup>13</sup> so successful at NOAA and elsewhere for the determination of ion-molecule reaction rate coefficients. The SIFT apparatus constructed at the Air Force Geophysics Laboratory and used in this work is similar to the variable-temperature version currently in use at the University of Birmingham.

Briefly, the AFGL SIFT apparatus consists of a 108 cm long, 7.3 cm i.d. stainless steel tube through which a helium carrier gas is flowed at low pressure ( $\sim 0.3$  Torr) and high speed ( $\sim 10^4$  cm/s). Ions are created in an electron-impact ion source, mass analyzed, and a current of  $\sim 10^{-10}$  A of a single desired ion species is injected into the flow tube at low energies ( $\sim 20$  eV) in the center of an annular jet of the helium carrier gas. The ions quickly thermalize in collisions with the helium gas and flow with the gas down the tube. The trip requires approximately 10 ms. At the end of this trip some of the ions pass through a small (0.03 cm) aperture leading into a second mass spectrometer maintained at high vacuum. The detected ion count rate is proportional to the original injected current.

The flow tube was cooled by bursts of liquid nitrogen through a copper jacket, or alternatively heated with car-

tridge heaters clamped to the jacket. The helium buffer gas was appropriately precooled or preheated.

In the ion source electrons were boiled off a thoriated-iridium filament in high vacuum and accelerated toward a gas cell where ions were produced by electron bombardment.  $N_2$  gas was used for  $N^+$  production, and  $CO_2$  gas was used for  $C^+$ ,  $CO^+$ , and  $CO_2^+$ . The electron acceleration potential was kept low enough that the apparent reaction rate coefficient was independent of the potential, i.e., presumably few metastable or doubly charged ions were produced. An acceleration potential of  $\sim 75$  V was used for the results reported here.

The reaction rate coefficient was determined from the attenuation  $I/I_0$  in the primary ion current reaching the detector, due to reactions with a density  $n$  of neutral reactants titrated into the helium buffer gas during a reaction time  $t$ :

$$k = \ln(I_0/I) / nt. \quad (1)$$

Experimentally, the small concentration  $n$  of the reactant gas was determined from the measured helium density  $N$  and the throughput  $Q_R$  of the reactant gas compared to the helium throughput  $Q_{He}$ . The reaction time  $t$  was determined from a direct measurement of the ion velocity  $v_i$  and of the distance  $d$  over which the reactant gas flows. Thus,

$$k = \frac{\ln(I_0/I)}{(Q_R N / Q_{He}) (d / v_i)}. \quad (2)$$

In practice, approximately ten values of  $I$  were obtained for corresponding values of  $n$  (or  $Q_R$ ), for each of three reaction distances  $d$  (74.0, 43.5, and 28.3 cm).

Product ion currents were also recorded for each value of  $n$  so that the product ion percentages could be determined (extrapolating to  $n = 0$ ). A product of a secondary reaction is easily identified with the procedure since the product percentage extrapolated to  $n = 0$  is zero for a secondary product.<sup>12</sup>

The ion velocity was determined from time-of-flight measurements. An electrode in a side port half-way down the flow tube could be biased so as to pull the ions out of the helium flow. When this electrode was pulsed to zero potential for 0.1 ms, a pulse of ions was allowed to pass down the second half of the flow tube and their arrival time at the detector was measured. It is known from the work of Ferguson *et al.*<sup>13</sup> and of Adams *et al.*<sup>14</sup> that the ion velocity  $v_i$  is greater than the bulk helium velocity  $v_{He}$  since the ion den-

<sup>a</sup>Permanent address: Department of Physics and Astronomy, University of Oklahoma, Norman, Oklahoma 73019.

sity is greatest near the flow tube axis where the helium velocity is greatest. In our apparatus  $v_i/v_{He} = 1.68$  above room temperature but the ratio increased with lower temperatures, reaching 2.1 at 90 K. These values are independent of the flow tube pressure.

We observed the effect of a shock wave associated with the helium injection. If the pressure drop across the injector was too large the resulting shock affected the helium flow far enough downstream to interfere with our measurements when using the longest reaction path. The disturbance of the helium flow could most easily be seen in the ion time-of-flight data. We were able to measure the ion flight time for both the first half of the flow tube and for the second half of the flow tube length. The former indicated a much larger ion speed near the helium injector and  $v_i/v_{He}$  depended on the pressure drop across the injector. The latter, used in our reaction data analysis, gave ratios  $v_i/v_{He}$  which were pressure independent. For low pressure differentials the ion speed was constant over the entire length of the flow tube. Our observations are consistent with the studies of Dupeyrat *et al.*,<sup>15</sup> who concluded as we do that the shock wave is inconsequential to the reaction data provided that adequate distance is allowed for equilibrium flow to be established. It is perhaps advantageous to have the helium injector not directed down the flow tube axis, but rather at an angle to the flow tube axis ( $\sim 30^\circ$  for Adams and Smith<sup>12</sup>), and to design the helium injector so as to have a small pressure drop across its opening.

We have uncertainties in the quantities in Eq. (2) as follows:  $\sim 2\%$  in each of  $d$ ,  $v_i$ , and  $Q_{He}$ , and  $\sim 10\%$  uncertainty in the capillary-flow calibration for  $Q_R$ . Considering statistical variations in the data, we believe that the reaction rate coefficients presented here are accurate to within 25%. The temperatures were measured with five platinum resistance thermometers along the flow tube and these generally agreed within 5 K. At our highest temperature a variation as great as 20 K was indicated because we were unable to pre-heat the helium buffer gas adequately.

### III. DISCUSSION OF RESULTS

#### A. $N^+ + CO$

The reaction  $N^+ + CO$  was found to have a reaction rate coefficient which decreases with temperature, as shown in Fig. 1. At our lowest temperature 88 K, the reaction rate coefficient is approximately equal to its gas kinetic value<sup>16</sup> of  $1.1 \times 10^{-9} \text{ cm}^3/\text{s}$ . At our highest temperature 454 K, the reaction rate coefficient has dropped to about half the gas kinetic value. Charge transfer dominates ( $\sim 85\%$ ) at all temperatures studied and is exoergic by 0.52 eV. The only other exoergic channel is  $N^+ + CO \rightarrow NO^+ + C + 0.65 \text{ eV}$ , which we observe taking place whenever charge transfer does not occur. Our product percentages are accurate only to within about five percentage points.

Our measured rate coefficients, in units of  $10^{-10} \text{ cm}^3/\text{s}$ , are: 9.7(88 K), 7.2(146 K), 6.7(196 K), 6.3(245 K), 5.9(296 K), 5.3(371 K), and 4.6(454 K). Our room temperature result is 31% higher than that of Smith *et al.*<sup>17</sup> and 8% lower than that of Lindinger,<sup>18</sup> 18% higher than that of Fehsenfeld *et*




FIG. 1. Measured reaction rate coefficients vs temperature. The vertical scale is broken as indicated to avoid overlapping of data points. The arrows give the gas kinetic rate coefficient (Langevin or average-dipole-orientation model) in each case, calculated using formulas in Ref. 16.

*et al.*,<sup>19</sup> and 37% higher than that of Tishy *et al.*,<sup>20</sup> but is within the combined uncertainties of the experiments. The flow-drift tube data of Lindinger<sup>18</sup> show a decrease in  $k$  with the center-of-mass kinetic energy ( $KE_{c.m.}$ ) which is consistent with the decrease in  $k$  vs temperature which we observe. It is, of course, difficult to make a direct comparison between temperature and relative energy in the flow-drift tube.

Frobin, Schlier, Strein, and Teloy<sup>1</sup> have studied this reaction by interacting a low-energy  $N^+$  beam with CO gas and collecting and mass analyzing all of the product ions. At sufficiently high energies (several electron volts) all possible product ions are observed by Frobin *et al.* The cross sections for the exoergic channels (producing  $NO^+$  and  $CO^+$  ions) show minima, apparently the result of a decrease in the very low energy "capture" cross section versus  $KE_{c.m.}$  and the rise from threshold of the "direct reaction" cross section. At their lowest energy ( $KE_{c.m.} = 0.13 \text{ eV}$ ) Frobin *et al.*<sup>1</sup> measure the product branching ratio as  $\sim 11\% NO^+$  and  $\sim 89\% CO^+$ , which agrees with our result and others.<sup>17,18</sup>

Guyer, Hüwel, and Leone<sup>21</sup> have used a crossed-beam apparatus to study the partition of energy in the  $N^+ + CO \rightarrow CO^+ + N$  reaction. The 0.52 eV exoergicity allows the product  $CO^+$  to be vibrationally excited up to  $v'' = 2$ . The  $CO^+$  vibrational excitation observed by Guyer *et al.* in this reaction is approximately that predicted by a Franck-Condon analysis. Guyer *et al.* found that for  $v'' = 0$

the rotational temperature was only 410 K, implying that almost all of the exoergicity of the reaction went into translational energy of the products.

### B. $C^+ + NO$

The reaction  $C^+ + NO$  proved to be near its gas kinetic rate of about  $1.0 \times 10^{-9} \text{ cm}^3/\text{s}$  for temperatures of 90–300 K, but decreased slowly at higher temperatures. Charge transfer is exoergic by 2.00 eV and we find it to dominate (92%) at all temperatures studied. There are two other exoergic reaction channels.  $C^+ + NO \rightarrow CO^+ + N$  is exoergic by 1.87 eV and we observe it in about 8% of the reactions occurring.  $C^+ + NO \rightarrow CO + N^+$  is energetically allowed (by 1.35 eV), but we see only trace amounts of  $N^+$  (0.1%) which although correlated with NO concentration could not be shown to be a definite product of the  $C^+ + NO$  reaction. The few  $N^+$  ions observed might well be produced through impurities or in surface reactions on the walls of the flow tube.

Our rate coefficients in units of  $10^{-10} \text{ cm}^3/\text{s}$ , are: 9.8(88 K), 9.8(148 K), 10.2(196 K), 9.3(247 K), 9.4(297 K), 8.7(377 K), and 8.3(453 K). The only other thermal-energy studies of  $C^+ + NO$  of which we know are those of Anicich *et al.*,<sup>9</sup> Tichy *et al.*,<sup>20</sup> and the preliminary work of St. Niccolini *et al.*<sup>11</sup> Anicich *et al.*<sup>9</sup> find a reaction rate coefficient of  $8.5 \times 10^{-10} \text{ cm}^3/\text{s}$  in an ion cyclotron resonance apparatus with ion energies  $< 0.1 \text{ eV}$ . Tichy *et al.*<sup>20</sup> used a SIFT apparatus and measured a rate coefficient of  $6.9 \times 10^{-10} \text{ cm}^3/\text{s}$  with charge transfer occurring in most of their reactions. Tichy *et al.* did not observe the  $CO^+ + N$  channel. We have carefully studied the product distribution measurements and cannot explain this discrepancy. St. Niccolini *et al.*<sup>11</sup> used a drift tube apparatus and reported a value of  $5.7 \times 10^{-10} \text{ cm}^3/\text{s}$  at room temperature, with  $NO^+$  as the major product. The reaction rate coefficient was found by St. Niccolini *et al.* to decrease slowly with  $KE_{c.m.}$ .

Cahnbley, Schlier, and Teloy<sup>2</sup> have studied this reaction in the beam-gas apparatus mentioned earlier. As before, at sufficiently high energies they observe all possible ion products and have obtained the individual reaction cross sections. Their experiment extends to low enough energies ( $KE_{c.m.} = 0.04 \text{ eV}$ ) that the capture cross sections for the exoergic channels are significant. Cahnbley *et al.* find the branching ratio at very low energies to be 98%  $NO^+$ , 1.5%  $CO^+$ , and 0.16%  $N^+$ , with the  $CO^+$  fraction consistently decreasing with decreasing  $KE_{c.m.}$ . Cahnbley *et al.*<sup>2</sup> point out that a crossing between the  $NO + C^+$  and  $NO^+ + C$  potential curves can account for the dominance of the charge transfer channel in their experiments.

We observe a larger fraction of the  $CO^+$  product ( $8 \pm 5\%$ ) in our experiment at thermal energies than one would expect from the results of the beam experiment, and we see essentially no change in the branching ratio over 90–450 K. A significant  $CO^+$  product fraction is somewhat surprising since there is no adiabatic correlation<sup>1,2</sup> between the reactant  $C^+ + NO$  and the product  $N + CO^+$  (for ground-state  $CO^+$ , the only exoergic possibility). Frobin *et al.*<sup>1</sup> found that the results of their study of  $N^+ + CO$  collisions could be explained qualitatively on the basis of adiabatic correla-

tions of collinear potential surfaces. An exception was the charge transfer channel, for which a diabatic connection between  $N^+ + CO$  and  $N + CO^+$  was required. In their analysis of the reaction  $C^+ + NO \rightarrow N + CO^+$ , Cahnbley *et al.*<sup>2</sup> speculate that the low energy production of  $CO^+$  is possibly due to the same  $N^+ + CO \rightarrow N + CO^+$  diabatic transition. (At higher energies, excited  $CO^+$  is adiabatically allowed.)

### C. $C^+ + O_2$

The interaction  $C^+ + O_2$  proved to be relatively independent of temperature over the range 90–450 K and only slightly lower than the calculated gas kinetic (Langevin) rate coefficient of  $1.0 \times 10^{-9} \text{ cm}^3/\text{s}$ . Simple charge transfer is endoergic (by 0.80 eV) in this case. Two other reaction channels were observed, however. The first,  $C^+ + O_2 \rightarrow O^+ + CO$  is exoergic by 3.64 eV, and is favored (62%) at all temperatures used. Although the neutral CO product is not observed directly, the energetics of the reaction require that it be bound. The second channel observed is  $C^+ + O_2 \rightarrow CO^+ + O$ , which is exoergic by 3.24 eV.

Our results for the reaction rate coefficient are shown in Fig. 1 and, in units of  $10^{-10} \text{ cm}^3/\text{s}$ , are: 8.3(96 K), 8.5(150 K), 8.8(200 K), 9.0(250 K), 9.3(300 K), 8.9(395 K), and 8.5(450 K). Our 300 K datum may be compared to the value obtained by Adams and Smith,<sup>4</sup>  $9.9 \times 10^{-10} \text{ cm}^3/\text{s}$ , and to that obtained by Rakshit *et al.*,<sup>3</sup>  $9.0 \times 10^{-10} \text{ cm}^3/\text{s}$ . Rakshit *et al.* also found  $9.0 \times 10^{-10} \text{ cm}^3/\text{s}$  at 100 K. Tichy *et al.*<sup>20</sup> found a lower rate coefficient,  $7.4 \times 10^{-10} \text{ cm}^3/\text{s}$  at 300 K, and nearly equal product percentages for  $O^+ + CO$  and  $CO^+ + O$ . Our product distribution is the same as that reported by Adams and Smith.<sup>4</sup> St. Niccolini *et al.*<sup>11</sup> have given preliminary results which likewise show the same product distribution at 300 K. By increasing  $KE_{c.m.}$  with an electric field they find that the product distribution gradually approaches 50%  $O^+$  and 50%  $CO^+$  at relative energies of  $\sim 1 \text{ eV}$ .

Anicich *et al.*<sup>9</sup> found  $12.2 \times 10^{-10} \text{ cm}^3/\text{s}$  for this reaction in an ion cyclotron resonance apparatus with ion energies  $< 0.1 \text{ eV}$ . Oddly, the product distributions reported by Anicich *et al.* are exactly the opposite of what we and others<sup>4,11</sup> have found. (They reported 35%  $O^+$  and 64%  $CO^+$  products, with an estimated uncertainty of three percentage points.) This discrepancy cannot be explained by the presence of nonthermal ions in the ICR cell because St. Niccolini *et al.*<sup>11</sup> show the  $O^+$  product to be the dominant one at energies up to 1 eV, after which the 50–50 ratio obtains.

### D. $CO^+ + O_2$

The rate coefficient for the charge transfer reaction  $CO^+ + O_2 \rightarrow CO + O_2^+$  was found to increase with decreasing temperature, approaching the calculated gas kinetic rate of  $7.7 \times 10^{-10} \text{ cm}^3/\text{s}$  at very low temperatures. This reaction is exoergic by 1.95 eV.

Our results for the reaction rate coefficient, in units of  $10^{-10} \text{ cm}^3/\text{s}$ , are: 3.1(92 K), 2.3(150 K), 2.0(200 K), 1.7(250 K), 1.4(300 K), 1.0(388 K), and 0.82(450 K). Our 300 K datum may be compared with the value obtained by Adams, Smith, and Grief,<sup>6</sup>  $1.2 \times 10^{-10} \text{ cm}^3/\text{s}$ .

Coefficients

or

E.  $\text{CO}_2^+ + \text{O}_2$ 

The rate coefficient for the charge transfer reaction  $\text{CO}_2^+ + \text{O}_2 \rightarrow \text{CO}_2 + \text{O}_2^+$  was likewise found to increase with decreasing temperature, approaching the calculated gas kinetic rate of  $6.9 \times 10^{-10} \text{ cm}^3/\text{s}$  at very low temperatures. Our data join smoothly onto the high temperature data of Lindinger *et al.*,<sup>7</sup> which extend to 900 K and show that the rate coefficient passes through a minimum in the neighborhood of 680 K, implying that other reaction channels (probably due to vibrational excitation) become effective at higher temperatures. The charge transfer reaction is exoergic by 1.71 eV. All rearrangement reaction channels are forbidden energetically in our temperature range.

Our results for the reaction rate coefficients, in units of  $10^{-10} \text{ cm}^3/\text{s}$ , are: 2.2(94 K), 1.4(148 K), 1.0(198 K), 0.80(250 K), 0.63(300 K), 0.40(391 K), and 0.36(450 K).

## ACKNOWLEDGMENTS

We are grateful to Ch. Schlier for providing us with unpublished results from the Freiburg experiments and for many helpful remarks on the interpretation of the gas phase and beam data. We thank our colleagues Fred Dale, Tim Thomas, and Michael Henschman, with whom we had valuable discussions throughout the course of this work. We appreciate the help of David Smith and Nigel Adams of the University of Birmingham with the SIFT apparatus, and the advice of David Fahey and Fred Fehsenfeld of NOAA in regard to the effects of the SIFT helium injector.

Two of the authors (TMM and REW) thank the Air Force Office of Scientific Research for support during May-August 1982 under a contract to the Southeastern Center for Electrical Engineering Education, and for continued support on this work under Grant No. AFOSR-83-0177.

<sup>1</sup>W. Frobin, Ch. Schlier, K. Strein, and E. Teloy, *J. Chem. Phys.* **67** 5506 (1978).

<sup>2</sup>D. Cahnbley, Ch. Schlier, and E. Teloy (to be published). See Ch. Schlier, *Contributions to the Symposium on Atomic and Surface Physics, Salzburg, Austria, February 1982*, (publisher, city, year), p. 251.

<sup>3</sup>We also made measurements for  $\text{O}^+ + \text{O}$ , but found the results dependent on the electron energy in the ion source, implying that excited states of  $\text{O}^+$  might be involved. Reactions involving metastable states of  $\text{O}^+$  have been studied by R. Johnson and M. A. Biondi, *J. Chem. Phys.* **73**, 190 (1980); B. R. Rowe, D. W. Fahey, F. C. Fehsenfeld, and D. L. Albritton, *ibid.* **73**, 194 (1980).

<sup>4</sup>N. G. Adams and D. Smith, *J. Phys.* **B9**, 1439 (1976).

<sup>5</sup>A. B. Rakshit, H. M. P. Stock, D. P. Wareing, and N. D. Twiddy, *J. Phys.* **B11**, 4237 (1978).

<sup>6</sup>N. G. Adams, D. Smith, and D. Grief, *Int. J. Mass Spectrom. Ion Phys.* **26**, 405 (1978).

<sup>7</sup>W. Lindinger, F. C. Fehsenfeld, A. L. Schmeltekopf, and E. E. Ferguson, *J. Geophys. Res.* **79**, 4112 (1974). See also W. Lindinger, M. McFarland, F. C. Fehsenfeld, D. L. Albritton, A. L. Schmeltekopf, and E. E. Ferguson, *J. Chem. Phys.* **63**, 2175 (1975).

<sup>8</sup>F. C. Fehsenfeld, A. L. Schmeltekopf, and E. E. Ferguson, *J. Chem. Phys.* **45**, 23 (1966).

<sup>9</sup>V. G. Anicich, W. T. Huntress, and J. H. Futrell, *Chem. Phys. Lett.* **40**, 233 (1976).

<sup>10</sup>R. S. Hensworth, R. C. Bolden, M. J. Shaw, and N. D. Twiddy, *Chem. Phys. Lett.* **5**, 237 (1970).

<sup>11</sup>St. Niccolini, D. Bassi, S. Fannotta, H. Villinger, R. Richter, A. Saxer, and W. Lindinger, *Contributions to the Symposium on Atomic and Surface Physics, Salzburg, Austria, February 1982* (publisher, city, year), p. 291.

<sup>12</sup>N. G. Adams and D. Smith, *Int. J. Mass Spectrom. Ion Phys.* **21**, 349 (1976).

<sup>13</sup>E. E. Ferguson, F. C. Fehsenfeld, and A. L. Schmeltekopf, *Adv. At. Mol. Phys.* **5**, 1 (1969).

<sup>14</sup>N. G. Adams, M. J. Church, and D. Smith, *J. Phys.* **D 8**, 1409 (1975).

<sup>15</sup>G. Dupeyrat, B. R. Rowe, D. W. Fahey, and D. L. Albritton, *Int. J. Mass Spectrom. Ion Phys.* **44**, 1 (1982).

<sup>16</sup>The gas kinetic reaction rate coefficients in this article have been calculated using the Langevin or ADO expressions. See T. Su and M. T. Bowers, *Int. J. Mass Spectrom. Ion Phys.* **12**, 347 (1973).

<sup>17</sup>D. Smith, N. G. Adams, and T. M. Miller, *J. Chem. Phys.* **69**, 308 (1978).

<sup>18</sup>W. Lindinger (private communication).

<sup>19</sup>F. C. Fehsenfeld, A. L. Schmeltekopf, and E. E. Ferguson, *J. Chem. Phys.* **46**, 2019 (1967).

<sup>20</sup>M. Tichy, A. B. Rakshit, D. G. Lister, and N. D. Twiddy, *Int. J. Mass Spectrom. Ion Phys.* **20**, 231 (1979).

<sup>21</sup>D. R. Guyel, L. Hüwel, and S. R. Leone, *J. Chem. Phys.* **79**, 1259 (1983).

Universität Innsbruck, Innsbruck, 1982

LIMIT OF TEXT

END

FILMED



DUPIC

