



.

MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS - 1963 - A



AD-A142 351

FILE COPY

JIIO

and the second se

INSTITUTE REPORT NO. 171

ACUTE ORAL TOXICITY OF 1-ACETYLOCTAHYDRO-3, 5, 7-TRINITRO-1, 3, 5, 7-TETRAZOCINE (SEX) IN MALE AND FEMALE RATS

CRAIG W. WHITE, DVM, CPT VC and EVELYN M. ZIMMERMAN, SP5

TOXICOLOGY GROUP, DIVISION OF RESEARCH SUPPORT

....; <u>.</u>ts

MAY 1984

Toxicology Series 67

J

21

LETTERMAN ARMY INSTITUTE OF RESEARCH PRESIDIO OF SAN FRANCISCO, CALIFORNIA 94129

84 . 06 21 011

Acute oral toxicity of 1-acetyloctahydro-3,5,7-trinitro-1,3,5,7-tetrazocine in male and female rats--White and Zimmerman Toxicology Series 67

Reproduction of this document in whole or in part is prohibited except with the permission of the Commander, Letterman Army Institute of Research, Presidio of San Francisco, California 94129. However, the Defense Technical Information Center is authorized to reproduce the document for United States Government purposes.

Destroy this report when it is no longer needed. Do not return it to the originator.

Citation of trade names in this report does not constitute an official endorsement or approval of the use of such items.

In conducting the research described in this report, the investigation adhered to the "Guide for the Care and Use of Laboratory Animals," as promulgated by the Committee on Revision of the Guide for Laboratory Animal Facilities and Care, Institute of Laboratory Animal Resources, National Research Council.

This material has been reviewed by Letterman Army Institute of Research and there is no objection to its presentation and/ or publication. The opinions or assertions contained herein are the private views of the author(s) and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense. (AR 360-5)

(Signature and date) 2 May 1984

This document has been approved for public release and sale; its distribution is unlimited.

UNCLASSIFIED SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered) READ INSTRUCTIONS **REPORT DOCUMENTATION PAGE** BEFORE COMPLETING FORM 1. REPORT NUMBER 2. GOVT ACCESSION NO. 3 RECIPIENTIS CATALOS NUMBER LAIR Institute Report No. 171 5 TYPE OF REPORT & PERIOF COVERED 4. TITLE (and Subtitle) Acute Oral Toxicity of 1-Acetyloctahys: 5-3,5,7-Final Trinitro-1,3,5,7-Tetrazocine in Male and Female 14 Apr - 10 May 1983 6 PERFORMING ORG. REFORT NUMBER Rats 7. AUTHOR(1) 8. CONTRACT OR GRANT NUMBER(A) Craig W. White, DVM, CPT VC Evelyn M. Zimmerman, SP4 9. PERFORMING ORGANIZATION NAME AND ADDRESS TA'SK 10 PROGRAM ELEMENT PROJECT, AREA & WORK UNIT NUMBERS Toxicology Group, Division of Research Support 612720.835AA Letterman Army Institute of Research Presidio of San Francisco, CA 94129 11. CONTROLLING OFFICE NAME AND ADDRESS 12. REPORT DATE May 1984 US Army Medical Research and Development Command Fort Detrick 13. NUMBER OF PAGES Frederick, MD 21701 25 14 MONITORING AGENCY NAME & ADDRESS(If different from Controlling Office) 15. SECURITY CLASS (of this report, UNCLASS1F1ED 158 DECLASSIFICATION DOWNGRADING SCHEDULE 16. DISTRIBUTION STATEMENT (of this Report) THIS DOCUMENT HAS BEEN CLEARED FOR PUBLIC RELEASE AND SALE: ITS DISTRIBUTION IS UNLIMITED. 17 DISTRIBUTION STATEMENT (of the obstract entered in Block 20, If different from Report) 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Octahydro-1-(N)-Acetyl-3,5,7-Trinitro-1,3,5,7-Tetrazocine, 1,3,5,7-Tetrazocine, 1-Acetyloctahydro-3,5,7-Trinitro, SEX, Nitramines, Holston Army Ammunition Plant, Acute Toxicity, CAS Res. No. 13980-00-2 20. ABSTRACT (Continue as reverse side if necessary and identify by block number) The acute oral toxicity potential of the explosives by-product, 1-acetyloctahvdro-3, 5, 7-trinitro-1, 3, 5, 7-tetrazocine (SEX) was determined in male and female albino Fisher 334 rats by using a single dose, free choice feeding method. The study was conducted in compliance with the Good Laboratory Practice Regulations. No compound related mortality was observed at a limit dose of 5.0 g/kg. 1

DO 1 JAN 73 1473 EDITION OF ! NOV 65 IS OBSOLETE

UNCLASSIFIED SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

ABSTRACT

Ŕ

The Revenues

The acute oral toxicity potential of the explosives by-product, lacetyloctahydro-3,5,7-trinitro-1,3,5,7-tetrazocine (SEX), was determined in male and female albino Fisher 334 rats by using a single dose, free-choice feeding method. The study was conducted in compliance with the Good Laboratory Practice Regulations. No compound related mortality was observed at a limit dose of 5.0 g/kg.



i

PREFACE

TYPE REPORT: Acute Oral Toxicity GLP Study Report

TESTING FACILITY: US Army Medical Research and Development Command Letterman Army Institute of Research Division of Research Support Presidio of San Francisco, CA 94129

SPONSOR: US Army Medical Research and Development Command US Army Medical Bioengineering Research and Development Laboratory Fort Detrick, MD 21/01

PROJECT: 612720.835AA Acute Mammalian Toxicology Testing, APC, TL06

GLP STUDY NUMBER: 82005

STUDY DIRECTOR: COL John T. Fruin, DVM, PhD, VC, Diplomate of American College of Veterinary Preventive Medicine

PRINCIPAL INVESTIGATOR: Craig W. White, DVM, CPT VC

CO-PRINCIPAL INVESTIGATOR: Evelyn M. Zimmerman, SP5

PATHOLOGIST: MAJ Glen E. Marrs Jr., DVM, MS, VC, Diplomate of American College of Veterinary Pathologists

REPORT AND DATA MANAGEMENT: A copy of the final report, study protocols, raw data, retired SOPs and an aliquot of the test compound will be retained in the LAIR Archives.

TEST SUBSTANCE: 1-Acetyloctahydro-3,5,7-Trinitro-1,3,5,7-Tetrazocine (SEX)

INCLUSIVE STUDY DATES: 14 April - 10 May 1983

OBJECTIVE: To determine the acute oral toxicity potential of 1-acetyloctahydro-3,5,7-trinitro-1,3,5,7-tetrazocine (SEX) in male and female, albino, Fisher 344 rats.

ACKNOWLEDGMENTS

The authors wish to thank SP5 Leonard J. Sauers, MS, and SP5 Thomas P. Kellner, BS, for their assistance in the conduct of this study. In addition, we wish to thank Jesse Barkley Jr., US Army Medical Bioengineering Research and Development Laboratory, for his assistance as Project Consultant.

SIGNATURES OF PRINCIPAL SCIENTISTS AND MANAGERS INVOLVED IN THE STUDY:

We, the undersigned, believe the study number 82005 described in this report to be scientifically sound and the results in this report and interpretation to be valid. The study was conducted to comply, to the best of our ability, with the Good Laboratory Practice Regulations for Non-clinical Laboratory Studies, outlined by the Food and Drug Administration.

MAJ. MS Study Director

CRAIG W. WHITE CPT. VC Principal Investigator

lin E. Man, J. 22 aug 83 E. MARRS / DATE

GLEN E. MAJ, VC Pathologist

SP4. USA Co-Principal Investigator

83 Carolyn M. LEWIS, MS / DATE DAC

Data Manager



DEPARTMENT OF THE ARMY

LETTERMAN ARMY INSTITUTE OF RESEARCH PRESIDIO OF SAN FRANCISCO, CALIFORNIA 94129

REPLY TO ATTENTION OF

SGRD-ULZ-QA

3 April 1984

MEMORANDUM FOR RECORD

SUBJECT: Report of GLP Compliance

I hereby certify that in relation to LAIR GLP study 82005 the following inspections were made:

15 Apr 83 26 Apr 83 (1000 hours) 26 Apr 83 (1030 hours) 2 May 83

The report and raw data for this study were audited on 21 Mar 84.

Routine inspections with no adverse findings are reported quarterly, thus these inspections are also included in the Jul 83 report to Management and the Study Director.

Non 1 View

NELSON R. POWERS, Ph.D. DAC Chief, Quality Assurance Unit

Abstracti
Prefaceiii
Acknowledgmentsiv
Signature of Principal Scientistsv
Report of Quality Assurance Unitvi
Table of Contentsvii
BODY OF REPORT
INTRODUCTION
Objective of Studyl
METHODS
Test Substance
Mortality
BECOMMENDATION /
RECORDER DATION
APPENDICES
Appendix A, Chemical Data

OFFICIAL DISTRIBUTION LIST.....

Acute Oral Toxicity of 1-Acetyloctahydro-3,5,7-Trinitro-1,3,5,7-Tetrazocine--White and Zimmerman

The manufacture of the explosives, hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) and octahydro-1,3,5,7-tetranitro-1,3,5,7tetrazocine (HMX), at the Holston Army Ammunition Plant (HSAAP) results in the formation of a by-product, 1-acetyloctahydro-3,5,7trinitro-1,3,5,7- tetrazocine (SEX). It is formed by the nitrolysis and acetylation of hexamine. As a result, quantities of SEX are discharged from HSAAP. HSAAP is the only known producer of SEX. Its discharge, while partially mitigated by pollution abatement facilities at HSAAP, will continue and could increase during times of mobilization. Information on the chemical, physical, and toxicological properties of SEX is limited. Many of its properties can only be inferred by comparisons with RDX and HMX. Although no specific data are available, SEX, based on structural comparisons, appears to be more hydrophilic than either RDX or HMX and thus, potentially, a more serious toxicological threat than RDX or HMX to the aquatic life in the Holston River. This report summarizes the results from one of a series of studies being conducted at the Letterman Army Institute of Research (LAIR) to assess the toxicological hazards of SEX (1-3).

Objective of Study

The objective of the study is to determine the acute oral toxicity of l-acetyloctahydro-3,5,7-trinitro- 1,3,5,7-tetrazocine (SEX) in male and female, albino, Fisher 344 rats.

METHODS

Test Substance

The SEX was received from SRI-International (333 Ravenswood Ave, Menlo Park, CA 94025) on 7 January 1983. The bulk of the material (approximately 800 g) was stored at the Presidio Central Magazine Storage Facility. The magazine bunker is an underground, reinforced concrete structure used for the storage of explosive materials. Approximately 200 g was stored at the laboratory in a flame proof cabinet at room temperature. Chemical name:

l-Acetyloctahydro-3,5,7-Trinitro-1,3,5,7-Tetrazocine Chemical Abstract Service Registry Number: 13980-00-2 Structural formula:



Empirical formula: C₆H₁₁N₇O₇

Other test substance information appears in Appendix A.

Animal Data

A total of 44 (22 males, 22 females) Fisher 344 rats were received from Charles River Breeding Laboratory (Kingston, NY). Additional animal data are found in Appendix B.

Environmental Conditions

A commercially available certified rodent ration and tap water were provided <u>ad libitum</u> for the animals during the study. Appendix C is a listing of the environmental conditions of this study.

Dosing

Animals were randomly assigned to three study groups consisting of seven male and seven female rats per group. The study groups were designated as vehicle-control, cage-control and 5.0-g/kg limit- dose groups. The 5.0 g/kg dose is considered a no mortality limit dose (4). The limit dose was based upon the results of a pilot study performed between 14 March and 31 March 1983. Commercial grade peanut butter was used as the vehicle because of the insolubility and poor suspension properties of SEX. The limit-dose group and the vehicle control group initials were conditioned to consume b-6 g of peanat butter daily for b days before the day of dosing. All test animals were weighed on the day before dosing and the weight-based dosage was calculated. All limit-dose and vehicle control group animals were dosed on 26 April 1963. The calculated dose of the test compound was weighed on a balance, wixed with 5-0 g peanut butter, and placed in the individual rat cages in disposable petri dishes. The peanut butter/dose mixture was lift in the cages until it was consumed by the rats.

The dosing procedure was conducted without animal sedation or anesthesia.

Observations

Animals were observed daily throughout the quarantine/acclimation period. During the course of the study, recorded observations were conducted once daily. Additional observations were made on the day of dosing. Animals were observed daily after dosing, undisturbed in cages, outside of cages, and after their return to cages. The tirsuregularly scheduled observation period commenced at 1200 hours on the day of dosing.

Duration of Study

The study period was 15 days with a 13-day quarantine/acclimation period before the study was begun.

Historical study events are listed in Appendix D.

Deviation from Original Protocol

The room temperature elevated to 26.7 °C (86 °F) during the quarantine period; however, this was not considered to impact the study results.

One animal from the 5.0 g/kg dose group was removed from this study because the rat did not consume the entire dose amount. This particular animal also diluted the remaining dosing material with water from the automatic watering system. This made it impossible to determine the percent of the dose the animal received.

RESULTS

Mortality

All animals allocated to this study survived the entire test period. No compound-related deaths occurred at a limit dose of p.e. g/kg.

Clinical Observations

The only clinical sign observed in this study was that of irritability which was observed in three of the fourteen animals dosed. No animal exhibited signs of acute toxicity and/or mortality at the limit dose of 5.0 g/kg.

Gross Pathological Observations

No gross lesions attributable to SEX were observed at necropsy. A report of gross pathological observations appears in Appendix E.

DISCUSSION

The test compound, l-acetyloctahydro-3,5,7-trinitro-1,3,5,7tetrazocine (SEX) should be classified as relatively non-toxic by the oral route of administration (5). This, in all probability, can be attributed to the insolubility of the test compound. It is highly probable that the compound has little bioavailability as it is insoluble in aqueous and lipid solvent systems (3).

CONCLUSION

The test compound l-acetyloctahydro-3,5,7-trinitro-1,3,5,7tetrazocine (SEX) was classified as non-toxic, when administered orally, based upon this acute-limit test.

RECOMMENDATION

Any further safety testing planned for SEX should take into account the low oral toxicity of SEX which is probably attributable to its low solubility.

REFERENCES

- Tyson CA, Dilley JV, Sasmore DP, Spanggord RJ, Newell GW, Dacre JC. Single-dose and repeated-exposure toxicity of a complex wastewater from munitions manufacturing plants. J Toxicol Environ Health 1982;9:545-564.
- Dilley JV, Tyson CA, Spanggord RJ, Sasmore DP, Newell GW, Dacre JC. Short-term oral toxicity of a 2,4,6-trinitrotoluene and hexahydro-1,3,5-trinitro 1,3,5-triazine mixture in mice, rats, and dogs. J Toxicol Environ Health 1982; 9:587-610.
- 3. Bedford CD, Deas BD, Broussard MM, Geigl MA, Marynowski CW. Preparation and purification of multigram quanitities of TAX and SEX. Third Phase, Final Report, Fort Detrick, Maryland: US Army Medical Research and Development Command, December 1981.
- 4. Federal Register, Part IV, Environmental Protection Agency, Vol 44 No. 145 (para 72.112.21) July 1979.
- Doull J, Klaassen CD, Amdur MO, eds. Casarett and Doull's toxicology. 2nd ed. New York: MacMillan Publishing Co., Inc., 1980:12,18-22.

٠.

Page

Appendix A	Chemical Data9
Appendix B	Animal Datall
Appendix C	Environmental Conditions
Appendix D	Historical Listing of Study Events15
Appendix E	Pathology Report17

e - e 1

ei ei

5

Ľ٦

APPENDICES

CHEMICAL DATA

1. Chemical name: l-Acetyloctahydro-3,5,7-Trinitro-1,3,5,7-Tetrazocine (SEX) l-Acetyloctahydro-3,5,7-Trinitro-1,3,5,7-Tetrazocine; l,3,5,7-Tetrazocine,l-Acetyloctohydro-3,5,7-Trinitro (CA Name); Octahydro-1-Acetyl-3,5,7-Trinitro-S-Tetramine; l-Aceto-3,5,7-Trinitro-1,3,5,7-Tetrazacyclooctane; l-(N)-Acetyl-3,5,7-Trinitrocyclotetramethylenetetramine

Chemical Abstract Service Registry No.: 13980-00-2

Structural formula:



Empirical formula: C₆H₁₁N₇O₇ Molecular weight: 293.2 g/mole (calculated). Physical state: Solid at 20°C Melting Point: 224.2 - 224.7°C Density: 1.785 g/cc at 21°C pH: N/A nonaqueous Compound Refractory Index: Unknown Stability: After 48 hours at 75°C, there was no change (NMR, IR, color, or weight) in SEX samples tested. Purity: 99.9% Manufacturer: SRI International Menlo Park, CA 94205

APPENDIX A

4

2. Name: Peanut Butter

Ingredients: Peanuts, Dextrose, Salt, Hydrogenated Palm Oil, Mono and Diglycerides.

Physical state: Light brown, creamy style.

Name of contaminants percentages: None known.

Manufacturer: Golden State Brand Laura Scudder Snack Foods Division IC Industries Company Anaheim, CA

Lot Number: 30 6 1

APPENDIX A (concluded)

ANIMAL DATA

2. Animals were trained to consume 5-6 g of peanut butter from 21 Apr to 25 Apr 83.

Justification: Rats are a proven sensitive animal model for this test.

APPENDIX B

ENVIRONMENTAL CONDITIONS

Caging: Number/cage = 1; Type of cage = stainless steel, wire mesh bottom, battery type, no bedding, automatic flush.

Diet: Purina Certified Rodent Chow No. 5002 ad lib,

Water: Central line to cage battery with automatic lick dispensers

Temperature: 22 + 1°C

Relative Humidity: 50% + 10%

Photoperiod: 0530 - 2000 hours per day

APPENDIX C

White of a

HISTORICAL LISTING OF STUDY EVENUS

Date	Events
14-31 Mar 83	Pilot study accomplished.
13 Apr 83	Animals arrive at LAIR. They were sexed, observed for illness, ear tagged, weighed and caged in the GLP Suite.
14-25 Apr 83	Animals checked once daily.
22 Apr 83	Animals weighed and randomized into dose groups.
21-25 Apr 83	Animals weighed daily and conditioned to dosing procedure by feeding and recording peanut butter intake.
25 Apr 83	Animals weighed and dose calculated. Feed removed at 1630 hours.
26 Apr 83	Animals dosed by mixing test compound with 5-6 g peanut butter. Observations conducted one to two hours after dosing and at 1600 hours. Animals observed for clinical signs which were recorded. Rat chow made available as soon as the peanut butter was consumed.
27 Apr - 9 May 83	Animals observed for clinical signs at 0930 hours.
29 Apr - 2,6 May 83	All animals weighed.
9 May 83	Food removed at 1600 hours.
10 May 83	Animals observed for clinical signs at 0700 hours and weighed. Animals delivered to the PSG Necropsy Suite for sacrifice and gross necropsy by

0830 hours.

APPENDIX D

PATHOLOGY REPORT

GLP Study 82005

Acute Oral Toxicity Limit Study in Male Rats of Octahydro-1-(N)acetyl-3,5,7-trinitro-1,3,5,7-tetrazine (SEX), (CAS #13980-00-2)

History: The male Fisher 344 rats in this study were divided into 3 groups. All groups but the cage controls were fed 5-6 e of peanut butter vehicle or 5-6 g of peanut butter vehicle mixed with test compound at a dose rate of 5.0 g/kg body weight. The material dosed with and the number of rats in each group were as follow:

Group 1 (Cage controls) - 7 rats Group 2 (Vehicle controls) - 7 rats

Group 3 (5.0 g/kg SEX) - 7 rats

One of 7 rats in group 3 was misdosed and was removed from the study. All other rats survived until completion of the study, 14 days after dosing. The rats were killed by exsanguination from severed axillary vessels while under anesthesia produced by intraperitoneal injection of pentobarbital.

Gross necropsy findings: Necropsies revealed no test compound related gross lesions in male rats that were kiled at completion of the study. Seven of 7 rats in group 1, 7/7* rats in group 2, and 6/6 rats in group 3 had bilaterally enlarged inguinal (prepucial) glands that were turgid and filled with yellow-green granular material that was most likely inspissated secretory material. External examination revealed a subcutaneous nodule** in the inguinal area of 1/7 rats in group 1 and 1/6 rats in group 3. The skin over the nodule in the rat in group 3 was umbilicated and ulcerated. The eyes of 1/7 rats in group 2 had red-black optic disks and red retinal foci and the right eye of 1/6 rats in group 3 had a red retinal focus. The red-black optic disks and the red retinal foci may have represented hemorrhage. A cause for the lesions in the prepucial glands and the eyes was not determined.

*Number of rats affected/Number of rats in the group.

**Microscopic examination of the subcutaneous nodules in the inguinal area of the rat in group 1 and the rat in group 3 revealed ruptured prepucial glands that had spilled secretion material into adjacent subcutis. Both the subcutis and the lumen of the gland contained abundant pyohistiocytic inflammatory cell infiltrate and cell debris.

APPENDIX E

Summary: No test related gross or microscopic lesions were observed in male Fisher 344 rats that were cage controls, vehicle controls, or dosed with 5.0 mg of SEX.

Alm E. Mann J.

GLEN E. MARRS, JR., DVM, MS Diplomate, A.C.V.P. MAJ, VC Assistant Chief, Pathology Services Group Division of Research Support

9 August 1983

TABLE I

Acute Oral Toxicity Limit Study in Male Rats of SEX, (CAS #13980-00-2) - GLP Study 82005

	Group 1								Group 2							Group 3						
ID# 83D00xxx	2 9 5	3 0 2	3 1 0	3 1 3	3 2 1	3 2 2	3 2 9	2 9 6	2 9 7	3 0 8	3 1 5	3 2 3	3 2 4	3 2 6	2 9 7	2 8 9	2 9 2	2 9 3	304	3 1 2	3 1 6	
Survived to Completion	+	+	+	+	+	+	+	+	+	+	+	÷	+	+	+	+	+	+		+	+	
Removed - misdose																			+			
Inguinal (Prepucial) Glands, enlarged	+	+	+	+	+	+	+	÷	+	+	+	+	+	+	+	+	+	÷		+	+	
Ckin:																						
(1) Subcutaneous inguinal nodule		÷															+					
(2) 'Imbilicated & ulcerate over inguinal nodule	edi																+	,				
∑ye:																						
(1) Optic disk, red-black									•	F												
(2) Retina, red fous/foli									4	ł								+	•			
															Α	ΡP	EN	ID I	Х	E	(cont.)

PATHOLOGY REPORT

GLP Study 82005

Acute Oral Toxicity Limit Study in Female Rats of Octahydro-1-(1)acetyl-3,5,7-trinitro-1,3,5,7-tetrazine (SEX), (CAS #13980-00-2)

History: The female Fisher 344 rats in this study were divided into 3 groups. All groups but the cage controls were fed 5-6 g of peanut butter vehicle or 5-6 g of peanut butter vehicle mixed with test compound at a dose rate of 5.0 g/kg body weight. The material dosed with and the number of rats in each group were as follow:

Group 1 (Cage controls) - 7 rats

Group 2 (Vehicle controls) - 7 rats

Group 3 (5.0 g/kg SEX) - 7 rats

All rats survived until completion of the study, 14 days after dosing. The rats were killed by exsanguination from severed axillary vessels while under anesthesia produced by intraperitoneal injection of pentobarbital.

Gross necropsy findings: Necropsies revealed no test compound related gross lesions in female rats that were kiled at completion of the study. One of 7 rats in group 3 had a fluid filled cyst in one ovary and $1/7^*$ rats in group 1, 1/7 rats in group 2, and 1/7rats in group 3 had bilaterally enlarged inguinal (clitoral) glands that were turgid and were filled with yellow-green granular material. The ovarian cyst was considered to be an incidental finding. A cause for the enlargement of the clitoral glands was not determined.

Summary: No test related gross lesions were observed in female Fisher 344 rats that were cage controls, vehicle controls, or dosed with 5.0 mg/kg of SEX.

*Number of rats affected/Number of rats in the group

Vien E. Man, Dr.

GLEN E. MARRS, JR., DVM, MS Diplomate, A.C.V.P. MAJ, VC Assistant Chief, Pathology Services Group Division of Research Support

9 August 1983

APPENDIX E (cont.)

and the second states of the second second

1.41.4 A 2.4 A

TAPLE I

Acute Tral Coxicity Limit Study in Female Rats of SEX, (CAS #13980-00-2) - FLP Study B2005

· • .-

	Group 1								Group 2								Group 3						
10#-83000xxx	ч ч б	7 1 5	א ה 2	3 5 7	3 5 1	5. 6 2	3 7 9	5 3 2	3 3 4	373	3 7 5	375	3 7 8	33 30 0	3 4 3	3 4 5	3 5 4	355	3 5 5	371	K. T - C.		
Survived to Completion	+	+	÷	+	+	+	+	+	٠	+	+	+	÷	+	+	+	+	+	٠	+	+		
No lesions recognized	+	+	+		+	+	+	+	+	ł	÷		+	÷	+	+	+	+		÷			
Inguinal (Clitoral) glands, enlarged				+								+									٠		
Ovary:																							

Syst. fluid-filled. unilateral

APF "DIX E (concluded)

OFFICIAL DISTRIBUTION LIST

.

Commander US Army Medical Research and Development Command AFTN - SGRD-RMS Mrs. Madigan Fort Petrick, Frederick MD 21701

Defense Technical Information Center ATTN DTIC-DDA (12 copies) Cameron Station Alexandria VA 22314

Director of Detense Research and Engineering ATTN - Assistant Director, Environmental and Life Sciences Washington DC 20301

The Surgeon General ATTN DASG-TLO Wishington DC 20314

HQ DA (DASG-ZXA) WASH DC 20310

Commandant Academy of Health Sciences ATTN: HSHA-CDM Fort Sam Houston TX 78234

Assistant Dean Institute and Research Support Uniformed Services University of Health Sciences 6917 Arlington Road Bethesda MD 20014

Commander US Army Environmental Hygiene Agency Aberdæn Proving Ground MD 21070

US Army Research Office ATTN Chemical and Biological Sciences Division P.O. Box 1221 Research Triangle Park NC 27709

Biological Sciences Division Office of Naval Research Arlington VA 22217

Director of Life Sciences USAF Office of Scientific Research (AFSC) Bolling AFB Wishington DC 20332 Director Walter Rend Army Estimate of Hele $\tau_{\rm Cl}$ Washington DC -2030

Commander US Army Medical Research Institute of Infective 3 Discusses Fort Detrick, Frederick MD 21761

Commander US Army Research Institute of Environmental Medicine Natick MA 01760

Commander US Army Institute of Surgicul Research Brooke Army Medicel Center Fort Sam Houston TX "8234

Commander US Army Medical Bioengineering Research and Development Laboratory Fort Detrick, Frederick MD 21701

Commander US Army Aeromedical Research Laboratory Fort Rucker AL 36362

Commander US Army Research Institute of Chemical Defense Aberdeen Proving Ground Edgewood Arsenal MD 21010

Commander Naval Medical Research Institute National Naval Medical Center Betnesda MD 20014

Commander USAF School of Aerospace Medicine Acrospace Medical Division Brooks Air Force Base TX 78235

