F	AD-R142 772 SOURCES OF ERROR IN OBJECTIVE ANALYSIS(U) NAVAL 1/1 POSTGRADUATE SCHOOL MONTEREY CA R FRANKE MAY 84 NP553-84-0003										'1		
UNCLASSIFIED F/G 12/1									NL				
			*			•							
			-										

1.	•.
3	-
14	
ð.	
5	
52	
2.5	
3.1	
3.2	
1.1	
3	
102	
1	
1.9	
1.2	
	· · · · · · · · · · · · · · · · · · ·
4	
24	
$a_{\rm eff} = 1$	1/1 3.6
- X	
¥.	
14	
19	
× *	
Ť.	

MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS-1963-A

.



AD-A142 772

# NAVAL POSTGRADUATE SCHOOL Monterey, California





05

07

107

SOURCES OF ERROR IN OBJECTIVE ANALYSIS by

Richard Franke

May 1984

Technical Report For Period

October 1983 - March 1984

Approved for public release; distribution unlimited

Prepared for: Naval Environmental Prediction Research Cacility Monterey, CA 93943 NAVAL POSTGRADUATE SCHOOL MONTEREY CALIFORNIA 93943

R. H. Shumaker Commodore, U. S. Navy Superintendent D. A. Schrady Provost

This work was funded by the Naval Environmental Prediction Research Facility, Monterey, CA under Program Element 61153N, Project (none), "Interpolation of Scattered Meteorological Data".

Reproduction of all or part of this report is authorized.

Richard Franke Associate Professor of Mathematics

Reviewed by:

Gordon E. Latta, Chairman Department of Mathematics

Released by:

Kneale T. Marshall Dean of Information and Policy Sciences

REPORT DOCUMENTATION PAGE	READ INSTRUCTIONS
1. REPORT NUMBER	D. 3. RECIPIENT'S CATALOG NUMBER
NPS-53-84-0003	77.2
4. TITLE (and Subtitie)	S. TYPE OF REPORT & PERIOD COVERED
Sources of Error in Objective Analysis	Technical Report
	Interim, FY 1984
	6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(a)	8. CONTRACT OR GRANT NUMBER(*)
Richard Franke	
. PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TASK
Naval Postgraduate School	
Monterey, CA 93943	Program element 61153N
1. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE
Naval Environmental Prediction Research Facility	May 1984
Monterey, CA 93943	13. NUMBER OF PAGES
14. MONITORING AGENCY NAME & ADDRESS(If different from Controlling Office)	18. SECURITY CLASS. (of this report)
Naval Air Systems Command	Unclassified
Washington, DC 20361	154. DECLASSIFICATION/DOWNGRADINA
Approved for public release. distribution unlimit	ted
·····	
<ul> <li>DISTRIBUTION STATEMENT (of the abstract entered in Block 20, 11 different in Block</li></ul>	cen Report)
<ul> <li>DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different fr</li> <li>SUPPLEMENTARY NOTES</li> <li>KEY WORDS (Continue on reverse elde if necessary and identify by block number Objective analysis Barnes' method Optimum interpolation Splines</li> </ul>	oen Report)
<ul> <li>7. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different fr</li> <li>8. SUPPLEMENTARY NOTES</li> <li>8. KEY WORDS (Continue on reverse eide if necessary and identify by block number Objective analysis Barnes' method Optimum interpolation Splines Thin plate splines Statistical inter</li> </ul>	oen Report)
<ul> <li>DISTRIBUTION STATEMENT (of the obstract entered in Block 20, 11 different in</li> <li>SUPPLEMENTARY NOTES</li> <li>KEY WORDS (Continue on reverse elde 11 necessary and identify by block number Objective analysis Barnes' method Optimum interpolation Splines Thin plate splines Statistical inter</li> <li>ABSTRACT (Continue on reverse elde 11 necessary and identify by block number) The error in objective analysis methods that to a first guess field is considered. An expres decomposition of the error into three independen To test the magnitudes of the contribution of ea computer simulations was conducted. Grid-to-obs schemes considered ranged from simple piecewise accurate spline functions. The observation-to-g</li> </ul>	polation t are based on corrections sion that gives a t components is derived. ch component a series of ervation point interpolation linear functions to highly rid interpolation methods
<ul> <li>DISTRIBUTION STATEMENT (of the abstract entered in Block 20, 11 different 20, 11 different 20, 11 different 20, 20, 20, 20, 20, 20, 20, 20, 20, 20,</li></ul>	polation t are based on corrections sion that gives a t components is derived. ch component a series of ervation point interpolation linear functions to highly rid interpolation methods

# UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

considered included most of those in present meteorological use, such as optimum interpolation and successive corrections, as well as proposed schemes such as thin plate splines, and several variations of these schemes. The results include an analysis of cost versus skill; this information is summarized in plots for most combinations. The degradation in performance due to inexact parameter specification in statistical observation-to-grid interpolation schemes is addressed. The efficacy of the mean squared error estimates in this situation is also explored.



Acces	ssion For	r /
NTIS	GRANI	<u>H</u>
DTIC	TAB	
Unan:	pecariot	
Just	<i>fication</i>	2
By		
Distr	ibution/	/
Avai	lability	Codes
·····	Avail a	ud/or
Dist	Specia	al
A /		
4~I		
Diry Person		
Adia OFY FETED		
Adia OFT FETRD		

S/N 0102- LF- 014- 6601

The second s

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

### 1.0 Introduction

The purpose of this investigation was to test the relative importance of various aspects of correcting predicted values on a grid by incorporating information from observed values at scattered data points. Grid and observation configurations were patterned after those routinely available over North America. Although investigations were limited to the univariate objective analysis methods, I believe the results are indicative of those that would be achieved in the more general case.

Previous investigations on the error contribution of various steps in the objective analysis process are limited. Koehler (1979) separately studied the errors of a number of grid-toobservation and observation-to-grid interpolation (approximation) He noted that although little attention is typically routines. paid to the grid-to-observation interpolation process significant errors may be caused by this phase of objective analysis. While this may be a surprise since these errors are usually small compared to the first-guess errors at the grid points, my results further demonstrated that the contribution to overall error made by the grid-to-observation interpolation process should not be ignored. This investigation complements recent work by Seaman (1983) regarding the accuracy of statistical and successive correction schemes. His work provides expected mean squared error estimates for these schemes. His work is very thorough in that it provides estimates of the analysis error as the parameters of the first-quess error are varied while holding the assumed values constant, and vice-versa.

In Section 2 I derive a generalized expression for the overall error in objective analysis which leads to several observations. In Section 3 I describe the simulation method and the various options which can be easily handled. In Section 4 I present the results of the simulations and discuss their implications with regard to the observations made in Section 2.

2.0 The Form of the Error Term in Objective Analysis

My setting for study of the objective analysis process assumes the following:

(i) The true field (function) to be analyzed is H.

(ii) H is known imperfectly at grid points through a "firstguess" which is in error by an amount to be denoted by g. The error is a normally distributed stationary random function which has a certain spatial correlation and standard deviation.

(iii) H is imperfectly measured at observation points yielding values with errors o. These errors are independent and normally distributed with certain standard deviation.

The nature of the errors makes it only possible to evaluate g at grid points, and o at observation points, although it is sometimes convenient to think of them as functions rather than as sets of errors. The objective analysis process consists of interpolation of the first-guess values from the grid to the observation points (by a linear operator designated M) followed by interpolation of the difference between the observed and firstguess values back to the grid point (by a linear operator designated L) as a correction to the first-guess values. Denote the error in the entire process by E, then the final approximation is

H + E = H + g + L(H + o - M(H + g)).

Let m(H) represent the error in the approximation of H by H(H), then M(H) = H - m(H). Rearranging and simplifying the above, leads to

E = g + L(H + o - M(H) - M(g))= g + L(H + o - H + m(H) - M(g)) = g + L(o + m(H)) - LM(g) ,

and finally,

E = L(o) + Lm(H) + (g - LM(g)) . (1)

Thus the error is made up of three parts. The term L(o) is dependent on the 'function' o, which describes instrumentation error and is typically not controlable. It is obviously advantageous to have o small. Since the values of o are assumed independent and random it is desirable for L to be a smoothing operator. The second part, Lm(H) is within our control and the grid-to-observation point interpolations error should be made small. If it is, then interpolation of the error back to the grid points by L is also small, assuming this smoothing operator is typical and does not magnify the error. The third part (q -LM(g)) is the error in interpolation of the first-quess error at the grid points to the observation locations by M, then back to the grid points by L. While it is possible that a certain symbiosis between parts could occur, the goal is certainly for each interpolation process to have small errors. Ideally the operator L should be a left inverse of the operator M, although this is almost certainly impossible.

Partitioning the error in this way shows, for example, that

using a better interpolation process from the grid to the observation points should decrease the overall analysis error. In certain realizations, of course, the errors may tend to cancel. Since the three terms represent uncorrelated errors, the total error variance over many realizations will tend to be the sum of the individual variances. Thus, decreasing any one will lead to statistically smaller error variances.

# 3.0 The Computer Simulation Methods

In order to simulate the behavior of the overall error under various interpolation processes and first-guess error assumptions, a modular computer program was written to give several options for the different processes. This made it possible to test a large number of combinations of methods and assumptions.

In general terms, the process simulated consists of the following steps:

 (i) An underlying mathematically defined function describing the field to be analyzed is evaluated on a grid of points.

(ii) "First-guess" error is generated from normal random deviates with a pre-specified standard deviation and spatial correlation.

(iii) "Observed values" are generated by evaluating the field to be analyzed at the observation points, and adding normally distributed uncorrelated random deviates to these values.

(iv) The first-guess values at the observation points are obtained by one of several interpolation schemes.

(v) Based on the difference between first-guess and observed values at the observation locations, "corrected" values at the grid points are obtained. I will refer to the corrected values as the analysis values.

Most of the simulations were done with two different grids and observation point sets. One was based on a 2.5° grid covering 112.5° W to 82.5° W and 30° N to 50° N, with 117 = 13x9 grid points and 36 observation points within the grid, as shown in Figure 1. The other was based on a 5° grid covering  $125^{\circ}$  W to 75° W, and 25° N to 50° N, with 88 = 11x8 grid points, and 57 observation points within the grid. This grid and the observation locations are shown in Figure 2. All the simulations used were univariate analysis methods on a two-dimensional field. This simplification was necessary for two reasons. The first reason is that the generation of error with a specified spatial correlation required factorization of the correlation matrix into the product of a lower triangular matrix and its transpose. The correlation matrix is of order equal to the number of grid points, and it is not particularly well conditioned. Incorporation of multiple levels, a large grid, or correlated multiple variables was therefore not possible. The other reason is that statistical results required that numerous realizations be simulated, thereby limiting the time available to do the computations.

The underlying mathematically defined field can be any specified function. The height field test function used is the one given by Koehler (1979) and also described in Wahba and Wendelberger (1930). The input parameters,  $\theta_0$  (the location of

the longitudinal wave),  $\Delta \Theta$ , (amount part of the field is skewed logitudinally), and p (the pressure for the height field) are easily varied. The experiments simulated the 500 mb height field, using fixed or randomly varying  $\theta_0$  and  $\Delta \theta_0$ . A typical field of height contours generated by this function is shown in Figure 3. First-guess errors had a nominal standard deviation,  $r_{\alpha}$ , of 30 m. The spatial correlation function was modeled using  $\exp((-d/c_d))^2$ , where d is distance (on the degree grid), and  $c_d$ is a correlation distance, specified as 10°. I have used degree measure for distance rather than true distance, to maintain a rectangular grid of first-guess points. This resulted in a distortion of the distance varying with location. The observation errors had a nominal standard deviation, ro, of 10 m. The observation locations approximately correspond to the North American radiosonde network within the grids being used. They are shown, along with the grids, in Figures 1 and 2.

The output consisted of mean, root-mean-square, and maximum errors over each data set (first-guess at grid points, firstguess at observation locations, observation at observation locations, and analysis values at grid points) for each realization. The first and third of these mainly served as a chack on the psuedo-random number generator (IMSL subroutines GGNSM and GGNML). The output also gave summaries of the same errors over all realizations as well as the mean and standard deviation of the root-mean-square errors over the realizations. Interpolation processes are sometimes ill-behaved around boundaries. Since in the global problem this can be avoided, the effects were minimized here by tabulating error only over the interior grid

points. Thus the results are over 77 grid points on the 2.5° grid and 54 grid points on the 5° grid. The options simulated for each step are described below.

a. Grid-to-observation point interpolation

First-guess values at the observation points are obtained by interpolation from the first-guess grid values. I compared four schemes. Others could be easily included, however my results indicate it will probably not be fruitful to do so. The methods I have used are:

(i) Piecewise bilinear interpolation. As with any piecewise defined method, one must first determine the rectangle in which the avaluation point lies. Then, the evaluation is most easily seen as translation to the square  $[0,1]^2$ , followed by 3 one dimensional interpolations. This requires 8 operations, where an operation is defined as a multiplication or division followed by an addition or subtraction. Practically, the evaluation can be accomplished in 5 operations (and a couple of extra additions/subtractions). In my cost analysis I have used 6 operations; this cost is very low compared to that of other necessary calculations.

(ii) Bicubic spline interpolation. I used the LMSL subroutines IBCCCU and IBCEVL. Preprocessing for the spline coefficients on a  $N_{\Theta}xN_{\phi}$  grid requires  $12N_{\Theta}xN_{\phi}+27N_{\phi}+51N_{\Theta}-10$  operations. Evaluation requires 2 operations to translate to  $[0,1]^2$  and 5 cubic interpolations at 9 operations each for a total of 47 operations. The preprocessing operations involve solution of tridiagonal systems of equations which are amenable to vectoriza-

tion for pipeline computers.

(iii) Piecewise bicubic interpolation. My implementation of this scheme used 2 operations for a translation to  $[0,3]^2$ followed by 5 cubic interpolations, each costing 5 operations. In addition, a difference table was formed at a cost of several subtractions.

(iv) Bessel bicubic interpolation. My implementation of this scheme used 2 operations for a translation to [0,3]<sup>2</sup> followed by 5 cubic interpolations, each costing 5 operations. Because of default to parabolic interpolation in boundary regions, there were some additional tests. There were also a few subtractions to form the difference table.

## b. Observation-to-grid point interpolation

As in operational weather forecasting programs, the differences between first-guess and observed values at the observation points are used to correct the first-guess values on the grid to obtain analysis values on the grid. I have tested twelve schemes for performing this correction. I will give a brief description of each method and refer the reader elsewhere for complete details. The first-guess error at the observation location,  $P_k =$  $(\Theta_k, \phi_k)$ , is denoted by  $\Delta H_k$ ,  $k=1,\ldots,N_0$ . The number of grid points is  $N_\Theta N_{\Phi}$ . I want to evaluate the approximation at grid points, but will write it in terms of a generic point,  $P = (\Theta, \phi)$ . Recall that the standard deviation of the first-guess errors is  $r_{\alpha}$ , and the spatial covariance function is denoted by C(P, g).

An operation count has been made for each of the methods. I discuss briefly how various phases of the process contribute, and

summarize the results in Table 1, along with some representative numbers that arise from my simulations. I have described some schemes as local, implying that others are global. In the context of global objective analysis, all the schemes 1 consider are local; the schemes which are global for my simulation are less local than the ones I refer to as local.

(i) Optimum interpolation (OI). This scheme was introduced to the meteorological literature by Gandin (1953) and has received widespread attention in recent years, e.g. see Borgman (1979) and Lorenc (1981). The method in its proper form requires that the spatial covariance function of the first-guess errors and the standard deviation of the observation error be known. Since these are known for this simulation, I have used their properties. I have implemented the scheme as described in Franke and Gordon (1983), viewing the approximation as a linear combination of the covariance functions associated with the observation points. Thus we have

$$\Delta H(P) = \sum_{k=1}^{N_{O}} a_{k}C(P,P_{k}),$$

where C(P,Q) is as noted above. The  $a_k$  are determined from the system of equations

$$\left( C(P_{i}, P_{j}) + \delta_{ij} r_{o}^{2} \right) \quad \left( \begin{array}{c} a_{1} \\ \vdots \\ a_{N_{o}} \end{array} \right) = \left( \begin{array}{c} \Delta H_{1} \\ \vdots \\ \Delta H_{N_{o}} \end{array} \right)$$

where  $\Delta H_i$  is the difference between the first-guess and observed values at the i<sup>th</sup> observation point,  $r_o$  is the standard deviation of the observation error, and  $\delta_{ij}$  is the Kronecker delta.

The cost of (OI) consists of a preprocessing phase that

includes the generation and solution of the system of equations, followed by evaluation of the analysis at the grid points. For  $N_0$  observations, preprocessing is at a cost of  $N_0(N_0+1)/2$  function evaluations to generate the coefficient matrix and  $(N_0^3+6N_0^2-N_0)/6$  operations plus  $N_0$  square roots to perform Cholesky decomposition and solution of the system of equations for the  $a_k$ . Evaluation costs  $N_0$  covariance function evaluations and  $N_0$  operations to form the linear combination representing the value of the correction at each grid point.

(ii) Local optimum interpolation. In my version of this scheme, nominally only points within the surrounding  $10^{\circ}$  square are used; if fewer than 4 observations are available, the square is expanded to  $15^{\circ}$  and so on, by  $2.5^{\circ}$  increments in each direction until at least 4 observations are available. The costs of the search were not assessed. For each grid value correction, a system of equations must be formed and solved, and the corresponding correction computed. With n observations being used the expressions given for OI above apply with n replacing N<sub>o</sub>. This process was performed for each grid point, making the total cost the sum of these costs over all grid points.

(iii) Global Barnes' method. This type of scheme is described by Barnes (1973) and others. My scheme used the known correlation functions as the weights for the first pass. Thus, the approximation is

 $\Delta H_{\star,1}(P) = \sum_{k=1}^{N_o} \frac{W_o}{k} (P) \Delta H_k / \sum_{k=1}^{N_o} W_k(P),$ where  $W_k = \exp((-(||P-P_k||/c_d)^2))$ , and  $\Delta H_k$  is as before. For the

second pass the correction has the same form, but  $\Delta H_k$  is replaced by  $\Delta H_{k,1}$ , the difference between the corrected first-guess and the observations. The quantity  $c_d$  is replaced by  $c_d/3^{1/2}$  for the second pass. The total correction at the grid points is then the sum of the two corrections. For each grid point the cost of this method is N<sub>o</sub> weight function evaluations per pass and N<sub>o</sub>+1 operations per pass. In addition a separate interpolation from the grid points to the observation points is required before the second pass. This type of scheme has been defined and studied in a different context, without a change of weight functions between iterations, by Foley and Nielson (1989).

(iv) Local Barnes' method. The same localization process as used for the local OI scheme (ii) was used here. As for the global version, two passes were used. Hence the cost for an evaluation at a grid point with n neighboring points is the same expression as in the global scheme, but with n replacing  $N_0$ . In addition, there was the search cost to determine the nearby observations, which was not assessed. Costs of an interpolation from the grid points to the observation points between passes was included.

(v) Statistical interpolation  $(c_d = 14^\circ)$ . In practical applications of OI the error correlations and standard deviations cannot be modeled precisely. This has lead to the use of the name "statistical interpolation". Computationally the method is identical to the OI scheme (i). Here the only difference is the substitution of an inexact correlation distance,  $c_d = 14$ . The algorithm and costs are identical.

(vi) Statistical interpolation ( $c_d = 7^\circ$ ). Again this is

identical to (i) except that the inexact value substituted for  $c_a$  is 7.

(vii) Statistical interpolation (damped cosine correlation function). Once more this scheme is computationally identical to (i) except that the correlation function used is of the form  $exp((-(||P-Q||/c_d)^2)cos((||P-Q||/c_d)(/2))$ . I used the value  $c_d$ = 10.

(viii) Thin plate splines. This method is described by Wahba and Wendlelberger (1980) and others. The approximating function used by the scheme is

$$H(P) = \sum_{k=1}^{N_{o}} A_{k}B(P,P_{k}) + a \vartheta + b \phi + c,$$
  
$$k=1$$

where the basis function  $B(P,Q) = ||P-Q||^2 \log ||P-Q||$ . The  $A_k$ and a, b, and c, are obtained by solving the system of equations

$$N_{o}$$

$$\Sigma A_{k} (B(P_{i}, P_{j}) + \lambda N_{k} r_{o}^{2} i_{j}) + a \vartheta_{i} + b \varphi_{i} + c = \Delta H_{i}, i = 1, ..., N_{o}$$

$$j = 1$$

$$N_{o}$$

$$\Sigma A_{j} \vartheta_{j} = \emptyset$$

$$j = 1$$

$$N_{o}$$

$$\Sigma A_{j} \varphi_{j} = 0$$

$$j = 1$$

$$N_{o}$$

$$\Sigma A_{j} = \emptyset$$

$$j = 1$$

In the above,  $\lambda$  is a smoothing parameter. The smoothing parameter was chosen on the basis of a few trials with no attempt to optimize its choice for a particular data set, as can be done. Wendelberger (1931) describes a program that will automatically choose  $\lambda$  (and m as well, see next method), but I have not tested

it yet. This system of equations is symmetric, but not positive definite. I have used standard L-U decomposition routines to solve the system. Methods for symmetric indefinite systems use about half as many operations, however I observed greater numerical stability using the general decomposition process. There are  $N_0(N_0+1)/2$  basis function evaluations, and solution of the system of equations requires  $(N_0+3)(N_0^2+5N_0+3)/3 + (N_0+3)^2$  operations. Unlike symmetric positive definite systems, solution of these equations requires searching for a pivot and pivoting. Evaluation at each grid point then requires  $N_0$  basis function evaluations and  $N_0+2$  operations to form the sum.

(ix) Laplacian smoothing spline (m=3). This scheme is also described by Wahba and Wendelberger (1987), and is one of those available in the program by Wendelberger (1981). The thin plate spline method is a member of this family (with m=2), but also has the "thin plate" interpretation. The reason for inclusion of this method is that the results of Wahba and Wendelberger indicate that pressure height fields are better approximated using values of m = 3 or 4. I will not describe the method fully. It requires evaluation of  $N_0(N_0+1)/2$  basis functions and  $3N_0$  multiplications to set up the system of  $N_0+6$  equations to be solved. Then  $N_0+5$  operations would be required for evaluation at each grid point, along with the evaluation of  $N_0$  basis functions.

(x) Franke/Gordon. This scheme was suggested by Franke and Gordon (1983) as one which is an explicit scheme, similar to Barnes' method, but which when iterated converges to the OI interpolant. Three iterations, with the parameter = .85[[M]]

(in the notation of that report) were performed. The cost in operations is  $2N_O(N_O+1)$  plus  $3N_O$  for each grid point. The number of weight function evaluations is  $2N_O^2$  plus  $3N_O$  for each grid point.

(xi) Pseudo-Barnes' method. This method was described in Franke and Gordon (1983) and was at that time mistaken for Barnes' method. It differs in that the error at the second iteration is Barnes' approximation evaluated at the observation point minus the first-guess error, rather than the the corrected first-guess at the grid point interpolated to the observation point minus the observed value. The cost of this algorithm is evaluation of  $N_0^2$ weight functions plus  $2N_0$  for each grid point. It requires  $N_0(N_0+1)$  operations, plus  $2(N_0+1)$  for each grid point.

(xii) Local pseudo-Barnes' method. This is a local version of (xi), using the same "nearby" observation points as (ii) and (iv). A grid point with n nearby observation points requires evaluation of  $n^2+2n$  basis functions and  $n^2+3n+2$  operations.

#### 4.0. Results

The simulation program described in the previous section was run for a substantial number of different options. Each run consisted of 100 realizations of a test field each containing associated first-guess and observation errors. Table 2 gives the assumed parameter values for the various cases. Not all combinations of grid-to-observation point and observation-to-grid point interpolation schemes were used in every case. The tables detail the complete results and the entries indicate which combinations were computed. Each combination in a given table (3-14)

corresponds to the same set of realizations, but different tables depend on different realizations.

This investigation was designed to determine the influence of the grid-to-observation point interpolation scheme. This influence is seen by noting changes in error for a particular observation-to-grid point interpolation scheme as the grid-toobservation point interpolation scheme is varied. The rows of Tables 3-14 give this information. The bicubic spline interpolation produced significant improvement over piecewise bilinear interpolation. This verifies the smaller magnitude of the term Lm(H) in the error expression given by (1) for the spline method. For 2.5° grids the errors were no smaller for spline interpolation than for piecewise bicubic or Bessel bicubic interpolation. Evidently the grid spacing was small enough (for the test function used) that the interpolation error was not significant. Spline interpolation did show an improvement over piecewise bicubic and Bessel bicubic interpolation on the 5° grid. Spline interpolation and the cubic interpolation methods showed even greater improvement over piecewise linear interpolation on the  $5^{\circ}$  grid than on the 2.5° grid. Interestingly the first-guess errors at the observation points had greater rms values for cubic interpolation than they did for linear interpo-This occurs because linear interpolation inherently has lation. greater smoothing.

Most of the useful information given in Tables 3-14 can be more easily obtained from plots of the salient values. Figures 4-8 give plots of skill vs. cost of the algorithm in thousands of

operations per analysis. Here "skill" is defined to be 1 rmsa/r, where rmsa is the rms error in the analysis values. The skill with respect to bilinear and bicubic interpolation are each indicated, connected with a straight line to delineate the extent between the two. The results for only one of the statistical schemes, (vi), has been plotted since the others were nearly identical. For these purposes I counted an evaluation of a basis, weight, square root, or covariance function as 10 operations. The plots reveal that the statistical schemes, local OI, and thin plate splines all had close to the same accuracy and all were slightly less accurate than OI. The Barnes' schemes, the Franke/Gordon scheme, and Laplacian smoothing splines were least accurate. The poor performance of the Laplacian smoothing splines here, in contrast to the better performance obtained by Wahba and Wendelberger (1980) is probably due to the scheme being applied to the first-guess error function rather than to the underlying true height field. The degradation in the performance of the less than optimal statistical schemes is perhaps less drastic than one might expect. It does appear that it was better to underestimate the correlation distance than to overestimate it.

Figure 9 shows plots of the rms errors in the analysis values as a function of first-guess errors. The improvement in the Barnes' scheme as the first-guess errors decrease was rapid. The scheme gave results nearly as good as OI, the statistical schemes, and thin plate splines. This occurred because the principal problem became smoothing observation errors as the first-guess errors tended to zero. Figure 10 shows plots of the

rms errors in the analysis values as a function of observation errors. As observation errors go to zero the importance of modelling the first-guess error was more important than smoothing. Thus OI, the statistical schemes, and thin plate splines improved the most, while both Barnes' schemes improved little. Figures 11-13 show the rms errors in the analysis values when incorrect variances were specified for the interpolation routines. Methods not using these values were naturally unaffected so that changes in the rms errors in the analysis values for these methods only reflect the variability of the (different) realizations used in the various cases. The plots show that the use of incorrect values for the first-guess and observation error variances did not drastically affect the accuracy of the statistical methods. The interested reader is referred to Seaman (1983) for more extensive tests of the effects of incorrect parameter specification on the performance of statistical interpolation methods.

One of the attractive features of the statistical schemes is that they afford a calculation for the estimated mean squared error. These estimates do not depend on any particular realization, so they were not incorporated into the process. However, I did compute them as a side calculation for my grids and observation points. The results of these calculations are tabulated for the 2.5° grid, along with the empirical rms errors obtained during the simulations. Table 15 shows that the estimates given by OI were quite good; the estimated and empirical errors varied only a few percent. They also were accurate for local OI, as

they should be. On the other hand, the slight degradation in performance of statistical methods when incorrect correlations or variances were specified did not carry over to the error estimates. In fact the schemes that have their performance degraded the most (in this case, using too long a correlation distance) showed a decrease in the estimated error variance. Conversely, shortening the correlation distance in the statistical method increased the error estimate as well as the empirical error obtained, although the empirical error is underestimated. This indicates that one must not put too much faith in the error estimates when the actual covariance structure is not known, as in practice. It appears one could obtain just about any error estimate wished simply by specifying unrealistic parameters for the covariance structure.

The principal results of this study were as follows. The decomposition of the error into independent components in (1) identified possible ways to decrease the analysis error. This lead to the results showing the contribution of the grid-toobservation interpolation process, the necessity of smoothing in the observation-to-grid interpolation process, along with accuracy. The simulations provided confirmation of the above and yielded information concerning the sensitivity of statistical interpolation schemes to inexact parameter specification. The error estimates provided by statistical schemes were shown to be sensitive to inexact parameter specification.

13

STATE AND A STATE

# 5.0 Acknowlegdements

k

I would like to acknowledge useful discussions that I had with Dr. Edward Barker. In addition, his helpful comments on preliminary drafts of the the paper were responsible for many improvements in content and form.

#### References

- Barnes, S. L., 1973: Mesoscale objective map analysis using weighted time-series observations, NOAA Tech. Memo. ERL NSSL-62 (NTIS No. COM-73-10781)
- Bergman, Kenneth H., 1979: Multivariate analysis of temperatures and winds using optimum interpolation, Mon. Wea. Rev. 107, 1423-1444
- Foley, Thomas A., and Gregory M. Nielson, 1980: Multivariate interpolation to scattered data using delta iteration, in Approximation Theory III (E. W. Cheney, ed.), Academic Press, New York, pp. 419-424
- Franke, Richard, and William J. Gordon, 1983: The structure of optimum interpolation functions, Tech. Rep. No. NPS-53-0005, Naval Postgraduate School, Monterey, 23 pp. (NTIS No. AD-A126 958/8)
- Gandin, L. S., 1963: Objective analysis of meteorological fields [Translated from Russian by the Israeli Program for Scientific Translations, 1955], 242 pp.
- Koehler, Thomas Lee, 1979: A case study of height and temperature analysis derived from Nimbus-5 satellite soundings on a fine mesh model grid, Ph.D. Thesis, Dept. of Meteorology, University of Wisconsin, Madison, 1979, 185 pp.
- Séaman, R. S., 1983: Objective analysis accuracies of statistical interpolation and successive correction schemes, Aust. Met. Mag. 31, 225-240

- Wahba, Grace, 1982: Variational methods in simultaneous optimum interpolation and initialization, in The Interaction Between Objective Analysis and Initialization (Proceedings of the Fourteenth Stanstead Seminar held at Bishop's University, Lennoxville, Quebec, Canada), D. Williamson, ed., Publication in Met. 127, McGill University, Montreal, pp. 173-135
- Wahba, Grace, and James Wendelberger, 1980: Some new mathematical methods for variational objective analysis using splines and cross validation, Mon. Wea. Rev. 103, 1122-1143
- Wendelberger, James, 1982: The computation of Laplacian smoothing splines with examples, Dept. of Statistics, Tech. Rep. No. 548, University of Wisconsin, Madison, 55 pp.

[ əldel

Method	l: #Operations	2: #Bàsis	3: Other	Example # (1,2,3) 2.5 <sup>0</sup> grid, 35 obs 5 <sup>0</sup> grid, 57 obs
Opt Interp	(N <sup>3</sup> +5N <sup>2</sup> -N <sub>0</sub> )/5 +N <sub>0</sub> N <sub>0</sub> N <sub>0</sub>	no (no+1)/2+nongn&	N <sub>o</sub> sqrts	13278, 4878, 35 60501, 8174, 67
Local OI	Σ ((n <sup>3</sup> +5n <sup>2</sup> -n)/6+n)	Σ (n.(n+1)/2+n)	search, în sgrts	9555, 3155, 683 7333, 2400, 515
barnes'	2N <sub>0</sub> V <sub>6</sub> (N <sub>0</sub> +1)	<sup>2N</sup> o <sup>N</sup> B <sup>N</sup> A	grid-to-obs	8658, 8424, 972 11958,11792,1809
Local Barnes'	Σ (2n+2)	Σ 2n	search, grìd-to-obs	1500, 1355, 972 1208, 1032,1809
Thin Pl Spl	(N <sub>o</sub> +3) ((N <sub>o</sub> +3) <sup>2</sup> -1)/3 +(N <sub>o</sub> +3) <sup>2</sup> +(N <sub>o</sub> +2) N <sub>O</sub> N <sub>b</sub>	No (No+1)/2+N <sub>0</sub> N <sub>6</sub> N <sub>6</sub> N	-	25725, 4878, - 125231, 3174, -
Lapl Sm Spl	(N <sub>o</sub> +5) ( (N <sub>o</sub> +5) <sup>2</sup> -1) / ۲ + (N <sub>o</sub> +5) <sup>2</sup> + (N <sub>o</sub> +5) N <sub>O</sub> N <sub>b</sub>	<sup>м</sup> о (и <sub>о</sub> +1) / 2+и <sub>о</sub> и <sub>9</sub> и <sub>ф</sub>		31585, 4879, - 41590, 8174, -
Frnke/Grdn	$3N_{\Theta}N_{\Phi}(N_{O}+1)+2N_{O}(N_{O}+1)$	ф <sub>N</sub> eno <sub>ne</sub>	}	15551, 15288, - 27054, 25555, -
Pseudo- Barnes'	<sup>2N</sup> θ <sup>N</sup> ¢ (N <sub>0</sub> +1) +N <sub>0</sub> (N <sub>0</sub> +1)	2.νονθνά	:	9990, 9720, - 15524, 15221, -
Local P- Barnos'	Σ (n <sup>2</sup> +3n+2)	Σ (n <sup>2</sup> +2n)	search	5419, 5549, - 1175, 4281, -

21

s.

i.

Ĭ.

Table#	rg	ro	θ <sub>o</sub>	θ	grið	notes <sup>C</sup>
3	30	1Ø	101	Ø	13x9,2.5°	
4	30	10	100	13.775	13x9,2.5°	
5	20	10	100	Ø	13x9,2.5°	
5	30	5	100	Ø	13x9,2.5°	
7	30	1Ø	random <sup>a</sup>	random <sup>b</sup>	13x9,2.5 <sup>0</sup>	
8	30	13	random <sup>a</sup>	random <sup>b</sup>	13x9,2.5°	$(r_0)_i = 30$
9	20	10	random <sup>a</sup>	random <sup>b</sup>	13x9,2.5°	$(r_{g})_{i} = 23$
10	30	10	random <sup>a</sup>	random <sup>b</sup>	13x9,2.5°	$(r_{0})_{i} = 5$
11	3Ø	5	random <sup>a</sup>	random <sup>b</sup>	13x9,2.5°	$(r_0)_i = 10$
12	5	10	random <sup>a</sup>	random <sup>b</sup>	13x9,2.5°	
13	3Ø	Ø	random <sup>a</sup>	random <sup>b</sup>	13x9,2.5°	
14	3Ø	10	random <sup>a</sup>	random <sup>b</sup>	11x8,5 <sup>0</sup>	

Table 2

а

b

⊖ uniformly distributed in (-82.5°,112.5°) ∆⊖ uniformly distributed in (-15°,15°) The statistical interpolation routines were given incorrect variances, as indicated С

Entries: RMSE Analysis P = 500, Theta= 100, Celth = 0 Mean RMSE(StDev) rq = 3C, ro = 10Number of realizations = 100 13X9 grid of 2.5 degrees, 36 observation points PW Bicub Bsl Bicub Grid-to-cbs: PW linear Bicub Spl Obs-to-grid 6.09 6.64 6.09 6.09 Opt Interp 5.98(1.19) 5.97 (1.20) 5.98(1.19) (Cd = 10)6.53(1.18) 6.54 6.55 7.09 6.53 Local OI 6.99 (1.19) 6.42(1.19) 6.44 (1.19) 6.44(1.18) (Cd = 10)8,87 8.87 8.88 Barnes! 9.27 9.08 (1.03) 8.68 (1.82) 8.68 (1.82) 8.69(1.82) 2-Pass 7.95 7.96 7.96 Barnes! 8.42 7.79(1.56) 7.80 (1.56) 7.81 (1.56) (Lccal) 8.27 (1.57) 6.78 6.79 6.79 Stat Interp 7.28 6.67 (1.26) 6.68 (1.26) 7.23(1.22) 6.66(1.27) (Cd = 14)6.87 6.87 7.34 6.87 Stat Interp 6.76 (1.25) 6.76(1.25) 7.23(1.25) 6.75 (1.26) (Cd = 7)6.91 6.91 6.91 Stat Interp 7.37 7.26 (1.28) 6.79 (1.28) 6.79 (1.28) 6.79(1.28) (Dmpd Cos) Thin Pl Spl 7.12 6.59 6.60 6.60 6.47(1.32) (m = 2)7.00(1.30) 6.45 (1.33) 6.46(1.32) 10.25 10.25 10.54 10.25 Lapl Sm Spl 10.11(1.73) 10.10(1.73) 10.10(1.73) (m = 3)10.40(1.73) 11.76 11.76 Frnke/Grdn 12.02 11.75 11.45 (2.65) 11.45 (2.65) 11.45 (2.65) 11.72(2.65) (3 Fass) 8.88 8.87 8.87 PseudoBarnes<sup>1</sup> 9.28 9.10 (1.83) 8.68(1.82) 8.68(1.82) 8.69(1.82) (2 Fass) 7.70 7.71 7,72 PseudoBarnes<sup>®</sup> 8.20 8.06 (1.51) 7.55 (1.50) 7.57 (1.50) 7.57(1.50) (Local)

F = 500, Theta = 100, Delth = 13.775 Entries: RMSE Analysis rg = 3C, ro = 10Mean RMSE(StDev) Number of realizations = 100 1319 grid of 2.5 degrees, 36 Observation points PW Bicub Grid-to-obs: FW Linear Bicub Spl Bsl Bicub Obs-to-grid 6.19 6.20 6.21 Opt Interp 6.88 6.06(1.34) 6.05(1.34)6.06(1.34)(Cd = 10) $\epsilon.72(1.49)$ 6.76 7.40 6.77 Local OI 7.24 (1.51) 6.62(1.37) (Cd = 10)6.63(1.37) 9.73 9.29 Baires' 9.28 9.09(1.88) 9.53(1.96) 9.10(1.88) 2-Fass Barnes\* 8.29 7.79 7.77 7.67(1.36) 7.66 (1.36) (Lccal) 8.16(1.48) 7.08 7.09 Stat Interp 7.71 (Cd = 14)7.56 (1.50) 6.95(1.35) 6.96 (1.35) 7.54 6.96 6.96 Stat Interp (Cd = 7)7.39 (1.50) 6.82(1.36)6.83(1.36) Stat Interp 7.37 6.91 6.91 6.91 (Derd Ccs) 7.26 (1.28) 6.79 (1.28) 6.79 (1.28) 6.79 (1.28) 7.45 6.80 Thin Pl Spl 6.81 (m = 2)7.29 (1.51) 6.66 (1.37) 6.67 (1.37) Lapl Sm Spl (m = 3)Frnke/Grdn (3 Fass) PseudoBarnes' 9.75 9.28 9.55 (1.97) 9.09(1.88) (2 Fass) PseudoBarnes' (Lccal)

F = 500, Theta = 100, Celth = 0 Entries: RMSE Analysis rq = 20, rc = 10Mean RMSE(StDev) Number of realizations = 100 13X9 grid of 2.5 degrees, 36 Observation points Grid-to-cbs: FW Linear Bicub Spl PW Bicub Bel Bicub Obs-to-grid 5.75 5.76 6.23 Opt Interp 5.62(1.22) 6.10(1.28) 5.63 (1.22) (Cd = 10)6.54 6.10 6.10 Local OI 5.97 (1.22) 5.98 (1.22) 6.41(1.28) (Cd = 10)7.18 6.85 6.86 Barnes' 6.71(1.47) 2-Pass 7.03 (1.47) 6.71 (1.41) Barnes! 7.19 6.77 6.76 7.08(1.22) 6.68(1.11) 6.67 (1.10) (Lccal) Stat Interp 6.70 6.30 6.31 6.58(1.27) 6.19(1.19) 6.19(1.19) (Cd = 14)6.71 6.24 6.25 Stat Interr 6.12(1.25) 6.58 (1.30) 6.12(1.24) (cd = 7)Stat Interp 6.78 6.31 6.32 6.18(1.28) (Dmpd Cos) 6.65 (1.33) 6.18(1.28) Thir Pl Spl 6.66 6.20 6.20 6.06 (1.29) (m = 2)6.52(1.34) 6.07 (1.29) 11.05 Larl Sm Spl 10.71 10.71 (m<sup>-</sup>= 3) 10.84 (2.16) 10.50 (2.12) 10.50 (2.12) Frnke/Grdn 8.72 8.56 8.56 8.54 (1.78) 8.39 (1.70) (3 Pass) 8.39 (1.70) PseudoBarnes' 7.19 6.85 6.86 (? Fass) 7.03(1.47) 6.85(1.41) 6.71 (1.41) PseudoBarnes' 6.78 6.37 6.38 6.66(1.28) 6.25(1.24) 6.26(1.23)(Local)

TABLE 5

F = 500, Theta = 100, Celth = 0 Entries: RMSE Analysis Mean RMSE(StDev) 10 = 30, 10 = 5Number of realizations = 100 13X9 grid of 2.5 degrees, 36 Observation points Grid-to-cbs: FW Linear Bicut Spl Bel Bicub PW Bicub Obs-to-grid 3.77 3.76 4.57 Opt Interp 3.70 (0.69) 3.70 (0.69) 4.50 (0.83) (Cd = 10)4.27 5.03 4.26 Local OI 4.20 (0.77) 4.19 (0.77) 4.95 (0.91) (Cd = 10)8.49 8.48 8.88 Barnes' 8.29 (1.76) 8.30 (1.77) 8.69 (1.85) 2-Fass 5.89 5 - 90 6.39 Barnes! 5.78 (1.11) 5.79(1.11) 6.30 (1.10) (Lccal) 4.57 4.57 5.26 Stat Interp 4.50 (0.83) 4.49(0.83) 5.17(0.92) (Cd = 14)4.28 4.28 5.02 Stat Interr 4.21 (0.74) 4.21(0.74) 4.95 (0.82) (Cd = 7)4.22 4.22 4.96 Stat Interp 4.16(0.75) 4.16 (0.75) 4.89 (0.83) (Dmpd Cos) 4.16 4.15 4.92 Thin Pl Spl 4.09 (0.72) 4.10(0.72) 4.85 (0.84) (m = 2)5.43 5.43 Lagl Sm Spl 6.08 5.33(1.06) 5.33(1.06) 5.98 (1.05) (m = 3)11.61 11.61 11.82 Frnke/Grdn 11.34 (2.50) 11.34 (2.51) 11.55 (2.55) (3 Pass) 8,49 8.48 Pseudo Barnes' 8.89 8.30 (1.77) 8.29(1.76) 8.70 (1.85) (2 Pass) 6.76 6.75 PseudoBarnes 7.30 6.62(1.41)6.60(1.40) 7.15(1.47) (Local)

TABLE 6

P = 500, Theta = RANDCH, Delth = RANDOM, Entries: RMSE Analysis rg = 30, ro = 10Mean RMSE(StDev) Number cf realizations = 100 13X9 grid of 2.5 degrees, 36 Observation points Grid-to-cbs: FW Linear Bicub Spl PW Bicub Bsl Bicub Obs-to-grid 7.00 6.40 Opt Interp 6.85(1.44) 6.27 (1.30) (Cd = 1C)7.48 6.90 Local OI 7.33(1.49) 6.90 (1.36) (Cd = 10)9.92 Barnes\* 9.52 2-Fass 9.70(2.07) 9.33 (1.89) Barnes<sup>\*</sup> 8.34 7.89 8.21(1.46) 7.76 (1.39) (Lccal) 7.41 Stat Interp 7.92 7.77 (1.56) 7.27 (1.43) (Cd = 14)Stat Interp 7.56 7.01 7.41(1.51) (Cd = 7)6.88(1.36) Stat Interp 7.58 7.04 (Dard Ccs) 7.43(1.53) 6.90 (1.38) 7.63 7.06 Thin Pl Spl 7.47(1.58) 6.92(1.42) (n = 2)Lapl Sm Spl (m = 3)Frnke/Grdn (3 Fass) PseudoBarnes\* (2 Fass) PseudoBaines\* (Lccal)

alalalalala alalalalalalaralaralarin ala

F = 500, Theta = RANDCM, Delth = RANDOM, Entries: RMSE Analysis rg = 30, rc = 10, rg(lie) = 20Mean RMSE(StDev) Number of realizations = 100 13X9 grid of 2.5 degrees, 36 Observation points Grid-to-cbs: FW Linear Bicub Spl PW Bicub Bel Bicub Obs-to-grid 6.47 7.14 Opt Interp (Cd = 10)6.97 (1.52) 6.32(1.35) Local OI 7.56 6.92 (Cd = 10)7.39(1.62) 6.77 (1.41) 9.64 Barnes' 9.21 2-Fass 9.42(2.04) 9.01(1.90) Barnes' 8.30 7.73 7.04(1.43) 7.62(1.33) (Local) Stat Interp (Cd = 14)Stat Interp (Cd = 7)Stat Interp (Dard Ccs) Thin Pl Spl 7.19 6.52 (m = 2)7.04(1.45) 6.39(1.26) Lapl Sm Spl (m = 3)Frnke/Grdn (3 Fass) PseudoBarnes\* (2 Fass) PseudoBaines' (Lccal)

#### TABLE 8

and the second secon

F = 500, Theta = RANDCM, Delth = RANDOM, Entries: RMSE Analysis rg = 2C, rc = 10, rg(lie) = 30Mean RMSE(StDev) Number of realizations = 100 13X9 grid of 2.5 degrees, 36 Observation points PW Bicub Bel Bicub Grid-to-cbs: FW Linear Bicub Spl Obs-to-grid 6.32 5.72 Opt Interp 6.17(1.36) 5.60 (1.21) (Cd = 10)Local OI 6.60 6.05 6.47(1.33) 5.94 (1.16) (Cd = 10)6.69 Barnes\* 7.10 6.55 (1.39) 2-Fass 6.95(1.45) Baines' 7.09 6.54 6.98 (1.22) 6.45 (1.08) (Lccal) Stat Interp (Cd = 14)Stat Interp (Cd = 7)Stat Interp (Drrd Ccs) Thin Pl Spl 6.31 5.75 6.17(1.30) 5.63 (1.16) (m = 2) Lapl Sm Spl (<u>m</u> = 3) Frnke/Grdn (3 Fass) PseudoBaines! (2 Fass) PseudoBarnes! (Lccal)

TABLE 9

F = 500, Theta = RANDCH, Delth = RANDOM, Entries: RMSE Analysis rg = 3C, rc = 10, ro(lie) = 5Mean RMSE(StDev) Number of realizations = 100 13X9 grid of 2.5 degrees, 36 Observation points Grid-to-cbs: FW Linear Bicub Spl PW Bicub Bsl Bicub Obs-to-grid 6.88 7.51 Opt Interp 7.37(1.43) 6.74 (1.38) (cd = 10)7.96 7.37 Local OI (Cd = 10)7.26 (1.27) 7.85(1.36) Barnes' 9.81 9.28 2-Fass 9.60(2.01)9.06 (1.99) 8.47 7.91 Barnes' 7.80 (1.31) 8.36 (1.32) (Local) Stat Interp (Cd = 14)Stat Interp (Cd = 7)Stat Interp (Derd Ccs) 6.94 Thin Pl Spl 7.65 7.51(1.43) 6.80 (1.39) (m = 2) Lapl Sm Spl (n = 3)Frnke/Grdn (3 Fass) PseudoBaines\* (2 Fass) PseudoBarnes' (Lccal)

F = 500, Theta = RANDCM, Delth = RANDON, Entries: RMSE Analysis rg = 30, ro = 5, ro(lie) = 10Mean RMSE(StDev) Number of realizations = 100 13X9 grid of 2.5 degrees, 36 Observation points Grid-to-cbs: PW Linear Bicub Spl PW Bicub Bsl Bicub Obs-to-grid Opt Interp 5.18 4.19 (Cd = 10)5.10(0.91) 4.13(0.71) Local OI 5.66 4.76 5.56(1.04) 4.69(0.81) (Cd = 10)9.09 Barnes' 8.62 8.47 (1.61) 8.93(1.69) 2-Fass Barnes! 6.71 6.06 6.63 (1.07) (Lccal) 5.97 (1.04) Stat Interp (Cd = 14)Stat Interp (Cd = 7)Stat Interp (Dard Ccs) 6.13 Thin Pl Spl 5.34 (m = 2)6.02(1.15) 5.25 (0.95) Lapl Sm Spl (m = 3)Frnke/Grdn (3 Fass) PseudoBaines\* (2 Fass) PseudoBarnes' (Lccal)

P = 50C, Theta = RANDCH, Delth = RANDON, Entries: RMSE Analysis rg = 5, rc = 10Mean RMSE(StDev) Number cf realizations = 100 13X9 grid of 2.5 degrees, 36 Observation points Grid-to-cbs: FW Linear Bicub Spl PW Bicub Bsl Bicub Obs-to-grid Opt Interp 3.58 3.32 (Cd = 10)3.44 (1.01) 3.18(0.94) 3.77 3.53 Local OI (Cd = 10)3.63(1.02) 3.40 (0.9769 3.98 4.44 Barnes\* 4.32(1.01) 3.87 (0.90) 2-Fass Barnes' 6.26 5.73 6.15(1.16) 5.64 (1.00) (Local) Stat Interp 3.61 3.41 (Cd = 14)3.46 (1.03) 3.27 (0.96) Stat Int∈rp 3.65 3.39 3.25 (0.97) (Cd = 7)3.50(1.03)3.78 Stat Interp 3.51 3.62(1.07)3.36 (1.02 (Derd Ccs) Thin Pl Spl 4.00 3.85 (n = 2)3.84(1.13)3.70(1.07) Lapl Sm Spl (m = 3)Frnke/Grðn (3 Fass) PseudoBarnes\* (2 Fass) PseudoBarnes' (Lccal)

F = 500, Theta = RANDCH, Delth = RANDOM, Entries: RMSE Analysis ig = 30, ic = 0Mean RMSE(StDev) Number of realizations = 100 13X9 grid of 2.5 degrees, 36 Observation points Grid-to-cts: FW Linear Bicut Spl PW Bicub Bsl Bicub Obs-to-grid 0.74 6.06 Opt Interp 0.69(0.27)(Cd = 10)5.84 (1.62) Local OI 3.86 2.40 2.2.30(0.69 (Cd = 10)3.79 (0.76) Barnes' 9.38 8.96 8.74 (1.98) 2-Pass 9.16 (2.05) Barnes! 6.27 5.63 5.49(1.23) (Lccal) 6.15(1.21) Stat Interp 7.62 1.23 (Cd = 14)7.10(2.74) 1.16(0.40)Stat Interp 3.70 1.29 (Cd = 7)3.64 (0.68) 1.20 (0.48) Stat Interp 1.03 4.48 4.37 (0.96) 0.97 (0.36 (Dmpd Cos) Thir Pl Srl 3.74 2.18 3.66 (0.78) (m = 2)2.02(0.83) Larl Sm Spl (m = 3) Frrke/Grdn (3 Pass) Pseudo Earnes' (2 Fass) Pseudo Barnes' (Lccal)

F = 500, Theta = 100, Celth = 0 RMSE Analysis Entries: rq = 30, ro = 10Mean RMSE(StDev) Number of realizations = 100 11 BY 8 grid of 5 degrees, 67 Observation points Grid-to-obs: FW Linear Bicut Spl PW Bicub Bsl Bicub Obs-to-grid 7.62 7.93 8.11 Opt Interp 12.84 7.53(1.19) 12.74 (1.62) 7.84 (1.20) 8.02(1.19) (Cd = 10)8.44 8.74 8.92 Local OI 13.33 (Cd = 10)13.22 (1.73) 8.33(1.33) 8.63(1.35) 8.82(1.32) 10.95 Barnes\* 14.33 10.62 10.82 14.21 (1.85) 10.49 (1.70) 10.68 (1.70) 10.81 (1.71) 2-Pass 9.40 Barnes' 14.00 8.82 9.12 13.91(1.55) 8.71(1.36) 9.79 (1.35) 9.31 (1.33) (Lccal) 13.75 8.80 9.02 9.20 Stat Interp 13.25(1.57) 8.70(1.33) 8.92(1.31) 9.10(1.32) (Cd = 14)8.31 8.79 Stat Interp 13.44 8.62 8.70(1.19) 13.35 (1.63) 8.23(1.18) 8.53 (1.21)  $(\dot{C}d = 7)$ 8.47 8.78 8.95 Stat Interp 13.57 13.47 (1.65) 8.38 (1.25) 8.68(1.27) 8.86 (1.25) (Dmpd Cos) 8.08 8.47 Thin Pl Spl 13.17 8.36 (m = 2)13.07 (1.59) 7.99(1.17) 8.27 (1.19) 8.39 (1.19) Lapl Sm Spl 17.12 11.87 12.08 12.16 17.01(1.86) 11.76(1.60) 11.97 (1.57) 12.05 (1.58) (m = 3)Frnke/Grdn 17.29 15.23 15.31 15.36 17.14(2.29) 15.04 (2.35) 15.13(2.34) 15.17 (2.37) (3 Fass) PseudoBarnes'14.27 10.62 10.82 10.95 (2 Fass) 14.14 (1.87) 10.49 (1.70) 10.69(1.70) 10.82(1.71) 9.73 10.00 10.17 PseudoBarnes 13.89 13.77 (1.82) 9.60 (1.58) 9.87 (1.58) 10.05(1.57) (Lccal)

	IG: IO:	5 10	20 10	30 10	30 5	30 0
method						
Opt Interr		3.30	5.62	6.29	3.78	0.65
(cd = 10)		(3.32)	(5.76)	(6.40)	(3.77)	(0.74)
Local OI		3.55	5.96	6.80	4.39	2.21
(cd = 10)		(3.53)	(6.10)	(6.90)	(4.27)	(2.40)
Stat Interp		2.90	4.60	5.07	2.95	0.11
(cd = 14)		(3.41)	(6.31)	(7.41)	(4.57)	(1.23)
Stat Interp		3.76	7.14	8.26	5.46	2.90
(cd = 7)		(3.39)	(6.25)	(7.01)	(4.28)	(1.29)
Stat Interp		3.82	7.00	7.94	5.03	2.06
(Dmpd Cos)		(3.51)	(6.32)	(7.04)	(4.22)	(1.03)

Estimated and (empirical) RMS errors for statistical methods







.







Figure 4



Figure 5







T7: RG = 30, RO = 10, 13X9 2.5 DEGREE GRID



and and a second second second a second

1933-3-3- (D.22)



RO - 10, 13X9 2.5 DEGREE GRID



Figure 9



Figure 10



Figure 11

# PERFORMANCE DEGRADATION, RG = 30, RO = 5



Figure 12

.



Figure 13

#### DISTRIBUTION LIST

CHIEF OF NAVAL RESEARCH (2)

LIBRARY SERVICES, CODE 784

ASST. FOR ENV. SCIENCES ASST. SEC. OF THE NAVY (R&D) ROOM 5E731, THE PENTAGON WASHINGTON, DC 20350

COMMANDING OFFICER NORDA NSTL, MS 39529

NAVAL POSTGRADUATE SCHOOL METEOROLOGY DEPT. MONTEREY, CA 93943 ARLINGTON, VA 22217 COMMANDING OFFICER FLENUMOCEANCEN

MONTEREY, CA 93940

BALLSTON TOWER #1

800 QUINCY ST.

NAVAL POSTGRADUATE SCHOOL OCEANOGRAPHY DEPT. MONTEREY, CA 93943

COMMANDER (2) NAVAIRSYSCOM ATTN: LIBRARY (AIR-7226) WASHINGTON, DC 20361

USAFETAC/TS SCOTT AFB, IL 62225 SUPERINTENDENT ATTN: USAFA (DEG)

USAF ACADEMY, CO 80840

NAVAIRSYSCOM (AIR-330)

WASHINGTON, DC 20361

COMMANDER

AFGL/LY HANSCOM AFB, MA 01731 AFGL/OPI HANSCOM AFB, MA 01731

COMMANDER & DIRECTOR ATTN: DELAS-D U.S. ARMY ATMOS. SCI. LAB WHITE SAND MISSILE RANGE WHITE SANDS, NM 88002

DIRECTOR NATIONAL METEORO. CENTER NWS, NOAA WWB W32, RM 204 WASHINGTON, DC 20233 DIRECTOR (2) DEFENSE TECH. INFORMATION CENTER, CAMERON STATION ALEXANDRIA, VA 22314

ACQUISITIONS SECT. IRDB-D823 LIBRARY & INFO. SERV., NOAA 6009 EXECUTIVE BLVD. ROCKVILLE, MD 20852 OFFICE OF NAVAL RESEARCH CODE 422AT ARLINGTON, VA 22217

SUPERINTENDENT LIBRARY REPORTS U.S. NAVAL ACADEMY ANNAPOLIS, MD 21402

LIBRARY (2) NAVAL POSTGRADUATE SCHOOL MONTEREY, CA 93943

COMMANDER NAVOCEANSYSCEN DR. J. RICHTER, CODE 532 SAN DIEGO, CA 92152

AFGWC/DAPL OFFUTT AFB, NE 68113

OFFICER IN CHARGE SERVICE SCHOOL COMMAND DET. CHANUTE/STOP 62 CHANUTE AFB, IL 61868

NOAA-NESDIS LIAISON ATTN: CODE SC2 NASA-JOHNSON SPACE CENTER HOUSTON, TX 77058

DIRECTOR OFFICE OF PROGRAMS RX3 NOAA RESEARCH LAB BOULDER, CO 80302

NATIONAL WEATHER SLAVE F WORLD WEATHER BLDG., RM 307 5200 AUTH ROAD CAMP SPRINGS, MD 20023

DIRECTOR TECHNIQUES DEVELOPMENT LAB GRAMAX BLDG. 8060 13TH ST. SILVER SPRING, MD 20910

COLORADO STATE UNIVERSITY ATMOSPHERIC SCIENCES DEPT. ATTN: DR. WILLIAM GRAY FORT COLLINS, CO 80523

CHAIRMAN, METEOROLOGY DEPT. CALIFORNIA STATE UNIVERSITY SAN JOSE, CA 95192

UNIVERSITY OF WASHINGTON ATMOSPHERIC SCIENCES DEPT. SEATTLE, WA 98195

UNIVERSITY OF HAWAII METEOROLOGY DEPT. 2525 CORREA ROAD HONOLULU, HI 96822

UNIVERSITY OF MARYLAND METEOROLOGY DEPT. COLLEGE PARK, MD 20742

CHAIRMAN, METEOROLOGY DEPT. UNIVERSITY OF UTAH SALT LAKE CITY, UT 84112 DIPECTOR NATIONAL SEVERE STORMS LAB 1313 HALLEY CIRCLE NORMAN, OK 73069

LABORATORY FOR ATMOS. SCI. NASA GODDARD SPACE FLIGHT CEN. GREENBELT, MD 20771

ATMOSPHERIC SCIENCES DEPT. UCLA 405 HILGARD AVE. LOS ANGELES, CA 90024

COLORADO STATE UNIVERSITY ATMOSPHERIC SCIENCES DEPT. ATTN: LIBRARIAN FT. COLLINS, CO 80523

CHAIRMAN, METEOROLOGY DEPT. PENNSYLVANIA STATE UNIV. 503 DEIKE BLDG. UNIVERSITY PARK, PA 16802

DIRECTOR COASTAL STUDIES INSTITUTE LOUISIANA STATE UNIVERSITY ATTN: O. HUH BATON ROUGE, LA 70803

CHAIRMAN ATMOS. SCIENCES DEPT. UNIVERSITY OF VIRGINIA CHARLOTTESVILLE, VA 22903

ATMOSPHERIC SCIENCES CENTER DESERT RESEARCH INSTITUTE P.O. BOX 60220 RENO, NV 89506 DIRECTOR GEOPHYS. FLUID DYNAMICS LAB NOAA, PRINCETON UNIVERSITY P.O. BOX 308 PRINCETON, NJ 08540

EXECUTIVE SECRETARY, CAO SUBCOMMITTEE ON ATMOS. SCI. NATIONAL SCIENCE FOUNDATION RM. 510, 1800 G. STREET, NW WASHINGTON, DC 20550

CHAIRMAN, METEOROLOGY DEPT. UNIVERSITY OF OKLAHOMA NORMAN, OK 73069

NATIONAL CENTER FOR ATMOS. RSCH., LIBRARY ACQUISITIONS P.O. BOX 3000 BOULDER, CO 80302

FLORIDA STATE UNIVERSITY ENVIRONMENTAL SCIENCES DEPT. TALLAHASSEE, FL 32306

ATMOSPHERIC SCIENCES DEPT. OREGON STATE UNIVERSITY CORVALLIS, OR 97331

CHAIRMAN METEOROLOGY DEPT. MASSACHUSETTS INSTITUTE OF TECHNOLOGY CAMBRIDGE, MA 02139

ATMOSPHERIC SCI. RSCH. CENTER NEW YORK STATE UNIV. 1400 WASHINGTON AVE. ALBANY, NY 12222

SYSTEMS & APPLIED SCI. CORP. ATTN: LIBRARY, SUITE 500 6811 KENILWORTH AVE. RIVERDALE, MD 20840

METEOROLOGY INTL., INC. P.O. BOX 22920 CARMEL, CA 93922

OCEAN DATA SYSTEMS, INC. 2460 GARDEN ROAD MONTEREY, CA 93940

DEPT. OF MATHEMATICS NAVAL POSTGRADUATE SCHOOL MONTEREY, CA 93943

DR. RICHARD LAU OFFICE OF NAVAL RESEARCH 1030 EAST GREEN ST. PASADENA, CA 91106

LIBRARY, FLEET NUMERICAL(2) OCEANOGRAPHY CENTER MONTEREY, CA 93943

PROFESSOR W.J. GORDON CENTER FOR SCIENTIFIC COMPUTATION & INTERACTIVE GRAPHICS DREXEL UNIVERSITY PHILADELPHIA, PA 19104

PROFESSOR DONALD GAVER DEPARTMENT OF OPERATIONS RESEARCH NAVAL POSTGRADUATE SCHOOL MONTEREY, CA 93943

CENTER FOR ENV. & MAN. INC. RESEARCH LIBRARY 275 WINDSOR ST. HARTFORD, CT 06120

CONTROL DATA CORP. METEOROLOGY DEPT. RSCH. DIV. 2800 E. OLD SHAKOPEE RD. BOX 1249 MINNEAPOLIS, MN 55440

EUROPEAN CENTRE FOR MEDIUM RANGE WEATHER FORECASTS SHINFIELD PARK, READING BERKSHIRE RG29AX, ENGLAND

PROFESSOR G. LATTA, CODE 53LZ CHRMN, DEPARTMENT OF MATH NAVAL POSTGRADUATE SCHOOL MONTEREY, CA 93943

PROFESSOR R.E. BARNHILL DEPARTMENT OF MATHEMATICS UNIVERSITY OF UTAH SALT LAKE CITY, UT 84112

(10) DR. EDWARD BARKED RESEARCH FACILITY MONTEREY, CA 93943

MS. LINDA THIEL DEPARTMENT OF MATHEMATICAL SCIENCES DREXEL UNIVERSITY PHILADELPHIA, PA 19104

PROFESSOR P.A. JACOBS DEPARTMENT OF OPERATIONS RESEARCH NAVAL POSTGRADUATE SCHOOL MONTEREY, CA 93943

METEOROLOGY RESEARCH, INC. 464 W. WOODBURY RD. ALTADENA, CA 91001

SCIENCE APPLICATIONS. INC. 205 MONTECITO AVENUE MONTEREY, CA 93940

(2)DEAN OF RESEARCH CODE 012 NAVAL POSTGRADUATE SCHOOL MONTEREY, CA 93943

PROFESSOR R. FRANKE, CODE 53FE(15) DEPARTMENT OF MATHEMATICS NAVAL POSTGRADUATE SCHOOL MONTEREY, CA 93943

PROFESSOR G.M. NIELSON DEPARTMENT OF MATHEMATICS ARIZONA STATE UNIVERSITY TEMPE, ARIZONA 85281

DR. THOMAS ROSMOND NAVAL ENVIRONMEN.... PREDICTION NAVAL ENVIRONMENTAL PREDICTION **RESEARCH FACILITY** MONTEREY, CA 93943

> PROFESSOR GR. CE WAHBA DEPARTMENT OF STATISTICS UNIVERSITY OF WISCONSIN MADISON, WI 53705

PROFESSOR H. JEAN THIEBAUX DEPARTMENT OF MATHEMATICS. STATISTICS & COMPUTING SCIENCE DALHAUSIE UNIVERSITY HALIFAX, NOVA SCOTIA, CANADA 133H 4H8

DR. R.S. SEAMAN AUSTRALIAN NUMERICAL METEOROLOGY RESEARCH CENTRE P.O. BOX 5089AA MELBOURNE, VICTORIA, AUSTRALIA, 3001

PROFESSOR PETER ALFELD DEPARTMENT OF MATHEMATICS UNIVERSITY OF UTAH SALT LAKE CITY, UT 84112 PROFESSOR MARK E. HAWLEY DEPARTMENT OF ENVIRONMENTAL SCIENCES UNIVERSITY OF VIRGINIA CHARLOTTESVILLE, VA 22903

DR. LAWRENCE BREAKER DEPARTMENT OF OCEANOGRAPHY NAVAL POSTGRADUATE SCHOOL MONTEREY, CA 93943

