

MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A148 995

4

OFFICE OF NAVAL RESEARCH

Contract No. N00014-83-K-0611

Task No. NR 053-842

TECHNICAL REPORT NO. UK/DC/TR-4

Structures of Three Pyrazaboles: 1,3,5,7-Tetramethylpyrazabole;
4,4-Bis(pyrazol-1'-yl)pyrazabole; 4,4,8,8-Tetrakis(pyrazol-1'-yl)pyrazabole

by

C. P. Brock, K. Niedenzu, E. Hanecker, and H. Noeth

Prepared for publication in

ACTA CRYSTALLOGRAPHICA

University of Kentucky
Department of Chemistry
Lexington, KY 40506

November 1984

DTIC
ELECTE
JAN 7 1985
B

Reproduction in whole or in part is
permitted for any purpose of the United States Government

This document has been approved for public
release and sale; its distribution is unlimited

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER UK/DC/TR-4	2. GOVT ACCESSION NO. AD-A148 495	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) STRUCTURES OF THREE PYRAZABOLES: 1,3,5,7-TETRA-METHYLPYRAZABOLE; 4,4-BIS(PYRAZOL-1'-YL)PYRAZABOLE; AND 4,4,8,8-TETRAKIS(PYRAZOL-1'-YL)PYRAZABOLE		5. TYPE OF REPORT & PERIOD COVERED Interim Technical Report, 1984
7. AUTHOR(s) C. P. Brock, K. Niedenzu, E. Hanecker,* and H. Noeth* (*University of Muenchen, West Germany)		6. PERFORMING ORG. REPORT NUMBER 4
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Kentucky, Department of Chemistry Lexington, KY 40506-0055		8. CONTRACT OR GRANT NUMBER(s) N00014-83-K-0611
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-842
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE November 1984
		13. NUMBER OF PAGES 21
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Prepared for publication in ACT CRYSTALLOGRAPHICA		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) pyrazaboles x-ray crystal structures		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The structures of three pyrazaboles have been determined by single crystal x-ray diffraction. 1,3,5,7-Tetramethylpyrazabole: $M_r = 215.9$, monoclinic, $P2_1/c$, $a = 7.552(1)$, $b = 12.047(4)$, $c = 13.814(6)$ Å, $\beta = 91.27(2)^\circ$, $v = 1256.5(7)$ Å ³ at $T = 295$ K, $Z = 4$, $D_x = 1.141$ g cm ⁻³ , MoK α , graphite monochromator, $\lambda = 0.71069$ Å, u		

= 0.65 cm^{-1} , $F(000) = 464$, final R of 0.055 for 199 variables and 1640 reflections. 4,4-Bis(pyrazol-1'-yl)pyrazabole: $M_r = 291.9$, triclinic, $P\bar{1}$, $a = 8.721(2)$, $b = 9.351(2)$, $c = 9.407(2) \text{ \AA}$, $\alpha = 93.75(2)$, $\beta = 100.75(2)$, $\gamma = 92.73(2)^\circ$, $V = 750.7(2) \text{ \AA}^3$ at $T = 295 \text{ K}$, $Z = 2$, $D_x = 1.291 \text{ g cm}^{-3}$, $\text{MoK}\alpha$, graphite monochromator, $\lambda = 0.71069 \text{ \AA}$, $\mu = 0.79 \text{ cm}^{-1}$, $F(000) = 304$, final R of 0.035 for 255 variables and 1957 reflections.

4,4,8,8-Tetrakis(pyrazol-1'-yl)pyrazabole: $M_r = 424.0$, monoclinic, $P2_1/n$, $a = 8.849(2)$, $b = 21.765(2)$, $c = 11.672(1) \text{ \AA}$, $\beta = 109.51(1)^\circ$, $V = 2118.9(5) \text{ \AA}^3$ at 296 K , $Z = 4$, $D_x = 1.329 \text{ g cm}^{-3}$, $\text{MoK}\alpha$, graphite monochromator, $\lambda = 0.71069 \text{ \AA}$, $\mu = 0.82 \text{ cm}^{-1}$, $F(000) = 880$, final R of 0.036 for 362 variables and 1880 reflections. In all three structures the pyrazabole ring adopts a pronounced boat conformation. A comparison with the nine other known pyrazabole structures, however, shows that the central B_2N_2 ring is quite flexible, and that its observed conformation is sensitive to substituent and packing effects.

✓

A-1



Structures of Three Pyrazaboles: 1,3,5,7-Tetramethylpyrazabole
($C_{10}H_{18}B_2N_4$); 4,4-Bis(pyrazol-1'-yl)pyrazabole ($C_{12}H_{14}B_2N_8$); and
4,4,8,8-Tetrakis(pyrazol-1'-yl)pyrazabole ($C_{18}H_{18}B_2N_{12}$).

C. P. Brock and K. Niedenzu

Department of Chemistry, University of Kentucky, Lexington, KY 40506-0055, USA

and

E. Hanecker and H. Nöth*

Institute for Inorganic Chemistry, University of Munich, D-8000 Munich 2,
Federal Republic of Germany

(Received _____; Accepted _____)


Abstract. 1,3,5,7-Tetramethylpyrazabole (hereafter, ME4PZB): $M_r = 215.9$, monoclinic, $P2_1/c$, $a = 7.552(1)$, $b = 12.047(4)$, $c = 13.814(6)$ Å, $\beta = 91.27(2)^\circ$, $V = 1256.5(7)$ Å³ at $T = 295$ K, $Z = 4$, $D_x = 1.141$ g cm⁻³, MoK α , graphite monochromator, $\lambda = 0.71069$ Å, $\mu = 0.65$ cm⁻¹, $F(000) = 464$, final R of 0.055 for 199 variables and 1424 reflections. 4,4-Bis(pyrazol-1'-yl)pyrazabole (hereafter, BISPZB): $M_r = 291.9$, triclinic, $P\bar{1}$, $a = 8.721(2)$, $b = 9.351(2)$, $c = 9.407(2)$ Å, $\alpha = 93.75(2)$, $\beta = 100.75(2)$, $\gamma = 92.73(2)^\circ$, $V = 750.7(2)$ Å³ at $T = 295$ K, $Z = 2$, $D_x = 1.291$ g cm⁻³, MoK α , graphite monochromator, $\lambda = 0.71069$ Å, $\mu = 0.79$ cm⁻¹, $F(000) = 304$, final R of 0.035 for 255 variables and 1957 reflections. 4,4,8,8-Tetrakis(pyrazol-1'-yl)pyrazabole (hereafter, TETPZB): $M_r = 424.0$, monoclinic, $P2_1/n$, $a = 8.849(2)$,

*Contribution No. 151 to the series "Chemistry of Boron"

$\underline{b} = 21.765(2)$, $\underline{c} = 11.672(1)$ Å, $\underline{\beta} = 109.51(1)^\circ$, $\underline{V} = 2118.9(5)$ Å³ at 296 K, $Z = 4$, $D_x = 1.329$ g cm⁻³, MoK α , graphite monochromator, $\lambda = 0.71069$ Å, $\underline{\mu} = 0.82$ cm⁻¹, $\underline{F}(000) = 880$, final \underline{R} of 0.036 for 362 variables and 1880 reflections.

In all three structures the pyrazabole adopts a pronounced boat conformation. A comparison with the nine other known pyrazabole structures, however, shows that the central B₂N₄ ring is quite flexible, and that its observed conformation is sensitive to substituent and packing effects.

Introduction. The pyrazaboles, which have the general structure and numbering scheme shown below, are a class of heterocycles containing four-coordinate boron and exhibiting remarkable chemical and thermal stability. Recent



efforts by one of us to exploit this stability (Clarke, Niedenzu, Niedenzu & Trofimenko, 1984) have focussed on synthesizing polymeric pyrazaboles containing boron atoms as spiro centers. In such a polymer the B_2N_4 ring must be approximately planar; however, if an analogy is made with 9,10-dihydroanthracene, the pyrazabole skeleton might just as well be expected to adopt a boat conformation (for discussion of the potential energy surface of 9,10-dihydroanthracene see Rabideau, 1978; Lipkowitz, Rabideau, Raber, Hardee, Schleyer, Kos & Kahn, 1982). The three structures reported here were undertaken in order to increase the information available about the pyrazabole ring system and its conformational flexibility.

Experimental. The structures of ME4FZB and BISPZB were determined at the University of Munich using a Syntex P3 diffractometer and the SHELXNTL program

package. Crystals of ME4PZB (Trofimenko, 1967a) grown from CH_2Cl_2 as plates having faces of the forms {100}, {010}, and {001}. (Needles are obtained from toluene). Crystal 0.3 x 0.2 x 0.5 mm, ω -2 θ scans, $2\theta \leq 45^\circ$, 3785 reflections (2 quadrants), after averaging 1424 with $F^2 > 3\sigma(F^2)$ treated as observed, 2 standard reflections, no absorption correction, decomposition (corrected) less than 2% overall, 21 reflections for measuring lattice parameters; $-8 \leq h \leq 8$, $k \leq 12$, $-14 \leq l \leq 14$. All C, B, and N atoms refined anisotropically; H atoms refined with fixed isotropic U 's of 0.070 \AA^2 . There was some difficulty in refining the methyl H atoms. Final values of agreement indices [refinement on F , $w^{-1} = \sigma^2(F_o) + (0.04 F_o)^2$] are $R = 0.055$, $wR = 0.055$ for 199 variables; error in an observation of unit weight is 1.94; most important features in a final difference-Fourier map have heights +0.17 and -0.19 $e^{-\sigma^{-3}}$. Positional parameters given in Table 1 and some bond lengths and angles in Tables 2 and 3; the molecular structure is illustrated in Figure 1. Crystals of BISPZB (Niedenzu & Niedenzu, 1984) grown from $\text{CHCl}_3/\text{C}_6\text{H}_{12}$ as prisms. Crystal 0.2 x 0.3 x 0.25 mm, ω -2 θ scans, $2\theta \leq 45^\circ$, 4226 reflections (full sphere), after averaging 1957 with $F^2 > 3\sigma(F^2)$ treated as observed, 2 standard reflections, no absorption correction, decomposition (corrected) less than 1.8% overall, 14 reflections for measuring lattice parameters, $-9 \leq h \leq 9$, $-10 \leq k \leq 10$, $l \leq 10$. All C, B, and N atoms refined anisotropically; H atoms refined isotropically. Final values of agreement indices [refinement on F , $w^{-1} = \sigma^2(F_o) + (0.02 F_o)^2$] are $R = 0.035$, $wR = 0.037$ for 255 variables; error in an observation of unit weight is 1.89; most important features in a final difference-Fourier map have heights +0.09 and -0.21 $e^{-\sigma^{-3}}$. Positional parameters given in Table 4 and some bond lengths and angles in Tables 2 and 3; the molecular structure is illustrated in Figure 2.

The structure of TETPZE was determined at the University of Kentucky using an Enraf-Nonius CAD4/F diffractometer and a program package described previously (Brock & Webster, 1976). Crystals of TETPZE (Trofimenko, 1967b) grown from $\text{CHCl}_3/\text{C}_6\text{H}_{12}$ as prisms elongated along \underline{a} and bounded by faces of the forms $\{010\}$, $\{011\}$, and $\{10\bar{1}\}$. Crystal 0.15 x 0.22 x 0.38 mm (all thicker crystals examined were twinned), θ - 2θ scans, $2\theta \leq 55^\circ$, 4839 reflections, 1880 with $F^2 > 2\sigma(F^2)$ treated as observed, 3 standard reflections, no absorption correction, correction for decomposition of 1.7% overall, 24 reflections having $22 < 2\theta < 24^\circ$ for measuring lattice parameters; $0 \leq h \leq 11$, $0 \leq k \leq 28$, $-15 \leq l \leq 15$. All C, B, and N atoms refined anisotropically, H atoms isotropically; an extinction factor (Zachariasen, 1967), final value $2.76(11) \times 10^{-5}$, improved agreement significantly. Final agreement indices [refinement on F , $w = 4F_o^2/\sigma^2(F_o^2)$ where $\sigma^2(F_o^2)$ includes a term $(.02 F_o^2)^2$] are $R = 0.036$, $wR = 0.033$ for 362 variables; error in an observation of unit weight is 1.38; most important features in a final difference-Fourier map have heights $\pm 0.17 e^{-\rho^{-3}}$. Positional parameters given in Table 5 and some bond lengths and angles in Tables 2 and 3; the molecular structure is illustrated in Figure 3.

Refined atomic displacement parameters and observed and calculated structure-factor amplitudes are available for all three structures.*

Discussion. If an analogy is made with 9,10-dihydroanthracene, the six-membered B_2N_4 ring of the pyrazabole would be expected to adopt either a boat or a planar conformation; however, the first structure of a member of this class [1,3,5,7,-tetramethyl-4,8-bis(3,5-dimethylpyrazol-1'-yl)pyrazabole;

*The refined atomic displacement parameters and observed and calculated structure-factor amplitudes have been deposited with the British Library Lending Division as Supplementary Publication No. SUP _____ (_____ pp.). Copies may be obtained through The Executive Secretary, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

Alcock & Sawyer, 1974] revealed a flattened chair conformation. The authors invoked steric interactions to explain this observation, and it is true that no other pyrazabole with comparable crowding has been studied crystallographically. The second structure done (2,6-dibromo-4,4,8,8-tetraethylpyrazabole; Holt, Tebben, Holt & Watson, 1977) contains a somewhat asymmetrical B_2N_4 ring that is approximately planar.

Quite recently ten more pyrazabole structures have been determined (Niedenzu & Nöth, 1983; Hanecker, Hodgkins, Niedenzu & Nöth, 1984; this work) and it is now possible to compare B_2N_4 ring conformations across a variety of substituents and crystal packing arrangements. Two dihedral angles have been chosen to describe ring geometries: an angle θ_1 between the plane of the four N atoms and a plane defined by a N-B-N grouping, and an angle θ_2 between the plane of the four N atoms and a bridging pyrazolyl ring. In the absence of imposed symmetry each molecule has two θ_1 and two θ_2 values, but in all cases but one (see below) the two corresponding values differ by at most a few degrees. The average value of the angle θ_1 measures the deviation of the B_2N_4 ring from planarity, and the average value of θ_2 describes the extent to which the bridging pyrazolyl rings are folded back in a butterfly arrangement. In all twelve structures the four N atoms of the B_2N_4 ring and the five atoms of the N_2C_3 rings form good planes; most deviations are much smaller than the maximum value of 0.02 Å.

A plot of θ_2 vs. θ_1 for eleven of the twelve structures (Figure 4) shows that a wide range of ring conformations has been observed. The interconversion of boat and chair forms must require only a small amount of energy, and must therefore be sensitive to substituent and packing effects. The boat form is, however, observed most frequently, and so is probably the

favored conformation.* The plot also confirms the expected correlation between θ_1 and θ_2 : the more pronounced the boat conformation, the greater the bending back of the pyrazolyl rings. Evidence from molecular mechanics and quantum mechanical calculations for the related hydrocarbon 9,10-dihydroanthracene (Lipowitz et al., 1982) and from various studies of its derivatives (Rabideau, 1978) suggests that the energy difference between the boat and planar conformations in such ring systems is very small, perhaps less than 0.5 kcal/mol for a θ_2 value of 10 deg. In the pyrazaboles, which may be even more flexible than the hydrocarbon, the chair conformation is also accessible.

The one structure omitted from Figure 4 is that of 2,6-dibromo-4,4,8,8-tetraethylpyrazabole (Holt, Tabben, Holt & Watson, 1977). Its two angles θ_2 are similar at 1.2 and 3.2 deg, but the two values of θ_1 , -0.5 and 12.2 deg, vary considerably. One end of the B_2N_4 ring is nearly planar, while the B atom at the other end is 0.24 Å out of the plane of the four N atoms. None of the other eleven structures even approaches this degree of asymmetry.

Differences in bond lengths and angles between bridging and terminal pyrazolyl rings are best examined in the structure of TETPZB. Not only is extensive averaging possible over the chemically equivalent bonds, but the internal consistency of the four (or in two cases, two) independent length and angle determinations is excellent. Figure 5 shows some of the average values; bond angles in the B_2N_4 ring are in general sensitive to ring conformation and so are not included. There is clear evidence of localization of double bonds in the terminal rings. Although the differences within the pairs of N-C and C-C average bond lengths are small, they bracket the corresponding average

*The recently completed structure of 4,4,8,8-tetrafluoropyrazabole (Nöth & Hanecker, 1984) also contains boat-shaped molecules with $\theta_1 = 30^\circ$.

values for the bridging rings. The same is true for the bond angles. Although a correction for the effects of thermal motion might increase the absolute values of these averages by ca. 0.003-0.006 Å, the pattern shown is probably insensitive to that kind of systematic error. A very similar pattern is also seen in the structure of BISPZ3. The structures of BISPZB and TETPZB show that a B-N bond to a terminal ring is about 0.040(4) Å shorter than a B-N bond to a bridging ring.

The B_2N_4 rings in both BISPZB and TETPZB adopt pronounced boat conformations (see Fig. 4); the arrangements of the two terminal rings on each B atom are also very similar (see Figs. 2 and 3). The orientations of the terminal rings in 4,8-diphenyl-4,8-bis(pyrazol-1'-yl)pyrazabole (Niedenzu & Nöth, 1983; site symmetry $\bar{1}$) are also related to those of BISPZB and TETPZB. The planes of the equatorial phenyl rings are almost perpendicular to the plane of the four N atoms and approximately bisect the N-B-N angles. The planes of the axial pyrazolyl rings are about perpendicular to both the equatorial rings and the central N-atom plane. The appearance of this same general arrangement of the terminal rings across three structures, two boat forms and one chair, suggests that it is the favored conformation for cis disubstitution by aromatic rings. In both BISPZB and TETPZB the uncoordinated N atoms of the pseudo-equatorial pyrazolyl rings are oriented so that the N-N bonds approximately eclipse the B-B vectors.

Acknowledgment. This work was supported by the Office of Naval Research (K.N.) and the Fonds der Chemischen Industrie (H.N.).

References

- ALCOCK, N.W. & SAWYER, J.F. (1974). Acta Cryst. B30, 2899-2901.
- BROCK, C.P. & WEBSTER, D.F. (1976). Acta Cryst. B32, 2089-2094.
- CLARKE, C.M., NIEDENZU, K., NIEDENZU, P.M. & TROFIMENKO, S. (1984). In Preparation.
- HANECKER, E., HODGKINS, T.G., NIEDENZU, K. & NÖTH, H. (1984). Inorg. Chem., in press.
- HOLT, E.M., TEBBEN, S.L., HOLT, S.L. & WATSON, K.J. (1977). Acta Cryst. B33, 1986-1988.
- LIPKOWITZ, K.B., RABIDEAU, P.W., RABER, D.J., HARDEE, L.E., SCHLEYER, P.v.R., KOS, A.J. & KAHN, R.A. (1982). J. Org. Chem. 47, 1002-1005.
- NIEDENZU, K. & NIEDENZU, P.M. (1984). Inorg. Chem., in press.
- NIEDENZU, K. & NÖTH, H. (1983). Chem. Ber. 116, 1132-1153.
- NÖTH, H. & HANECKER, E. (1984). In preparation.
- RABIDEAU, P.W. (1978). Acc. Chem. Res. 11, 141-147.
- TROFIMENKO, S. (1967a). J. Amer. Chem. Soc. 89, 3165-3170.
- TROFIMENKO, S. (1967b). J. Amer. Chem. Soc. 89, 4948-4952.
- ZACHARIASEN, W.H. (1967). Acta Cryst. 23, 558-564.

Table 1. Atom Coordinates ($\times 10^4$) and Temperature Factors for
1,3,5,7-Tetramethylpyrazabole (ME4PZB)

Atom	x	y	z	$B_{eq}, \text{\AA}^2$
B(1)	256(4)	898(3)	7196(2)	4.2
B(2)	-857(4)	2778(3)	6045(2)	4.1
N(1)	1682(2)	1732(2)	6842(1)	3.7
N(2)	1158(2)	2624(2)	6294(1)	3.8
N(3)	-1526(2)	1593(2)	5750(1)	3.5
N(4)	-999(2)	701(2)	6300(1)	3.4
C(1)	3451(3)	1783(2)	6971(2)	4.3
C(2)	4042(3)	2719(3)	6512(2)	5.1
C(3)	2594(3)	3232(2)	6086(2)	4.5
C(4)	-2569(3)	1228(2)	5016(2)	3.6
C(5)	-2705(3)	92(2)	5098(2)	3.7
C(6)	-1705(3)	-218(2)	5906(2)	3.4
C(11)	4454(5)	931(4)	7536(3)	6.3
C(31)	2458(4)	4266(3)	5505(3)	6.6
C(41)	-3364(4)	1983(3)	4275(2)	5.0
C(61)	-1416(4)	-1339(2)	6328(2)	4.7
H(11)	874(35)	133(20)	7446(18)	*
H(12)	-566(33)	1337(20)	7814(17)	*
H(21)	-1501(34)	3074(21)	6696(19)	*
H(22)	-995(34)	3322(21)	5444(19)	*
H(2)	5093(32)	2982(19)	6441(17)	*
H(5)	-3311(33)	-346(20)	4682(17)	*
H(11A)	5590(39)	1107(23)	7497(20)	*
H(11B)	4451(36)	197(25)	7228(19)	*
H(11C)	4229(38)	921(23)	8181(23)	*
H(31A)	3579(34)	4524(19)	5409(17)	*
H(31B)	1945(33)	4145(20)	4852(18)	*
H(31C)	1793(33)	4840(21)	5795(18)	*
H(41A)	-3870(37)	1612(22)	3769(21)	*
H(41B)	-2656(36)	2595(22)	4089(20)	*
H(41C)	-4339(33)	2381(21)	4512(18)	*
H(61A)	-2213(38)	-1494(23)	6785(21)	*
H(61B)	-346(37)	-1444(22)	6619(20)	*
H(61C)	-1593(36)	-1870(22)	5898(21)	*

*Fixed at 5.5\AA^2 .

Table 2. Selected Bond Lengths (\AA).

	ME4PZB	BISPZB	TETPZB
B(1)-N(1)	1.560(4)	1.549(2)	1.545(3)
B(1)-N(4)	1.561(3)	1.558(2)	1.551(3)
B(2)-N(2)	1.563(4)	1.563(2)	1.555(3)
B(2)-N(3)	1.565(4)	1.564(2)	1.557(3)
B(1)-N(5)		1.515(2)	1.516(3)
B(1)-N(7)		1.519(2)	1.510(3)
B(2)-N(9)			1.513(3)
B(2)-N(11)			1.519(3)
N(1)-N(2)	1.368(3)	1.368(1)	1.370(2)
N(3)-N(4)	1.369(3)	1.358(1)	1.369(2)
N(5)-N(6)		1.373(2)	1.367(3)
N(7)-N(8)		1.368(1)	1.364(3)
N(9)-N(10)			1.362(3)
N(11)-N(12)			1.372(3)
N(1)-C(1)	1.345(3)	1.346(2)	1.340(3)
N(2)-C(3)	1.345(3)	1.335(2)	1.342(3)
N(3)-C(4)	1.344(3)	1.336(2)	1.337(3)
N(4)-C(6)	1.339(3)	1.342(2)	1.331(3)
N(5)-C(7)		1.355(2)	1.356(3)
N(7)-C(10)		1.357(2)	1.354(3)
N(9)-C(13)			1.349(3)
N(11)-C(16)			1.351(3)
N(6)-C(9)		1.322(2)	1.326(3)
N(8)-C(12)		1.331(2)	1.324(3)
N(10)-C(15)			1.318(3)
N(12)-C(18)			1.338(4)
C(1)-C(2)	1.373(4)	1.370(2)	1.361(4)
C(2)-C(3)	1.377(4)	1.369(2)	1.360(4)
C(4)-C(5)	1.378(4)	1.372(2)	1.367(4)
C(5)-C(6)	1.385(4)	1.364(2)	1.364(4)
C(1)-C(11)	1.486(5)		
C(3)-C(31)	1.488(4)		
C(4)-C(41)	1.486(4)		
C(6)-C(61)	1.485(4)		
C(7)-C(8)		1.358(2)	1.358(4)
C(10)-C(11)		1.362(2)	1.352(5)
C(13)-C(14)			1.359(4)
C(16)-C(17)			1.362(4)
C(8)-C(9)		1.385(2)	1.379(4)
C(11)-C(12)		1.382(2)	1.370(5)
C(14)-C(15)			1.367(4)
C(17)-C(18)			1.364(5)
B(1)-H(11)	1.086(25)		
B(1)-H(12)	1.190(24)		
B(2)-H(21)	1.091(27)	1.151(13)	
B(2)-H(22)	1.061(26)	1.067(13)	

Table 3. Selected Bond Angles ($^{\circ}$).

	ME4PZB	BISPZB	TETPZB
N(1)-B(1)-N(4)	105.1(2)	105.6(1)	105.6(2)
N(2)-B(2)-N(3)	104.8(2)	103.6(1)	105.3(2)
N(1)-B(1)-N(5)		111.2(1)	109.4(2)
N(4)-B(1)-N(5)		109.2(1)	110.8(2)
N(1)-B(1)-N(7)		109.7(1)	110.7(2)
N(4)-B(1)-N(7)		110.5(1)	110.0(2)
N(2)-B(2)-N(9)			109.3(2)
N(3)-B(2)-N(9)			109.6(2)
N(2)-B(2)-N(11)			110.9(2)
N(3)-B(2)-N(11)			109.5(2)
N(5)-B(1)-N(7)		110.5(1)	110.2(2)
N(9)-B(2)-N(11)			112.0(2)
B(1)-N(1)-N(2)	119.1(2)	119.6(1)	121.2(2)
B(1)-N(4)-N(3)	118.9(2)	121.0(1)	121.6(2)
B(2)-N(2)-N(1)	118.9(2)	121.0(1)	122.3(2)
B(2)-N(3)-N(4)	118.9(2)	119.7(1)	121.9(2)
H(11)-B(1)-H(12)	112.1(18)		
H(21)-B(2)-H(22)	113.9(19)	115.7(10)	

Table 4. Atom Coordinates ($\times 10^4$) and Temperature Factors for
4,4-Bis(pyrazol-1'-yl)pyrazobole (BISPZB).

Atom	x	y	z	B_{eq} or $B, \text{\AA}^2$
B(1)	6515(2)	2269(2)	7138(2)	3.2
B(2)	8782(2)	4539(2)	6679(2)	4.4
N(1)	6286(1)	2975(1)	5674(1)	3.5
N(2)	7339(1)	4037(1)	5456(1)	3.8
N(3)	8094(1)	4599(1)	8101(1)	3.9
N(4)	7094(1)	3515(1)	8327(1)	3.5
N(5)	4988(1)	1586(1)	7392(1)	3.5
N(6)	3642(1)	3213(1)	7231(1)	4.4
N(7)	7734(1)	1159(1)	7163(1)	3.5
N(8)	8486(1)	0723(1)	8457(1)	4.4
C(1)	5161(2)	2788(2)	4474(1)	4.2
C(2)	5498(2)	3720(2)	3494(2)	4.9
C(3)	6854(2)	4486(2)	4148(2)	4.7
C(4)	8290(2)	5548(2)	9254(2)	4.9
C(5)	7427(2)	5084(2)	10230(2)	5.4
C(6)	6690(2)	3812(2)	9618(2)	4.5
C(7)	4747(2)	0317(2)	7957(2)	4.3
C(8)	3248(2)	0210(2)	8165(2)	5.2
C(9)	2611(2)	1463(2)	7690(2)	4.5
C(10)	8005(2)	0272(1)	6048(2)	4.3
C(11)	8944(2)	-0753(2)	6619(2)	5.2
C(12)	9208(2)	-0432(2)	8103(2)	4.8
H(21)	9673(15)	3667(13)	6693(13)	4.3(3)
H(22)	9202(16)	5594(14)	6514(14)	5.2(3)
H(1)	4263(15)	2079(13)	4457(14)	4.8(3)
H(2)	4870(18)	3782(15)	2585(17)	5.4(3)
H(3)	7417(18)	5285(16)	3866(17)	6.5(4)
H(4)	8954(18)	6369(16)	9228(16)	6.2(3)
H(5)	7352(19)	5581(17)	11097(18)	7.2(4)
H(6)	5962(17)	3181(15)	9949(15)	4.8(3)
H(7)	5550(17)	-0298(14)	8136(14)	4.3(3)
H(8)	1553(17)	1736(15)	7655(15)	5.6(3)
H(9)	2738(19)	-0553(17)	8497(17)	6.3(4)
H(10)	7585(16)	0442(14)	5069(14)	5.2(3)
H(11)	9342(18)	-1445(15)	6085(16)	6.2(3)
H(12)	9857(18)	-0884(16)	8878(17)	5.9(3)

TABLE 5. POSITIONAL AND THERMAL PARAMETERS FOR THE ATOMS OF
4,4,8,8-TETRAKIS(PYRAZOL-1'-YL)PYRAZABOLE (TEFPZB)

ATOM	A X	Y	Z	B or B_{eq} , \AA^2	ATOM	X	Y	Z	B or B_{eq} , \AA^2
N(1)	0.2693(2)	0.44465(9)	-0.08711(16)	2.9	C(14)	0.5064(5)	0.36251(16)	0.4061(3)	5.8
N(2)	0.3173(2)	0.43257(9)	0.03503(16)	2.9	C(15)	0.6871(4)	0.37568(15)	0.3426(3)	5.3
N(3)	0.3921(2)	0.32562(9)	-0.00359(17)	2.9	C(16)	0.0790(4)	0.29476(15)	0.0345(3)	4.1
N(4)	0.3342(2)	0.33808(9)	-0.12546(17)	3.1	C(17)	-0.0576(4)	0.29467(20)	0.0640(3)	5.4
H(5)	0.0471(2)	0.37466(10)	-0.20291(17)	3.3	C(18)	-0.0463(4)	0.3455(2)	0.1346(3)	5.7
H(6)	-0.0083(3)	0.31939(11)	-0.25603(19)	4.3	D(1)	0.2200(4)	0.39235(14)	-0.1818(3)	3.1
N(7)	0.2405(2)	0.41195(10)	-0.29988(17)	3.5	B(2)	0.3286(3)	0.36608(14)	-0.0860(2)	3.1
N(8)	0.3874(3)	0.42733(11)	-0.30525(20)	4.6	H(1)	0.245(3)	0.5225(10)	-0.181(2)	3.9(6)
N(9)	0.4552(2)	0.36386(9)	0.21086(16)	3.4	H(2)	0.327(3)	0.5733(12)	0.025(2)	3.6(6)
N(10)	0.6107(3)	0.37695(11)	0.22437(19)	4.8	H(3)	0.383(3)	0.4855(10)	0.177(2)	3.8(6)
N(11)	0.1664(2)	0.34424(10)	0.08784(17)	3.5	H(4)	0.512(3)	0.2598(10)	0.097(2)	3.6(6)
N(12)	0.0091(3)	0.37726(12)	0.1518(2)	5.1	H(5)	0.540(3)	0.2223(12)	-0.100(2)	4.6(7)
C(1)	0.2696(3)	0.50575(13)	-0.1012(3)	3.6	H(6)	0.378(3)	0.2960(11)	-0.266(2)	4.9(6)
C(2)	0.3166(4)	0.53323(15)	0.0099(3)	4.1	H(7)	-0.053(3)	0.4488(13)	-0.152(2)	4.5(7)
C(3)	0.3466(3)	0.48689(13)	0.0927(3)	3.6	H(8)	-0.315(4)	0.3888(13)	-0.235(3)	6.9(9)
C(4)	0.4722(3)	0.27481(13)	0.0169(3)	3.7	H(9)	-0.226(3)	0.2846(12)	-0.304(2)	5.0(7)
C(5)	0.4831(4)	0.25451(15)	-0.0909(3)	4.3	H(10)	0.019(4)	0.4114(15)	-0.416(3)	8.7(11)
C(6)	0.3965(3)	0.29495(13)	-0.1772(3)	3.9	H(11)	0.150(4)	0.4413(17)	-0.577(3)	10.8(11)
C(7)	-0.0746(4)	0.40801(16)	-0.1883(3)	4.1	H(12)	0.452(4)	0.4547(14)	-0.443(3)	7.1(9)
C(8)	-0.2113(4)	0.37448(17)	-0.2319(3)	4.6	H(13)	0.337(3)	0.3473(12)	0.325(2)	5.6(8)
C(9)	-0.1639(4)	0.32060(17)	-0.2720(3)	4.8	H(14)	0.613(4)	0.3581(14)	0.406(3)	7.5(9)
C(10)	0.1277(5)	0.41810(19)	-0.4110(3)	6.2	H(15)	0.807(4)	0.3849(12)	0.370(2)	6.1(8)
C(11)	0.2014(5)	0.4371(2)	-0.4892(3)	8.0	H(16)	0.113(3)	0.2681(13)	-0.020(2)	6.3(8)
C(12)	0.3601(5)	0.44229(17)	-0.4203(3)	6.0	H(17)	-0.142(4)	0.2661(14)	0.034(3)	7.2(9)
C(13)	0.4395(5)	0.35468(15)	0.3206(3)	4.9	H(18)	-0.117(4)	0.3630(13)	0.175(2)	6.8(8)

A ESTIMATED STANDARD DEVIATIONS IN THE LEAST SIGNIFICANT FIGURE(S) ARE GIVEN IN PARENTHESES IN THIS AND ALL SUBSEQUENT TABLES.
B THE EQUIVALENT DIS. ARE CALCULATED FROM THE ANISOTROPIC THERMAL PARAMETERS.

Figure Captions

Figure 1. Perspective drawing of the molecular structure of 1,3,5,7-tetramethylpyrazabole (ME4PZB). In this and the following drawings the shapes of the ellipsoids for the non-H atoms correspond to 50% probability contours of atomic displacement.

Figure 2. Perspective drawing of the molecular structure of 4,4-bis(pyrazol-1'-yl)pyrazabole (BISPZB).

Figure 3. Perspective drawing of the molecular structure of 4,4,8,8-tetrakis(pyrazol-1'-yl)pyrazabole (TETPZB).

Figure 4. A scatter plot showing the relationship between the dihedral angles θ_1 of the planes (B1,N1,N4) and (B2,N2,N3) with the plane (N1,N2,N3,N4), and the angles θ_2 of the planes (N1,N2,C1,C2,C3) and (N3,N4,C4,C5,C6) with the plane (N1,N2,N3,N4) for 11 pyrazabole structures (see Fig. 1-3 for the atom-numbering scheme). The structures are (1) 1,3,5,7-tetramethyl-4,8-bis(3,5-dimethylpyrazol-1'-yl)pyrazabole (Alcock & Sawyer, 1974); (2) 4,4,8,8-bis(ethylenedithio)pyrazabole (Niedenzu & Nöth, 1983); (3) 4,4,8,8-tetrakis(methylthio)pyrazabole (Niedenzu & Nöth, 1983); (4) 4,4,8,8-tetrachloropyrazabole (Niedenzu & Nöth, 1983); (5) 4,8-diphenyl-4,8-bis(pyrazol-1'-yl)pyrazabole (Niedenzu & Nöth, 1983); (6) 4,8-dibromo-2,6-dichloropyrazabole (Niedenzu & Nöth, 1983); (7) pyrazabole (Hanecker, Hodgkins, Niedenzu & Nöth, 1984); (8) 4,4,8,8-tetrabromopyrazabole (Hanecker, Hodgkins, Niedenzu & Nöth, 1984); (9) 1,3,5,7-tetramethylpyrazabole (this paper); (10)

4,4-bis(pyrazol-1'-yl)pyrazabole (this paper); and (11)
4,4,8,8-tetrakis(pyrazol-1'-yl)pyrazabole (this paper). The
rectangle for each structure covers the range of the two θ_1 and two
 θ_2 values that are observed in the absence of imposed symmetry (see
text).

Figure 5. Diagram showing some average bond lengths and angles for
4,4,8,8-tetrakis(pyrazabole-1'-yl)pyrazabole. The numbers in
parentheses are the esd's of the means. The esd's for the angles
range from 0.1 to 0.4 deg, and average 0.2 deg.

Figure 1.





Figure 2.






Figure 3.

Figure 4.





Figure 5.



END

FILMED

2-85

DTIC