

AD-A155 763

FINITE TEMPERATURE STABILIZATION OF THE GRADIENT DRIFT  
INSTABILITY IN BARIUM CLOUDS(U) NAVAL RESEARCH LAB  
WASHINGTON DC J F DRAKE ET AL. 21 MAY 85 NRL-MR-5566

1/1

UNCLASSIFIED


F/G 20/9

NL

END

FORM

DTIC



NATIONAL BUREAU OF STANDARDS  
MICROCOPY RESOLUTION TEST CHART

# Finite Temperature Stabilization of the Gradient Drift Instability in Barium Clouds

J. F. DRAKE

*Science Applications International Corp.  
McLean, VA 22102*

J. D. HUBA AND S. T. ZALESK

*Geophysical and Plasma Dynamics Branch  
Plasma Physics Division*

May 21, 1985

This research was sponsored by the Defense Nuclear Agency under Subtask QIEQMXBB, work unit 00005 and work unit title "Plasma Structure Evolution."



DTIC  
ELECTE  
JUN 25 1985  
S B D

NAVAL RESEARCH LABORATORY  
Washington, D.C.

Approved for public release; distribution unlimited.

AD-A155 763

DTIC FILE COPY

## REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>		1b. RESTRICTIVE MARKINGS	
2a SECURITY CLASSIFICATION AUTHORITY		3 DISTRIBUTION / AVAILABILITY OF REPORT <b>Approved for public release; distribution unlimited.</b>	
2b DECLASSIFICATION / DOWNGRADING SCHEDULE			
4 PERFORMING ORGANIZATION REPORT NUMBER(S) <b>NRL Memorandum Report 5566</b>		5 MONITORING ORGANIZATION REPORT NUMBER(S)	
5a. NAME OF PERFORMING ORGANIZATION <b>Naval Research Laboratory</b>	6b OFFICE SYMBOL (if applicable) <b>Code 4780</b>	7a NAME OF MONITORING ORGANIZATION	
6c ADDRESS (City, State, and ZIP Code) <b>Washington, DC 20375-5000</b>		7b ADDRESS (City, State, and ZIP Code)	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION <b>Defense Nuclear Agency</b>	8b. OFFICE SYMBOL (if applicable) <b>RAAE</b>	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c ADDRESS (City, State, and ZIP Code) <b>Washington, DC 20305</b>		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO. <b>62715H</b>	PROJECT NO. <b></b>
		TASK NO. <b></b>	WORK UNIT ACCESSION NO. <b>DN580-072</b>
11 TITLE (Include Security Classification) <b>Finite Temperature Stabilization of the Gradient Drift Instability in Barium Clouds</b>			
12 PERSONAL AUTHOR(S) <b>Drake, J.F.,* Huba, J.D. and Zalesak, S.T.</b>			
13a TYPE OF REPORT <b>Interim</b>	13b TIME COVERED FROM <b>10/84</b> TO <b>10/85</b>	14. DATE OF REPORT (Year, Month, Day) <b>1985 May 21</b>	15 PAGE COUNT <b>42</b>
16 SUPPLEMENTARY NOTATION <b>*Science Applications International Corporation, McLean, VA 22102</b> (Continues)			
17 COSATI CODES		18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
		Gradient drift instability Ionospheric structure	
		Barium clouds	
19 ABSTRACT (Continue on reverse if necessary and identify by block number) <p>We present a relatively simple analysis of the gradient drift instability in barium clouds which includes the effects of both finite temperature and finite parallel length. It is found that short wavelength modes are stabilized as the electrons redistribute parallel to the magnetic field and neutralize the charge imbalance set up by the instability. An analytical expression for the critical wavenumber for stabilization is given, as well as numerical results. We discuss the application of these results to the structuring of barium clouds.</p>			
20 DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> OTIC USERS		21 ABSTRACT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>	
22a NAME OF RESPONSIBLE INDIVIDUAL <b>J. D. Huba</b>		22b TELEPHONE (Include Area Code) <b>(202) 767-3630</b>	22c OFFICE SYMBOL <b>Code 4780</b>

**16. SUPPLEMENTARY NOTATION (Continued)**

This research was sponsored by the Defense Nuclear Agency under Subtask QIEQMXBB, work unit 00005 and work unit title "Plasma Structure Evolution."

CONTENTS

I. INTRODUCTION ..... 1

II. GENERAL EQUATIONS ..... 3

III. EQUILIBRIUM ..... 7

IV. LINEARIZED EQUATIONS AND DISPERSION EQUATION ..... 11

V. NUMERICAL RESULTS ..... 21

VI. CONCLUDING REMARKS ..... 22

ACKNOWLEDGMENT ..... 24

REFERENCES ..... 27

**S** DTIC  
 ELECTE **D**  
 JUN 25 1985

B



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

# FINITE TEMPERATURE STABILIZATION OF THE GRADIENT DRIFT INSTABILITY IN BARIUM CLOUDS

## I. INTRODUCTION

The study of the dynamic evolution of artificial plasma clouds (e.g., barium clouds) in the earth's ionosphere and magnetosphere has been an active area of research for more than two decades. The initial motivation for these active space experiments was to use the plasma cloud as a diagnostic to determine ambient plasma conditions (e.g., electric fields, neutral winds). ~~However~~ it was ~~soon~~ discovered that the interaction of an artificial plasma cloud and the near-earth space plasma was very complex, and that plasma clouds undergo a complicated and dynamic evolution. One of the prominent features of cloud evolution in the ionosphere is the development of field-aligned striations on the cloud's steepened "backside". These striations have been attributed to the onset of the gradient drift instability. ~~(Linson and Workman, 1970)~~ This instability can occur in a weakly collisional plasma which contains a density gradient and a neutral wind (or ambient electric field). ~~(Simon, 1963; Hoh, 1963)~~

A substantial amount of theoretical and computational research has been invested in understanding the linear and nonlinear development of the gradient drift instability and its relevance to plasma cloud structure (Volk and Haerendel, 1971; Perkins et al., 1973; Zabusky et al., 1973; Shiau and Simon, 1972; Perkins and Doles, 1975; Scannapieco et al., 1976; Chaturvedi and Ossakow, 1979; Keskinen et al., 1980; McDonald et al., 1980; McDonald et al., 1981; Huba et al., 1983). The bulk of analyses to date have neglected the effects of plasma dynamics parallel to the ambient magnetic field  $B_0$  (i.e., considered perturbations only in the plane transverse to  $B_0$ ) or have incorporated parallel effects in a crude way.

Manuscript approved February 22, 1985.

Francis and Perkins (1975), for example, assume that the ambient and perturbed potentials map uniformly along  $B_0$  thereby connecting the cloud dynamics to that of the conducting background at different altitudes. In this model finite thermal effects have a stabilizing influence on short wavelength modes. However, several studies have been performed which attempt to include parallel dynamics self-consistently in the stability analysis (Goldman et al., 1976; Sperling and Glassman, 1984; Sperling et al., 1984). These studies incorporate the parallel length of the cloud along the ambient field into the stability analysis and have shown that parallel effects can play an important role in the development of the gradient drift instability as it relates to ionospheric plasma clouds.

In particular, Sperling et al. (1984) have recently shown that when the finite size of the cloud along  $B_0$  is self-consistently incorporated in the linear stability analysis, the long wavelength modes tend to have much smaller growth rates than short wavelength modes. This reduction in growth due to finite cloud length occurs because the integrated Pedersen conductivity of the background plasma over the extent of the mode along the field becomes greater than the cloud. The distance the mode extends along the field is proportional to the perpendicular wavelength of the mode so that long wavelength modes are more strongly affected. However, this analysis neglected the effects of finite temperature. Sperling and Glassman (1984) included finite temperature in the analysis and found that short wavelength modes became propagating rather than purely growing modes. Moreover, for sufficiently short wavelengths there was some evidence that the modes were be completely stabilized. A drawback of this work though is that a relatively complex, second-order differential equation is solved numerically to obtain results. The underlying physics of the stabilization mechanism is therefore somewhat obscure and not



addressed in the paper. Nonetheless, it is evident that finite temperature and finite cloud length effects can impact the development of the gradient drift instability.

The purpose of this paper is to present a linear stability analysis of the gradient drift instability which incorporates both finite temperature and finite parallel length effects. A simple plasma model is used (similar to the one used in Sperling et al. (1984)) which permits an analytical solution to the dispersion equation. It is found that the short wavelength modes are stabilized by redistribution of electrons parallel to  $B_0$  (i.e., parallel electron diffusion or parallel electron response to the perturbed fields) which neutralize the charge imbalance set up by the instability. An analytical expression for the critical wavenumber for the stabilization of instability is given, as well as numerical results.

The organization of the paper is as follows. In the next section we derive a set of general nonlinear equations which describe the evolution of a three dimensional plasma cloud. In Section III we discuss the equilibrium to be used in the instability analysis. In Section IV we derive a general dispersion equation, and present both analytical and numerical results. Finally, in the last section we summarize our findings and discuss the relevance of our results to cloud structure.

## II. GENERAL EQUATIONS

We first derive a set of nonlinear equations to describe the evolution of a warm plasma cloud in a uniform magnetic field  $\underline{B} = B_0 \hat{z}$  with a background neutral wind  $\underline{V}_n = V_n \hat{x}$  [see Fig. 1a]. For simplicity we consider only low frequency  $\partial/\partial t \ll v_\alpha$  motion of the cloud and take the electron collisions to be sufficiently weak so that  $v_e/\Omega_e \ll 1$  but allow  $v_i/\Omega_i$  to be arbitrary. The collision frequency and gyrofrequency of

the  $\alpha$  species are given by  $v_\alpha$  and  $\Omega_\alpha$ , respectively. Both electrons and ions are taken to be warm and for simplicity we consider the isothermal limit. In this case the fundamental equations of our analysis are continuity, momentum transfer, charge neutrality and Ampere's law:

$$\frac{\partial n}{\partial t} + \nabla \cdot (n \mathbf{v}_e) = 0 \quad (1)$$

$$0 = -e \mathbf{E} - \frac{e}{c} \mathbf{v}_e \times \mathbf{B} - m_e \nu_{en} (\mathbf{v}_e - \mathbf{v}_n) - m_e \nu_{ei} (\mathbf{v}_e - \mathbf{v}_i) - \frac{T_e}{n} \nabla n \quad (2)$$

$$0 = e \mathbf{E} + \frac{e}{c} \mathbf{v}_i \times \mathbf{B} - m_i \nu_{in} (\mathbf{v}_i - \mathbf{v}_n) - m_i \nu_{ie} (\mathbf{v}_i - \mathbf{v}_e) - \frac{T_i}{n} \nabla n \quad (3)$$

$$\nabla \cdot \mathbf{J} = \nabla \cdot [n(\mathbf{v}_i - \mathbf{v}_e)] = 0 \quad (4)$$

$$\nabla \times \mathbf{B} = \frac{4\pi}{c} \mathbf{J} \quad (5)$$

where  $v_\alpha$  and  $T_\alpha$  are the fluid velocity and temperature of species  $\alpha$ , respectively,  $\nu_{\alpha n}$  is the  $\alpha$  specie-neutral collision frequency,  $\nu_{ei}$  is the electron-ion collision frequency,  $\nu_{ie}$  is the ion-electron collision frequency, and  $\alpha$  refers to electrons (e) or ions (i). We take the electric and magnetic fields to be represented by potentials as

$$\mathbf{E} = -\nabla\phi - \frac{1}{c} \frac{\partial A}{\partial t} \hat{z} \quad (6)$$

and

$$\mathbf{B} = B_0 \hat{z} + \nabla A_z \times \hat{z} \quad (7)$$

where  $\phi$  is the electrostatic potential and  $A_z$  is the vector potential associated with the magnetic field produced by the self-consistent plasma currents. We consider only  $A_z$  since  $J_{\parallel} \gg J_{\perp}$  and assume  $|\nabla A_z \times \hat{z}| \ll B_0$ .

The electron cross-field motion is given by

$$\tilde{v}_{e\perp} = -\frac{c}{B} \nabla_{\perp} \phi \times \hat{z} + \frac{cT_e}{eBn} \nabla_{\perp} n \times \hat{z} \quad (8)$$

while the parallel motion is given by

$$v_{\parallel e} = -\frac{[eE_{\parallel} + T_e \nabla_{\parallel} \ln(n) + f(v_{e\perp}/v_{en}) T_e \nabla_{\parallel} \ln(n)]}{[m_e (v_e + f v_{e\perp})]} \quad (9)$$

with

$$E_{\parallel} = -\nabla_{\parallel} \phi - \frac{1}{c} \frac{\partial A_z}{\partial t}, \quad (10)$$

where  $v_e = v_{e\perp} + v_{en}$ ,  $\nabla_{\parallel} = \hat{b} \cdot \nabla$ ,  $\hat{b} = B/|B_0|$ , and  $T = T_e + T_i$ .

The assumption that  $v_e/\Omega_e \ll 1$  while  $v_{in}/\Omega_i \sim 1$  requires that

$$f \equiv m_e v_{en}/m_i v_{in} \ll 1.$$

The ion cross-field motion is given by

$$\begin{aligned} \tilde{v}_{i\perp} = & \delta_i \left[ -\frac{c}{B} \nabla_{\perp} \phi \times \hat{z} + \frac{v_{in}}{\Omega_i} \nabla_{\perp} n \times \hat{z} - \frac{cT_i}{eBn} \nabla_{\perp} n \times \hat{z} \right. \\ & \left. - \frac{v_{in}}{\Omega_i} \frac{c}{B} \nabla_{\perp} \phi + \left( \frac{v_{in}}{\Omega_i} \right)^2 \nabla_{\perp} n - \frac{v_{in}}{\Omega_i} \frac{cT_i}{eBn} \nabla_{\perp} n \right], \end{aligned} \quad (11)$$

where  $\delta_i = (1 + v_{in}^2/\Omega_i^2)^{-1}$ , and the parallel motion is given by

$$v_{\parallel i} = \frac{[eE_{\parallel} + T_e \nabla_{\parallel} \ln(n)] v_{en} - v_e T_e \nabla_{\parallel} \ln(n)}{[m_i v_{in} (v_e + f v_{e\perp})]}. \quad (12)$$

In (11) we have included both the Pedersen and Hall responses to the electric field, neutral wind and pressure.

Substituting (8)-(12) into (1), (4) and (5) we find that

$$\frac{dn}{dt} - \frac{c}{B} \nabla \hat{\phi} \times \hat{z} \cdot \nabla n + \nabla_{\parallel} \left( \frac{c}{4\pi e} \nabla_{\perp}^2 A_z - D_{\parallel} \nabla_{\parallel} n \right) = 0, \quad (13)$$

$$\begin{aligned} \delta_{\perp} \frac{c}{B} \frac{v_{in}}{\Omega_i} \nabla_{\perp} \cdot n \nabla_{\perp} \hat{\phi} + \delta_{\perp} \frac{c}{B} \frac{v_{in}^2}{\Omega_i^2} \hat{z} \times \nabla \hat{\phi} \cdot \nabla n + \frac{v_{in}}{\Omega_i} \hat{z} \times \nabla n \cdot \nabla n \\ + D_{\perp} \nabla_{\perp}^2 n + \frac{c}{4\pi e} \nabla_{\parallel} \nabla_{\perp}^2 A_z = 0, \end{aligned} \quad (14)$$

$$\nabla_{\perp}^2 A_z = \frac{4\pi}{c\eta_e} \left( \nabla_{\parallel} \hat{\phi} + \frac{1}{c} \frac{dA_z}{dt} \right), \quad (15)$$

where  $\eta_e = m_e v_e / ne^2$  is the parallel resistivity,  $D_{\parallel} = T/m_i v_{in}$  and  $D_{\perp} = \delta_{\perp} (v_{in}/\Omega_i) cT/eB$  are the parallel and perpendicular ion transport coefficients,  $T = T_e + T_i$ , and

$$\hat{\phi} \equiv \phi - \frac{T_e}{e} \ln(n) - \frac{B}{c} \frac{v_{in}}{\Omega_i} \nabla n \cdot \hat{x}, \quad (16)$$

$$\frac{d}{dt} \equiv \frac{\partial}{\partial t} + \frac{v_{in}}{\Omega_i} \hat{z} \times \nabla n \cdot \nabla, \quad (17)$$

$$\nabla_{\parallel} \equiv \frac{\partial}{\partial z} + B_0^{-1} \nabla A_z \times \hat{z} \cdot \nabla. \quad (18)$$

Equation (13) is the electron continuity equation, (14) is the charge conservation equation and (15) is Ampere's law. Note that the electron pressure has been absorbed into  $\hat{\phi}$  in (16) and that terms of order  $f \ll 1$  have been discarded compared with those of order unity. Equations (13)-(18) constitute a complete description of the evolution of a three-dimensional warm plasma cloud.

### III. EQUILIBRIUM

We will consider the linear stability of a two-dimensional cloud which is localized both along and across the magnetic field  $B_0$ :  $n_c = n_c(x, z)$  with  $n_c \neq 0$  for  $|x| \lesssim x_c$  and  $|z| \lesssim z_c$ . The background plasma is taken to be uniform throughout the region  $|z| < z_b$  between two insulating plates at  $z = \pm z_b$ . The location of the plates, enables us to control the ratio of the total magnetic-field-line integrated Pedersen conductivity of the cloud to that of the background,  $n_c z_c / (n_c z_c + n_b z_b)$ . This ratio is an important parameter of the equilibrium configuration.

The equations describing the two dimensional plasma cloud are given by

$$\frac{\partial n}{\partial t} - D_{\parallel} \frac{\partial^2 n}{\partial z^2} + \frac{c}{4\pi e} \frac{\partial}{\partial z} \frac{\partial^2 A_z}{\partial x^2} = 0, \quad (19)$$

$$\delta_{\perp} \frac{c}{B} \frac{v_{in}}{\Omega_i} \frac{\partial}{\partial x} n \frac{\partial}{\partial x} \hat{\phi} + D_{\perp} \frac{\partial^2 n}{\partial x^2} + \frac{c}{4\pi e} \frac{\partial}{\partial z} \frac{\partial^2 A_z}{\partial x^2} = 0, \quad (20)$$

$$\frac{\partial^2 A_z}{\partial x^2} = \frac{4\pi}{c n_e} \left( \frac{\partial \hat{\phi}}{\partial z} + \frac{1}{c} \frac{\partial A_z}{\partial t} \right). \quad (21)$$

When there are no density gradients in the x direction, the solution to these equations is  $\hat{\phi} = - (B/c) (v_{in}/\Omega_i) v_n x$  so that

$$\phi = (T_e/e) \ln [n(z)/n_b] \quad (22)$$

with  $A_z = 0$ , i.e., the potential adjusts itself so that the electrons are in force balance along the magnetic field as the cloud diffuses along  $B_0$ . In writing the solution for  $\phi$  in (22) we have required  $\phi \rightarrow 0$  for  $|z| > z_c$  since we have assumed that only the charge accumulations associated with the motion of the plasma cloud itself are responsible for the development of  $\phi$ . There is no charge at  $z = \pm z_b$ . The density satisfies

$$\partial n / \partial t - D_{\parallel} \partial^2 n / \partial z^2 = 0 \quad (23)$$

so that  $n$  diffuses along  $B_0$ . In the opposite limit, where there are no density gradients in the  $z$  direction ( $\partial/\partial z = 0$ ),  $A_z = 0$  and

$$\hat{\phi} = - (T/e) \ln(n) - (B/c) (v_{in}/\Omega_i) v_n \int_{-\infty}^x dx (n_b/n) \text{ so that}$$

$$\phi = - \frac{T_i}{e} \ln\left[\frac{n(x)}{n_b}\right] + \frac{B}{c} \frac{v_{in}}{\Omega_i} v_n \int_{-\infty}^x dx \left(1 - \frac{n_b}{n}\right), \quad (24)$$

where we have required  $\phi \rightarrow 0$  as  $|x| \rightarrow \infty$ . In Eq. (24) the potential adjusts itself so that the perpendicular ion pressure gradient is balanced by the electrostatic field. In comparing the expressions for  $\phi$  in (22) and (24), it is important to note that the electron and ion pressures push the potential in the opposite direction (compare the signs of the terms proportional to  $T_e$  and  $T_i$ ).

We now return to the more general two dimensional equations in (19)-(21). These equations describe three time scales: the resistive flux diffusion time,  $\tau_r = 4\pi x_c^2 / \eta_e c^2$ ; the parallel diffusion time,  $\tau_{\parallel} = z_c^2 / D_{\parallel}$ ; and the perpendicular diffusion time,  $\tau_{\perp} = x_c^2 / D_{\perp}$ . We assume that the flux diffusion time is the shortest time scale so that inductive effects are not important in the equilibrium, i.e., in Eq. (21)  $\partial A_z / \partial t \ll c \partial \hat{\phi} / \partial z$ . The equations then simplify to

$$\frac{\partial n}{\partial t} - D_{\parallel} \frac{\partial^2 n}{\partial z^2} - D_{\perp} \frac{\partial^2 n}{\partial x^2} - \delta_i \frac{c}{B} \frac{\partial}{\partial x} n \frac{\partial \hat{\phi}}{\partial x} = 0, \quad (25)$$

$$\delta_i \frac{c}{B} \frac{v_{in}}{\Omega_i} \frac{\partial}{\partial x} n \frac{\partial \hat{\phi}}{\partial x} + \frac{1}{en_e} \frac{\partial^2 \hat{\phi}}{\partial z^2} + D_{\perp} \frac{\partial^2 n}{\partial x^2} = 0. \quad (26)$$

## 1. CONCLUDING REMARKS

We have presented a relatively simple analysis of the gradient drift instability in barium clouds which includes both finite temperature and finite parallel length effects. We have derived a general set of nonlinear equations which describe the evolution of a three dimensional plasma cloud in the ionosphere. We then investigate the stability of a two dimensional plasma cloud in which the density varies in the direction of the neutral wind and along the ambient magnetic field  $B_0$ . By modeling the density variation along  $B_0$  as a waterbag, we are able to obtain an analytic dispersion relation for the gradient drift instability. There is no static equilibrium since the cloud diffuses both perpendicular and parallel to  $B_0$  so that we restrict our analysis to time scales short compared to the cloud diffusion time scales.

In recent work Sperling et al. (1984) found that the finite length of the cloud suppressed the growth of long wavelength modes because of "good" coupling to the background plasma. However, the growth rate of short wavelength modes is unaffected by the finite size of the cloud since they do not couple to the background in a zero temperature plasma. In contrast to this result, we find that in a finite temperature plasma the short wavelength modes do couple to the background plasma and that for sufficiently short wavelengths the modes are completely stabilized. Stabilization results as the electrons redistribute parallel to  $B_0$  and neutralize the charge imbalance set up by the instability. The parallel electron motion results from diffusion and/or the response to the perturbed fields. We have also derived a simple analytic expression (see (54)) for the wavenumber corresponding to marginal stability (i.e.,  $\gamma = 0$ ). Our results are consistent with previous numerical computations of Sperling and

## V. NUMERICAL RESULTS

We now present quantitative results for the wave frequency of the gradient drift instability by solving (43) numerically. In Fig. 2 we plot  $\gamma/\gamma_0$  (solid curve) and  $\omega_r/\gamma_0$  (dashed curve) vs.  $k_y \rho_s$  where  $\omega = \omega_r + i\gamma$  is the wave frequency. We consider the following typical parameters for a barium cloud at 180 km:  $n_c/n_b = 10.0$ ,  $T_e = T_i = 0.1$  eV,  $m_i = 16 m_p$  ( $O^+$  background),  $z_c = 10$  km,  $B = 0.3$  G,  $\alpha = 10^6$ ,  $D_{\perp} = 100$  m<sup>2</sup>/sec,  $V_n = V_{\text{eff}} = 20$  m/sec and  $L_n = 1$  km. From these values we note that  $c_s = 10^3$  m/sec and  $\rho_s = 6$  m. The main features of Fig. 2 are described as follows. First, the growth rate has a maximum value at  $k_y \rho_s = k_{ym} \rho_s \approx 0.08$ . For  $k_y < k_{ym}$  the growth rate decreases because of coupling to the background plasma as described in Sperling et al. (1984). For  $k_y > k_{ym}$  the growth rate decreases rapidly and becomes stable (i.e.,  $\gamma < 0$ ). This is due to parallel electron motion as described in the preceding section and is the dominant finite temperature effect. Second, the critical wavenumber for stabilization as given by (54) is denoted by the arrow along the  $k_y \rho_s$  axis ( $k_y \rho_s \approx 0.26$ ). It is seen that (54) gives a very good approximation to the critical wavenumber obtained numerically ( $k_y \rho_s \approx 0.28$ ). For the parameters used the critical wavelength is given by  $\lambda_c \approx 135$  m. Third, the real frequency is linear in  $k_y$  and is proportional to the diamagnetic drift velocity  $V_d = (cT/eB)(n_c'/n_c)$ .

The results shown in Fig. 2 appear to be in qualitative agreement with the numerical work of Sperling and Glassman (1984). They consider a plasma cloud with a Gaussian distribution along the magnetic field rather than a waterbag model. The shape of the curve of  $\gamma$  vs  $k_y$  shown in Fig. 2 is similar to corresponding curves presented in their report. Moreover, they also find that  $\omega_r \approx k_y$  for large  $k_y$ .



with

$$\omega_r = \omega_* \left( 1 + \frac{n_c^2}{n_b^2} \frac{2\gamma_0 z_c^2}{\alpha D_1} \right)^{-1} \quad (55)$$

where  $L_n = - (n_c^- / n_c)^{-1}$  and  $\rho_s = c_s / \Omega_i$  is the Larmor radius based on the sound speed  $c_s = (T/m_i)^{1/2}$ . Note from (54) that in the limit  $T \rightarrow 0$ , then  $k_c \rightarrow \infty$  so the mode is not stabilized for a zero temperature plasma [Sperling et al. (1984)]. Equation (54) also appears to imply that  $k_c$  becomes very large as the integrated cloud density  $n_c$  increases. This result is misleading. In a cloud which is finite in the  $y$  direction, the neutral wind velocity  $V_n$  in (54) should be replaced by the effective slip velocity  $V_{\text{eff}}$  of the cloud and the neutral wind. For a circular cloud, the effective velocity is given by [Ossakow and Chaturvedi, 1978]

$$V_{\text{eff}} = 2V_n n_b z_b / (n_b z_b + n_c z_c). \quad (56)$$

In the limit of very large  $n_c z_c$ , Eq. (54) yields the critical wavelength

$$k_c^2 L_n^2 = \frac{4V_n^2 z_b^2}{2D_1^2} \left( 1 + \frac{4z_b^2 V_n^2}{\alpha \rho_s^2 c_s^2} \right), \quad (57)$$

which is independent of  $n_c z_c$ . Thus, short wavelength modes are stable even in large ionospheric clouds.

Two significant features of (52) are first that  $\gamma_0$  does not enter the equation, and second that there appears to be an instability even in the absence of  $\gamma_0$ . However, the unstable root is invalid. In order for  $\phi$  to remain bounded as  $|z|$  becomes large, we require  $\text{Re}(k_b) > 0$ . Thus, for  $\bar{\gamma} = i\omega_*$  we find

$$k_b = (\omega_*/\alpha D_{\perp})^{1/2} e^{i\pi/4}. \quad (53)$$

The root with the opposite sign must be discarded. Inserting this expression for  $k_b$  into (46), we obtain the damped root in (52). Thus, when thermal effects are retained, the gradient-drift mode is stable for large  $k_y$ .

The stabilization mechanism can be understood in terms of our earlier discussion of the local dispersion relation. When  $\omega_*$  is discarded in (45) and (46), the parallel wavevector  $k_b$  remains real (the mode is evanescent in the outer region) and the instability survives even as  $k_y \rightarrow \infty$ . However, when  $\omega_*$  is retained, the parallel wavevector  $k_b$  becomes complex (the mode is evanescent and propagating in the outer region) and the mode is stable for large  $k_y$ . In other words,  $\omega_*$  causes the mode to become oscillatory along  $B_z$  and the electrons can now move along rather than across  $B$  to neutralize the ions; the mode is therefore stable.

To obtain an expression for the critical wavelength  $k_c$  where the mode becomes stable, we take  $\bar{\gamma} = -i\omega_r$  and solve (46) directly. The critical wavelength is given by

$$k_c^2 \frac{\rho_s^2}{c_s^2} = \frac{v_n^2}{c_s^2} \left[ \left( 1 + \frac{2n_c^2 z_c^2}{n_b^2} \frac{v_n}{\alpha D_{\perp} L_n} \right)^2 + \frac{4n_c^2 z_c^2}{n_b^2} \frac{\rho_s^2 c_s^2}{\alpha D_{\perp} L_n^2} \right] \quad (54)$$

$$(\gamma_0 - \bar{\gamma})^2 = \alpha \frac{n_b^2 D_r}{n_c^2 z_c^2} \bar{\gamma}, \quad (47)$$

and is independent of the plasma temperature. For  $z_c^2 \gg \alpha D_r n_b^2 / n_c^2 \gamma_0$ ,

$$\bar{\gamma} = \gamma_0 \quad (48)$$

while in the opposite limit  $z_c^2 \ll \alpha D_r n_b^2 / n_c^2 \gamma_0$ ,

$$\bar{\gamma} = \frac{\gamma_0^2 n_c^2 z_c^2}{\alpha n_b^2 D_r}. \quad (49)$$

These results have been obtained previously [Sperling et al. (1984)].

When  $k_y$  becomes very large,  $k_b^2 = \bar{\gamma} / \alpha D_\perp$  and the dispersion relation becomes

$$(\bar{\gamma} - i\omega_* - \gamma_0)^2 = \alpha D_\perp \frac{n_b^2}{n_c^2 z_c^2} \bar{\gamma}. \quad (50)$$

For large  $k_y$ , we have  $\omega_* \gg \gamma_0$  since  $\omega_* \propto k_y$  and therefore to lowest order

$$\bar{\gamma} = i\omega_*, \quad (51)$$

the mode simply propagates at the diamagnetic frequency. To investigate whether the mode is stable or unstable in this limit, corrections to this result must be calculated. These corrections arise from the right side of (50) and yield the eigenvalue

$$\bar{\gamma} = i\omega_* \pm \frac{n_b}{n_c z_c} (\alpha D_\perp \omega_*^2)^{1/2} e^{i\pi/4}. \quad (52)$$

where

$$\frac{k_b}{k_c} = \left[ \frac{n_b}{n_c} \frac{\bar{\gamma}}{\gamma_0 - \bar{\gamma} + i\omega_*} \right]^{1/2}. \quad (44)$$

In Sperling et al. (1984), the dispersion relation was carefully solved in all possible limits of the various parameters. In order to clearly emphasize the role of finite temperature in stabilizing short wavelength modes, we will focus our analysis primarily on modes with growth rates less than  $\gamma_0$  for which  $k_b/k_c \ll 1$ . In this limit (43) becomes

$$k_c^2 z_c^2 = k_b^2 \quad (45)$$

or

$$n_c z_c (\bar{\gamma} - i\omega_* - \gamma_0) + n_b k_b^{-1} \bar{\gamma} = 0. \quad (46)$$

In the form shown in (46), the nature of the dispersion relation can be simply understood. The first (second) term is the magnetic field line integrated contribution from the cloud (background). The background contribution arises only from the region  $|z| < k_b^{-1}$  since the perturbed potential  $\tilde{\phi}$  is small outside of this interval. The result in (46) can be directly obtained by integrating (33) for  $\tilde{\phi}$  along  $z$ .

The dispersion relation in (45) or (46) can be solved analytically for various values of  $k_y$ . In the limit  $k_y \rightarrow 0$ ,  $k_b^2 = \bar{\gamma}/\alpha D_r$  and the dispersion relation is given by

discontinuity in the density at  $z = \pm z_c$ , we find that  $\tilde{\phi}$  and  $\tilde{A}_z$  must be continuous. For the even  $\tilde{\phi}$  solution ( $\tilde{\phi}_c^2 = 0$ ), we obtain the dispersion equation

$$k_c z_c = \tan^{-1} \left[ \frac{k_b \bar{\gamma} + k_y^2 D_{rc}}{k_c \bar{\gamma} + k_y^2 D_{rb}} \right] + m\pi \quad (42)$$

where  $m$  is an integer. The dispersion relation for the odd  $\tilde{\phi}$  mode ( $\tilde{\phi}_c^1 = 0$ ) is similar to (42) except  $\tan^{-1}$  is replaced by  $-\cot^{-1}$ . The result in (42) is identical to the dispersion relation obtained previously by Sperling et al. (1984) except that the expressions for  $k_b$  and  $k_c$  given in (40d) and (41d) now contain thermal effects which were previously neglected.

In general, the dispersion equation (42) has an infinite number of solutions for a given set of physical parameters, corresponding to eigenmodes with an increasing number of nodes ( $m$ ) along  $z$ . The dispersion equation (42) can be solved numerically for arbitrary values of the background and cloud parameters. However, to gain an understanding of the general scaling of the growth rate  $\gamma$  with the parallel extent of the cloud and temperature, it is useful to solve (42) analytically. To do this we make a number of simplifying assumptions. We consider only the lowest order mode (i.e.,  $m = 0$  which implies  $0 < k_c z_c < \pi/2$ ; it is easily shown that this mode has the largest growth rate); take  $v_{in}/\Omega_i$  to be small so that take all parameters but the density to be the same inside and outside of the cloud; assume  $n_c \gg n_b$ ; and  $v_{ei} \gg v_{en}$  so that the resistive diffusion coefficient is continuous across the boundary  $z = z_c$  (i.e.  $D_{rb} = D_{rc}$ ). With these assumptions the dispersion equation becomes

$$k_c z_c = \tan^{-1}(k_b/k_c) \quad (43)$$

with

$$(\bar{\gamma}_b + k_y^2 D_{rb}) \tilde{A}_{zb} = k_b c \tilde{\phi}_b \quad (40c)$$

$$k_b^2 = \frac{\bar{\gamma}(\bar{\gamma} + k_y^2 D_r)}{\alpha D_r (\bar{\gamma} + k_y^2 D_l)} \Big|_b \quad (40d)$$

where the subscript b on a given parameter indicates that it is to be evaluated in the region  $|z| > z_c$ . Note that we have assumed  $k_b z_b \gg 1$  and that solutions which diverge as  $z$  becomes large have been omitted from (40). This assumption is consistent with the approximation  $z_b \ll \alpha^{1/2} x_c$  made in deriving the equilibrium potential  $\phi$  as long as  $k_b z_b \gg 1$ .

In the region  $|z| < z_c$  the solutions for  $\hat{\phi}$  and  $\tilde{A}_z$  are

$$\tilde{A}_z = \tilde{A}_{zc}^1 \sin(k_c z) - \tilde{A}_{zc}^2 \cos(k_c z) \quad (41a)$$

$$\tilde{\phi} = \tilde{\phi}_c^1 \cos(k_c z) + \tilde{\phi}_c^2 \sin(k_c z) \quad (41b)$$

with

$$(\bar{\gamma}_c + k_y^2 D_{rc}) \tilde{A}_{zc}^{1,2} = k_c c \tilde{\phi}_c^{1,2} \quad (41c)$$

$$k_c^2 = - \frac{(\bar{\gamma} - i\omega_* - \gamma_0)(\bar{\gamma} + k_y^2 D_r)}{\alpha D_r (\bar{\gamma} + k_y^2 D_l)} \Big|_c \quad (41d)$$

We note that in writing (40d) and (41d) we have assumed  $\bar{\gamma} \gg k_y \bar{v}_1$  for simplicity.

To complete the dispersion relation, we must match the various plane wave solutions at  $z = \pm z_c$ . The appropriate matching conditions are obtained from (32) and (33). Integrating these two equations across the

where  $D_{\perp i} = \delta_i (v_{in}/\Omega_i) c T_i / e B$ . In the limit  $k_z = 0$ ,  $\tilde{n}_e = (i k_y c n' / \gamma B) \tilde{\phi}$  and the usual growth rate of the instability is obtained by equating  $\tilde{n}_e$  and  $\tilde{n}_i$ . In this case the density perturbation  $\tilde{n}_e$  and the electric field perturbation,  $\tilde{E}_y = -i k_y \tilde{\phi}$ , are in phase and the usual physical description of instability applies. However, in the limit of large  $k_z$  with  $T_e \neq 0$ ,  $\tilde{n}_e = n_0 e \tilde{\phi} / T_e$ , i.e., the electrons are adiabatic and neutralize the ions by moving along  $\underline{B}$  rather than across  $\underline{B}$ . In this case  $\tilde{n}_e$  is out of phase with  $\tilde{E}_y$  and there is no instability. When  $T_e = 0$  and  $k_z$  is large, the mechanism is somewhat different although the mode is also stable. In this case, the electrons bunch parallel to  $\underline{B}$ , and  $\tilde{n}_e = (k_z^2 / e n_e \gamma) \tilde{\phi}$ . For  $k_z$  large the ions can only neutralize the electrons if  $\gamma = -k_y^2 D_{\perp i}^2 + i k_y v_{in} / \Omega_i$ . Namely, the electrons bunch parallel to  $\underline{B}$  and the ions diffuse across  $\underline{B}$  to neutralize the charge imbalance. Finally, we note that while electron parallel diffusion or ion perpendicular diffusion stabilizes the mode when the perturbation is periodic along  $\underline{B}$ , when the spatial dependence of the mode is exponential, i.e.,  $\tilde{p}(z) \sim \tilde{p} \exp(\pm k_z z)$ , the thermal effects are not stabilizing. The distinction between a periodic and a nonperiodic solution is important in interpreting the dispersion relation which is obtained for the equilibrium shown in Fig. 1b.

We now solve (32) and (33) for the profile given by (19). The boundary conditions used are  $\tilde{\phi}, \tilde{A}_z \rightarrow 0$  as  $|z| \rightarrow z_m$ . For the step profile for  $n(x, z)$  the solutions to (32) and (33) in the region  $|z| > z_c$  can be written as plane waves

$$\tilde{A}_z = \tilde{A}_{zb} \exp[-k_b |z|] \quad (40a)$$

$$\tilde{\phi} = \tilde{\phi}_b \exp[-k_b |z|] \quad (40b)$$

$$\gamma = \left( \gamma_0 - \frac{k_z^2 T}{m_e v_e} \right) \left( \frac{k_y^2 D_r}{k_y^2 D_r + k_z^2 v_A^2 / v_{in} \delta_i} \right) \quad (36)$$

The Alfvén wave has a stabilizing influence on the instability in the electromagnetic limit but thermal effects do not affect the growth rate.

Of course, the gradient drift instability is not stable when  $k_z v_A^2 / v_{in} \delta_i > \gamma_0$  as might be construed from (35) since the expression for  $\gamma$  given in (35) breaks down when  $\gamma < k_y^2 D_r$ . Thermal effects can have a significant influence on the growth rate when  $\gamma < k_z^2 D_r$  (electrostatic limit). Namely, for

$$k_z^2 T / m_e v_e < \gamma_0, \quad (37)$$

the electrons can move a wavelength along  $\underline{B}$  during the growth time of the instability, and the mode is stable. In the limit of  $k_z \rightarrow \infty$  in (36),

$$\gamma = -k_y^2 D_l \quad (38)$$

so that the mode damps at the ion diffusion rate.

The stabilization mechanism can be understood by examining the electron and ion density perturbations in the electrostatic limit,

$$\tilde{n}_e = \frac{(ik_y c n / B + k_z^2 / e n_e)}{\gamma + k_z^2 T_e / m_e v_e} \tilde{\phi} \quad (39a)$$

$$\tilde{n}_i = \frac{[ik_y c n / B - k_y^2 c (v_{in} / \Omega_i) n / B]}{\gamma + k_y^2 D_{li} - ik_y v_n v_{in} / \Omega_i} \tilde{\phi} \quad (39b)$$



In deriving (32) and (33) we have neglected parallel ion diffusion as discussed previously and assumed  $n_c \gg n_b$ . The primes denote a derivative with respect to  $x$ . The important finite temperature effects that appear in (32) and (33) are the diamagnetic drift frequency ( $\omega_*$ ), perpendicular ion diffusion ( $D_\perp$ ), and modification of the equilibrium ( $\hat{\phi}_0$ ). In the limit  $T_i = T_e = 0$ , Eqs. (32) and (33) reduce to those previously derived by Sperling et al. (1984). The eigenvalue  $\bar{\gamma}$  is in a frame of reference moving with the electron fluid.

Prior to solving (32) and (33) for the density profile shown in Fig. 1b, we first consider a cloud of infinite extent [ $z_c + \infty$ ] and Fourier expand in the direction parallel to  $B_0$ , i.e.,  $\tilde{p}(z) \sim \tilde{p} \exp [ik_z z]$ . This allows insight into the influence of finite temperature and parallel dynamics on the instability. The local dispersion equation is given by

$$(\bar{\gamma} - i\omega_* - \gamma_0)(\bar{\gamma} + k_y^2 D_r) = -\frac{k_z^2 V_A^2}{v_{in} \delta_i} (\bar{\gamma} - ik_y \bar{v}_i + k_y^2 D_\perp), \quad (34)$$

where  $V_A = B/(4\pi m_i)^{1/2}$  is the Alfvén velocity. This dispersion relation illustrates the coupling between the gradient drift instability and the Alfvén wave when  $k_z \neq 0$ .

For simplicity, we neglect the terms in (34) which cause the mode to propagate ( $\omega_r$ ,  $k_y \bar{v}_i \rightarrow 0$  and  $\bar{\gamma} \rightarrow \gamma$ ) and note that  $D_r \gg D_\perp$  in ionospheric applications. The growth rate can be easily obtained in two limits: in the electromagnetic limit ( $\gamma \gg k_y^2 D_r$ ),

$$\gamma = \gamma_0 - k_z^2 V_A^2 / v_{in} \delta_i, \quad (35)$$

while in the electrostatic limit ( $\gamma \ll k_y^2 D_r$ )

When the electron parallel conductivity is large,  $k_{\parallel}$  is typically quite small so that these inequalities are easily satisfied. Finally, the time evolution of the equilibrium should have a negligible influence on the stability analysis providing

$$|\gamma\tau_{\parallel}|, |\gamma\tau_{\perp}| \ll 1$$

where  $\gamma$  is the growth (damping) rate of the mode of interest.

#### IV. LINEARIZED EQUATIONS AND DISPERSION EQUATION

To determine the influence of finite temperature and parallel dynamics on the gradient drift instability we linearize (13)-(15) using the equilibrium discussed in the preceding section. We assume perturbed quantities to vary as  $\tilde{p} \sim \tilde{p}(z) \exp(\gamma t + ik_y y)$ . After eliminating  $\tilde{n}$  algebraically from (13)-(15), we obtain two coupled differential equations for  $\tilde{\phi}$  and  $\tilde{A}_z$ :

$$(\bar{\gamma} + k_y^2 D_r) \tilde{A}_z = -c \frac{\partial \tilde{\phi}}{\partial z} \quad (32)$$

$$(\bar{\gamma} - i\omega_* - \gamma_0) \tilde{\phi} = -\alpha D_r (\bar{\gamma} - ik_y \bar{v}_i + k_y^2 D_{\perp}) \frac{1}{c} \frac{\partial \tilde{A}_z}{\partial z} \quad (33)$$

where  $\bar{\gamma} = \gamma + i(c/B)k_y \hat{\phi}'_0$ ,  $\tilde{\phi} = \tilde{\phi} - (T_e/e) \ln \tilde{n}$ ,  $D_r = v_e c^2 / \omega_{pe}^2$ ,  $\bar{v}_i = (v_{in}/\Omega_i) \delta_i \bar{v}_n$ ,  $\alpha = (\Omega_e/v_e)(\Omega_i/v_{in}) \delta_i^{-1}$ ,  $\omega_* = -k_y (cT/eB)(n'_c/n_c)$ ,  $\bar{v}_n = v_n + (c/B)(v_{in}/\Omega_i) \hat{\phi}'_0$ ,  $\gamma_0 = -(n'_c/n_c) \bar{v}_n$ , and

$$\hat{\phi}'_0 = \left( \frac{B}{c} \frac{v_{in}}{\Omega_i} v_n - \frac{T}{e} \frac{n'_c}{n_c} \right) \frac{n_c z_c}{n_c z_c + n_b z_b}.$$

$$\frac{\partial \Delta n}{\partial t} - D_{\parallel} \frac{\partial^2 \Delta n}{\partial z^2} - D_{\perp} \frac{\partial^2 \Delta n}{\partial x^2} - \delta_i \frac{c}{B} \frac{\partial}{\partial x} \Delta n \frac{\partial \bar{\phi}}{\partial x} = 0,$$

$$\frac{1}{e n_e} \frac{\partial^2 \hat{\Delta \phi}}{\partial z^2} = - D_{\perp} \frac{\partial^2 \Delta n}{\partial x^2} - \delta_i \frac{c}{B} \frac{v_i}{\Omega_i} \frac{\partial}{\partial x} \Delta n \frac{\partial \bar{\phi}}{\partial x},$$

where  $\hat{\Delta \phi} \ll \bar{\phi}$ . The equation for  $\Delta n$  has no steady state solution so  $\Delta n$  will generally evolve on a time scale of order  $\tau_{\parallel} \sim \tau_{\perp}$ . The  $z$  dependent potential  $\hat{\Delta \phi}$  drives an equilibrium current  $J_z = n_e^{-1} \partial \hat{\Delta \phi} / \partial z$  which is required so that  $\nabla \cdot \underline{J} = 0$ . The time variation of the equilibrium and the equilibrium parallel current  $J_z$  will be neglected when we carry out the linear stability analysis.

Specifically, in the linear stability analysis we formally take the limit  $D_{\parallel} \rightarrow 0$  and choose

$$n(x, z) = \begin{cases} n_c(x) + n_b & |z| < z_c \\ n_b & z_b > |z| > z_c \end{cases} \quad (31)$$

as shown in Fig. 1b. The perpendicular diffusion coefficient is retained in the stability analysis. The step function model for the density is valid as long as the parallel wavelengths of the modes of interest are much longer than the actual parallel equilibrium density scale length, i.e.,

$$k_{\parallel}^2 z_c^2 \ll 1.$$

The neglect of the equilibrium parallel current and parallel diffusion can similarly be justified for

$$k_{\parallel} J_z / n_e \ll \omega$$

$$k_{\parallel}^2 D_{\parallel} \ll \omega,$$

Equation (25) governs the time evolution of  $n$  while (26) determines  $\hat{\phi}$ . For  $z_b^2 \ll x_c^2 \Omega_e \Omega_i / v_e v_{in}$ , the second term in (26) is much larger than the remaining terms unless  $\partial \hat{\phi} / \partial z = 0$ . Thus,  $\hat{\phi}$  is nearly constant along  $B_0$ . An equation for  $\bar{\phi}$  can then be obtained by averaging (25) and (26) along  $B_0$ . The resulting equations are

$$\partial \bar{n} / \partial t = 0 \quad (27)$$

$$\frac{\partial}{\partial x} \bar{n} \frac{\partial \bar{\phi}}{\partial x} + \frac{T}{e} \frac{\partial^2 \bar{n}}{\partial x^2} = 0 \quad (28)$$

where the average  $\bar{p}$  of a function  $p(z)$  is defined as

$$\bar{p} = \int_{-z_b}^{z_b} dz p(z) / 2z_b. \quad (29)$$

Thus, the integrated density  $\bar{n}$  is a constant in time. From (28), we obtain an expression for the potential,

$$\phi = \frac{T_e}{e} \ln \left( \frac{n}{n_b} \right) - \frac{T_i}{e} \ln \left( \frac{\bar{n}}{n_b} \right) + \frac{B}{c} \frac{v_{in}}{\Omega_i} v_n \int_{-\infty}^x dx \left( 1 - \frac{n_b}{n} \right). \quad (30)$$

The first and second terms in (30) compete in forcing the potential to balance the parallel electron pressure and ion perpendicular pressure, respectively. In the limit in which  $n$  is constant along  $B_0$ ,  $\bar{n} = n$  and  $\phi \sim - (T_i/e) \ln (n/n_b)$  so that  $\phi$  balances the ion pressure. In opposite limit, where  $z_b/z_c \rightarrow \infty$ ,  $\bar{n} = n_b$  and  $\phi \sim (T_e/e) \ln (n/n_b)$  so  $\phi$  balances the electron pressure.

Finally, subtracting the field line average of (25) and (26) from itself, we obtain an equation for  $\Delta n = n - \bar{n}$  and  $\Delta \hat{\phi} = \hat{\phi} - \bar{\phi}$ ,

Glassman (1984) who also noted the importance of finite temperature on the stability of the gradient drift mode.

The extent to which the work described here may have relevance to the barium cloud "freezing" phenomenon deserves some discussion. Briefly, it is observed that barium ion clouds released at ionospheric altitudes will, through the nonlinear evolution of the gradient drift instability, break into smaller pieces. This process, known as bifurcation, continues until a certain minimum transverse dimension is reached, at which point further bifurcation ceases. This frozen scale size is observed to be approximately 400 m. Simple two dimensional models of barium cloud evolution fail to explain frozen scale sizes this large [McDonald et al, 1981]. One explanation which has been advanced to explain freezing is that of end-shortening [Francis and Perkins, 1975; Zalesak et al., 1984] which takes into account the distribution of plasma along magnetic field lines, finite plasma temperature, and parallel currents, as we have done here. However end-shortening depends on the "mapping" of transverse electric fields over relatively large distances along  $B$ ; this becomes increasingly more difficult as  $k$  increases. Indeed, as shown here, and in Sperling et al. (1984), end-shortening fails to provide stabilization of the very high  $k$  modes (at least for the simple profiles of  $n_c$ ,  $n_b$  and  $v_{in}$  considered). If nothing else, the present work shows the existence of a viable stabilization mechanism (i.e., finite temperature) in the gap missed by the end-shortening mechanism. It should also be pointed out, however, that we are not that far removed (135 m stabilization wavelength) from offering a mechanism that by itself may stabilize a 400 m diameter barium striation.

Given the uncertainties in applying a linear theory to explain a complex nonlinear phenomena, the stabilization of the gradient drift instability by parallel electron diffusion must be considered as a strong candidate to explain striation freezing.

ACKNOWLEDGMENT

This research has been supported by the Defense Nuclear Agency.




Fig. 1 Slab geometry and plasma model used in the analysis.

(a) Geometry and plasma configuration.

(b) Cloud and background density profile.




Fig. 2 Plot of  $\omega_r/\omega_0$  (dashed curve) and  $\gamma/\gamma_0$  (solid curve) vs.  $k_y \rho_s$  where  $\omega = \omega_r + i\gamma$ . Parameters used are described in text.



## REFERENCES

- Chaturvedi, P.K., and S.L. Ossakow, Nonlinear stabilization of the  $E \times B$  gradient drift instability in ionospheric plasma clouds, J. Geophys. Res., 84, 419, 1979.
- Francis, S.H., and F.W. Perkins, Determination of striation scale sizes for plasma clouds in the ionosphere, J. Geophys. Res., 80, 3111, 1975.
- Goldman, S.R., L. Baker, S.L. Ossakow, and A.J. Scannapieco, Striation formation associated with barium clouds in an inhomogeneous ionosphere, J. Geophys. Res., 81, 5097, 1976.
- Hoh, F.C., Instability of Penning-type discharges, Phys. Fluids, 6, 1184, 1963.
- Huba, J.D., S.L. Ossakow, P. Satyanarayana, and P.N. Guzdar, Linear theory of the  $E \times B$  instability with an inhomogeneous electric field, J. Geophys. Res., 88, 425, 1983.
- Keskinen, M.J., S.L. Ossakow, and P.K. Chaturvedi, Preliminary report of numerical simulations of intermediate wavelength  $E \times B$  gradient drift instability in ionospheric plasma clouds, J. Geophys. Res., 85, 3485, 1980.
- Linson, L.M., and J.B. Workman, Formation of striations in ionospheric plasma clouds, J. Geophys. Res., 75, 321, 1970.
- McDonald, B.E., M.J. Keskinen, S.L. Ossakow, and S.T. Zalesak, Computer simulation of gradient drift instability processes in Operation Avesfia, J. Geophys. Res., 85, 2143, 1980.
- McDonald, B.E., S.L. Ossakow, S.T. Zalesak, and N.J. Zabusky, Scale sizes and lifetimes of F region plasma cloud striations as determined by the condition of marginal stability, J. Geophys. Res., 86, 5775, 1981.

- Ossakow, S.L. and P.K. Chaturvedi, Morphological studies of rising equatorial spread F bubbles, J. Geophys. Res., 83, 2085, 1978.
- Perkins, F.W., and J.H. Doles III, Velocity shear and the  $E \times B$  instability, J. Geophys. Res., 80, 211, 1975.
- Perkins, F.W., N.J. Zabusky, and J.H. Doles III, Deformation and striation of plasma clouds in the ionosphere, 1, J. Geophys. Res., 78, 697, 1973.
- Scannapieco, A.J., S.L. Ossakow, S.R. Goldman, and J.M. Pierre, Plasma cloud late time striation spectra, J. Geophys. Res., 81, 6037, 1976.
- Shiau, J.N., and A. Simon, Onset of striations in barium clouds, Phys. Rev. Lett., 29, 1664, 1972.
- Simon, A., Instability of a partially ionized plasma in crossed electric and magnetic fields, Phys. Fluids, 6, 382, 1963.
- Sperling, J.L., and A.J. Glassman, Striation eigenmodes along the geomagnetic field and eigenvalues in the limit of strong ion-neutral collisions, JAYCOR Report J530-83-135, JAYCOR, San Diego, CA, 1983.
- Sperling, J.L., J.F. Drake, S.T. Zalesak, and J.D. Huba, "The role of finite parallel length on the stability of barium clouds," J. Geophys. Res., 89, 10913, 1984.
- Volk, H.J., and G. Haerendel, Striation in ionospheric clouds, J. Geophys. Res., 76, 454, 1971.
- Zabusky, N.J., J.H. Doles III, and F.W. Perkins, Deformation and striation of plasma clouds in the ionosphere, 2, Numerical simulation of a nonlinear two-dimensional model, J. Geophys. Res., 78, 711, 1973.

Zalesak, S.T., P.K. Chaturvedi, S.L. Ossakow, and J.A. Fedder, Finite temperature effects on the evolution of ionospheric barium clouds in the presence of a conducting background ionosphere I. A high altitude incompressible background ionosphere, Memo Report 5346, Naval Res. Lab., Washington, D.C., July 1984.

DISTRIBUTION LIST

DEPARTMENT OF DEFENSE

ASSISTANT SECRETARY OF DEFENSE  
COMM, CMD, CONT 7 INTELL  
WASHINGTON, D.C. 20301

DIRECTOR  
COMMAND CONTROL TECHNICAL CENTER  
PENTAGON RM BE 685  
WASHINGTON, D.C. 20301  
01CY ATTN C-650  
01CY ATTN C-312 R. MASON

DIRECTOR  
DEFENSE ADVANCED RSCH PROJ AGENCY  
ARCHITECT BUILDING  
1400 WILSON BLVD.  
ARLINGTON, VA. 22209  
01CY ATTN NUCLEAR  
MONITORING RESEARCH  
01CY ATTN STRATEGIC TECH OFFICE

DEFENSE COMMUNICATION ENGINEER CENTER  
1860 WIEHLE AVENUE  
RESTON, VA. 22090  
01CY ATTN CODE R410  
01CY ATTN CODE R812

DEFENSE TECHNICAL INFORMATION CENTER  
CAMERON STATION  
ALEXANDRIA, VA. 22314  
02CY

DIRECTOR  
DEFENSE NUCLEAR AGENCY  
WASHINGTON, D.C. 20305  
01CY ATTN STVL  
04CY ATTN TITL  
01CY ATTN DDST  
03CY ATTN RAAE

COMMANDER  
FIELD COMMAND  
DEFENSE NUCLEAR AGENCY  
KIRTLAND, AFB, NM 87115  
01CY ATTN FCPR

DEFENSE NUCLEAR AGENCY  
SAO/DNA  
BUILDING 20676  
KIRTLAND AFB, NM 87115  
01CY D.C. THORNBURG

DIRECTOR  
INTERSERVICE NUCLEAR WEAPONS SCHOOL  
KIRTLAND AFB, NM 87115  
01CY ATTN DOCUMENT CONTROL

JOINT CHIEFS OF STAFF  
WASHINGTON, D.C. 20301  
01CY ATTN J-3 WWMCCS EVALUATION  
OFFICE

DIRECTOR  
JOINT STRAT TGT PLANNING STAFF  
OFFUTT AFB  
OMAHA, NB 68113  
01CY ATTN JSTPS/JLKS  
01CY ATTN JPST G. GOETZ

CHIEF  
LIVERMORE DIVISION FLD COMMAND DNA  
DEPARTMENT OF DEFENSE  
LAWRENCE LIVERMORE LABORATORY  
P.O. BOX 808  
LIVERMORE, CA 94550  
01CY ATTN FCPRL

COMMANDANT  
NATO SCHOOL (SHAPE)  
APO NEW YORK 09172  
01CY ATTN U.S. DOCUMENTS OFFICER

UNDER SECY OF DEF FOR RSCH & ENGRG  
DEPARTMENT OF DEFENSE  
WASHINGTON, D.C. 20301  
01CY ATTN STRATEGIC & SPACE  
SYSTEMS (OS)

WWMCCS SYSTEM ENGINEERING ORG  
WASHINGTON, D.C. 20305  
01CY ATTN R. CRAWFORD

COMMANDER/DIRECTOR  
ATMOSPHERIC SCIENCES LABORATORY  
U.S. ARMY ELECTRONICS COMMAND  
WHITE SANDS MISSILE RANGE, NM 88002  
O1CY ATTN DELAS-EO, F. NILES

DIRECTOR  
BMD ADVANCED TECH CTR  
HUNTSVILLE OFFICE  
P.O. BOX 1500  
HUNTSVILLE, AL 35807  
O1CY ATTN ATC-T MELVIN T. CAPPS  
O1CY ATTN ATC-O W. DAVIES  
O1CY ATTN ATC-R DON RUSS

PROGRAM MANAGER  
BMD PROGRAM OFFICE  
5001 EISENHOWER AVENUE  
ALEXANDRIA, VA 22333  
O1CY ATTN DACS-BMT J. SHEA

CHIEF C-E- SERVICES DIVISION  
U.S. ARMY COMMUNICATIONS CMD  
PENTAGON RM 18269  
WASHINGTON, D.C. 20310  
O1CY ATTN C- E-SERVICES DIVISION

COMMANDER  
FRADCOM TECHNICAL SUPPORT ACTIVITY  
DEPARTMENT OF THE ARMY  
FORT MONMOUTH, N.J. 07703  
O1CY ATTN DRSEL-NL-RD H. BENNET  
O1CY ATTN DRSEL-PL-ENV H. BOMKE  
O1CY ATTN J.E. QUIGLEY

COMMANDER  
U.S. ARMY COMM-ELEC ENGRG INSTAL AGY  
FT. HUACHUCA, AZ 85613  
O1CY ATTN CCC-EMEO GEORGE LANE

COMMANDER  
U.S. ARMY FOREIGN SCIENCE & TECH CTR  
220 7TH STREET, NE  
CHARLOTTESVILLE, VA 22901  
O1CY ATTN DRXST-SD

COMMANDER  
U.S. ARMY MATERIAL DEV & READINESS CMD  
5001 EISENHOWER AVENUE  
ALEXANDRIA, VA 22333  
O1CY ATTN DRCLDC J.A. BENDER

COMMANDER  
U.S. ARMY NUCLEAR AND CHEMICAL AGENCY  
7500 BACKLICK ROAD  
BLDG 2073  
SPRINGFIELD, VA 22150  
O1CY ATTN LIBRARY

DIRECTOR  
U.S. ARMY BALLISTIC RESEARCH  
LABORATORY  
ABERDEEN PROVING GROUND, MD 21005  
O1CY ATTN TECH LIBRARY,  
EDWARD BAICY

COMMANDER  
U.S. ARMY SATCOM AGENCY  
FT. MONMOUTH, NJ 07703  
O1CY ATTN DOCUMENT CONTROL

COMMANDER  
U.S. ARMY MISSILE INTELLIGENCE AGENCY  
REDSTONE ARSENAL, AL 35809  
O1CY ATTN JIM GAMBLE

DIRECTOR  
U.S. ARMY TRADOC SYSTEMS ANALYSIS  
ACTIVITY  
WHITE SANDS MISSILE RANGE, NM 88002  
O1CY ATTN ATAA-SA  
O1CY ATTN TCC/F. PAYAN JR.  
O1CY ATTN ATTA-TAC LTC J. HESSE

COMMANDER  
NAVAL ELECTRONIC SYSTEMS COMMAND  
WASHINGTON, D.C. 20360  
O1CY ATTN NAVALEX 034 T. HUGHES  
O1CY ATTN PME 117  
O1CY ATTN PME 117-T  
O1CY ATTN CODE 5011

COMMANDING OFFICER  
NAVAL INTELLIGENCE SUPPORT CTR  
4301 SUITLAND ROAD, BLDG. 5  
WASHINGTON, D.C. 20390  
O1CY ATTN MR. DUBBIN STIC 12  
O1CY ATTN NISC-50  
O1CY ATTN CODE 5404 J. GALET

COMMANDER  
NAVAL OCEAN SYSTEMS CENTER  
SAN DIEGO, CA 92152  
O1CY ATTN J. FERGUSON

NAVAL RESEARCH LABORATORY

WASHINGTON, D.C. 20375  
01CY ATTN CODE 4700 S. L. Ossakow  
26 CYS IF UNCLASS. 1 CY.  
IF CLASS)  
01CY ATTN CODE 4701 I Vitkovitsky  
01CY ATTN CODE 4780 J. Huba (50  
CYS IF UNCLASS, 1 CY IF CLASS)  
01CY ATTN CODE 7500  
01CY ATTN CODE 7550  
01CY ATTN CODE 7580  
01CY ATTN CODE 7551  
01CY ATTN CODE 7555  
01CY ATTN CODE 4730 E. MCLEAN  
01CY ATTN CODE 4108  
01CY ATTN CODE 4730 B. RIPIN  
20CY ATTN CODE 2628

COMMANDER

NAVAL SPACE SURVEILLANCE SYSTEM  
DAHLGREN, VA 22448  
01CY ATTN CAPT J.H. BURTON

OFFICER-IN-CHARGE

NAVAL SURFACE WEAPONS CENTER  
WHITE OAK, SILVER SPRING, MD 20910  
01CY ATTN CODE F31

DIRECTOR

STRATEGIC SYSTEMS PROJECT OFFICE  
DEPARTMENT OF THE NAVY  
WASHINGTON, D.C. 20376  
01CY ATTN NSP-2141  
01CY ATTN NSSP-2722 FRED WIMBERLY

COMMANDER

NAVAL SURFACE WEAPONS CENTER  
DAHLGREN LABORATORY  
DAHLGREN, VA 22448  
01CY ATTN CODE DF-14 R. BUTLER

OFFICER OF NAVAL RESEARCH

ARLINGTON, VA 22217  
01CY ATTN CODE 465  
01CY ATTN CODE 461  
01CY ATTN CODE 402  
01CY ATTN CODE 420  
01CY ATTN CODE 421

COMMANDER

AEROSPACE DEFENSE COMMAND/DC  
DEPARTMENT OF THE AIR FORCE  
ENT AFB, CO 80912  
01CY ATTN DC MR. LONG

COMMANDER

AEROSPACE DEFENSE COMMAND/XPD  
DEPARTMENT OF THE AIR FORCE  
ENT AFB, CO 80912  
01CY ATTN XPDQQ  
01CY ATTN XP

AIR FORCE GEOPHYSICS LABORATORY

HANSCOM AFB, MA 01731  
01CY ATTN OPR HAROLD GARDNER  
01CY ATTN LKB  
KENNETH S.W. CHAMPION  
01CY ATTN OPR ALVA T. STAIR  
01CY ATTN PHD JURGEN BUCHAU  
01CY ATTN PHD JOHN P. MULLEN

AF WEAPONS LABORATORY

KIRTLAND AFT, NM 87117  
01CY ATTN SUL  
01CY ATTN CA ARTHUR H. GUENTHER  
01CY ATTN NTYCE 1LT. G. KRAJEI

AFTAC

PATRICK AFB, FL 32925  
01CY ATTN TN

AIR FORCE AVIONICS LABORATORY

WRIGHT-PATTERSON AFB, OH 45433  
01CY ATTN AAD WADE HUNT  
01CY ATTN AAD ALLEN JOHNSON

DEPUTY CHIEF OF STAFF

RESEARCH, DEVELOPMENT, & ACQ  
DEPARTMENT OF THE AIR FORCE  
WASHINGTON, D.C. 20330  
01CY ATTN AFRDQ

HEADQUARTERS

ELECTRONIC SYSTEMS DIVISION  
DEPARTMENT OF THE AIR FORCE  
HANSCOM AFB, MA 01731  
01CY ATTN J. DEAS

HEADQUARTERS

ELECTRONIC SYSTEMS DIVISION/YSEA  
DEPARTMENT OF THE AIR FORCE  
HANSCOM AFB, MA 01732  
01CY ATTN YSEA

HEADQUARTERS

ELECTRONIC SYSTEMS DIVISION/DC  
DEPARTMENT OF THE AIR FORCE  
HANSCOM AFB, MA 01731  
01CY ATTN DCKC MAJ J.C. CLARK

COMMANDER  
FOREIGN TECHNOLOGY DIVISION, AFSC  
WRIGHT-PATTERSON AFB, OH 45433  
01CY ATTN NICD LIBRARY  
01CY ATTN ETDP B. BALLARD

COMMANDER  
ROME AIR DEVELOPMENT CENTER, AFSC  
GRIFFISS AFB, NY 13441  
01CY ATTN DOC LIBRARY/TSLD  
01CY ATTN OCSE V. COYNE

SAMSO/SZ  
POST OFFICE BOX 92960  
WORLDWAY POSTAL CENTER  
LOS ANGELES, CA 90009  
(SPACE DEFENSE SYSTEMS)  
01CY ATTN SZJ

STRATEGIC AIR COMMAND/XPFS  
OFFUTT AFB, NB 68113  
01CY ATTN ADWATE MAJ BRUCE BAUER  
01CY ATTN NRT  
01CY ATTN DOK CHIEF SCIENTIST

SAMSO/SK  
P.O. BOX 92960  
WORLDWAY POSTAL CENTER  
LOS ANGELES, CA 90009  
01CY ATTN SKA (SPACE COMM SYSTEMS)  
M. CLAVIN

SAMSO/MN  
NORTON AFB, CA 92409  
(MINUTEMAN)  
01CY ATTN MNML

COMMANDER  
ROME AIR DEVELOPMENT CENTER, AFSC  
HANSCOM AFB, MA 01731  
01CY ATTN EEP A. LORENTZEN

DEPARTMENT OF ENERGY  
LIBRARY ROOM G-042  
WASHINGTON, D.C. 20545  
01CY ATTN DOC CON FOR A. LABOWITZ

DEPARTMENT OF ENERGY  
ALBUQUERQUE OPERATIONS OFFICE  
P.O. BOX 5400  
ALBUQUERQUE, NM 87115  
01CY ATTN DOC CON FOR D. SHERWOOD

EG&G, INC.  
LOS ALAMOS DIVISION  
P.O. BOX 809  
LOS ALAMOS, NM 85544  
01CY ATTN DOC CON FOR J. BREEDLOVE

UNIVERSITY OF CALIFORNIA  
LAWRENCE LIVERMORE LABORATORY  
P.O. BOX 808  
LIVERMORE, CA 94550  
01CY ATTN DOC CON FOR TECH INFO  
DEPT  
01CY ATTN DOC CON FOR L-389 R. OTT  
01CY ATTN DOC CON FOR L-31 R. HAGER

LOS ALAMOS NATIONAL LABORATORY  
P.O. BOX 1663  
LOS ALAMOS, NM 87545  
01CY ATTN DOC CON FOR J. WOLCOTT  
01CY ATTN DOC CON FOR R.F. TASCHEK  
01CY ATTN DOC CON FOR E. JONES  
01CY ATTN DOC CON FOR J. MALIK  
01CY ATTN DOC CON FOR R. JEFFRIES  
01CY ATTN DOC CON FOR J. ZINN  
01CY ATTN DOC CON FOR P. KEATON  
01CY ATTN DOC CON FOR D. WESTERVELT  
01CY ATTN D. SAPPENFIELD

SANDIA LABORATORIES  
P.O. BOX 5800  
ALBUQUERQUE, NM 87115  
01CY ATTN DOC CON FOR W. BROWN  
01CY ATTN DOC CON FOR A.  
THORNBROUGH  
01CY ATTN DOC CON FOR T. WRIGHT  
01CY ATTN DOC CON FOR D. DAHLGREN  
01CY ATTN DOC CON FOR 3141  
01CY ATTN DOC CON FOR SPACE PROJECT  
DIV

SANDIA LABORATORIES  
LIVERMORE LABORATORY  
P.O. BOX 969  
LIVERMORE, CA 94550  
01CY ATTN DOC CON FOR B. MURPHEY  
01CY ATTN DOC CON FOR T. COOK

OFFICE OF MILITARY APPLICATION  
DEPARTMENT OF ENERGY  
WASHINGTON, D.C. 20545  
01CY ATTN DOC CON DR. YO SONG

OTHER GOVERNMENT

INSTITUTE FOR TELECOM SCIENCES  
NATIONAL TELECOMMUNICATIONS & INFO  
ADMIN  
BOULDER, CO 80303  
01CY ATTN A. JEAN (UNCLASS ONLY)  
01CY ATTN W. UTLAUT  
01CY ATTN D. CROMBIE  
01CY ATTN L. BERRY

NATIONAL OCEANIC & ATMOSPHERIC ADMIN  
ENVIRONMENTAL RESEARCH LABORATORIES  
DEPARTMENT OF COMMERCE  
BOULDER, CO 80302  
01CY ATTN R. GRUBB  
01CY ATTN AERONOMY LAB G. REID

DEPARTMENT OF DEFENSE CONTRACTORS

AEROSPACE CORPORATION  
P.O. BOX 92957  
LOS ANGELES, CA 90009  
01CY ATTN I. GARFUNKEL  
01CY ATTN T. SALMI  
01CY ATTN V. JOSEPHSON  
01CY ATTN S. BOWER  
01CY ATTN D. OLSEN

ANALYTICAL SYSTEMS ENGINEERING CORP  
5 OLD CONCORD ROAD  
BURLINGTON, MA 01803  
01CY ATTN RADIO SCIENCES

AUSTIN RESEARCH ASSOC., INC.  
1901 RUTLAND DRIVE  
AUSTIN, TX 78758  
01CY ATTN L. SLOAN  
01CY ATTN R. THOMPSON

BERKELEY RESEARCH ASSOCIATES, INC.  
P.O. BOX 983  
BERKELEY, CA 94701  
01CY ATTN J. WORKMAN  
01CY ATTN C. PRETTIE  
01CY ATTN S. BRECHT

BOEING COMPANY, THE  
P.O. BOX 3707  
SEATTLE, WA 98124  
01CY ATTN G. KEISTER  
01CY ATTN D. MURRAY  
01CY ATTN G. HALL  
01CY ATTN J. KENNEY

CHARLES STARK DRAPER LABORATORY, INC.  
555 TECHNOLOGY SQUARE  
CAMBRIDGE, MA 02139  
01CY ATTN D.B. COX  
01CY ATTN J.P. GILMORE

COMSAT LABORATORIES  
LINTHICUM ROAD  
CLARKSBURG, MD 20734  
01CY ATTN G. HYDE

CORNELL UNIVERSITY  
DEPARTMENT OF ELECTRICAL ENGINEERING  
ITHACA, NY 14850  
01CY ATTN D.T. FARLEY, JR.

ELECTROSPACE SYSTEMS, INC.  
BOX 1359  
RICHARDSON, TX 75080  
01CY ATTN H. LOGSTON  
01CY ATTN SECURITY (PAUL PHILLIPS)

EOS TECHNOLOGIES, INC.  
606 Wilshire Blvd.  
Santa Monica, Calif 90401  
01CY ATTN C.B. GABBARD  
01CY ATTN R. LELEVIER

ESL, INC.  
495 JAVA DRIVE  
SUNNYVALE, CA 94086  
01CY ATTN J. ROBERTS  
01CY ATTN JAMES MARSHALL

GENERAL ELECTRIC COMPANY  
SPACE DIVISION  
VALLEY FORGE SPACE CENTER  
GODDARD BLVD KING OF PRUSSIA  
P.O. BOX 8555  
PHILADELPHIA, PA 19101  
01CY ATTN M.H. BORTNER  
SPACE SCI LAB

GENERAL ELECTRIC COMPANY  
P.O. BOX 1122  
SYRACUSE, NY 13201  
01CY ATTN F. REIBERT

GENERAL ELECTRIC TECH SERVICES  
CO., INC.  
HMES  
COURT STREET  
SYRACUSE, NY 13201  
01CY ATTN G. MILLMAN



EOPHYSICAL INSTITUTE  
UNIVERSITY OF ALASKA  
AIRBANKS, AK 99701  
(ALL CLASS ATTN: SECURITY OFFICER)  
01CY ATTN T.N. DAVIS (UNCLASS ONLY)  
01CY ATTN TECHNICAL LIBRARY  
01CY ATTN NEAL BROWN (UNCLASS ONLY)

STATE SYLVANIA, INC.  
ELECTRONICS SYSTEMS GRP-EASTERN DIV  
17 A STREET  
LEEDEDHAM, MA 02194  
01CY ATTN DICK STEINHOF

ISS, INC.  
12 ALFRED CIRCLE  
BEDFORD, MA 01730  
01CY ATTN DONALD HANSEN

ILLINOIS, UNIVERSITY OF  
107 COBLE HALL  
150 DAVENPORT HOUSE  
CHAMPAIGN, IL 61820  
(ALL CORRES ATTN DAN MCCLELLAND)  
01CY ATTN K. YEH

INSTITUTE FOR DEFENSE ANALYSES  
1801 NO. BEAUREGARD STREET  
ALEXANDRIA, VA 22311  
01CY ATTN J.M. AEIN  
01CY ATTN ERNEST BAUER  
01CY ATTN HANS WOLFARD  
01CY ATTN JOEL BENGSTON

INTL TEL & TELEGRAPH CORPORATION  
500 WASHINGTON AVENUE  
MUTLEY, NJ 07110  
01CY ATTN TECHNICAL LIBRARY

JAYCOR  
11011 TORREYANA ROAD  
P.O. BOX 85154  
SAN DIEGO, CA 92138  
01CY ATTN J.L. SPERLING

JOHNS HOPKINS UNIVERSITY  
APPLIED PHYSICS LABORATORY  
JOHNS HOPKINS ROAD  
LAUREL, MD 20810  
01CY ATTN DOCUMENT LIBRARIAN  
01CY ATTN THOMAS POTEMRA  
01CY ATTN JOHN DASSOULAS

KAMAN SCIENCES CORP  
P.O. BOX 7463  
COLORADO SPRINGS, CO 80933  
01CY ATTN T. MEAGHER

KAMAN TEMPO-CENTER FOR ADVANCED  
STUDIES  
816 STATE STREET (P.O. DRAWER QQ)  
SANTA BARBARA, CA 93102  
01CY ATTN DASIAC  
01CY ATTN WARREN S. KNAPP  
01CY ATTN WILLIAM MCNAMARA  
01CY ATTN B. GAMBILL

LINKABIT CORP  
10453 ROSELLE  
SAN DIEGO, CA 92121  
01CY ATTN IRWIN JACOBS

LOCKHEED MISSILES & SPACE CO., INC  
P.O. BOX 504  
SUNNYVALE, CA 94088  
01CY ATTN DEPT 60-12  
01CY ATTN D.R. CHURCHILL

LOCKHEED MISSILES & SPACE CO., INC.  
3251 HANOVER STREET  
PALO ALTO, CA 94304  
01CY ATTN MARTIN WALT DEPT 52-12  
01CY ATTN W.L. IMHOF DEPT 52-12  
01CY ATTN RICHARD G. JOHNSON  
DEPT 52-12  
01CY ATTN J.B. CLADIS DEPT 52-12

MARTIN MARIETTA CORP  
ORLANDO DIVISION  
P.O. BOX 5837  
ORLANDO, FL 32805  
01CY ATTN R. HEFFNER

M.I.T. LINCOLN LABORATORY  
P.O. BOX 73  
LEXINGTON, MA 02173  
01CY ATTN DAVID M. TOWLE  
01CY ATTN L. LOUGHLIN  
01CY ATTN D. CLARK

ONNEL DOUGLAS CORPORATION  
11 BOLSA AVENUE  
MINGTON BEACH, CA 92647  
01CY ATTN N. HARRIS  
01CY ATTN J. MOULE  
01CY ATTN GEORGE MROZ  
01CY ATTN W. OLSON  
01CY ATTN R.W. HALPRIN  
01CY ATTN TECHNICAL  
LIBRARY SERVICES

SSION RESEARCH CORPORATION  
5 STATE STREET  
ATA BARBARA, CA 93101  
01CY ATTN P. FISCHER  
01CY ATTN W.F. CREVIER  
01CY ATTN STEVEN L. GUTSCHE  
01CY ATTN R. BOGUSCH  
01CY ATTN R. HENDRICK  
01CY ATTN RALPH KILB  
01CY ATTN DAVE SOWLE  
01CY ATTN F. FAJEN  
01CY ATTN M. SCHEIBE  
01CY ATTN CONRAD L. LONGMIRE  
01CY ATTN B. WHITE  
01CY ATTN R. STAGAT

SSION RESEARCH CORP.  
20 RANDOLPH ROAD, S.E.  
BUQUERQUE, NEW MEXICO 87106  
01CY R. STELLINGWERF  
01CY M. ALME  
01CY L. WRIGHT

ITRE CORPORATION, THE  
.O. BOX 208  
EDFORD, MA 01730  
01CY ATTN JOHN MORGANSTERN  
01CY ATTN G. HARDING  
01CY ATTN C.E. CALLAHAN

ITRE CORP  
ESTGATE RESEARCH PARK  
320 DOLLY MADISON BLVD  
CLEAN, VA 22101  
01CY ATTN W. HALL  
01CY ATTN W. FOSTER

ACIFIC-SIERRA RESEARCH CORP  
2340 SANTA MONICA BLVD.  
OS ANGELES, CA 90025  
01CY ATTN E.C. FIELD, JR.

PENNSYLVANIA STATE UNIVERSITY  
IONOSPHERE RESEARCH LAB  
318 ELECTRICAL ENGINEERING EAST  
UNIVERSITY PARK, PA 16802  
(NO CLASS TO THIS ADDRESS)  
01CY ATTN IONOSPHERIC RESEARCH LAB

PHOTOMETRICS, INC.  
4 ARROW DRIVE  
WOBURN, MA 01801  
01CY ATTN IRVING L. KOFSKY

PHYSICAL DYNAMICS, INC.  
P.O. BOX 3027  
BELLEVUE, WA 98009  
01CY ATTN E.J. FREMOUW

PHYSICAL DYNAMICS, INC.  
P.O. BOX 10367  
OAKLAND, CA 94610  
ATTN A. THOMSON

R & D ASSOCIATES  
P.O. BOX 9695  
MARINA DEL REY, CA 90291  
01CY ATTN FORREST GILMORE  
01CY ATTN WILLIAM B. WRIGHT, JR.  
01CY ATTN WILLIAM J. KARZAS  
01CY ATTN H. ORY  
01CY ATTN C. MACDONALD  
01CY ATTN R. TURCO  
01CY ATTN L. DeRAND  
01CY ATTN W. TSAI

RAND CORPORATION, THE  
1700 MAIN STREET  
SANTA MONICA, CA 90406  
01CY ATTN CULLEN CRAIN  
01CY ATTN ED BEDROZIAN

RAYTHEON CO.  
528 BOSTON POST ROAD  
SUDBURY, MA 01776  
01CY ATTN BARBARA ADAMS

RIVERSIDE RESEARCH INSTITUTE  
330 WEST 42nd STREET  
NEW YORK, NY 10036  
01CY ATTN VINCE TRAPANI

SCIENCE APPLICATIONS, INC.  
1150 PROSPECT PLAZA  
LA JOLLA, CA 92037

01CY ATTN LEWIS M. LINSON  
01CY ATTN DANIEL A. HAMLIN  
01CY ATTN E. FRIEMAN  
01CY ATTN E.A. STRAKER  
01CY ATTN CURTIS A. SMITH

SCIENCE APPLICATIONS, INC  
1710 GOODRIDGE DR.  
MCLEAN, VA 22102

01CY J. COCKAYNE  
01CY E. HYMAN

SRI INTERNATIONAL  
333 RAVENSWOOD AVENUE  
MENLO PARK, CA 94025

01CY ATTN J. CASPER  
01CY ATTN DONALD NEILSON  
01CY ATTN ALAN BURNS  
01CY ATTN G. SMITH  
01CY ATTN R. TSUNODA  
01CY ATTN DAVID A. JOHNSON  
01CY ATTN WALTER G. CHESNUT  
01CY ATTN CHARLES L. RINO  
01CY ATTN WALTER JAYE  
01CY ATTN J. VICKREY  
01CY ATTN RAY L. LEADABRAND  
01CY ATTN G. CARPENTER  
01CY ATTN G. PRICE  
01CY ATTN R. LIVINGSTON  
01CY ATTN V. GONZALES  
01CY ATTN D. MCDANIEL

TECHNOLOGY INTERNATIONAL CORP  
75 WIGGINS AVENUE  
BEDFORD, MA 01730  
01CY ATTN W.P. BOQUIST

TOYON RESEARCH CO.  
P.O. Box 6890  
SANTA BARBARA, CA 93111  
01CY ATTN JOHN ISE, JR.  
01CY ATTN JOEL GARBARINO

TRW DEFENSE & SPACE SYS GROUP  
ONE SPACE PARK  
REDONDO BEACH, CA 90278  
01CY ATTN R. K. PLEBUCH  
01CY ATTN S. ALTSCHULER  
01CY ATTN D. DEE  
01CY ATTN D/ STOCKWELL  
SNTF/1575

VISIDYNE  
SOUTH BEDFORD STREET  
BURLINGTON, MASS 01803  
01CY ATTN W. REIDY  
01CY ATTN J. CARPENTER  
01CY ATTN C. HUMPHREY

UNIVERSITY OF PITTSBURGH  
PITTSBURGH, PA 15213  
01CY ATTN: N. ZABUSKY

DIRECTOR OF RESEARCH  
U.S. NAVAL ACADEMY  
ANNAPOLIS, MD 21402  
02CY

**DEPARTMENT OF THE NAVY**

**NAVAL RESEARCH LABORATORY**  
Washington, D.C. 20375-5000

**OFFICIAL BUSINESS**  
PENALTY FOR PRIVATE USE, \$300

**POSTAGE AND FEES PAID**  
**DEPARTMENT OF THE NAVY**  
DoD-316  
**THIRD CLASS MAIL**



**END**

**FILMED**

**8-85**

**DTIC**