



MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS-1963-A

a series and the series of the s



Information in Censored Models

1.1

by

Myles Hollander, Frank Proschan and James Sconing

FSU-Statistics Report M701, 71

AFOSR Technical Report No. 85-177

June, 1985

The Florida State University The Department of Statistics Tallahassee, Florida 32306

F.49620-85-C-0007

Research sponsored/by the Air Force Office of Scientific Research, AFSC, USAF, under Grant-AFCEN-95-5-5567. The U.S. Government is authorized to reproduce and distribute reprints for Governmental purposes notwithstanding any copyright notation thereon.

AMS (1981) Subject Classification: Primary 62B10; secondary 62B15

Key words and Phrases: Information, randomly censored model, majorization.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC) NOTICE OF INTEGRATION TO LETER This technic and a science approved Distribute a science to to MATTHEW J. KERTLE Chief, Technical Information Division

AF ACI

ELECTE

AUG 2 9 1985

G

DISTRIBUTION STATEMENT A Approved for public release

Distribution Unlimited

2.

الاجربيين ويستجاب فيرتجد والمستخفي وتوجيعا الموجوب فيتقابل والأراز		· AD -	A158	644		
	REPORT DOCUM	ENTATION PAGE				
14. REPORT SECURITY CLASSIFICATION		16. RESTRICTIVE MARKINGS				
UNCIASSITIED 28. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT				
		Approved for public release; distribution				
b. DECLASSIFICATION/DOWNGRADING SCHEDULE		unlimited				
. PERFORMING ORGANIZATION REPORT NUMBER(S)		5. MONITORING ORGANIZATION REPORT NUMBER(S)				
		AFOSR	-TR- 8	5 - 0 6 1	7	
NAME OF PERFORMING ORGANIZATION	66. OFFICE SYMBOL	78. NAME OF MONIT	ORING ORGANI	ZATION		
Elonida Stato University		AFOCD				
<u>Florida State University</u>		AFOSR 7b. ADDRESS (City, State and ZIP Code)				
Tallahassee, FL 32306		B1dg. 410				
		Bolling AFB	3, D.C. 203	32-6448		
B. NAME OF FUNDING/SPONSORING Bb. OFFICE SYMBOL		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER				
AFOSR	NM	F49620-85-0	-0007	,		
. ADDRESS (City, State and ZIP Code)		10. SOURCE OF FUNDING NOS.				
B1dg. 410		PROGRAM ELEMENT NO	PROJECT	TASK NO.	WORK UNIT	
Bolling AFB, D.C. 20332-6448		61102F	2204	Λ.Ε.		
1. TITLE (Include Security Classification) INTOY	mation in	01102r	2304	CA		
2. PERSONAL AUTHORIS	- <u></u>					
Myles Hollander, Frank Prosch	nan, and James S	Sconing				
JA TYPE OF REPORT 13b. TIME COVERED Technical FROM TO		14. DATE OF REPORT (Yr., Mo., Day) 15. PAGE COUNT June 1985 20				
S. SUPPLEMENTARY NOTATION						
7. COSATI CODES	18. SUBJECT TERMS (Continue on reverse if ne	cessary and identi	ly by block numbe	r)	
FIELD GROUP SUB. GR.	Information,	, randomly cens	ored model,	majorizati	on	
		,				
. ABSTRACT (Continue on reverse if necessary and	d identify by block numb	er;				
Customer and developed for	e include an ex	ation in the r	andomly rig nnon's entr	nt-censored	easures	
-Criteria are developed for me Measures which are appropriat		UCHSION OF SHA	A Alex was an	cound coop	ic	
-Criteria are developed for me Measures which are appropriat are seen to satisfy some fund	lamental theorem	including (i) the uncen	soreu case	13	
-Criteria are developed for me Measures which are appropriat are seen to satisfy some fund always at least as invormativ ing increases stochastically.	lamental theorem e as any censor and (iii) the	red model, (ii)	informatio	n decreases	as censor	
-Criteria are developed for me Measures which are appropriat are seen to satisfy some fund always at least as invormativ ing increases stochastically,	lamental theorem re as any censor and (iii) the	ns including (i red model, (ii) information ga	informatio	n decreases nally decre	as censor	
Criteria are developed for me Measures which are appropriat are seen to satisfy some fund always at least as invormativ ing increases stochastically,	lamental theorem re as any censor and (iii) the	information ga	information in is marging Satistica	n decreases nally decre	as censor	
Criteria are developed for me Measures which are appropriat are seen to satisfy some fund always at least as invormativ ing increases stochastically,	lamental theorem we as any censor and (iii) the	information ga	informatio in is margi Sataluca	n decreases nally decre	as censor- asing.	
-Criteria are developed for me Measures which are appropriat are seen to satisfy some fund always at least as invormativ ing increases stochastically,	lamental theorem re as any censor and (iii) the	including (i red model, (ii) information ga	informatio in is margi Satatia	n decreases nally decre	as censor asing.	
Criteria are developed for me Measures which are appropriat are seen to satisfy some fund always at least as invormativ ing increases stochastically,	lamental theorem re as any censor and (iii) the	information ga	informatio in is margi Satistica	n decreases nally decre	as censor asing.	
Criteria are developed for me Measures which are appropriat are seen to satisfy some fund always at least as invormativ ing increases stochastically,	lamental theorem re as any censor and (iii) the	including (i red model, (ii) information ga	informatio in is margi Sataluca	n decreases nally decre	as censor asing.	
Criteria are developed for me Measures which are appropriat are seen to satisfy some fund always at least as invormativ ing increases stochastically,	lamental theorem re as any censor and (iii) the	21. ABSTRACT SEC) the uncen informatio in is margi Satistica	n decreases nally decre	as censor	
Criteria are developed for me Measures which are appropriat are seen to satisfy some fund always at least as invormativ ing increases stochastically,	CT	21. ABSTRACT SECULUTION) the uncen informatio in is margi Satistica URITY CLASSIFIC ed	n decreases nally decre	as censor	
Criteria are developed for me Measures which are appropriat are seen to satisfy some func always at least as invormativ ing increases stochastically,	CT	21. ABSTRACT SECULUTION Unclassifi) the uncen informatio in is margi Salishia URITY CLASSIFIC ed UMBER	CATION	ABOL	
Criteria are developed for me Measures which are appropriat are seen to satisfy some func always at least as invormativ ing increases stochastically,	CT	21. ABSTRACT SECU Unclassifi 22b. TELEPHONE N (202)767-50) the uncen informatio in is margi Satalica UNITY CLASSIFIC ed UMBER MER 27	CATION	ABOL	
 Criteria are developed for me Measures which are appropriat are seen to satisfy some func always at least as invormativ ing increases stochastically, DISTRIBUTION/AVAILABILITY OF ABSTRACE DISTRIBUTION/AVAILABILITY OF ABSTRACE NCLASSIFIED/UNLIMITED SAME AS APT. NAME OF RESPONSIBLE INDIVIDUAL Brian W. Woodruff, Maj, USAF D FORM 1473 R3 APR 	CT EDITION OF 1 JAN 27	21. ABSTRACT SECU Unclassifi 22b. TELEPHONE N (Include Area Co (202)767-50) the uncen informatio in is margi Satatua URITY CLASSIFIC ed UMBER Sde) 27	CATION 22c. OFFICE SYN NM Unclassi	ABOL	

Information in Censored Models

by

Myles Hollander, Frank Proschan, and James Sconing Florida State University

Abstract

Criteria are developed for measuring information in the randomly rightcensored model. Measures which are appropriate include an extension of Shannon's entropy. The measures are seen to satisfy some fundamental theorems including (i) the uncensored case is always at least as informative as any censored model, (ii) information decreases as censoring increases stochastically, and (iii) the information gain is marginally decreasing.

Accession For NTIS GRA&I DTIC TAB Unannounsed Justification By_ Distribution/ Availability Codes Avail and/or Dist Special

1. Introduction.

Let X_1, X_2, \ldots, X_n be independent and identically distributed positive random variables corresponding to the true lifetimes of some items on test. With every X_i there is a corresponding Y_i , independent of X_i . The Y_i 's are also independent and identically distributed on the positive real line. Y_i is said to be the <u>censoring variable</u>. The observations consist of the iid pairs (Z_i, δ_i) , $i = 1, \ldots, n$, where $Z_i = \min(X_i, Y_i)$, $\delta_i = I(X \le Y)$, and I(A) denotes the indicator function for the set A. This is the randomly right-censored model. Typically the goal is to make inferences about some property of the distribution of X. The censoring variable can be thought of as a confounding variable which inhibits the ability to see X. Suppose it is desired to compare experiments where different types of censoring may take place and then decide which experiment is preferred. One approach is to use the "information" in the experiment as a basis for decision.

The term information was first used by Fisher (1925) to describe the efficiency of an estimator of some parametric component of the unknown distribution function.

A more common usage of the term is in the field of communication theory pioneered by Shannon (1948). Shannon's information can be viewed as a measure of uncertainty as to the outcome of a random variable. In our paper Shannon's measure is extended to provide a comparison of experiments in the censored model. In extending Shannon's measure to the censored case and developing other suitable measures of information, we find that the notion that more censoring should yield less information is fundamental. This property should hold for any satisfactory measure of information.

The property of decreasing information as censoring increases has also been

studied by Lindley (1956), Brooks (1982), and Barlow and Hsiung (1983), all in the Bayesian context where information is given in terms of expected risk. The connection between our approach and the Bayesian approach is given by Bernardo (1979).

Some satisfactory notions of information are developed in Sections 2, 3, and 4. Each information measure advanced is shown to satisfy Theorems 1.1 and 1.2 below.

Theorem 1.1. E[Information (X)] \geq E[Information (Z, δ)] for every X and Y.

<u>Theorem 1.2</u>. E[Information (Z_1, δ_1)] $\leq E$ [Information (Z_2, δ_2)] for every X, where (Z_i, δ_i) is the censored variable associated with Y_i , i = 1, 2, and $Y_1 \stackrel{\text{st}}{\leq} Y_2$.

The form of the information measure is utilized in the proof. In the discrete case, information takes the form $E_{v}[G]$ where

G(i) = Information (X = j | Y = i, i > j) + Information (X > i | Y = i).

The first term represents the information in observing the X variable directly. The second term is the "partial" information in observing only that X is larger than the observed variable Y. Information is given by taking the expectation over the Y variable. Theorems 1.1 and 1.2 are proved by first showing G(i)< G(i+1), for every i > 0. This says that information is increased if the experiment is observed for the additional time from i to i+1. With this preliminary lemma the theorems follow directly.

While information increases as censoring decreases there are limits to this increase. Barlow and Hsiung (]983) state "it would be interesting to see when this (information) gain is marginally decreasing." This leads to the following theorem.

<u>Theorem 1.3</u>. Let $X^{(i)}$ be the lifetime variable which is censored deterministically at time i (Type I censoring). Then for sufficiently large i, E[Information $X^{(i)}$] is a concave increasing function of i.

In Sections 2, 3, and 4 various regularity conditions are imposed to obtain versions of Theorem 1.3 for the particular information measures considered.

In Section 2 Shannon's original measure, entropy, is defined and extended to the censored case and the three fundamental theorems are proved. In Section 3 a more general class of measures is developed based on the theory of majorization. Once again the three basic theorems are proved. In Section 4 Shannon's measure is shown to be inadequate in the continuous case. Several measures are developed based on the variance of the lifetime variable X and the fundamental theorems are proved.

2. Information in the discrete case.

Shannon (1948) axiomatically derived an information measure which satisfies some intuitive requirements. Suppose a variable X takes on only two values with probabilities p and 1 - p. If an information measure is denoted H(p) (or H(X)) then it should satisfy the following requirements:

(i) $H(p) \ge 0$ for all p, $0 \le p \le 1$,

(ii) $H(\frac{1}{2}) = 1$, H(1) = H(0) = 0,

(iii) H(X, Y) = H(X|Y) + H(Y) for all (X, Y),

where H(X, Y) is the information in the joint experiment (X, Y) and H(X|Y) is the conditional information in the experiment X, given the outcome of experiment Y. Imposing (i) - (iii) leads to the definition of information as

(2.1)
$$H(\underline{p}) = -\sum_{i=1}^{n} p_i \log_2 p_i,$$

- 3 -

where $0\log_2 0 \stackrel{\text{def}}{=} 0$ and $\underline{p} = (p_1, p_2, \dots, p_n)$ with $P(X = i) = p_i$. The choice of the base of the logarithm is unimportant and henceforth will be defined as the base of the natural logarithm. The definition can also be extended to the case where X has a countably infinite number of support points. This measure is termed <u>entropy</u>.

For the censored model where X and Y have discrete distributions, let $p_i = P(X = i), q_i = P(Y = i)$, then extend (2.1) to:

Definition 2.1. The information in the discrete experiment (X, Y) is

(2.2) $H(\underline{p}, \underline{q}) = -\sum_{i=1}^{n} q_{i} \left[\sum_{j \leq i}^{n} p_{j} \log p_{j} + \overline{P}_{i+1} \log \overline{P}_{i+1} \right],$

where $\overline{P}_{i} = \sum_{j \ge i} p_{j}$.

Our definition of information in the discrete censored case can be interpreted as follows. Suppose the censoring variable takes the value i. Then the information in our observed variable Z is full information, $-p_j \log p_j$, if a death occurs prior to the censoring time. Otherwise we receive partial information, $-\overline{P}_{i+1} \log \overline{P}_{i+1}$. Note that if a death and a censorship occur at the same time we say that a death is observed. The definition follows by averaging over the censoring times. It is interesting to note that (2.2) is equivalent to Shannon's mutual information, $H(X) - H(X|Z, \delta)$. To see this write $H(X) - H(X|Z, \delta)$ as $-\sum_{i=1}^{n} \log p_i - \sum_{i=1}^{n} \sum_{j=0}^{n} p_{i|j} \log p_{i|j}$ where r_j is the probability that (Z, δ) = $j = (j_1, j_2)$ where $j_1 = 1, 2, \ldots$ and $j_2 = 0, 1$. Also $p_{i|j}$ is the conditional probability that X = i given that (Z, δ) = (j_1, j_2) . The mutual information can be rewritten as $\sum_{i=1}^{n} p_{ij} \log (p_{ij}/p_{irj})$ where p_{ij} is the joint probability that $[X = i, (Z, \delta) = (j_1, j_2)]$. Note that $P[X = i, Z = j, \delta = 0] = p_i q_i$ if i > j,

0 otherwise. Also $P[X=i, Z=j, \delta=1] = p_i \overline{Q}_j$ if i=j, 0 otherwise. Finally $P[Z=j, \delta=0] = q_j \overline{P}_{j+1}$, $P[Z=j, \delta=1] = p_j \overline{Q}_j$. With these probabilities (2.2) follows from straightforward calculations.

Now with this definition for information in the discrete censored case we show that the three basic theorems stated in Section 1 hold.

<u>Theorem 2.2.</u> $H(p) \ge H(p, q)$ for all probability vectors p and q.

The theorem states that any amount of censoring reduces information. In order to prove this we first prove Lemma 2.3. This lemma has appeared in the literature in several different forms. Dobrušin (1963) showed that $H(p) \ge H(f(p))$ with equality if and only if f(p) is a one-to-one function. Khinchin (1957) showed that if $UA_i = A$ and $A_i \cap A_j = \emptyset$, $i \ne j$, then $H(p(A_i)) \ge H(p(A))$, where A_i represents a set of support points for X and $p(A_i) = P(X \in A_i)$. We prove the lemma directly.

Lemma 2.3.
$$-\sum_{j=1}^{j} \log p_{j} \ge -\sum_{j=1}^{j} p_{j} \log p_{j} - \overline{P}_{i+1} \log \overline{P}_{i+1}$$
, for every i.

<u>Proof.</u> Since log x is an increasing function, then $\log \overline{P}_{i+1} \ge \log p_j$, $j \ge i+1$. Hence, $\sum_{j j} \log p_j \le \sum_{j \le i} p_j \log p_j + \sum_{j \ge i} p_j \log \overline{P}_{i+1} = \sum_{j \le i} p_j \log p_j + \overline{P}_{i+1} \log \overline{P}_{i+1}$.

We now proceed with the proof of Theorem 2.2.

Proof of Theorem 2.2. $H(\underline{p}, \underline{q}) \approx -\sum_{i=1}^{n} q_{i} (\sum_{j \leq i}^{p} p_{j} \log p_{j} + \overline{P}_{i+1} \log \overline{P}_{i+1})$ $\leq \sum_{i=1}^{n} q_{i} (-\sum_{j=1}^{p} p_{j} \log p_{j}) = H(\underline{p}) \text{ (from Lemma 2.3.). } \|$ Next we compare amounts of information available in two models with different censoring distributions, one of which is stochastically larger than the other. First we prove the following lemma.

Lemma 2.4. Let
$$G_i = \sum_{j \leq i} p_j \log p_j + \overline{P}_{i+1} \log \overline{P}_{i+1}$$
. Then $G_i \geq G_{i+1}$, for $i = 1, 2, ...$

<u>Proof.</u> $G_i - G_{i+1} = -p_{i+1} \log p_{i+1} + \overline{P}_{i+1} \log \overline{P}_{i+1} - \overline{P}_{i+2} \log \overline{P}_{i+2}$

$$= p_{i+1}(\log \overline{P}_{i+1} - \log p_{i+1}) + \overline{P}_{i+2}(\log \overline{P}_{i+1} - \log \overline{P}_{i+2}) \ge 0,$$

since $\overline{P}_{i+1} \ge p_{i+1}$ and $\overline{P}_{i+1} \ge \overline{P}_{i+2}$.

We are now ready to prove an analogue of Theorem 1.2.

Theorem 2.5. Let $Y_1 \stackrel{\text{st}}{\leq} Y_2$. Let Y_i have outcome probability vector \underline{q}_i , i = 1, 2. Then $H(\underline{p}, \underline{q}_1) \leq H(\underline{p}, \underline{q}_2)$ for every life distribution vector \underline{p} .

<u>Proof.</u> From (2.2) we see that $H(\underline{p}, \underline{q_i}) = \underline{F_Y}_i$ (-G), i = 1, 2, where G is the function defined by $G(i) = G_i$ as in Lemma 2.4. From Lemma 2.4 G_i is increasing (non-decreasing) in i. Thus $E_{Y_1}(-G) \le E_{Y_2}(-G)$.

Thus we see that our intuition has been justified in the simple case where Shannon's entropy is the measure of information. Theorem's 2.2 and 2.5 should represent a sort of "acid test" for the applicability of any measure of information.

The condition of stochastic domination of the censoring variables is also necessary. If stochastic domination does not occur then there exists an interval where $Y_1 \stackrel{st}{\leq} Y_2$ and another interval where $Y_2 \stackrel{st}{\leq} Y_1$. By defining X to have support only on one of these intervals and applying Theorem 2.5, a contradiction arises. In a similar fashion stochastically decreasing the lifetime variable X also yields more information. Note that (2.1) and (2.2) are scale invariant. By relabeling the axis after stochastically decreasing X and then applying Theorem 2.5 we get analogous results.

We now establish a parallel for Theorem 1.3.

<u>Theorem 2.6.</u> If there exists a k such that for all i > k, $p_i > p_{i+1}$, and $\overline{P}_k < e^{-1}$, then $G_i - G_{i+1}$ is decreasing (nonincreasing) in i, i > k.

$$\begin{array}{l} \underline{\operatorname{Proof}} & \text{It is sufficient to show } (\operatorname{G}_{i-1} - \operatorname{G}_{i}) - (\operatorname{G}_{i} - \operatorname{G}_{i+1}) \geq 0. \\ (\operatorname{G}_{i-1} - \operatorname{G}_{i}) - (\operatorname{G}_{i} - \operatorname{G}_{i+1}) &= -\operatorname{p}_{i} \operatorname{logp}_{i} + \operatorname{p}_{i+1} \operatorname{logp}_{i+1} + 2\overline{\operatorname{P}}_{i+1} \operatorname{log\overline{P}}_{i+1} - \overline{\operatorname{P}}_{i} \operatorname{log\overline{P}}_{i} - \overline{\operatorname{P}}_{i+2} \operatorname{log\overline{P}}_{i+2} \\ &\geq 2\overline{\operatorname{P}}_{i+1} \operatorname{log\overline{P}}_{i+1} - \overline{\operatorname{P}}_{i} \operatorname{log\overline{P}}_{i} - \overline{\operatorname{P}}_{i+2} \operatorname{log\overline{P}}_{i+2} \text{ (since } \overline{\operatorname{P}}_{k} < e^{-1}) \\ &\geq 2[\overline{\operatorname{P}}_{i+1} \operatorname{log\overline{P}}_{i+1} - (\frac{1}{2})(\overline{\operatorname{P}}_{i} + \overline{\operatorname{P}}_{i+2}) \operatorname{log}\{(\frac{1}{2})(\overline{\operatorname{P}}_{i} + \overline{\operatorname{P}}_{i+2})\}] \geq 0. \end{array} \right\| \end{array}$$

The conditions of the theorem assure that the index i is far enough in the right-hand tail for the marginally decreasing property to take hold.

3. Majorization and information.

We wish to consider generalizations of Shannon's entropy measure. In particular requirement (iii) of Section 2 is not universally accepted, and it is this requirement which leads to the specific functional form for Shannon's entropy. By relaxing this assumption we can generalize Shannon's measure. Note that the information in an event is governed solely by the probability of that event. Thus information in the event labeled i is given by $f(p_i)$. What types of measures perform satisfactorily as measures of information? To answer this question the theory of the majorization ordering is utilized. Majorization is a powerful tool, useful in proving inequalities. The standard reference on majorization is Marshall and Olkin (1979). First we state some definitions and preliminary theorems.

Definition 3.1. Let $\underline{x}, \underline{y} \in \mathbb{R}^n$, n-dimensional Euclidean space. We say \underline{y} majorizes \underline{x} $(\underline{y} \stackrel{\times}{>} \underline{x})$ if

(1)
$$\sum_{i=1}^{k} y_{[i]} \ge \sum_{i=1}^{k} x_{[i]}, k=1, ..., n-1;$$

and,

(2)
$$\sum_{i=1}^{n} y_{i} = \sum_{i=1}^{n} x_{i}$$
,

where $x_{[1]} \ge x_{[2]} \ge \dots \ge x_{[n]}$, $y_{[1]} \ge y_{[2]} \ge \dots \ge y_{[n]}$ are the decreasing rearrangements for <u>x</u> and <u>y</u> respectively.

An equivalent definition of majorization is given by:

<u>Definition 3.2</u>. Let $\underline{x}, \underline{y} \in \mathbb{R}^n$. Then $\underline{y} \xrightarrow{\mathbb{R}} \underline{x}$ if and only if there exists a doubly stochastic matrix P, such that $\underline{x} = \underline{y}P$.

A matrix is <u>doubly stochastic</u> if each row sum and each column sum equals one. This definition illuminates why majorization is particularly useful in the study of information under censoring. The <u>y</u> vector can be thought of as the probabilities when a censorship has occurred, that is, $(p_1, p_2, \ldots, p_i,$ $\overline{P}_{i+1}, 0, \ldots)$. The <u>x</u> vector is the vector of probabilities of the life distribution (p_1, p_2, \ldots) . Note that

(3.1)
$$(p_1, p_2, \ldots, p_i, \overline{P}_{i+1}, 0, \ldots) \stackrel{w}{>} (p_1, p_2, \ldots)$$

for every i. The doubly stochastic matrix in this case consists of the

conditional probabilities of surviving to time j given that the item was censored at time i, i < j.

Functions which preserve (reverse) the ordering of majorization are called Schur-convex (Schur-concave).

Definition 3.3. A function f: $\mathbb{R}^n \to \mathbb{R}^1$ is said to be <u>Schur-convex</u> (Schur-concave) if $\underline{y} \xrightarrow{\mathbb{R}} \underline{x}$ implies $f(\underline{y}) \ge (\le) f(\underline{x})$.

Theorem 3.4. (Schur, 1923, Ostrowski, 1952). A permutation invariant function ϕ is Schur-convex (concave) if and only if $(z_i - z_j)(\phi_{(i)}(\underline{z}) - \phi_{(j)}(\underline{z})) \ge (\le) 0$, where $\phi_{(i)}(\underline{z})$ is the partial derivative of ϕ with respect to z_i .

It is useful to identify specific types of functions which can represent the average information in a random variable.

<u>Theorem 3.5.</u> (Schur, 1923). Let $\phi(\underline{x}) = \sum_{i=1}^{n} f(x_i)$ where f: $\mathbb{R}^1 + \mathbb{R}^1$. Then ϕ is Schur-convex (concave) if and only if f is a convex (concave) function.

This provides a basis for constructing information measures with a general function f. Let the information in the occurrence of a death at time i be represented by $f(p_i)$. Two possibilities for classes of information measures can be obtained as follows. Define $A = \{f: f(x) \text{ is decreasing and } f(x)/x \text{ is concave}\}$ and $B = \{g: g(x) \text{ is concave and } g(x)/x \text{ decreasing}\}$. Then there are the following two candidates for general information measures,

(3.2)
$$H_{f}(\underline{p}) = \sum_{i} p_{i} f(p_{i}), f \in A,$$

or

$$(3.3) \qquad H_{g}(\underline{p}) = \sum g(\underline{p}_{i}), \ g \in B.$$

Solomon (1979) uses a measure similar to (3.3) to measure ecological diversity. The measure given by (3.2) will be adopted here as it represents the average information in an experiment and it facilitates the proof of Theorem 3.12.

<u>Definition 3.6.</u> Let <u>p</u> be the vector of probabilities associated with a life variable X. Then the "f-type" information measure in X is given by (3.2).

Note that from the definition of the class A, $H_{f}(\underline{p})$ is Schur-concave.

<u>Definition 3.7</u>. Let \underline{p} and \underline{q} be the probability vectors associated with X and Y. Then the amount of information in the censored model is defined to be

(3.4)
$$H_{\mathbf{f}}(\underline{p}, \underline{q}) = \sum_{i=1}^{n} \left[\sum_{j \leq i}^{n} P_{j} f(p_{j}) + \overline{P}_{i+1} f(\overline{P}_{i+1}) \right],$$

where $f \in A$.

Choosing $f(x) = -\log x$ gives (2.2), however (3.4) cannot be obtained as a measure of Shannon's mutual information. Henceforth, $Of(0) \stackrel{\text{def}}{=} 0$ to fix the location.

Lemma 3.8.
$$\sum_{j=1}^{n} f(p_j) \ge \sum_{j \le i}^{n} p_j f(p_j) + \overline{P}_{i+1} f(\overline{P}_{i+1})$$
 for every i.

<u>Proof</u>. This follows immediately from (3.1) and the fact that $H_{f}(\underline{p})$ is Schurconcave.

<u>Theorem 3.9.</u> $H_f(p) \ge H_f(p, q)$ for every p and q.

Proof. The proof is analogous to that of Theorem 2.2.

Lemma 3.10. Let G(f) be a function defined by $G(f(i)) = G_f(i) = \sum_{\substack{j \le i \\ j \le i \le 1}} p_j f(p_j) + \overline{P_{i+1}}f(\overline{P_{i+1}})$. Then $G_i(f) \le G_{i+1}(f)$, i = 1, 2,

<u>Proof.</u> Note that $(p_1, p_2, \dots, p_{i+1}, \overline{P}_{i+2}, 0, \dots)$ $\stackrel{\text{m}}{\leftarrow} (p_1, p_2, \dots, p_i, \overline{P}_{i+1}, 0, \dots)$ and that $H_f(p)$ is Schur-concave. \parallel

<u>Theorem 3.11</u>. Let Y_1 and Y_2 be censoring variables with probability vectors \underline{q}_1 and \underline{q}_2 respectively. Let $Y_1 \stackrel{\text{st}}{\leq} Y_2$. Then for every \underline{p} , $\underline{H}_f(\underline{p}, \underline{q}_1) \leq \underline{H}_f(\underline{p}, \underline{q}_2)$.

Proof. With Lemma 3.10 the proof is analogous to that of Theorem 2.5.

<u>Theorem 3.12</u>. If there exists a k such that for all i > k, $p_i > p_{i+1}$, then $G_i(f)$ is a concave function of i, i > k.

Proof. The proof is similar to the proof in Theorem 2.6.

Thus the "f-type" information measures are suitable for the discrete censored model.

4. Information in the continuous case.

Sec. Carl

Our goal is to extend our definition to include life distributions which are continuous. The obvious analogue of Definition 2.1 would be to define $H(p(x)) = -\int p(x) \log p(x) dx$. However, Example 4.1 shows that such a definition is unsatisfactory.

Example 4.1. Let
$$p(x) = \begin{cases} \lambda e^{-\lambda x}, & 0 < x < \infty, \\ \lambda > 0, \\ 0 & \text{otherwise.} \end{cases}$$

Then $H(p(x)) = -\int_0^\infty \lambda e^{-\lambda x} [-\lambda x + \log \lambda] dx = 1 - \log \lambda$.

From Example 4.1 it is seen that $H(p(x)) \ge 0$ if and only if $\lambda > e$. Thus the base of the logarithm is crucial in determining a key property of information. Furthermore H(p(x)) does not have the scale invariant property present in the discrete case. Finally note that if $\lambda < e$, then $H(p(x)) \le 0$ so that an observation will decrease our knowledge. All these properties run counter to the properties which measures of information should possess. Thus H(p(x)) as defined above is unsatisfactory for defining information.

In order to find a new measure of information, recall the properties Shannon used to define entropy: (i) $H(p) \ge 0$, (ii) $H(\frac{1}{2}, \frac{1}{2}) = 0$ and H(1, 0) = 1 and (iii) H(X, Y) = H(X|Y) + H(Y). The first two requirements simply fix the scale and thus are not crucial. It is the third requirement, the so-called additivity criterion, that is crucial in defining entropy. It is desirable to retain this crucial property in the continuous case. Restricting consideration to functions of the form $\vartheta(X, EX)$, where $\vartheta(\cdot, \cdot)$ is a metric, leads to $H(X) = E(X - EX)^2 = \sigma_X^2$ (Blyth, 1959). This suggests:

<u>Definition 4.2</u>. Let X be a continuous random variable on the positive real line with p.d.f. f(x) and finite variance. Then the information in X is defined to be $H(X) = H(f) = \int_0^\infty (x - \mu)^2 f(x) dx = \sigma_X^2$, where $\mu = \int_0^\infty x f(x) dx$.

Note that information, in any sense, measures the spread of the distribution. From this it seems unreasonable to expect any measure of information to be scaleinvariant in the continuous case. Thus when comparing measures of information care must be taken to use the same scale of measurement. Definition 4.2 gives a measure of information in the uncensored case. Recall that in the discrete case there is full information if death occurs prior to censorship, and only partial information, $-\overline{P}_{i+1}\log\overline{P}_{i+1}$, otherwise. In the case of censoring the only constraint is that the remaining probabilities sum to \overline{P}_{i+1} . Note that among all discrete probability distributions which have probability \overline{P}_{i+1} remaining, the one that gives the least amount of information is that which puts its entire remaining mass at a single point. This would yield information $-\overline{P}_{i+1}\log\overline{P}_{i+1}$. Thus $-\overline{P}_{i+1}\log\overline{P}_{i+1}$ can be viewed as a type of "worst case" under the probability constraint. With this "worst case" type of reasoning for the variance measure, information measures can be developed.

In the continuous case, minimizing information is equivalent to minimizing $\partial(X, EX)$. Given that the censorship takes place at time c, the constraint is that the remaining probability, $\overline{F}(c)$, must be placed in the set $A = \{x: x > c\}$. It is easy to show that if $c \leq EX$, then $\partial(X, EX)$ is minimized by placing all the remaining mass at EX. If c > EX, then $\partial(X, EX)$ is minimized by placing all the mass at c. We now give a definition for information in the continuous censored case.

<u>Definition 4.3</u>. Let X be a lifetime variable with p.d.f. f(x) and finite variance. Let Y be a censoring variable with p.d.f. g(y). Let Z = {min(X, Y), I(X ≤ Y)} be the observed variable. Then the information in Z is defined to be:

 $H^{(1)}(X, Y) = H^{(1)}(f, g) = \int_0^\infty g(c) \left[\int_0^c (x - \mu)^2 f(x) dx + (c - \mu)^2 \overline{F}(c) I(c > \mu) \right] dc;$

equivalently,

 $H^{(1)}(f, g) = \int_0^\infty g(c) \int_0^c (x - \mu)^2 f(x) dx + \int_{\mu}^\infty g(c) (c - \mu)^2 \overline{F}(c) dc.$

From this definition results analogous to those of the discrete case are obtained.

Lemma 4.4. Let $k_c^{(1)} = \int_0^c (x - \mu)^2 f(x) dx + (c - \mu)^2 \overline{F}(c) I(c > \mu)$. Then for every c > 0, $\sigma_x^2 \ge k_c^{(1)}$.

Proof.
$$\sigma_{\chi}^2 - k_c^{(1)} = \int_c^{\infty} (x - \mu)^2 f(x) dx - (c - \mu)^2 \overline{F}(c) I(c > \mu).$$

Case 1. If $c < \mu$, the second term is zero, and $\sigma_{\chi}^{2} - k_{c}^{(1)} = \int_{c}^{\infty} (x - \mu)^{2} f(x) dx \ge 0$. Case 2. If $c \ge \mu$, then $\sigma_{\chi}^{2} - k_{c}^{(1)} \ge (c - \mu)^{2} \int_{c}^{\infty} f(x) dx - (c - \mu)^{2} \overline{F}(c) = 0$. || <u>Theorem 4.5.</u> $H^{(1)}(X, Y) \le H(X)$. <u>Proof.</u> $H^{(1)}(X) = \int_{0}^{\infty} (x - \mu)^{2} f(x) dx = \int_{0}^{\infty} g(c) [\int_{0}^{\infty} (x - \mu)^{2} f(x) dx] dc$ $\ge \int_{0}^{\infty} g(c) [k_{c}^{(1)}] dc = H^{(1)}(X, Y)$. || <u>Lemma 4.6.</u> $k_{c}^{(1)}$ is increasing in c. <u>Proof.</u> $dk_{c}^{(1)}/dc = (c - \mu)^{2} f(c) - (c - \mu)^{2} f(c) I(c > \mu) + 2\overline{F}(c) (c - \mu)^{2} I(c > \mu)$ $= \begin{cases} (c - \mu)^{2} f(c) & \text{if } c \le \mu, \\ 2\overline{F}(c) (c - \mu) & \text{if } c > \mu, \end{cases}$

and each expression is positive.

<u>Theorem 4.7</u>. Suppose that Y_1 and Y_2 are censoring variables with d.f.'s G_1 and G_2 respectively. Suppose $Y_1 \stackrel{\text{st}}{\leq} Y_2$. Then $H^{(1)}(X, Y_1) \leq H^{(1)}(X, Y_2)$.

<u>Proof.</u> Define a function $k^{(1)}$ by $k^{(1)}(c) = k_c^{(1)}$ as defined in Lemma 4.4. Then $H^{(1)}(X, Y_i) = E_{Y_i}(k^{(1)})$. From Lemma 4.6 the conclusion follows. ||

<u>Definition 4.8.</u> X is said to have an <u>increasing failure rate</u> (IFR) if $r(t) = f(t)(\overline{F}(t))^{-1}$ is increasing in t.

<u>Theorem 4.9</u>. Let censoring be deterministic at time c and let X be an IFR variable. If there exists a value A such that f(x) is decreasing for x > A, then for c sufficiently large, $H^{(1)}(X, c)$ is a concave increasing function of c. <u>Proof.</u> $H^{(1)}(X, c) = k_c^{(1)}$ which is increasing from Lemma 4.6. Computing

d

$$lk_{c}^{(1)}/dc = \begin{cases} 2(c - \mu)f(c) + (c - \mu)^{2}f'(c) & \text{for } c \leq \mu, \\ 2\overline{F}(c) - 2(c - \mu)f(c) & \text{for } c > \mu. \end{cases}$$

The first term is negative if f is decreasing; thus we need only consider the second term. We have $2\overline{F}(c) - 2(c - \mu)f(c) \ge 0$ if and only if $(c - \mu)^{-1} \le r(c)$. But $(c - \mu)^{-1} \Rightarrow 0$ as $c \Rightarrow \infty$. Thus if X is IFR and c is sufficiently large, then the inequality holds. ||

Theorem 4.9 shows that more censoring yields less information; however, this relationship is not as strong as one would like. Consider two censoring distributions G_1 and G_2 , where G_1 is stochastically larger than G_2 up to time μ and equal thereafter. Then the difference in information reduces to $\int_{0}^{\infty} (x-\mu)^{2} f(x) (\overline{G}_{1}(x) - \overline{G}_{2}(x)) dx.$ This term is positive from Theorem 4.7 but it merely reflects the information in those observations where a death occurred under Model 1 and a censorship occurred under Model 2. The difference for the censored observations is zero even though they are stochastically larger in one case than in the other. This occurs because all censored observations which occur prior to time μ are shifted to μ , regardless of when they actually occur. An alternate measure is sought which will more carefully distinguish among censored observations. This can be achieved by a constraint which was previously ignored, that corresponding to the value of the mean of the distribution, μ . Agair, the "worst case" will be used under this new set of restrictions. Given that censorship takes place at time c, consider a new variable, x^0 , with p.d.f. $f^{0}(x)$, which equals f(x) for x < c, and minimizes $\int_{c}^{\infty} (x-\mu)^{2} f^{0}(x) dx$, under the restrictions that $\int_{c}^{\infty} f^{0}(x) = \overline{F}(c)$ and $\int_{c}^{\infty} x f^{0}(x) dx = \int_{c}^{\infty} x f(x) dx$. It can be shown that $f^{0}(x)$ must put all its mass at the point $\alpha(c) = (\overline{F}(c))^{-1} \int_{c}^{\infty} xf(x) dx$. This gives a new definition for information.

- 15 -

Definition 4.10. Let X, Y, Z be defined as in Definition 4.3. Then the information in the random variable Z is defined by $H^{(2)}(X, Y) = \int_0^\infty g(c) [\int_0^C (x - \mu)^2 f(x) dx + (\alpha(c) - \mu)^2 \overline{F}(c)] dc.$

Lemma 4.11. Let $k_c^{(2)} = \int_0^c (x - \mu)^2 f(x) dx + (\alpha(c) - \mu)^2 \overline{F}(c)$. Then for every c > 0, $\sigma_v^2 \ge k_c^{(2)}$.

Proof.
$$\sigma_{X}^{2} - k_{c}^{(2)} = \int_{c}^{\infty} (x - \mu)^{2} f(x) dx - (\alpha(c) - \mu)^{2} \overline{F}(c)$$

= $\int_{c}^{\infty} (x - \alpha(c))^{2} f(x) dx + 2(\alpha(c) - \mu) \int_{c}^{\infty} (x - \alpha(c)) f(x) dx - \int_{c}^{\infty} (x - \alpha(c))^{2} f(x) dx \ge 0.$ ||
Theorem 4.12. $H(X) \ge H^{(2)}(X, Y)$, for every X, Y.

<u>Proof.</u> From Lemma 4.11 the proof is the same as that of Theorem 4.5. $\|$ Lemma 4.13. $k_c^{(2)}$ is increasing in c.

<u>Proof</u>. Direct calculations show that $dk_c^{(2)}/dc = f(c)(c - \mu)^2 \ge 0$.

<u>Theorem 4.14</u>. Let X, Y₁, Y₂ be as in Theorem 4.7. Then $H^{(2)}(X, Y_1) \le H^{(2)}(X, Y_2)$ for every X.

<u>Proof.</u> From Lemma 4.13 the proof follows along the lines of the proof of Theorem 4.7.

<u>Definition 4.15</u>. A random variable is said to have <u>increasing (decreasing) mean</u> residual life IMRL(DMRL), if $g(y) = (\overline{F}(y))^{-1} \int_0^{\infty} \overline{F}(y+t) dt$ is increasing (decreasing) in y.

<u>Theorem 4.16</u>. Suppose censoring is deterministic at time c and X is a DMRL variable. If there exists a number A such that f(x) is decreasing for all x > A then, for sufficiently large c, $H^{(2)}(X, c)$ is a concave increasing function of c. <u>Proof.</u> $H^{(2)}(X, c) = k_c^{(2)}$ which is increasing by Lemma 4.13. Also $d^2k_c^{(2)}/dc^2$ = $2f(c)(c - a(c))[1 - (\overline{F}(c))^{-2}(-cf(c)\overline{F}(c) + a(c)f(c)\overline{F}(c))] + (c - a(c))^2 f'(c)$. The second term is negative beyond the point A. The first term is negative if $f(c)(\overline{F}(c))^{-1} \le (a(c) - c)^{-1}$. We now use the following identity of Meilijson (1971), g(c)r(c) = 1 + g'(c), where r(c) is the failure rate at c, and g(c) = a(c) - c is the mean residual life function at c. If X is a DMRL variable then $g'(c) \le 0$. Thus $g(c)r(c) \le 1$ and the conclusion follows.

The discrete case can be paralleled in one more fashion. The "worst case" scenario is no longer used. Now the remaining mass $\overline{F}(c)$ is simply moved to the point of censoring. Note that this is not the same as in Definition 4.3. There, mass was sometimes displaced to the right. Here, it is always displaced to the left.

<u>Definition 4.17</u>. Let X, Y, and Z be as in Definition 4.3. Then information in the variable Z is $H^{(3)}(X, Y) = \int_0^{\infty} g(c) [\sigma_X^2 \overline{F}(c) + \int_0^c (x - \mu_c)^2 f^*(x) dx] dc$, where μ_c^* = $(\overline{F}(c))^{-1} \int_0^c x f^*(x) dx$ and $f^*(x) = f(x)$ for x < c, = $\overline{F}(x)$ for x = c.

Lemma 4.18. Let σ_c^2 denote the variance of the truncated density f^* . Then $\sigma_c^2 \leq \sigma_{\chi}^2$.

<u>Proof.</u> Let X_1 , X_2 be iid copies of X with p.d.f f(x). Then $2\sigma_X^2 = E(X_1 - X_2)^2$ and $2\sigma_c^2 = E(X_1^c - X_2^c)$ where X_1^c is the truncated version of X_1 , i = 1, 2. Then letting $A = \{X_1 < c, X_2 < c\}$, $B = \{X_1 < c, X_2 \ge c\}$, $C = \{X_1 \ge c, X_2 < c\}$, and $D = \{X_1 \ge c, X_2 \ge c\}$ we have $2\sigma^2 \ge \iint_A (x_1 - x_2)^2 f(x_1) f(x_2) dx_1 dx_2 + \iint_B (x_1 - x_2)^2 f(x_1) f(x_2) dx_1 dx_2 + \iint_C (x_1 - x_2)^2 f(x_1) f(x_2) dx_1 dx_2 + \iint_C (x_1 - x_2)^2 f(x_1) f(x_2) dx_1 dx_2 + \iint_C (x_1 - x_2)^2 f(x_1) f(x_2) dx_1 dx_2 + \iint_C (x_1 - x_2)^2 f(x_1) f(x_2) dx_1 dx_2 + \iint_C (x_1 - x_2)^2 f(x_1) f(x_2) dx_1 dx_2 + \iint_C (x_1 - c)^2 f(x_1) F(c) dx_1$

+ $\int_{0}^{c} (c - x_{2})^{2} f(x_{2}) \overline{F}(c) dx_{2} = 2\sigma_{0}^{2}$. ||

<u>Theorem 4.19</u>. $H^{(3)}(X, Y) \le H(X)$.

Proof. From Lemma 4.18 we have

$$H^{(3)}(X, Y) = \int_0^\infty g(c) [F(c)\sigma_X^2 + \overline{F}(c)\sigma_c^2]$$

$$\leq \int_0^\infty g(c) [F(c)\sigma_X^2 + \overline{F}(c)\sigma_X^2] = \sigma_X^2 = H(X). \parallel$$

Lemma 4.20. σ_c^2 is increasing in c.

<u>Proof.</u> Let $c_1 < c_2$. It is enough to show $\sigma_{c_1}^2 \le \sigma_{c_2}^2$. Denote the two random variables as X_{c_1} and X_{c_2} , then X_{c_1} can be obtained from X_{c_2} by truncating X_{c_2} at c_1 . The desired result follows from Lemma 4.18. ||

<u>Lemma 4.21</u>. Let $L_c = F(c)\sigma_X^2 + \overline{F}(c)\sigma_c^2$. Then L_c is increasing in c.

<u>Proof.</u> $\partial L_c/dc = f(c)(\sigma_X^2 - \sigma_c^2) + \overline{F}(c)(d\sigma_c^2/dc)$. Now, from Lemma 4.18 and Lemma 4.20, $dL_c/dc \ge 0$. ||

Theorem 4.22. Let X, Y_1 , and Y_2 be defined as in Theorem 4.7. Then $H^{(3)}(X, Y_1) \le H^{(3)}(X, Y_2)$.

Proof. From Lemma 4.21, the conclusion follows as in Theorem 4.7.

<u>Theorem 4.23</u>. Let X be an IFR variable. Suppose censoring is deterministic at time c. Suppose there exists a value A such that f(x) is decreasing for all x > A. Then for c sufficiently large, $H^{(3)}(X, C)$ is a concave, increasing function of c.

<u>Proof</u>. $H^{(3)}(X, c) = L_c$, which is increasing by Lemma 4.21. Also, $d^2L_c/dc^2 \le -2f(c)(d\sigma_c^2/dc) + \overline{F}(c)(d^2\sigma_c^2/dc^2)$. Thus $d^2L_c/dc^2 \le 0$ if $f(c)(\overline{F}(c))^{-1} \ge F(c)[3\int_0^c (c-y)^2 f(y)dy]^{-1}$. The term on the right decreases to zero. Hence since X is an IFR variable, the result holds.

It is interesting to note that Rao (1983) also suggests variance as a measure of ecological diversity. He considers measures of the form $\iint k(X, Y) dP_x dP_y$, where $k(\cdot, \cdot)$ is a kernel measuring the distance between X and Y. Taking $k(X, Y) = (X - Y)^2$ gives the variance measure.

We also note that alternate proofs of some of our results can be obtained by using Blackwell's (1951) method for comparing two experiments. For example, to show that the uncensored case is always at least as informative as any censored model, let P denote the distribution of the lifetime variable X, Q the distribution of the independent censoring variable Y. Transform X to (Z, δ) by $(Z = X, \delta = 1)$ if $X \le Y^*$, $(Z = Y^*, \delta = 0)$ if $X > Y^*$, where Y^* is independent of X and has the distribution Q.

<u>Acknowledgments</u>. We are grateful to Ian McKeague for helpful comments and to Prem Goel for a useful discussion concerning the relationship of our approach to David Blackwell's method of comparing two experiments.

References

- Barlow, R. and Hsiung, J. (1983). Information in a Life Test Experiment. <u>The Statistician 32</u>, 35-45.
- Bernardo, J. (1979). Expected Information as Expected Utility. <u>Ann. Statist. 7</u>, 686-690.
- Blackwell, D. (1951). Comparison of Experiments. Proc. Second Berkeley Symp. Math. Statist. Prob. Univ. California Press. 93-102.

Blyth, C. (1959). Note on Estimating Information. Ann. Math. Stat. 30, 71-79.

Brooks, R. J. (1982). On the Loss of Information through Censoring. <u>Biometrika</u> 69, 137-144.

Dobrušin, R. L. (1963). General Formulation of Shannon's Main Theorem. Amer. Math. Soc. Transl. 33, 323-428.

- Fisher, R. A. (1925). Theory of Statistical Estimation. Proc. Cambridge Philos. Soc. 22, 700-725.
- Khinchin, A. I. (1957). <u>Mathematical Foundations of Information Theory</u>. Dover, New York.
- Lindley, D. L. (1956). On a Measure of Information Provided by an Experiment. Ann. Math. Stat. 27, 986-1005.

Marshall, A. and Olkin, I. (1979). Inequalities: Theory of Majorization and its Applications. Academic Press, New York.

- Meilijson, I. (1972). Limiting Properties of the Mean Residual Lifetime Function. Ann. Math. Stat. 43, 354-357.
- Ostrowski, A. M. (1952). Sur Quelques Applications des Convexes et Concaves au Sens de I. Schur. J. Pures Appl. 31, 253-292.

Rao, C. R. (1983). Convexity Properties of Entropy Functions and Analysis of Diversity. University of Pittsburgh Technical Report 83-11.

Schur, I. (1923). Über eine Klasse von Mittelbildungen mit Anwendungen die Dieterminaten. Theorie Sitzungsber Berlin Math. Gesellschaft 22, 9-20.

Shannon, C. E. (1948). A Mathematical Theory of Communication. <u>Bell System</u> Technical Journal 27, 379-423.

Solomon, D. L. (1979). A Comparative Approach to Species Diversity. <u>Ecological</u> <u>Diversity in Theory and Practice</u> (eds. Grassle, *et al.*). International <u>Cooperative Publishing</u>.

CONTRACTOR OF THIS PAGE				
REPORT DOCUMENTAT	I (III) PAGE			
REPORT NUMBER 2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG HUBBER			
AFOSR 85-177 4D-H155644				
TITLE (and subtitle)	5. TYPE OF REPORT & PERIOD COVERED			
Information in Concernal Malala	Technical			
Information in Censored Models	6. PERFORMING ORG. REPORT NUMBER			
AUTHOR(s)	8. CONTRACT OR GRANT NUMBER(s)			
Myles Hollander, Frank Proschan, and	AEOSP E49620-85-C-0007			
James Sconing	Arosk 149020-85-C-0007			
PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TASK			
Department of Statistics	AREA, & MORK UNIT NIMBERS			
Florida State University				
Tallahassee, FL 32306				
CONTROLLING OFFICE MAME AND ADDRESS	12. REPORT DATE			
The U.S. Air Force	June, 1985			
Air Force Office of Scientific Research	13. HURBER OF PAGES			
Bolling Air Force Base, DC 20332	20			
HUNITORING AGENCY NAME & ADDRESS (IF	15. SECURITY CLASS. (of this report)			
different from Controlling Office)				
	15a. DECLASSIFICATION/DOUNGRADING SCHEDULE			
DISTRIBUTION STATEMENT (of this report)	l			

distribution unlimited

DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from report)

SUPPLEMENTARY MOTES

KEY MORDS

Information, randomly censored model, majorization

ABSTRACT (Continue on reverse side if necessary and identify by block number)

Criteria are developed for measuring information in the randomly rightcensored model. Measures which are appropriate include an extension of Shannon's entropy. The measures are seen to satisfy some fundamental theorems including (i) the uncensored case is always at least as informative as any censored model, (ii) information decreases as censoring increases stochastically, and (iii) the information gain is marginally decreasing.

END

FILMED

10-85

DTIC