


RD-A159 456 INTERACTION OF FRONTS WITH TOPOGRAPHY(U) NAVAL
POSTGRADUATE SCHOOL MONTEREY CA D A ZANKOFSKI JUN 85


1/1

UNCLASSIFIED

F/G 4/2

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS - 1963 - A

AD-A159 456

NAVAL POSTGRADUATE SCHOOL

Monterey, California



DTIC
SELECTED
SEP 30 1985
~~S~~ B ~~D~~

THESIS

INTERACTION OF FRONTS WITH TOPOGRAPHY

by

Deborah Ann Zankofski

June 1985

DTIC FILE COPY

Thesis Advisor:

R. T. Williams

Approved for public release; distribution is unlimited

85 09 30 00

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
		8159436
4. TITLE (and Subtitle) Interaction of Fronts with Topography	5. TYPE OF REPORT & PERIOD COVERED Master's Thesis June 1985	
7. AUTHOR(s) Deborah Ann Zankofski	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Postgraduate School Monterey, California 93943-5100	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Postgraduate School Monterey, California 93943-5100	12. REPORT DATE June 1985	
14. MONITORING AGENCY NAME & ADDRESS(if different from Controlling Office)	13. NUMBER OF PAGES 95	
	15. SECURITY CLASS. (of this report) Unclassified	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution is unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Front; Mountain; Deformation; Frontogenesis; Model; Boussinesq; Quasi-geostrophic; Finite Difference		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This study examines the formation of a front over a mountain due to a pre-existing deformation field. The dependent variables are assumed to be independent of x. The hydrostatic Boussinesq equations are used with no diffusion of heat or momentum. A modified coordinate system similar to Phillips' sigma system is used. The model is bounded at the top by a rigid plane and periodic boundary conditions are used in the horizontal. The		

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

#20 - ABSTRACT - (CONTINUED)

solutions show the formation of a front within a finite period of time that tilts toward the cold air. The frontogenesis experiments with a mountain of wavelength 3600 km showed negligible topographic effects.

S N 0102-LF-014-6601

Approved for public release; distribution unlimited

Interaction of Fronts with Topography

by

Deborah Ann Zankofski
Lieutenant, United States Navy
B.S., John Carroll University, 1977
M.S., The Ohio State University, 1979

Submitted in partial fulfillment of the
requirement for the degree of

MASTER OF SCIENCE IN METEOROLOGY AND OCEANOGRAPHY

from the

NAVAL POSTGRADUATE SCHOOL
June 1985

Author:

Deborah Ann Zankofski
Deborah Ann Zankofski

Approved by:

R T Williams
R. T. Williams, Thesis Advisor

M A Rennick
M. A. Rennick, Second Reader

R J Richard
R. J. Richard, Chairman,
Department of Meteorology


J N Dyer
John N. Dyer,
Dean of Science and Engineering

ABSTRACT

This study examines the formation of a front over a mountain due to a pre-existing deformation field. The dependent variables are assumed to be independent of x . The hydrostatic Boussinesq equations are used with no diffusion of heat or momentum. A modified coordinate system similar to Phillips' sigma system is used. The model is bounded at the top by a rigid plane and periodic boundary conditions are used in the horizontal. The solutions show the formation of a front within a finite period of time that tilts toward the cold air. The frontogenesis experiments with a mountain of wavelength 3600 km showed negligible topographic effects.

TABLE OF CONTENTS

I.	INTRODUCTION -----	12
II.	BASIC EQUATIONS -----	14
III.	FINITE DIFFERENCE FORMULATIONS -----	23
IV.	RESULTS -----	28
	A. MOUNTAIN WITH NO DEFORMATION FLOW -----	28
	B. DEFORMATION OVER FLAT TOPOGRAPHY -----	36
	C. MOUNTAIN WITH NON-MOVING DEFORMATION -----	41
	D. MOUNTAIN WITH MOVING DEFORMATION -----	53
	1. Maximum Deformation over the Valley -----	64
	2. Deformation Shifted One-Quarter Wavelength -----	73
	3. Reversed Deformation Field -----	73
	4. Deformation Reverse of Case 2 -----	73
V.	CONCLUSIONS AND RECOMMENDATIONS -----	90
	LIST OF REFERENCES -----	91
	INITIAL DISTRIBUTION LIST -----	92



LIST OF FIGURES

1.	Schematic diagram of the zeta-coordinate system ---	16
2.	The vertical grid, showing the indexing convention and staggering of the variables -----	24
3.	The horizontal grid, showing the indexing convention and staggering of the variables -----	25
4.	$\theta(K)$. Initial. Mountain only -----	31
5.	u (m/sec). Initial. Mountain only -----	32
6.	v (m/sec). Initial. Mountain only -----	33
7.	$\dot{\zeta} (\times 10^{-6} \text{ sec}^{-1})$. Initial. Mountain only -----	34
8.	Profile of the mountain with maximum height 375 m -	35
9.	$\theta(K)$. 72 hours. Mountain only -----	37
10.	u (m/sec). 72 hours. Mountain only -----	38
11.	v (m/sec). 72 hours. Mountain only -----	39
12.	$\dot{\zeta} (\times 10^{-6} \text{ sec}^{-1})$. 72 hours. Mountain only -----	40
13.	$\theta(K)$. Initial. Deformation only -----	42
14.	u (m/sec). Initial. Deformation only -----	43
15.	Deformation flow -----	44
16.	$\theta(K)$. 24 hours. Deformation only -----	45
17.	u (m/sec). 24 hours. Deformation only -----	46
18.	v (m/sec). 24 hours. Deformation only -----	47
19.	$\dot{\zeta} (\times 10^{-6} \text{ sec}^{-1})$. 24 hours. Deformation only -----	48
20.	$\theta(K)$. 48 hours. Deformation only -----	49
21.	u (m/sec). 48 hours. Deformation only -----	50
22.	v (m/sec). 48 hours. Deformation only -----	51

23.	$\dot{\zeta} (\times 10^{-6} \text{ sec}^{-1})$. 48 hours. Deformation only -----	52
24.	$\theta(K)$. Initial. Deformation and mountain -----	54
25.	$u (\text{m/sec})$. Initial. Deformation and mountain -----	55
26.	$\theta(K)$. 24 hours. Deformation and mountain -----	56
27.	$u (\text{m/sec})$. 24 hours. Deformation and mountain ---	57
28.	$v (\text{m/sec})$. 24 hours. Deformation and mountain ---	58
29.	$\dot{\zeta} (\times 10^{-6} \text{ sec}^{-1})$. 24 hours. Deformation and mountain -----	59
30.	$\theta(K)$. 48 hours. Deformation and mountain -----	60
31.	$u (\text{m/sec})$. 48 hours. Deformation and mountain ---	61
32.	$v (\text{m/sec})$. 48 hours. Deformation and mountain ---	62
33.	$\dot{\zeta} (\times 10^{-6} \text{ sec}^{-1})$. 48 hours. Deformation and mountain -----	63
34.	$\theta(K)$. 24 hours. Maximum deformation over valley -	65
35.	$u (\text{m/sec})$. 24 hours. Maximum deformation over valley -----	66
36.	$v (\text{m/sec})$. 24 hours. Maximum deformation over valley -----	67
37.	$\dot{\zeta} (\times 10^{-6} \text{ sec}^{-1})$. 24 hours. Maximum deformation over valley -----	68
38.	$\theta(K)$. 48 hours. Maximum deformation over valley -----	69
39.	$u (\text{m/sec})$. 48 hours. Maximum deformation over valley -----	70
40.	$v (\text{m/sec})$. 48 hours. Maximum deformation over valley -----	71
41.	$\dot{\zeta} (\times 10^{-6} \text{ sec}^{-1})$. 48 hours. Maximum deformation over valley -----	72
42.	Deformation when maximum is on the upslope side of the mountain -----	74
43.	$\theta(K)$. 24 hours. Deformation shifted 1/4 wavelength -----	75

44. u (m/sec). 24 hours. Deformation shifted 1/4 wavelength -----	76
45. v (m/sec). 24 hours. Deformation shifted 1/4 wavelength -----	77
46. $\dot{\zeta} (\times 10^{-6} \text{ sec}^{-1})$. 24 hours. Deformation shifted 1/4 wavelength -----	78
47. Deformation flow when maximum is over the ridge ---	79
48. θ (K). 24 hours. Reversed deformation field -----	80
49. u (m/sec). 24 hours. Reversed deformation field -	81
50. v (m/sec). 24 hours. Reversed deformation field -	82
51. $\dot{\zeta} (\times 10^{-6} \text{ sec}^{-1})$. 24 hours. Reversed deformation field -----	83
52. Deformation when maximum is on the downslope side of the mountain -----	84
53. θ (K). 24 hours. Deformation reverse of case 2 ---	85
54. u (m/sec). 24 hours. Deformation reverse of case 2 -----	86
55. v (m/sec). 24 hours. Deformation reverse of case 2 -----	87
56. $\dot{\zeta} (\times 10^{-6} \text{ sec}^{-1})$. 24 hours. Deformation reverse of case 2 -----	88

LIST OF SYMBOLS

a	One-half total horizontal temperature variation
B	Mountain amplitude
c_p	Specific heat of air at constant temperature
D	Average depth of atmosphere
D_d	Deformation scale
f	Coriolis parameter
g	Acceleration due to gravity
H	θZ
\hat{k}	Vertically directed unit vector
L	Domain scale
M_x	uZ
M_y	vZ
p	Pressure
p_0	Reference pressure
R	Gas constant for dry air
t	Time
T	Temperature
U_d	Deformation component in x-direction
u	Velocity component in x-direction
v	Mean flow in the y-direction
v_d	Deformation component in y-direction
v	Velocity component in y-direction
\vec{v}	Horizontal wind vector
w	Vertical velocity component in the z-coordinate system

x, y	Space coordinates
z	Height above sea level
z_s	Height of lower surface above sea level
D	Distance from the upper boundary to the lower boundary
θ_0	Constant reference potential temperature
θ	Departure of potential temperature from θ_0
θ_s	Mean θ as a function of z in the undisturbed atmosphere
ζ	$(z - z_s)/(D - z_s)$
$\dot{\zeta}$	Vertical velocity in the ζ -coordinate system
∇	Horizontal gradient operator
$\frac{\partial \theta_s}{\partial z}$	Constant lapse rate
Φ	$\theta_0 \left(c_p \left(\frac{p}{p_0} \right)^\kappa - \pi_0 \right)$
π_0	$- \frac{g}{\theta_0} z + c_p$
κ	$\frac{R}{c_p}$
α	$\frac{L}{f} \left(\frac{g}{\theta_0} \frac{\partial \theta_s}{\partial z} \right)^{1/2}$

ACKNOWLEDGEMENTS

The author wishes to express her appreciation to Dr. Roger Terry Williams for his technical assistance, recommendations, encouragement, and patience.

Thanks are also expressed to Dr. Mary Alice Rennick and Dr. James Boyle for answering my unending questions about the computer system.

I. INTRODUCTION

Williams and Plotkin (1968) have shown that atmospheric frontogenesis occurs for large time limits using the quasi-geostrophic equation. Their results are unrealistic in the respect that the frontal zone does not tilt with height. A comparison of the linear and nonlinear hydrostatic primitive equations by Williams (1972) shows that the linear solutions are close to the solutions of the quasi-geostrophic equations. The nonlinear solutions are more realistic and show the formation of frontal discontinuities in a finite period of time. Hoskins and Bretherton (1972) showed that dynamic processes in addition to horizontal deformation can produce a front. Bannon (1983) and (1984) investigated quasi-geostrophic and semi-geostrophic frontogenesis over topography. Monotonically sloping topography beneath a deformation leads to the formation of a surface discontinuity in the absence of an initial horizontal thermal gradient. A front associated with a translating deformation field over an isolated orographic feature is affected by the topographically forced divergence field as the front passes over the mountain. Convergence will increase frontogenesis and divergence will decrease it.

The purpose of this study is to investigate the effect of topography on a front which arises from deformation. Periodic boundary conditions are introduced because they

simplify the mountain flow simulation. The deformation is independent of height. The Boussinesq equations are used in which the dependent variables are independent of x . A coordinate system similar to that of Phillips (1957) is used. The equations are solved numerically and output is shown in the form of vertical cross-sections in the $y-\zeta$ plane.

II. BASIC EQUATIONS

The basic equations used in this study are the Boussinesq equations which neglect the compressibility of the atmosphere. The hydrostatic approximation is made and heating and friction are neglected. These equations will be applied to a model in which the flow is bounded above by a rigid horizontal plane.

The Boussinesq equations [Ogura and Phillips, 1962] may be expressed in the z-coordinate system as

$$\frac{\partial \vec{v}}{\partial t} + \vec{v} \cdot \nabla \vec{v} + w \frac{\partial \vec{v}}{\partial z} = -\nabla \phi - f \hat{k} \times \vec{v}, \quad (1)$$

$$\frac{\partial \theta}{\partial t} + \vec{v} \cdot \nabla \theta + w \frac{\partial \theta}{\partial z} = 0, \quad (2)$$

$$\frac{\partial \phi}{\partial z} = \frac{g}{\theta_0} \theta, \quad (3)$$

$$\nabla \cdot \vec{v} + \frac{\partial w}{\partial z} = 0, \quad (4)$$

where

$$\theta = T \left(\frac{P}{P_0} \right)^{\kappa} - \theta_0, \quad (5)$$

$$\phi = \theta_0 \left[c_p \left(\frac{P}{P_0} \right)^{\kappa} - \pi_0 \right], \quad (6)$$

and

$$\pi_0 = - \frac{g}{\theta_0} z + c_p . \quad (7)$$

In this model, the Coriolis parameter is constant, and the only quantity which varies with longitude is ϕ .

To simplify the finite differencing of the equations, a modified coordinate system is introduced in which the lower boundary surface becomes a coordinate surface. This system permits the application of the lower boundary condition at the surface of the mountain. The coordinate system is similar to the sigma system introduced by Phillips (1957).

Following Walton (1968) and DeBoer (1970), the vertical coordinate, zeta, is defined as

$$\zeta = \frac{z - z_s}{D} , \quad (8)$$

where D and z_s are defined in Fig. 1 and $Z = D - z_s$. With these definitions, $\zeta = 0$ at the lower boundary and $\zeta = 1$ at the upper boundary.

Transformation of equations (1) through (4) from the z -coordinate system into the ζ -coordinate system yields

$$\frac{\partial \vec{v}}{\partial t} + \vec{v} \cdot \nabla \vec{v} + \zeta \frac{\partial \vec{v}}{\partial \zeta} = - \nabla \phi - \frac{\partial \phi}{\partial \zeta} (\frac{\zeta-1}{z}) \nabla_h z_s - f \hat{k} \times \vec{v} , \quad (9)$$

$$\frac{\partial \theta}{\partial t} + \vec{v} \cdot \nabla \theta + \dot{\zeta} \frac{\partial \theta}{\partial \zeta} = 0 , \quad (10)$$

$$\frac{\partial \phi}{\partial \zeta} = \frac{g}{\theta_0} z \theta , \quad (11)$$




Fig. 1. Schematic diagram of the zeta-coordinate system

$$\nabla \cdot (Z \vec{v}) + Z \frac{\partial \zeta}{\partial \zeta} = 0 . \quad (12)$$

In equations (9) through (12), ∇ is the horizontal gradient operator on zeta surfaces.

It is desirable to put the equations in flux form in order to satisfy certain computational properties. Equations (9) through (12) become

$$\frac{\partial (M_x)}{\partial t} + \frac{\partial}{\partial y}(M_y u) + \frac{\partial}{\partial \zeta}(\dot{\zeta} u Z) = -Z \frac{\partial \Phi}{\partial x} + f M_y , \quad (13)$$

$$\frac{\partial (M_y)}{\partial t} + \frac{\partial}{\partial y}(M_y v) + \frac{\partial}{\partial \zeta}(\dot{\zeta} v Z) = -Z \frac{\partial \Phi}{\partial y} + \frac{g}{\theta_0} Z \theta (\zeta - 1) \frac{\partial Z}{\partial y} - f M_x , \quad (14)$$

$$\frac{\partial H}{\partial t} + \frac{\partial}{\partial y}(M_y \theta) + \frac{\partial}{\partial \zeta}(\dot{\zeta} \theta Z) = 0 , \quad (15)$$

$$\frac{\partial \Phi}{\partial \zeta} = \frac{g}{\theta_0} Z \theta , \quad (16)$$

$$\frac{\partial M_y}{\partial y} + Z \frac{\partial \dot{\zeta}}{\partial \zeta} = 0 \quad (17)$$

where $M_x = uZ$, $M_y = vZ$ and $H = \theta Z$.

The boundary conditions are

$$\dot{\zeta} = 0 \quad \text{at} \quad \zeta = 0 , \quad (18)$$

and

$$\dot{\zeta} = 0 \quad \text{at} \quad \zeta = 1 . \quad (19)$$

For this study, the flow will be independent of x and periodic in y . Φ is the only variable allowed to vary in the x -direction, so $\frac{\partial}{\partial x} = 0$ for all other variables.

In order to obtain an expression for Φ , equation (16) may be integrated with respect to ζ giving

$$\Phi = \int_0^\zeta \frac{q}{\theta} \theta z d\zeta + C . \quad (20)$$

The constant of integration, C , can be eliminated by taking the vertical average of equation (20) and subtracting from equation (20) which gives the following

$$\Phi = \frac{q}{\theta} \int_0^\zeta \theta z d\zeta - \overline{\left(\int_0^\zeta \theta z d\zeta \right)} + \overline{\Phi} . \quad (21)$$

In the above equation, $\overline{\cdot}$ represents the vertical average of a given variable.

A value for Φ must be found before a solution for Φ can be determined. First, take the vertical average of (17) and apply the boundary conditions (18) and (19) which gives

$$\frac{\partial}{\partial y} \overline{M_y} = - z \dot{\zeta} \Big|_0^1 = 0 . \quad (22)$$

This equation shows that $\overline{M_y}$ must be a constant in order to satisfy the boundary conditions. Next, take the vertical

average of equation (14) and differentiate with respect to y which yields

$$\frac{\partial^2}{\partial t \partial y} (\overline{M}_y) + \frac{\partial^2}{\partial y^2} (\overline{M}_y v) = \frac{\partial}{\partial y} (-z \frac{\partial \overline{\Phi}}{\partial y} + \frac{q}{\theta_0} \sqrt{(z\theta)(\zeta-1)} \frac{\partial \overline{Z}}{\partial y} - \overline{f M}_x) . \quad (23)$$

The first term in equation (23) is identically zero from (22) and the resulting equation becomes

$$\frac{\partial}{\partial y} (z \frac{\partial \overline{\Phi}}{\partial y}) = \frac{\partial}{\partial y} (- \frac{\partial}{\partial y} (\overline{M}_y v) + \frac{q}{\theta_0} \sqrt{(z\theta)(\zeta-1)} \frac{\partial \overline{Z}}{\partial y} - f z \overline{u}) . \quad (24)$$

This equation can be solved diagnostically for $\overline{\Phi}$. This equation must be applied in order to satisfy the boundary conditions (18) and (19).

To obtain an expression for $\frac{\partial \overline{\Phi}}{\partial x}$, which is a constant, equation (21) is differentiated with respect to x which yields

$$\frac{\partial \overline{\Phi}}{\partial x} = \frac{q}{\theta_0} z \left(\int_0^\zeta \frac{\partial \theta}{\partial x} d\zeta - \overline{\int_0^\zeta \frac{\partial \theta}{\partial x} d\zeta} \right) + \frac{\partial \overline{\Phi}}{\partial x} . \quad (25)$$

Since $\frac{\partial \theta}{\partial x} = 0$ in this study,

$$\frac{\partial \overline{\Phi}}{\partial x} = \frac{\partial \overline{\Phi}}{\partial x} \quad (26)$$

and this quantity is assumed to be independent of y.

Consider the domain averages of equations (13) and (14)
which are

$$\frac{d}{dt} (\overline{\overline{M_x}}) = - \bar{z} \frac{\partial \overline{\Phi}}{\partial x} + f \overline{\overline{M_y}}, \quad (27)$$

$$\frac{d}{dt} (\overline{\overline{M_y}}) = - \bar{z} \frac{\partial \overline{\Phi}}{\partial y} + \overline{\overline{\frac{g}{\theta_0} (z\theta)(\zeta-1) \frac{\partial z}{\partial y}}} - f \overline{\overline{M_x}}, \quad (28)$$

where $\overline{\overline{\quad}}$ represents the y average. Initially, $\frac{\partial \overline{\Phi}}{\partial x}$ is computed geostrophically as

$$\frac{\partial \overline{\Phi}}{\partial x} = \frac{f}{\bar{z}} \overline{\overline{M_y}}_{t=0}. \quad (29)$$

If this relationship is used for all time and if the mass flux in the x-direction is zero initially, then $\overline{\overline{M_x}}$ will be zero for all time. The first two terms on the right side of equation (28) represent the mountain drag. The sum of these two terms for mid-latitude flow has been shown to be negative in the mean. This will lead to a continual loss of y-momentum. This kind of loss would make a quasi-steady final state impossible. An artificial source of momentum is required to maintain the mean meridional flow. This artificial source of momentum is provided by holding $\frac{\partial \overline{\Phi}}{\partial x}$ constant as obtained in equation (29). In this case, if $\overline{\overline{M_y}}$ is decreased below the initial geostrophic value, $\overline{\overline{M_x}}$ will increase and restore the momentum loss. Thus, in the long term mean, the mountain

terms are balanced by the mean Coriolis force. However, for the horizontal scales to be considered in this study, this momentum source may not be needed since synoptic scale mountains cause very little wave drag.

The dependent variables are to be divided as follows:

$$u(x,y,\zeta,t) = u_d(x,y) + u'(y,\zeta,t) ,$$

$$v(x,y,\zeta,t) = v_d(x,y) + v'(y,\zeta,t) ,$$

$$\dot{\zeta}(y,\zeta,t) = \zeta'(y,\zeta,t) , \quad (30)$$

$$\theta(y,\zeta,t) = \theta'(y,\zeta,t) ,$$

$$\phi(x,y,\zeta,t) = \phi(x,y) + \phi'(y,\zeta,t) .$$

The basic wind deformation field is chosen to be periodic in y and to have zero vorticity and divergence. Thus, the deformation field is given by

$$u_d(x,y) = -D_d L \sinh(\frac{x}{L}) \sin(\frac{y}{L}) , \quad (31)$$

$$v_d(x,y) = -D_d L \cosh(\frac{x}{L}) \cos(\frac{y}{L}) , \quad (32)$$

where D_d and L are constants. This deformation field was used by Stone (1966) in his study of quasi-geostrophic

frontogenesis. Since the flow is assumed to be independent of x in this study, equations (31) and (32) are evaluated at $x = 0$ resulting in

$$u_d(0, y) = 0 , \quad (33)$$

$$v_d(0, y) = -D_d L \cos\left(\frac{y}{L}\right) . \quad (34)$$

Substituting the expressions in equations (30) through (34) into equations (13) through (15) and dropping the prime notation yields

$$\begin{aligned} z \frac{\partial u}{\partial t} + v_d z \frac{\partial u}{\partial y} + u \frac{\partial (v_d z)}{\partial y} + \frac{\partial}{\partial y}(v_z u) + \frac{\partial}{\partial \zeta}(\dot{\zeta} u z) \\ = -z \frac{\partial \phi}{\partial x} + f v z , \end{aligned} \quad (35)$$

$$\begin{aligned} z \frac{\partial v}{\partial t} + \frac{\partial}{\partial y}(v_z v) + v \frac{\partial (v_d z)}{\partial y} + v_d z \frac{\partial v}{\partial y} + \frac{\partial}{\partial \zeta}(\dot{\zeta} u z) \\ = -z \frac{\partial \phi}{\partial y} + \frac{g}{\theta_0} \theta z (\zeta - 1) \frac{\partial z}{\partial y} - fu z , \end{aligned} \quad (36)$$

$$z \frac{\partial \theta}{\partial t} + \frac{\partial}{\partial y}(v_z \theta) + \frac{\partial}{\partial \zeta}(\dot{\zeta} \theta z) = 0 , \quad (37)$$

where $v_d = -D_d L \cos\left(\frac{y}{L}\right)$. This last set of equations is the set that needs to be solved for u , v and θ . In this study, equation (24) does not have to be solved if M_y is restored to the initial constant value at every timestep.

III. FINITE DIFFERENCE FORMULATIONS

Staggered grids are used in both the horizontal and vertical. The vertical grid is divided into $K-1$ levels of constant $\Delta\zeta$ (Fig. 2). Quantities defined for the layers will be denoted by integer subscripts and the ζ coordinate value will be denoted by ζ_{I_k} . The quantities θ , u , v and ϕ are all defined on the integer levels. The vertical motion, $\dot{\zeta}$, on the other hand, is defined on the top, bottom and interface levels. These levels will be denoted with half-integer subscripts and the vertical coordinate will be identified by $\zeta_{k+1/2}$. The horizontal grid is divided into $J-1$ sections of constant resolution (Fig. 3). Momentum variables are defined on the integer subscripted points denoted by Y_j . The momentum variables include u , v , M_x , M_y and V_d . The mass variables, θ , ϕ , Z and $\dot{\zeta}$, are defined on the half-integer subscripted points. These points will be denoted as $Y_{j+1/2}$. Values needed off their respective grid points will be identified by a caret and will be defined as

$$\hat{s}_j = \frac{(Y_{j+1} - Y_j) s_{j+1/2} + (Y_j - Y_{j-1}) s_{j-1/2}}{2[Y_{j+1/2} - Y_{j-1/2}]}, \quad (38)$$

for quantities defined on the half-integer points and as

$$\zeta_{k_1+1/2} \bullet \dot{\zeta}$$

$$\zeta l_{k_1} \bullet u, v, \theta, \phi$$

$$\zeta_{k_1-1/2} \bullet \dot{\zeta}$$

$$\zeta_{k+1/2} \bullet \dot{\zeta}$$

$$\zeta l_k \bullet u, v, \theta, \phi$$

$$\zeta_{k-1/2} \bullet \dot{\zeta}$$

$$\zeta_{3/2} \bullet \dot{\zeta}$$

$$\zeta l_1 \bullet u, v, \theta, \phi$$

$$\zeta_{1/2} \bullet \dot{\zeta}$$

Fig. 2. The vertical grid, showing the indexing convention and staggering of the variables




Fig. 3. The horizontal grid, showing the indexing convention and staggering of the variables

$$\hat{s}_{j+1/2} = \frac{(YI_{j+3/2} - YI_{j+1/2})s_{j+1} + (YI_{j+1/2} - YI_{j-1/2})s_j}{2[Y_{j+1} - Y_j]}, \quad (39)$$

for quantities at the integer points. Periodic boundary conditions are used in this study for all quantities.

Equations (11), (13) through (15) and (17) are solved by using finite differences in y , ζ and t . The finite differencing scheme is chosen to conserve the sum of kinetic energy and potential energy.

The time differencing scheme uses a leapfrog scheme with a periodic use of the Euler backward scheme. The Euler backward scheme is used to eliminate the computational mode arising with the leapfrog scheme [Haltiner and Williams, 1980].

Equations (11), (17) and (13) through (15) may be written in finite difference form as

$$\begin{aligned} \hat{z}_j \frac{\partial u_{j,k}}{\partial t} &+ \frac{\left(\frac{M_{Y_{j+1,k}} + M_{Y_{j,k}}}{2}\right)\left(\frac{u_{j+1,k} + u_{j,k}}{2}\right) - \left(\frac{M_{Y_{j,k}} + M_{Y_{j-1,k}}}{2}\right)\left(\frac{u_{j,k} + u_{j-1,k}}{2}\right)}{YI_{j+1/2} - YI_{j-1/2}} \\ &+ \frac{(\dot{\zeta}z)_{j,k+1/2}\left(\frac{u_{j,k+1} + u_{j,k}}{2}\right) - (\dot{\zeta}z)_{j,k-1/2}\left(\frac{u_{j,k} + u_{k,k-1}}{2}\right)}{\zeta_{k+1/2} - \zeta_{k-1/2}} \\ &= - \hat{z}_j \frac{\partial \phi}{\partial x} + f v_{j,k} \hat{z}_j, \end{aligned} \quad (40)$$

$$\begin{aligned}
& \hat{z}_j \frac{\partial v_{j,k}}{\partial t} + \frac{\left(\frac{M}{2} \frac{y_{j+1,k} + M}{2} y_{j,k} \right) \left(\frac{v_{j+1,k} + v_{j,k}}{2} \right) - \left(\frac{M}{2} \frac{y_{j,k} + M}{2} y_{j-1,k} \right) \left(\frac{v_{j,k} + v_{j-1,k}}{2} \right)}{YI_{j+1/2} - YI_{j-1/2}} \\
& + \frac{(\dot{\zeta}z)_{j,k+1/2} \left(\frac{v_{j,k+1} + v_{j,k}}{2} \right) - (\dot{\zeta}z)_{j,k-1/2} \left(\frac{v_{j,k} + v_{j-1,k-1}}{2} \right)}{\zeta_{k+1/2} - \zeta_{k-1/2}} \\
= & - \hat{z}_j \frac{\phi_{j+1/2,k} - \phi_{j-1/2,k}}{YI_{j+1/2} - YI_{j-1/2}} + \frac{q}{\theta_0} \hat{z}_{j,k}^{(\zeta-1)} \frac{z_{j+1/2} - z_{j-1/2}}{YI_{j+1/2} - YI_{j-1/2}} \\
& - f u_{j,k} \hat{z}_{j,k} , \tag{41}
\end{aligned}$$

$$\begin{aligned}
z_{j+1/2} \frac{\partial \theta_{j+1/2,k}}{\partial t} + \frac{M}{2} \frac{y_{j+1,k}}{Y_{j+1} - Y_j} \left(\frac{\theta_{j+3/2,k} + \theta_{j+1/2,k}}{2} \right) - \frac{M}{2} \frac{y_{j,k}}{Y_j - Y_{j-1}} \left(\frac{\theta_{j+1/2,k} + \theta_{j-1/2,k}}{2} \right) \\
+ \frac{(\dot{\zeta}z)_{j+1/2,k+1/2} \left(\frac{\theta_{j+1/2,k+1} + \theta_{j+1/2,k}}{2} \right) - (\dot{\zeta}z)_{j+1/2,k-1/2} \left(\frac{\theta_{j+1/2,k} + \theta_{j+1/2,k-1}}{2} \right)}{\zeta_{k+1/2} - \zeta_{k-1/2}} \\
= 0 , \tag{42}
\end{aligned}$$

$$\frac{\phi_{j+1/2,k+1} - \phi_{j+1/2,k}}{\zeta_{k+1} - \zeta_k} = \frac{q}{\theta_0} z_{j+1/2} \frac{\theta_{j+1/2,k+1} + \theta_{j+1/2,k}}{2} , \tag{43}$$

and

$$\frac{M}{2} \frac{y_{j+1,k} - M}{2} y_{j,k} = - z_{j+1/2} \left(\frac{\dot{\zeta}_{j+1/2,k+1/2} - \dot{\zeta}_{j+1/2,k-1/2}}{\zeta_{k+1/2} - \zeta_{k-1/2}} \right) . \tag{44}$$

IV. RESULTS

All numerical results to be shown use the following numerical values for the constants:

$$\begin{aligned} a &= 12 \text{ K} \\ B &= 375 \text{ m} \\ D &= 9 \text{ km} \\ D_d &= 10^{-5} \text{ sec}^{-1} \\ f &= 10^{-4} \text{ sec}^{-1} \\ g &= 9.81 \text{ m/sec}^2 \\ L &= 3600 \text{ km} \\ \theta_0 &= 300 \text{ K} \\ \frac{\partial \theta_s}{\partial z} &= 4 \text{ K/km} \end{aligned} \tag{45}$$

The following finite-difference increments are used.

$$\begin{aligned} \Delta t &= 180 \text{ sec} \\ \Delta y &= 40 \text{ km} \\ \Delta \zeta &= 1/27 \end{aligned} \tag{46}$$

A. MOUNTAIN WITH NO DEFORMATION FLOW

The initial conditions are obtained from the quasi-geostrophic equations for steady state flow over topography [Walton, 1968; DeBoer, 1970]. The initial potential temperature field is given by

$$\theta(y, z, 0) = \theta_s + B \frac{\partial \theta}{\partial z} \frac{s \sinh[\alpha(z-D)]}{\sinh(\alpha D)} \cos\left(\frac{y}{L}\right) - \frac{2a}{\pi} \cos\left(\frac{y}{L}\right) . \quad (47)$$

The quantity $\frac{\partial \theta}{\partial z}$ is the constant initial lapse rate and

$$\alpha = \frac{1}{fL} \left(\frac{g}{\theta_0} \frac{\partial \theta}{\partial z} \right)^{1/2} . \quad (48)$$

Substituting equation (47) into the thermal wind equation yields

$$\frac{\partial u}{\partial z} = \frac{1}{f} \frac{g}{\theta_0} \frac{1}{L} [B \frac{\partial \theta}{\partial z} \frac{s \sinh[\alpha(z-D)]}{\sinh(\alpha D)} \sin\left(\frac{y}{L}\right) - \frac{2a}{\pi} \sin\left(\frac{y}{L}\right)] . \quad (49)$$

This equation can be integrated with respect to z to give

$$u(y, z, 0) = B \left(\frac{g}{\theta_0} \frac{\partial \theta}{\partial z} \right)^{1/2} \frac{\cosh[\alpha(z-D)]}{\sinh(\alpha D)} \sin\left(\frac{y}{L}\right) - \frac{2a}{\pi} \sin\left(\frac{y}{L}\right) (z - \frac{D}{2}) . \quad (50)$$

The initial v field can be shown to be

$$v(y, z, 0) = V + \alpha VB \frac{\cosh[\alpha(z-D)]}{\sinh(\alpha D)} \cos\left(\frac{y}{L}\right) . \quad (51)$$

The initial w and ϕ are obtained from equations (4) and (3).

It is necessary to rewrite the initial conditions in the zeta-coordinate system. From equation (6), equations (47), (50), and (51) can be rewritten as

$$\theta(y, \zeta, 0) = \theta_s + B \frac{\partial \theta}{\partial z} \frac{s \sinh[\alpha z(\zeta-1)]}{\sinh(\alpha D)} \sin\left(\frac{y}{L}\right) - \frac{2a}{\pi} \sin\left(\frac{y}{L}\right), \quad (52)$$

$$u(y, \zeta, 0) = B \left(\frac{q}{\theta_0} \frac{\partial \theta}{\partial z} \right)^{1/2} \frac{\cosh[\alpha z(\zeta-1)]}{\sinh(\alpha D)} \sin\left(\frac{y}{L}\right) - \frac{2a}{\pi} \sin\left(\frac{y}{L}\right) (\zeta - \frac{1}{2}) D, \quad (53)$$

and

$$v(y, \zeta, 0) = v + \alpha VB \frac{\cosh[\alpha z(\zeta-1)]}{\sinh(\alpha D)} \cos\left(\frac{y}{L}\right). \quad (54)$$

These initial conditions are shown in Figs. 4-7. ζ and ϕ are given by equations (10) and (9).

The value of θ_s was computed using the formula

$$\theta_s = \frac{\partial \theta}{\partial z} \left[\frac{D}{2} + z(\zeta-1) \right]. \quad (55)$$

The profile of the mountain used in this study is given by

$$z_s = B \cos\left(\frac{y}{L}\right). \quad (56)$$

and is shown in Fig. 8.

The initial conditions show anticyclonic vorticity over the ridge and cyclonic vorticity over the valley. There is divergence on the upslope and convergence on the downslope. These perturbations die off with height above the surface.




Fig. 4. θ (K). Initial. Mountain only




Fig. 5. u (m/sec). Initial. Mount in only




Fig. 6. v (m/sec). Initial. Mountain only




Fig. 7. ζ ($\cdot 10^{-6} \text{ sec}^{-1}$). Initial. Mountain only




Fig. 8. Profile of the mountain with maximum height 375 m

Figs. 9-12 show the results at $t = 72$ hours. Since the original fields were analytic, only a slight adjustment to the mountain can be seen in the v , and hence, the ζ fields.

B. DEFORMATION OVER FLAT TOPOGRAPHY

For this case, the initial potential temperature field contains a horizontal variation. The initial field is given by

$$\theta(y, z, 0) = \frac{\partial \theta}{\partial z} \left(z - \frac{D}{2} \right) - \frac{2a}{\pi} \cos \left(\frac{y}{L} \right) . \quad (57)$$

The initial u field is again derived using the thermal wind relationship and can be shown to be

$$u(y, z, 0) = - \frac{2a}{\pi} \sin \left(\frac{y}{L} \right) \left(z - \frac{D}{2} \right) . \quad (58)$$

The initial v field is identically zero and thus, w is forced to be zero by equation (3).

It is again necessary to write equations (48) and (49) in zeta-coordinates as

$$\theta(y, \zeta, 0) = \frac{\partial \theta}{\partial z} (\zeta - 1) D - \frac{2a}{\pi} \cos \left(\frac{y}{L} \right) , \quad (59)$$

and

$$u(y, \zeta, 0) = - \frac{2a}{\pi} \sin \left(\frac{y}{L} \right) \left(\zeta - \frac{1}{2} \right) D . \quad (60)$$




Fig. 9. θ (K). 72 hours. Mountain only




Fig. 10. u (m/sec). 72 hours. Mountain only




Fig. 11. v (m/sec). 72 hours. Mountain only




Fig. 12. $\dot{\gamma} (\times 10^{-6} \text{ sec}^{-1})$. 72 hours. Mountain only

v and ζ are zero initially and ϕ is given by equation (9).

Figures 13 and 14 show the initial conditions and Fig. 15 shows the deformation flow given by equation (34).

The front is developed by 24 hours as shown in Figs. 16-19. The frontal discontinuity forms along the point where $v = 0$ and tilts toward the colder air. Cyclonic shear can be seen along the frontal boundary. The relative vorticity is less than zero on the north side of the front and is greater than zero to the south.

By 48 hours, Figs. 20-23 show the strengthening of the frontal discontinuity. The tilt remains the same as at the 24 hour point. On the other side of the domain, the temperature gradient is weakening, which indicates frontolysis occurring in that region.

C. MOUNTAIN WITH NON-MOVING DEFORMATION

The previous two cases are combined in order to investigate the combined effects of a deformation flow and a mountain. The initial fields are given by

$$\theta(y, \zeta, 0) = \theta_s + B \frac{\partial \theta}{\partial z} \frac{s}{\sinh(\alpha D)} \frac{\sinh[\alpha Z(\zeta-1)]}{\sinh(\alpha D)} \sin\left(\frac{y}{L}\right) - \frac{2a}{\pi} \cos\left(\frac{y}{L}\right), \quad (61)$$

$$u(y, \zeta, 0) = B \left(\frac{\sigma}{\theta_0} \frac{\partial \theta}{\partial z} \right)^{1/2} \frac{\cosh[\alpha Z(\zeta-1)]}{\sinh(\alpha D)} \sin\left(\frac{y}{L}\right) - \frac{2a}{\pi} \sin\left(\frac{y}{L}\right), \quad (62)$$




Fig. 13. θ (K). Initial. Deformation only




Fig. 14. u (m/sec). Initial. Deformation only




Fig. 15. Deformation flow




Fig. 16. θ (K). 24 hours. Deformation only




Fig. 17. u (m/sec). 24 hours. Deformation only




Fig. 18. v (m/sec). 24 hours. Deformation only




Fig. 19. ζ ($\times 10^{-6} \text{ sec}^{-1}$). 24 hours. Deformation only




Fig. 20. θ (K). 48 hours. Deformation only




Fig. 21. u (m/sec). 48 hours. Deformation only




Fig. 22. v (m/sec). 48 hours. Deformation only




Fig. 23. ξ ($\times 10^{-6} \text{ sec}^{-1}$). 48 hours. Deformation only

$$v(y, \zeta, 0) = v + \alpha VB \frac{\cosh[\alpha Z(\zeta-1)]}{\sinh(\alpha D)} \cos \left(\frac{y}{L} \right), \quad (63)$$

and ζ is given by equation (10). These fields can be seen in Figs. 24, 25, 26 and 27. The deformation is the same as in the previous case.

As in the previous case, a tightening of the temperature gradient has occurred by the 24 hour point, although not as strong (Figs. 26-29). The front has moved farther to the north and the tilt is smaller than in Case B. Throughout the next 24 hours, the front continues to move to the north and passes over the mountain ridge (Figs. 30-33). The gradient weakens as the front passes over the ridge and the tilt is negligible. In this experiment, the front weakens because it is advected away from the frontogenetic part of the deformation field.

D. MOUNTAIN WITH MOVING DEFORMATION

The next four cases to be considered start with the same initial conditions as in Case C (Figs. 24, 25, 26 and 27). However, in these cases, the deformation moves to the north at a speed of 10 m/sec. Each case examines a different starting place for the maximum northward deformation flow. The cases have the maximum flow over the valley, on the upslope side of the mountain, directly over the ridge, and on the downslope side of the mountain.




Fig. 24. θ (K). Initial. Deformation and mountain




Fig. 25. u (m/sec). Initial. Deformation and mountain




Fig. 26. θ (K). 24 hours. Deformation and mountain




Fig. 27. u (m/sec). 24 hours. Deformation and mountain




Fig. 28. v (m/sec). 24 hours. Deformation and mountain




Fig. 29. : ($\cdot \cdot 10^{-6} \text{ sec}^{-1}$). 24 hours. Deformation and mountain




Fig. 30. θ (K). 48 hours. Deformation and mountain




Fig. 31. u (m/sec). 48 hours. Deformation and mountain




Fig. 32. v (m/sec). 46 hours. Deformation and mountain




Fig. 33. $\dot{\xi}$ ($\times 10^{-6} \text{ sec}^{-1}$). 48 hours. Deformation and mountain

1. Maximum Deformation over the Valley

The initial deformation is shown in Fig. 15. As in the previous cases, the front has formed by 24 hours (Figs. 34-37). There is a small tilt toward the colder air in the north and the front has moved farther up the mountain than in Case C. The upper level front has passed over the mountain ridge.

As the deformation flow continues to move northward, so does the front. During the next 24 hours, the speed of the front slows down, but the gradient in the temperature field tightens (Figs. 38-41) and forms a discontinuity. The cyclonic shear around the front has increased and there is strong vertical motion in that region.

A careful comparison of the potential temperatures at $t = 24$ hours between Figs. 24 and 34 shows very little change between the flat experiment and the mountain experiment, except for the translation in Fig. 34. The same conclusion holds at 48 hours. Bannon (1983,1984) has shown that mountain effects on frontogenesis will be important when the mountain forced convergence is of the same order as the frontal convergence. Comparison of v in Figs. 6 and 18 shows that the mountain forced convergence is only about 25 percent of the mountain forced convergence at 24 hours. This ratio is even smaller later in the frontal evolution. With the large scale mountain used in this and later experiments, the effects of the topography are unimportant.




Fig. 34. $\theta(K)$. 24 hours. Maximum deformation over valley




Fig. 35. u (m/sec). 24 hours. Maximum deformation over valley




Fig. 36. v (m/sec). 24 hours. Maximum deformation over valley




Fig. 37. $\dot{\zeta}$ ($\times 10^{-6} \text{ sec}^{-1}$). 24 hours. Maximum deformation over valley




Fig. 38. θ (K). 48 hours. Maximum deformation over valley




Fig. 39. u (m/sec). 48 hours. Maximum deformation over valley




Fig. 40. v (m/sec). 48 hours. Maximum deformation over valley




Fig. 41. $\dot{\zeta}$ ($\times 10^{-6} \text{ sec}^{-1}$). 48 hours. Maximum deformation over valley

2. Deformation Shifted One-Quarter Wavelength

In this experiment, the deformation field is shifted one-quarter of a wavelength to the north with respect to the initial fields (Fig. 42). The topography is also unimportant in this experiment. At $t = 24$ hours, there is only a slight indication of frontal formation in the temperature field, but a much stronger suggestion of the front in the u field. The u field, however, shows little tilt (Figs. 43-46). Williams (1968) has shown that this phasing of the deformation field is inefficient for frontogenesis.

3. Reversed Deformation Field

As is shown in Fig. 47, the initial deformation flow is in opposite phase with the original deformation field. The front can be seen at 24 hours in Figs. 48-51. As with previous cases, the front tilts toward the cold air, but in this case, the tilt is toward the south. This is a result of the reversed deformation field. The fields are mirror images of the fields in the basic experiment except for the shift.

4. Deformation Reverse of Case 2

The last case to be considered is the one in which the maximum deformation flow is the reverse of Case 2 (Fig. 52). The first indication of frontal activity can be seen 24 hours later (Figs. 53-56) especially in the u field. Strong cyclonic activity can be seen here with no evidence of a tilt. The temperature field shows only a small tightening




Fig. 42. Deformation when maximum is on the upslope side of the mountain




Fig. 43. θ (K). 24 hours. Deformation shifted $1/4$ wavelength




Fig. 44. u (m/sec). 24 hours. Deformation shifted $1/4$ wavelength




Fig. 45. v (m/sec). 24 hours. Deformation shifted $1/4$ wavelength




Fig. 46. $\dot{\xi}$ ($\times 10^{-6} \text{ sec}^{-1}$). 24 hours. Deformation shifted $1/4$ wavelength.




Fig. 47. Deformation flow when maximum is over the ridge




Fig. 48. θ (K). 24 hours. Reversed deformation field




Fig. 49. u (m/sec). 24 hours. Reversed deformation field




Fig. 50. v (m/sec). 24 hours. Reversed deformation field




Fig. 51. $\dot{\xi}$ ($\times 10^{-6} \text{ sec}^{-1}$). 24 hours. Reversed deformation field




Fig. 52. Deformation when maximum is on the down-slope side of the mountain




Fig. 53. θ (K). 24 hours. Deformation reverse of case 2




Fig. 54. u (m/sec). 24 hours. Deformation reverse of case 2




Fig. 55. v (m/sec). 24 hours. Deformation reverse of case 2




Fig. 56. $\dot{\xi}$ ($\times 10^{-6} \text{ sec}^{-1}$). 24 hours. Deformation reverse of case 2

of the gradient from the initial conditions. This is very similar to Case 2, but the maximum shear is at the lower boundary.

V. CONCLUSIONS AND RECOMMENDATIONS

In this study, a numerical model of frontogenesis was constructed which included surface topography. Periodic boundary conditions were included and a periodic deformation field was introduced. The model was successfully tested with frontogenesis with no topography and for stratified flow over a mountain with no deformation field. Combined experiments with topography and a moving deformation field were carried out. The experiments showed negligible effects on frontogenesis by the topography.

The next step in this research is to introduce smaller scale topography which should lead to a noticeable effect as suggested by the frontogenesis theories of Bannon (1983, 1984). Other effects including vertical wind shear could be included later.

LIST OF REFERENCES

- Bannon, P. J., 1983: Quasi-geostrophic frontogenesis over topography. J. Atmos. Sci., 40, 2266-2277.
- _____, 1984: A semi-geostrophic model of frontogenesis over topography. Beitr. Phys. Atmosph., 57, 393-408.
- DeBoer, J. K., 1970: Numerical Simulation of Atmospheric Flow over an Idealized Mountain. MS Thesis, Department of Meteorology, Naval Postgraduate School, 50 pp.
- Haltiner, G. J. and R. T. Williams, 1980: Numerical Prediction and Dynamic Meteorology, John Wiley and Sons, 477 pp.
- Hoskins, B. J. and F. P. Bretherton, 1972: Atmospheric frontogenesis models: Mathematical formulations and solution. J. Atmos. Sci., 29, 11-37.
- Ogura, Y. and Phillips, N. A., 1962: Scale analyses of deep or shallow convection in the atmosphere. J. Atmos. Sci., 19, 173-179.
- Phillips, N. A., 1957: A coordinate system having some special advantages for numerical forecasting. J. Meteor., 24, 184-185.
- Stone, P. H., 1966: Frontogenesis by horizontal wind deformation fields. J. Atmos. Sci., 23, 455-565.
- Walton, J. R., 1968: The Validity of Using Smoothed Topography for Numerical Weather Prediction. MS Thesis, Department of Meteorology, Univ. of Utah, 21 pp.
- Williams, R. T., 1968: A note on quasi-geostrophic frontogenesis. J. Atmos. Sci., 25, 1157-1159.
- _____, 1972: Quasi-geostrophic versus non-geostrophic frontogenesis. J. Atmos. Sci., 29, 3-10.
- _____, and J. Plotkin, 1968: Quasi-geostrophic frontogenesis, J. Atmos. Sci., 25, 201-206.

INITIAL DISTRIBUTION LIST

	No. Copies
1. Defense Technical Information Center Cameron Station Alexandria, Virginia 22306-6145	2
2. Library, Code 0142 Naval Postgraduate School Monterey, California 93943-5100	2
3. Meteorology Reference Center, Code 63 Department of Meteorology Naval Postgraduate School Monterey, California 93943-5100	1
4. Chairman, Code 63Rd Department of Meteorology Naval Postgraduate School Monterey, California 93943-5100	1
5. Chairman, Code 68Mr Department of Oceanography Naval Postgraduate School Monterey, California 93943-5100	1
6. Dr. Roger T. Williams Department of Meteorology Naval Postgraduate School Monterey, California 93943-5100	2
7. Director Naval Oceanography Division Naval Observatory 34th and Massachusetts Avenue NW Washington, D.C. 20390	1
8. Commander Naval Oceanography Command NSTL Station Bay St. Louis, Missouri 39522	1
9. Commanding Officer Naval Oceanographic Office NSTL Station Bay St. Louis, Missouri 39522	1

- | | | |
|-----|---|---|
| 10. | Commanding Officer
Fleet Numerical Oceanography Center
Monterey, California 93943-5100 | 1 |
| 11. | Commanding Officer
Naval Ocean Research and
Development Activity
NSTL Station
Bay St. Louis, Missouri 39522 | 1 |
| 12. | Commanding Officer
Naval Environmental Prediction
Research Facility
Monterey, California 93943-5100 | 1 |
| 13. | Chairman, Oceanography Department
U.S. Naval Academy
Annapolis, Maryland 21402 | 1 |
| 14. | Chief of Naval Research
800 N. Quincy Street
Arlington, Virginia 22217 | 1 |
| 15. | Office of Naval Research, Code 420
Naval Ocean Research and Development
Activity
NSTL Station
Bay St. Louis, Missouri 39522 | 1 |
| 16. | Commander (Air-370)
Naval Air Systems Command
Washington, D.C. 20360 | 1 |
| 17. | LT Deborah Ann Zankofski
USS Lexington (AVT 16)
Naval Air Station
Pensacola, Florida 32508 | 2 |
| 18. | Dr. C.-P Chang, Code 63Cp
Department of Meteorology
Naval Postgraduate School
Monterey, California 93943-5100 | 1 |
| 19. | Dr. R. L. Elsberry, Code 63Es
Department of Meteorology
Naval Postgraduate School
Monterey, California 93943-5100 | 1 |
| 20. | Dr. R. L. Haney, Code 63Hy
Department of Meteorology
Naval Postgraduate School
Monterey, California 93943-5100 | 1 |

- | | | |
|-----|---|---|
| 21. | Dr. B. J. Hoskins
Department of Geophysics
University of Reading
Reading, United Kingdom | 1 |
| 22. | Dr. E. C. Nickerson
NOAA, Atmospheric Physics &
Chemistry Laboratory
Boulder, Colorado 80302 | 1 |
| 23. | Dr. C. H. Wash, Code 63
Naval Postgraduate School
Monterey, California 93943-5100 | 1 |
| 24. | Dr. Andrew Staniforth
Recherche en Prevision Numerique
West Isle Office Tower, 5 ieme etage
2121 route Trans-Canada
Dorval, Quebec H9P1J3, Canada | 1 |
| 25. | Dr. J. Hovermale
Naval Environmental Prediction
Research Facility
Monterey, California 93943 | 1 |
| 26. | Dr. R. T. Pierrehumbert
Geophysical Fluid Dynamics Lab/NOAA
P.O. Box 308
Princeton, New Jersey 08540 | 1 |
| 27. | W. Blumen
Campus Box 391
University of Colorado
Boulder, Colorado 80309 | 1 |
| 28. | Dr. M. A. Rennick
Department of Meteorology
Naval Postgraduate School
Monterey, California 93943-5100 | 1 |
| 29. | Dr. M. Peng
Department of Meteorology
Naval Postgraduate School
Monterey, California 93943-5100 | 1 |
| 30. | Dr. C.-S. Liou, Code 63
Naval Postgraduate School
Monterey, California 93943-5100 | 1 |
| 31. | Dr. David Adamec, Code 63
Naval Postgraduate School
Monterey, California 93943-5100 | 1 |

32. Dr. R. B. Smith 1
Department of Geology and Geophysics
Yale University
P.O. Box 6666
New Haven, Connecticut 06511
33. Dr. Peter Bannon 1
Department of Geophysical Sciences
University of Chicago
Chicago, Illinois 60637
34. Dr. R. Gall, Code 63 1
Department of Meteorology
Naval Postgraduate School
Monterey, California 93943-5100
35. Mr. Lang C. Chou, Code 63 1
Department of Meteorology
Naval Postgraduate School
Monterey, California 93943-5100