



MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS - 1963 - A

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS
I. REPORT NUMBER	2. GOVT ACCESSION NO.	BEFORE COMPLETING FORM 3. RECIPIENT'S CATALOG NUMBER
5		
4. TITLE (and Subtitie)		S. TYPE OF REPORT & PERIOD COVE
SURFACE DYNAMICS OF RARE GAS	CRYSTALS	Interim Technical Repor
		6. PERFORMING ORG. REPORT NUMBE
7. AUTHORIE)		8. CONTRACT OR GRANT NUMBER(*)
K. D. Gibson and S. J. Sibener	<u>c</u>	N00014-77-C-0240
3. PERFORMING ORGANIZATION NAME AND The University of Chicago	ADDRESS	10. PROGRAM ELEMENT, PROJECT, TA AREA & WORK UNIT NUMBERS
The James Franck Institute 5640 South Ellis Avenue, Chica	ago, IL 60637	NR-392-023
1. CONTROLLING OFFICE NAME AND ADDR	[58	12. REPORT DATE
The Office of Naval Research		August, 1985
Physical Sciences Division (Co 800 North Oningy Str. Arling	oae 421) ton. VA 22217	13. NUMBER OF PAGES
14. NONITORING AGENCY NAME & AODRESS	It different from Controlling Office)	18. SECURITY CLASS. (of this report)
		unclassified
		154. DECLASSIFICATION DOWN GRADIN
10.15 document has been approved istribution is unlimited.	ed for public release a	Ind sale; its
Inis document has been approved distribution is unlimited. 17. DISTRIBUTION STATEMENT (of the observed) 18. SUPPLEMENTARY NOTES Prepared for Publication in J	ed for public release a te entered in Block 20, if different fre ournal of Vacuum Scient	The sale; its DTIC ELECTE OCT 2 1985 B ce and Technology.
19. SUPPLEMENTARY NOTES Prepared for Publication in J	ed for public release a t entered in Block 20, if different fre ournal of Vacuum Scient	Report) DTIC ELECTE OCT 2 1985 B ce and Technology.
In 15 document has been approved distribution is unlimited. 17. DISTRIBUTION STATEMENT (of the obstration 18. SUPPLEMENTARY NOTES Prepared for Publication in J Colored 19. Ally WORDS (Continue on reverse olds if new Surface phonon spectroscopy; films; inelastic single phonon surface energy exchange; rare	ed for public release a remered in Block 20, if different fre ournal of Vacuum Scient phonon dispersion curvi on scattering; physisor gas interaction poten	The sale; its The second seco
Inis document has been approved distribution is unlimited. 17. DISTRIBUTION STATEMENT (of the observed) 18. SUPPLEMENTARY NOTES Prepared for Publication in J Colored 19. AEY WORDS (Continue on reverse olds If new Surface phonon spectroscopy; films; inelastic single phono surface energy exchange; rare	ed for public release a t entered in Block 20, if different free ournal of Vacuum Scient phonon dispersion curvi on scattering; physisor e gas interaction poten	and sale; its Traperty DTIC ELECTE OCT 2 1985 B ce and Technology. Es; surface dynamics; thin ption interactions; gas- tials
 Inis document has been approved distribution is unlimited. 17. DISTRIBUTION STATEMENT (of the observed for Publication in J J J J J J J J J J J J J J J J J J	ed for public release a memory and location by block numbers phonon dispersion curve memory and location by block numbers phonon dispersion curve memory and location potent as interaction potent e gas crystals have be probe. Specifically, o and ke were grown on an physical properties o c diffraction and sele He-surface interaction culations with the expe and higher order terms	en explored using a He mole rdered monolayer, bilayer, Ag(11) substrate. This ha f these films evolve on a ctive adsorption experiment potential. Comparing the rimental data, both the in the energy expansion ca

unclassified

SECURITY CLASSIFICATION OF THIS PAGE (Then Date Entered)

be determined. To examine the dynamical properties of the adsorbed rare gases, angle and time resolved inelastic scattering was used. This technique made it possible to investigate the dynamics of the surface across the entire Brillouin zone. In this paper the properties of Ar(111) multilayers are specifically examined. These data are compared to the theoretical calculations of several other groups, testing the validity of several model potentials, and the importance of various interaction terms when constructing these potentials.

A COMPLETE COPY OF THE MANUSCRIPT WILL BE SENT UPON REQUEST.

S/N 0102- LF- 014- 6401

unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered

Surface Dynamics of Rare Gas Crystals

K. D. Gibson and S. J. Sibener⁺

Department of Chemistry and The James Franck Institute The University of Chicago 5640 South Ellis Avenue Chicago, IL 60637

Journal of Vacuum Science and Technology, in press

⁺Alfred P. Sloan Research Fellow PACS: 79.20.Rf; 63.20.Dj; 68.30.+z; 68.60.+q

ABSTRACT

The surface properties of rare gas crystals have been explored using a He molecular beam as the dynamical probe. Specifically, ordered monolayer, bilayer, trilayer, and bulk Ar, Kr, and Xe were grown on an Ag(111) substrate. This has allowed us to examine how the physical properties of these films evolve on a layer by layer basis. Elastic diffraction and selective adsorption experiments were used to investigate the He-surface interaction potential. Comparing the results of close-coupled calculations with the experimental data, both the laterally averaged potential and higher order terms in the energy expansion can be determined. To examine the dynamical properties of the adsorbed rare gases, angle and time resolved inelastic scattering was used. This technique made it possible to investigate the dynamics of the surface across the entire Brillouin zone. In this paper the properties of Ar(111) multilayers are specifically examined. These data are compared to the theoretical calculations of several other groups, testing the validity of several model potentials, and the importance of various interaction terms when constructing

these potentials. Originator Supplied Key a dosinclude:



Dric

In this paper we present preliminary results dealing with the surface dynamics of ordered rare gas overlayers physisorbed on an Ag(111) substrate. The overlayers examined were 1, 2, 3, and 25 layer (111) oriented films of Ar, Kr, and Xe. Experiments with physisorbed rare gas films on other metal surfaces have been reported by Mason and Williams [1]. However, in this paper we report for the first time, to our knowledge, how the surface phonon relations evolve on a layer-by-layer basis from monolayer to bulk across the Brillouin zone. Our experimental results were obtained using the angle resolved time of flight (TOF) of inelastically scattered He. Inelastic He scattering is a good technique for probing these weakly bound systems because of its surface sensitivity, excellent energy resolution (\sim 250µeV), and because the momentum of the incident beam is of the right order of magnitude to probe the entire Brillouin zone for the low energy modes of these materials. A complete description of the experimental procedure and apparatus will appear in forthcoming publications [2].

Thermodynamic and structural information for rare gas overlayers on Ag(111) was already available from LEED experiments [3,4] and theoretical calculations [5]. These form hexagonal structures that are azimuthally aligned but incommensurate with the surface. Overlayers were grown on a cryogenically cooled Ag(111) crystal with a low pressure dosing beam by controlling the crystal temperature, beam pressure, and dosing time. The ordering and lattice constants were checked using elastic He diffraction. The angular profiles obtained in these diffraction runs indicate that the coherence lengths of these overlayers are only slightly less than that of the Ag substrate (~ 100 Å). When an experiment on a particular overlayer was completed, the number of layers was confirmed by temperature programmed thermal desorption. By controlling the rate of the temperature ramp, the

monolayer desorption peak occurred separately from any of the multilayer peaks. Therefore, the coverage could be quantitatively determined by comparing the total integrated signal to that of the monolayer.

The data were analyzed by first fitting the TOF peaks with a nonlinear least squares routine. Examples of the data and the fits are shown in Figure 1. The arrow indicates the TOF of elastically scattered He. In general, there was always a peak at this position, which we attribute to incoherent scattering from crystal defects. The well resolved peaks show that with an 18 meV incident beam, single phonon events are the predominant inelastic scattering channel. To give an idea of the probabilities for inelastic scattering, the large energy loss peak of the monolayer has an intensity which is $\sim 1 \times 10^{-3}$ of the specular beam, and $\sim 1 \times 10^{-5}$ of the incident beam. The intensity of the principal modes observed do not vary by more than a factor of five from zone center to edge $(\Gamma \rightarrow M)$. The difference in the flight times for these peaks compared with the elastic TOF (determined at the specular angle) was used to determine the phonon energies. Using this information, and the incident and detector angles, the momentum exchange with the surface was calculated. Most of the experiments were done at an incident angle of 45 degrees, and the Brillouin zone was probed by varying the detector angle.

Figure 2 is a reduced zone plot of the dispersion relations for Ar in the $\overline{\Gamma} + \overline{M}$ direction. This figure clearly shows that inelastic He scattering experiments can be used to quantitatively probe how the surface vibrations of thin films evolve, on a layer-by-layer basis, from monolayer to bulk. This also indicates how the forces present at the surfaces of thin supported films vary with increasing distance of the surface from the substrate. The same qualitative features have also been seen for Kr and Xe.

The monolayer SP₁ mode (the only monolayer mode observed) is found to be dispersionless within experimental error. This indicates that the adsorbed atoms are behaving as independent Einstein oscillators, moving in an uncorrelated fashion in the surface potential well. Only the single phonon inelastic events corresponding to the energy loss and gain peaks adjacent to the elastic position in Figure 1 are plotted in Figure 2. This monolayer frequency gives the curvature at the equilibrium position, an essential piece of information for constructing the interaction potential. Temperature dependent studies show that the relative intensities of the single phonon creation and annihilation peaks vary in a manner consistent with statistical expectations. There are also two other peaks in the monolayer spectrum of Figure 1, corresponding to phonon creation, with energies twice and three times the fundamental. Due to the nearly harmonic nature of the rare gas-surface holding potential, we are at this time unable to determine the relative importance of overtone and multiphonon excitations in explaining these energy loss features. As further overlayers are added, the amount of dispersion increases as lateral interactions become systematically more important. Again, note that only the energy gain and loss features immediately adjacent to the elastic time of flight in Figure 1 were used in constructing the multilayer dispersion curves shown in Figure 2.

Lattice dynamics calculations have been carried out which well represent our experimental dispersion curves [6]. These calculations demonstrate the adequacy of using known gas phase pair potentials for modelling the interactions present in condensed rare gas films, and the importance of the substrate in modifying the phonon frequencies. Work is currently in progress to determine the inelastic scattering probabilities from the TOF spectra.

<u>Acknowledgments</u>: We would like to thank B. Hall, D. Mills, J. Black, M. Klein, and L. Bruch for their theoretical assistance. This work was supported in part by the ONR and the NSF-MRL at The University of Chicago. S. J. Sibener also acknowledges support from a Camille and Henry Dreyfus Foundation Young Faculty Grant.

References

- [1] B. F. Mason and B. R. Williams, Surf. Sci. 139 (1984) 173.
- [2] K. D. Gibson and S. J. Sibener, to be published.
- [3] J. Unguris, L. W. Bruch, E. R. Moog, and M. B. Webb, Surf. Sci. 87 (1979)
 415.
- [4] J. Unguris, L. W. Bruch, E. R. Moog and M. B. Webb, Surf. Sci. 109 (1981) 522.
- [5] L. W. Bruch, Surf. Sci. 125 (1983) 194.
- [6] K. D. Gibson, S. J. Sibener, B. Hall, D. L. Mills, and J. Black, J. Chem. Phys., to be published.

Figure Captions

- Figure 1 TOF spectra of the Ar overlayers studied at a surface temperature of 12 K and an incident beam energy of 18 meV. Crosses are the experimental data, solid lines are a least-squares fit of these data, and dashed lines are a computer simulation assuming single phonon processes and delta function phonon energy widths. Arrows indicate position of elastic incoherent scattering. The spectra were collected for 1 hour with 1 usec/channel resolution at an incident angle of 45 degrees. The detector angle was, from top to bottom, 42, 39, 40, and 50 degrees.
- Figure 2 Dispersion curves of the Ar overlayers from I-M. Open symbols are for energy loss of the beam, closed symbols are for energy gain. Representative error bars are shown for energy and momentum loss features (with respect to the He beam) which occurred in the first zone. Solid lines are polynomial fits to the data.





. .

FILMED

11-85

DTIC