

AD-A160 589

CHEMICAL APPLICATIONS OF TOPOLOGY AND GROUP THEORY 20
EIGHT-VERTEX POLYHEDRA (U) GEORGIA UNIV ATHENS DEPT OF
CHEMISTRY R B KING 14 OCT 85 TR-5 N00014-84-K-0365


1/1

UNCLASSIFIED

F/G 7/4

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

Unclassified

6

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 5	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) CHEMICAL APPLICATIONS OF TOPOLOGY AND GROUP THEORY. 20. EIGHT-VERTEX POLYHEDRA AND THEIR REARRANGEMENTS ¹		5. TYPE OF REPORT & PERIOD COVERED Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) R.B. King		8. CONTRACT OR GRANT NUMBER(s) N00014-84-K-0365
PERFORMING ORGANIZATION NAME AND ADDRESS University of Georgia Department of Chemistry Athens, GA 30602		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 051-861
CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, VA 22217		12. REPORT DATE October 14, 1985
1. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 21
		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES To be published in "Inorganic Chemistry"		DTIC SELECTED OCT 28 1985 B
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Polyhedra Diamond-square-diamond process Deltahedra Cube Rearrangements Eight-coordinate complexes		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) There are 257 combinatorially distinct eight-vertex polyhedra, of which 14 are deltahedra. Of the 14 deltahedra only the bisdisphenoid, commonly known as the D _{2d} or triangular dodecahedron, both lacks tetrahedral chambers and can be formed by the hybridization of only s, p, and d orbitals. Degenerate single and symmetrical parallel multiple diamond-square-diamond processes involving the 14 eight-vertex deltahedra are tabulated. Among the eleven such processes (six single, two symmetrical double, one fully symmetrical triple, and two fully (continued)		

AD-A160 589

DTIC FILE COPY

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Block #20.

symmetrical quadruple) those relating the bisdisphenoid to the 4,4-bicapped trigonal prism and square antiprism are of current chemical significance. Single, double, and quadruple dsd rearrangements of the bidisphenoid are depicted as topologically equivalent cubes with added diagonals so that the pivot faces are faces of the underlying cube and the dsd processes involve shifting only the positions of selected diagonals without disturbing the 12 edges of the underlying cube.

OFFICE OF NAVAL RESEARCH

Contract N00014-84-K-0365

Task No. NR 051-861

TECHNICAL REPORT NO. 5

Chemical Applications of Topology and Group Theory. 20

Eight-Vertex Polyhedra and their Rearrangements¹

by

R.B. King

Prepared for Publication

in

Inorganic Chemistry

**University of Georgia
Department of Chemistry
Athens, Georgia**

October 14, 1985

**Reproduction in whole or in part is permitted for
any purpose of the United States Government**

**This document has been approved for public release
and sale; its distribution is unlimited.**

111

85 10 25 039

Abstract

There are 257 combinatorially distinct eight-vertex polyhedra, of which 14 are deltahedra. Of the 14 deltahedra only the bisdisphenoid, commonly known as the D_{2d} or triangular dodecahedron, both lacks tetrahedral chambers and can be formed by the hybridization of only s, p, and d orbitals. Degenerate single and symmetrical parallel multiple diamond-square-diamond processes involving the 14 eight-vertex deltahedra are tabulated. Among the eleven such processes (six single, two symmetrical double, one fully symmetrical triple, and two fully symmetrical quadruple) those relating the bisdisphenoid to the 4,4-bicapped trigonal prism and square antiprism are of current chemical significance. Single, double, and quadruple dsd rearrangements of the bisdisphenoid are depicted as topologically equivalent cubes with added diagonals so that the pivot faces are faces of the underlying cube and the dsd processes involve shifting only the positions of selected diagonals without disturbing the 12 edges of the underlying cube.

Structure, Isomerism, Molecular Structure, Isomerism



PER CALL JC



A-1

in

1. Introduction

Previous papers in this series have discussed new mathematical approaches for the treatment of stereochemical non-rigidity in ML_n coordination complexes (M = central atom, generally a metal; L = ligands surrounding M) and metal clusters. Thus Gale transformations² allow all possible non-planar isomerization processes to be found for polyhedra having five and six vertices, i.e., corresponding to ML_5 and ML_6 complexes.³ This Gale transformation approach is no longer effective for polyhedra having seven or more vertices, since Gale transforms no longer reduce the dimensionality of the problem.⁴ However, the chemically based assumption⁴ of minimum pivot face size in intermediate polyhedra coupled with the still manageable number of combinatorially distinct seven-vertex polyhedra, namely 34 (ref. 5), allows an exhaustive study of chemically relevant diamond-square-diamond (dsd) processes⁶ in seven-vertex polyhedra.⁴


This paper presents a related treatment of polyhedral isomerizations in eight-coordinate complexes. Here the problem is considerably more complicated since there are 257 combinatorially distinct eight-vertex polyhedra (Table 1).⁵ In order to make the problem tractable, the following two assumptions are used:

- (1) Only dsd processes (i.e., 4-pyramidal rather than n -pyramidal ($n \geq 5$) processes³) having a quadrilateral pivot face⁴ will be considered. This assumption is justified on the energetic basis that polyhedra having one or more faces with five or more edges (i.e., pentagons, hexagons, heptagons, etc.) are unfavorable relative to polyhedra having only triangular and quadrilateral faces. This principle was already recognized in 1969 in the first paper of this series.⁷
- (2) Only symmetrical parallel multiple dsd processes will be considered. In a symmetrical multiple dsd process the quadrilateral pivot faces are equivalent because of the symmetry of the intermediate polyhedron.

In addition the dsd processes in only the 14 combinatorially distinct eight-vertex deltahedra (Table 2) are considered. In general, minimum energy coordination polyhedra are deltahedra or polyhedra derived from deltahedra through low energy processes. Furthermore, this paper shows that the two eight-vertex non-deltahedra of greatest significance as coordination polyhedra, namely the 4,4-bicapped trigonal prism and the square antiprism,^{8,9,10,11} appear as intermediate polyhedra in single and double dsd processes of the bisdisphenoid ("D_{2d}-dodecahedron"), the only one of the 14 eight-vertex deltahedra (Table 2) found in ML₈ coordination complexes not involving f orbital elements.^{8,9,10,11}

2. Background

Many aspects of the methods used in this paper resemble those used in the previous paper on seven-coordinate complexes.⁴ Thus we consider sequences of polyhedral isomerizations $P_1 \rightarrow P_2 \rightarrow P_3$ in which polyhedra P_1 and P_3 are combinatorially equivalent, i.e., $P_1 \rightarrow P_2 \rightarrow P_3$ is a degenerate polyhedral isomerization. The intermediate polyhedron P_2 has fewer edges than P_1 (or P_3). Since only dsd polyhedral isomerizations are considered, the polyhedral sequence $P_1 \rightarrow P_2 \rightarrow P_3$ has the following structure at the quadrilateral pivot face of P_2 :



Thus P_1 and P_3 can be generated from P_2 by drawing diagonals in two different ways across the quadrilateral pivot face of P_2 . In the case of symmetrical multiple dsd processes, the intermediate polyhedron P_2 has two or more quadrilateral pivot

faces equivalent by symmetry and the multiple diagonals are drawn across these faces to preserve the symmetry elements making these faces equivalent.


Using this method the problem of finding all possible degenerate dsd polyhedral isomerizations of eight-vertex deltahedra can be reduced to the following:

- (1) Find all eight-vertex polyhedra having only triangular faces except for a single quadrilateral face or several symmetrically equivalent quadrilateral faces. These polyhedra correspond to the intermediate polyhedron P_2 of the sequence (1) above.
- (2) Draw new polyhedra by adding diagonals in both ways across the quadrilateral pivot face of P_2 to generate the polyhedra P_1 and P_3 . If polyhedra P_1 and P_3 are combinatorially equivalent, then a degenerate eight-vertex deltahedral isomerization has been found. In the case of an intermediate polyhedron P_2 having two or more symmetrically equivalent quadrilateral faces, the diagonals to form P_1 and P_3 must be drawn so as to preserve the element of symmetry making equivalent the two or more quadrilateral faces of P_2 .

Federico⁵ has tabulated all combinatorially distinct polyhedra having eight faces. These are the duals of the desired polyhedra having eight vertices where a dual P^* of P is obtained as follows¹²:

- (1) The vertices of P^* are located at the midpoints of the faces of P .
- (2) Two vertices of P^* are connected by an edge if and only if the corresponding faces of P share an edge.

The Schlegel diagrams¹³ in Federico's paper are, of course, for the duals of the eight-vertex polyhedra of interest but can be used for this work if the dsd process of the sequence (1) is expressed in dual form, i.e.,



In the dual transformation of sequence (1) into sequence (2) the quadrilateral pivot face of P_2 becomes a degree four vertex in P_2^* and the two new triangular faces in P_1 and P_3 generated by diagonals across this quadrilateral face become pairs of degree three vertices in P_1^* and P_3^* .


3. Properties of the Eight-Vertex Polyhedra

Elementary topological considerations derivable from Euler's theorem⁷ show that eight-vertex polyhedra must have at least six faces but not more than twelve faces. The 257 combinatorially distinct eight-vertex polyhedra can be divided into 18 classes (Table 1) so that all members of a given class have the same set of polygon types for faces, i.e., the same face index.⁴ Of these 18 classes those classes having one or more faces with five or more edges (i.e., the classes C, E, F, H, I, J, K, M, N, O, and R in Table 1 containing a total of 74 polyhedra) are unfavorable energetically and are therefore excluded from further consideration in this paper. The remaining seven classes (A, B, D, G, L, P, and Q in Figure 1) correspond to the 183 eight-vertex polyhedra containing only triangular and quadrilateral faces. These are the polyhedra potentially involved in the rearrangements discussed in this paper.

The eight-vertex deltahedra are of particular chemical interest. Table 2 lists the 14 combinatorially distinct eight-vertex deltahedra according to the size of the largest cavity or (equivalently in this case) the number of tetrahedral chambers. Note that there are only two eight-vertex deltahedra without tetrahedral chambers, the bisdisphenoid (" D_{2d} dodecahedron") frequently occurring as an eight-vertex coordination polyhedron^{8,9,10,11} and the hexagonal bipyramid which is a "forbidden" eight-vertex polyhedron since the inversion center in its symmetry point group prevents it from being formed by the hybridization of only s, p, and d orbitals.¹⁴

The special role of the bisdisphenoid in eight-coordinate metal complexes is already apparent.

The 12 remaining eight-vertex deltahedra, i.e., those having one or more tetrahedral chambers, follow the expected pattern. Since all ten (triangular) faces of the pentagonal bipyramid are equivalent, there is only one distinct capped pentagonal bipyramid (Federico number 56 of its dual). The four eight-vertex deltahedra having octahedral cavities can be best considered as their duals, namely cubes in which two vertices are truncated. The two vertices being truncated can be at each ends of an edge (ortho: Federico number 51 of its dual), a face-diagonal (meta: Federico number 55 of its dual), or a body diagonal (para: Federico number 57 of its dual).



The fourth eight-vertex deltahedron having an octahedral cavity (Federico number 53 of its dual) has two layers of capping. First, one of the eight equivalent faces of an octahedron is capped giving three equivalent new triangular faces in place of the one face that was capped. A second cap is then placed on one of the three new triangular faces giving a "capped capped octahedron," i.e., a polyhedron with two levels of capping. Finally Table 2 summarizes the properties of the seven combinatorially distinct eight-vertex deltahedra formed by fusing five tetrahedra. One of these, the "tetracapped tetrahedron" (Federico number 49 of its dual), is distinctive in having relatively high T_d symmetry and only one level of capping relative to the central tetrahedron.

4. Rearrangements of Eight-Vertex Polyhedra

Table 3 summarizes all possible symmetrical dsd rearrangements of eight-vertex deltahedra in terms of the intermediate polyhedra (P_2 in sequence 1) containing the quadrilateral pivot faces. Note that only a small number of these dsd rearrangements are degenerate, i.e., have combinatorially equivalent starting and finishing deltahedra. The possible number of symmetrical dsd rearrangements becomes smaller as a larger number of symmetrically equivalent quadrilateral pivot faces become involved in parallel processes.

Table 3 indicates six possible degenerate single dsd processes for eight-vertex deltahedra. By far the most interesting such process is the degenerate dsd rearrangement of the chemically significant bisdisphenoid (Federico number 58 of its dual) through a 4,4-bicapped trigonal prism intermediate (Federico number 96 of its dual). The 4,4-bicapped trigonal prism intermediate polyhedron occasionally appears in eight-coordinate structures (e.g., terbium (III) chloride).¹⁵ The remaining five degenerate dsd processes involve eight-vertex deltahedra of low symmetry (C_1 , C_2 , and C_s) either consisting of five fused tetrahedra (Federico numbers 46, 47, and 52 of their duals) or the bilevel capped capped octahedron (Federico number 53 of its dual). These processes do not appear to be particularly significant chemically in eight-coordinate derivatives because of the unlikely polyhedra involved.

Table 3 indicates two possible degenerate symmetrical double dsd rearrangements of eight-vertex deltahedra. The chemically interesting such process involves the degenerate double dsd rearrangement of the bisdisphenoid through a square antiprism intermediate. The square antiprism is the eight-vertex non-deltahedron most frequently found in coordination compounds.^{8,9,10,11} Also the bisdisphenoid-square antiprism-bisdisphenoid double dsd rearrangement is included in a topological

representation of hyperoctahedrally restricted eight-coordinate polyhedral rearrangements discussed several years ago.¹⁶

Symmetrical triple dsd rearrangements are necessarily rare since a C_3 axis must be preserved throughout such processes. The single example of a degenerate eight-vertex such process (Table 3) involves rearrangement of a bicapped octahedron (Federico number 57 of its dual) through a 3,3-bicapped trigonal prism intermediate (Federico number 245 of its dual) in a process resembling a Bailer twist¹⁷ but with the two capping vertices not directly participating in the rearrangement.

Symmetrical quadruple dsd processes are more complicated since there are different symmetry-preserving ways of drawing diagonals across the four quadrilateral faces of the three eight-vertex polyhedra having four quadrilateral and four triangular faces and D_{2d} and D_{2h} symmetry (Federico numbers 282, 287, and 288 of their duals). Two generic degenerate symmetrical quadruple dsd processes are listed in Table 3. Again one of these processes involves the bisdisphenoid this time undergoing a degenerate rearrangement through one of the two polyhedral intermediates having four triangular and four quadrilateral faces (Federico numbers 282 or 288 of their duals).

Figure 1 depicts topologically single (top), double (middle), and quadruple (bottom) dsd processes of the bisdisphenoid representing all of the relevant eight-vertex polyhedra as topologically equivalent cubes with added diagonals. There are two such presentations of the bisdisphenoid as a cube with an added diagonal in each of the six square faces: the tetrahedral presentation (e.g., Figure 1: top left, center left, and center right) in which the four degree 5 vertices of the bisdisphenoid (circled in Figure 1) form the vertices of a tetrahedron and the rectangular presentation (e.g., Figure 1: top right, bottom left, and bottom right) in which the four degree 5 vertices of the bisdisphenoid are coplanar and form the vertices of a rectangle. Presentations of the bisdisphenoid are selected for

a given dsd process so that the pivot face(s) correspond to faces on the underlying cube and the dsd process involves shifting only the positions of selected diagonals without disturbing the 12 edges of the underlying cube.

The top rearrangement in Figure 1 is a single dsd rearrangement of the bisdisphenoid (Federico number 58 of its dual) through a 4,4-bicapped trigonal prism (Federico number 96 of its dual) intermediate using the front face of the underlying cube as the pivot face. This process interchanges the tetrahedral and rectangular presentations of the bisdisphenoid. The middle rearrangement in Figure 1 is a double dsd rearrangement of the bisdisphenoid through a square antiprism (Federico number 172 of its dual) intermediate using the front and back faces of the underlying cube as the two pivot faces. This process involves the tetrahedral presentation of the bisdisphenoid. The bottom rearrangement in Figure 1 is a quadruple dsd rearrangement of the bisdisphenoid through a polyhedron (Federico number 282 of its dual) having four triangular faces, four quadrilateral faces, four vertices of degree 4, four vertices of degree 3, and D_{2h} symmetry using the left, top, right, and bottom faces of the underlying cube as the four pivot faces. This process involves the rectangular presentation of the bisdisphenoid. The other quadruple dsd rearrangement of the bisdisphenoid through the D_{2d} gyrobifastigium (Federico number 288 of its dual) listed in Table 3 cannot be depicted using either the tetrahedral or rectangular presentations of the bisdisphenoid so that a face of the underlying cube is the pivot face and only diagonals are shifted.

The three types of dsd rearrangements of the bisdisphenoid depicted in Figure 1 represent all symmetrical possibilities involving the tetrahedral and rectangular presentations of the bisdisphenoid. The single dsd rearrangement through the 4,4-bicapped trigonal prism (Figure 1: top) interchanges the two presentations. The double dsd rearrangement through the square antiprism (Figure 1: middle)

requires the tetrahedral presentation but the quadruple dsd rearrangement (Figure 1: bottom) requires the rectangular presentation for both the starting and finishing bisdisphenoids in order for the pivot face of the intermediate polyhedron to be a face of the underlying cube. Thus a factor affecting the relative ease of the various dsd rearrangements of the bisdisphenoid can be the degree of distortion of an actual bisdisphenoid to conform to the presentations (i.e., tetrahedral or rectangular) required for the pivot face of the dsd rearrangement to be a face of an underlying cube. In this connection the double dsd process involving a square antiprism intermediate (Figure 1: middle) should be the most favorable since less distortion of an actual bisdisphenoid is required to give the tetrahedral presentation than the rectangular presentation on a cube because in an actual bisdisphenoid the four degree 4 vertices and the four degree 5 vertices are located at the vertices of subtetrahedra.⁸

5. Conclusions

This paper shows how treatment of polyhedral isomerizations in systems having increasing numbers of vertices requires increasingly restrictive assumptions in order to keep the number of possibilities tractable. Thus for five- and six-vertex polyhedra the dimensionality reduction afforded by Gale transformations³ allows all non-trivial (i.e., non-planar) rearrangements to be found without any restrictive assumptions. For seven-vertex polyhedra, the smallest number of vertices for which Gale transformation does not offer dimensionality reduction, the assumption of no pentagonal (or hexagonal) faces suggested by energetic considerations (thereby restricting polyhedral rearrangements to dsd processes) is sufficient to reduce the possibilities to a manageable number.⁴ The eight-vertex polyhedra treated

in this paper require symmetry considerations in addition to energetic considerations in order to select from the intractable number of possibilities those of greatest chemical interest. Extension of this work to polyhedra having nine or more vertices not only goes beyond the scope of Federico's tables of polyhedra⁵ but is also unattractive because of the intractably large number of possibilities and decreasing relevance to real chemical systems.

The analysis in this paper also underscores the special role of the bisdisphenoid (popularly known as the "D_{2d} dodecahedron") in eight-coordinate structures. The bisdisphenoid is the only eight-vertex deltahedron which has no tetrahedral chambers and which can be formed from sp³d⁴ hybrid orbitals. In addition, the bisdisphenoid is unique among eight-vertex deltahedra in undergoing single, symmetrical double, and symmetrical quadruple dsd processes as depicted in Figure 1. Furthermore, the intermediate (non-deltahedral) polyhedra in the single and symmetrical double dsd processes of the bisdisphenoid, namely the 4,4-bicapped trigonal prism and square antiprism, respectively, also occur in eight-coordinate complexes.^{8,9,10,11} For these reasons the bisdisphenoid has as fundamental a role in the chemistry of eight-coordinate complexes as the octahedron has in the chemistry of six-coordinate complexes. However, the bisdisphenoid is highly fluxional whereas the octahedron is relatively rigid.¹⁸

Acknowledgment: I am indebted to the Office of Naval Research for partial support of this work.

LITERATURE REFERENCES

- (1) For part 19 of this series see King, R. B.; Rouvray, D. H. Theor. Chim. Acta in press.
- (2) Grünbaum, B. "Convex Polytopes." Interscience, New York, 1967: Chapter 6.
- (3) King, R. B. Theor. Chim. Acta **1984**, 64, 439.
- (4) King, R. B. Inorg. Chem. **1985**, 24, 1716.
- (5) Federico, P. J. Geometriae Dedicata **1975**, 3, 469.
- (6) Lipscomb, W. N. Science **1966**, 153, 373.
- (7) King, R. B. J. Am. Chem. Soc. **1969**, 91, 7211.
- (8) Hoard, J. L.; Silverton, J. V. Inorg. Chem. **1963**, 2, 235.
- (9) Muetterties, E. L.; Wright, C. M. Quart. Revs. **1967**, 21, 109.
- (10) Lippard, S. J. Prog. Inorg. Chem. **1967**, 8, 109.
- (11) Kepert, D. L. Prog. Inorg. Chem. **1978**, 24, 179.
- (12) Reference 2, pp. 46-48.
- (13) Reference 2, pp. 42-46.
- (14) King, R. B. Theor. Chim. Acta **1984**, 64, 453.
- (15) Forrester, J. D.; Zalkin, A.; Templeton, D. H.; Wallmann, J. C. Inorg. Chem. **1964**, 3, 185.
- (16) King, R. B. Theor. Chim. Acta **1981**, 59, 25.
- (17) Bailar, J. C., Jr. J. Inorg. Nucl. Chem. **1958**, 8, 165.
- (18) King, R. B. Inorg. Chim. Acta **1981**, 49, 237.

TABLE 1
 THE 257 COMBINATORICALLY DISTINCT
 EIGHT-VERTEX POLYHEDRA DIVIDED INTO 18 CLASSES^a

<u>v</u>	<u>e</u>	<u>f</u>	Face Types					Number of Polyhedra	Class
			<u>f₇</u>	<u>f₆</u>	<u>f₅</u>	<u>f₄(=q)</u>	<u>f₃(=t)</u>		
8	18	12	0	0	0	0	12	14	A
8	17	11	0	0	0	1	10	38	B
8	16	10	0	0	1	0	9	12	C
8	16	10	0	0	0	2	8	64	D
8	15	9	0	1	0	0	8	3	E
8	15	9	0	0	1	1	7	24	F
8	15	9	0	0	0	3	6	47	G
8	14	8	1	0	0	0	7	1	H
8	14	8	0	1	0	1	6	2	I
8	14	8	0	0	1	2	5	20	J
8	14	8	0	0	2	0	6	2	K
8	14	8	0	0	0	4	4	17	L
8	13	7	0	1	0	2	4	2	M
8	13	7	0	0	1	3	3	5	N
8	13	7	0	0	2	1	4	2	O
8	13	7	0	0	0	5	2	2	P
8	12	6	0	0	0	6	0	1	Q
8	12	6	0	0	2	2	2	1	R

a) v=number of vertices; e=number of edges; f=number of faces; f_n=number of faces having n edges; f₃=t (triangle) and f₄=q (quadrilateral) for clarity.

TABLE 2

THE 14 COMBINATORICALLY DISTINCT EIGHT-VERTEX DELTAHEDRA

Federico Number of Duala	Symmetry	Vertex Types					Symmetrical dsd Levels ^b	Description ^b
		v7	v6	v5	v4	v3		
<u>A) No Tetrahedral Chambers</u>								
#58	D _{2d}	0	0	4	4	0	1,2,4	bisdiphenoid ("D _{2d} dodecahedron")
#54	D _{6h}	0	2	0	6	0	0	hexagonal bipyramid
<u>B) Pentagonal Bipyramidal Central Cavity: One Tetrahedral Chamber</u>								
#56	C _s	0	1	3	3	1	0	capped pentagonal bipyramid
<u>C) Octahedral Central Cavity: Two Tetrahedral Chambers</u>								
#57	D _{3d}	0	0	6	0	2	3	para-bicapped octahedron
#55	C _{2v}	0	1	4	1	2	0	meta-bicapped octahedron
#51	C _{2v}	0	2	2	2	2	1,4	ortho-bicapped octahedron
#53	C _s	0	2	1	4	1	1	capped capped octhedron (two levels of capping)
<u>D) No Cavity Larger Than a Tetrahedron (Bicapped Bicapped Tetrahedra): Five Tetrahedral Chambers</u>								
#45	C _{2v}	2	0	0	4	2	0	two 355 outer caps
#46	C ₁	1	1	2	1	3	1	445 + 355 outer caps
#47	C ₁	1	1	1	3	2	1	345 + 355 outer caps

TABLE 2 (Continued)

Federico Number of Duala	Symmetry	Vertex Types				Symmetrical dsd Levels ^b	Description ^b	
		v7	v6	v5	v4			v3
#48	C _s	1	0	3	2	2	0	two 345 outer caps
#49	T _d	0	4	0	0	4	0	two 445 outer caps (symmetrically tetracapped tetrahedron)
#50	C _s	0	3	1	1	3	0	345 + 445 outer caps
#52	C ₂	0	2	2	2	2	1,2	two 345 outer caps

a) See Federico, P.J. Geom. Ded. 1975, 3, 469.

b) See text for a description of the nomenclature.

TABLE 3
 SYMMETRICAL DSD-PROCESSES OF EIGHT-VERTEX
 DELTAHEDRA

<u>Intermediate Polyhedron^a</u>	<u>Deltahedra Formed by Diagonalization^a</u>	<u>Remarks</u>
<u>A) Single dsd Processes</u>		
#59 (11033-C ₁)	#45 (20042-C _{2v}) and #46 (11213-C ₁)	
#60 (11033-C ₁)	#45 (20042-C _{2v}) and #47 (11132-C ₁)	
#61 (10304-C ₅)	#46 (11213-C ₁)	degenerate
#62 (10223-C ₁)	#47(11132-C ₁) and #48 (10322-C ₅)	
#63 (10223-C ₁)	#46 (11213-C ₁) and #47 (11132-C ₁)	
#64 (10223-C ₅)	#46 (11213-C ₁)	degenerate
#65 (10142-C ₁)	#47 (11132-C ₁) and #48 (10322-C ₅)	
#66 (02204-C ₁)	#46 (11213-C ₁) and #50 (03113-C ₅)	
#67 (02204-C _{2v})	#49 (04004-T _d) and #51 (02222-C _{2v})	
#68 (02123-C ₁)	#46 (11213-C ₁) and #51 (02222-C _{2v})	
#69 (02123-C ₁)	#46 (11213-C ₁) and #53 (02141-C ₅)	
#70 (02123-C ₁)	#47 (11132-C ₁) and #50 (03113-C ₅)	
#71 (02123-C ₅)	#47 (11132-C ₁)	degenerate
#72 (02123-C ₁)	#47 (11132-C ₁) and #52 (02222-C ₂)	
#73 (02123-C ₅)	#50 (03113-C ₅) and #53 (02141-C ₅)	
#74 (02042-C _{2v})	#45 (20042-C _{2v}) and #54 (02060-D _{6h})	
#75 (02042-C ₁)	#47 (11132-C ₁) and #53 (02141-C ₅)	
#76 (01313-C ₁)	#46 (11213-C ₁) and #56 (01331-C ₅)	
#77 (01313-C ₁)	#48 (10322-C ₅) and #52 (02222-C ₂)	

TABLE 3 (Continued)

<u>Intermediate Polyhedron^a</u>	<u>Deltahedra Formed by Diagonalization^a</u>	<u>Remarks</u>
#78 (01313-C ₁)	#50 (03113-C ₃) and #55 (01412-C _{2v})	
#79 (01313-C ₂)	#50 (03113-C ₃) and #52 (02222-C ₂)	
#80 (01232-C ₁)	#47 (11132-C ₁) and #56 (01331-C _{2v})	
#81 (01232-C ₂)	#48 (10322-C ₃) and #55 (01412-C _{2v})	
#82 (01232-C ₁)	#48 (10322-C ₃) and #53 (02141-C ₃)	
#83 (01232-C ₃)	#53 (02141-C ₃)	degenerate
#84 (01232-C ₁)	#51 (02222-C _{2v}) and #56 (01331-C ₃)	
#85 (01232-C ₃)	#55 (01412-C _{2v}) and #56 (01331-C ₃)	
#86 (01232-C ₃)	#52 (02222-C ₂)	degenerate
#87 (01232-C ₁)	#52 (02222-C ₂) and #53 (02141-C ₃)	
#88 (01151-C ₃)	#54 (02060-D _{6h}) and #56 (01331-C ₃)	
#89 (01151-C ₁)	#53 (02141-C ₃) and #56 (01331-C ₃)	
#90 (00422-C _{2v})	#51(02222-C _{2v}) and #58 (00440-D _{2d})	
#91 (00422-C ₁)	#55 (01412-C _{2v}) and #56 (01331-C ₃)	
#92 (00422-C ₃)	#56 (01331-C ₃) and #57 (00602-D _{3d})	
#93 (00422-C ₂)	#52 (02222-C ₂) and #57 (00602-D _{3d})	
#94 (00341-C ₃)	#53 (02141-C ₃) and #58 (00440-D _{2d})	
#95 (00341-C ₃)	#56 (01331-C ₃) and #58 (00440-D _{2d})	
#96 (00260-C _{2v})	#58 (00440-D _{2d})	degenerate ^b

TABLE 3 (Continued)


<u>Intermediate Polyhedron^a</u>	<u>Deltahedra Formed by Diagonalization^a</u>	<u>Remarks</u>
<u>B) Symmetrical Double dsd Processes</u>		
#99 (10124-C ₅)	#45 (20042-C _{2v}) and #48 (10322-C ₅)	
#103 (02024-C ₂)	#45 (20042-C _{2v}) and #51 (02222-C _{2v})	
#105 (02024-C ₂)	#45 (20042-C _{2v}) and #52 (02222-C ₂)	
#117 (01214-C ₅)	#49 (04004-T _d) and #56 (01331-C ₅)	
#136 (01052-C _{2v})	#54 (02060-D _{6h}) and #55 (01412-C _{2v})	
#137 (01052-C ₅)	#51 (02222-C _{2v}) and #54 (02060-D _{6h})	
#139 (00404-C _{2v})	#52 (02222-C ₂)	degenerate
#140 (00404-D _{2d})	#49 (04004-T _d) and #58 (00440-D _{2d})	
#151 (00323-C ₅)	#50 (03113-C ₅) and #56 (01331-C ₅)	
#152 (00323-C ₅)	#51 (02222-C _{2v}) and #57 (00602-D _{3d})	
#159 (00242-C ₂)	#51 (02222-C _{2v}) and #58 (00440-D _{2d})	
#161 (00242-C ₂)	#54 (02060-D _{6h}) and #58 (00440-D _{2d})	
#162 (00242-C ₅)	#54 (02060-D _{6h}) and #55 (01412-C _{2v})	
#163 (00242-C _{2v})	#54 (02060-D _{6h}) and #57 (00602-D _{3d})	
#165 (00242-C ₅)	#53 (02141-C ₅) and #55 (01412-C _{2v})	
#167 (00242-C ₂)	#52 (02222-C ₂) and #57 (00602-D _{3d})	
#168 (00242-C ₂)	#57 (00602-D _{3d}) and #58 (00440-D _{2d})	
#171 (00161-C ₅)	#56 (01331-C ₅) and #58 (00440-D _{2d})	
#172 (00080-D _{4d})	#58 (00440-D _{2d})	degenerate ^c

TABLE 3 (continued)


<u>Intermediate Polyhedron^a</u>	<u>Deltahedra Formed by Diagonalization^a</u>	<u>Remarks</u>
<u>C) Symmetrical Triple dsd Processes</u>		
#191 (01034-C _{3v})	#49 (04004-T _d) and #54 (02060-D _{6h})	
#194 (00305-C _{3v})	#49 (04004-T _d) and #57 (00602-D _{3d})	
#245 (00062-D _{3h})	#57 (00602-D _{3d})	degenerate
<u>D) Symmetrical Quadruple dsd Processes^d</u>		
#282 (00044-D _{2h})	#51 (02222-C _{2v}) or #58 (00440-D _{2d})	degenerate (for #51 or #58)
#287 (00044-D _{2d})	#49(04004-T _d) and #58 (00440-D _{2d})	
#288 (00044-D _{2d}) (gyrobifastigium)	#58 (00440-D _{2d}) or #51 (02222-C _{2v})	degenerate (for #51 or #58)

- a) The numbers of the polyhedra correspond to the numbers of their dual polyhedra given in Table 1 of Federico, P.J. Geometriae Dedicata 1975, 3, 469. The vertex index $v_7v_6v_5v_4v_3$ and the symmetry point group are given in parentheses after the polyhedron dual number.
- b) This corresponds to the bisdisphenoid-4,4-bicapped trigonal prism-bisdisphenoid single dsd degenerate rearrangement.
- c) This corresponds to the bisdisphenoid-square antiprism-bisdisphenoid symmetrical double dsd degenerate rearrangement.
- d) Symmetrical quadruple dsd processes are more complicated since there are different symmetry-preserving ways of drawing diagonals across the four quadrilateral faces of polyhedra #282, #287, and #288.


Figure 1: Single (top), double (middle), and quadruple (bottom) dsd processes of the bisdisphenoid depicted as topologically equivalent cubes with added diagonals so that the pivot face(s) are faces of the underlying cube and the dsd processes involve shifting only the positions of selected diagonals without disturbing the 12 edges of the underlying cube. Degree 5 vertices are circled and degree 3 vertices are starred.




#58
Bisdisphenoid
(tetrahedral presentation)




#96
4,4-Bicapped
Trigonal Prism




#58
Bisdisphenoid
(rectangular presentation)




#58
Bisdisphenoid
(tetrahedral presentation)




#172
Square Antiprism




#58
Bisdisphenoid
(tetrahedral presentation)



#58
Bisdisphenoid
(rectangular presentation)



#282



#58
Bisdisphenoid
(rectangular presentation)

END

FILMED

12-85

DTIC