

MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963 A



National Défense
Defence nationale

UNCLASSIFIED



UNLIMITED
DISTRIBUTION

DRES

SUFFIELD MEMORANDUM

NO. 1150

AD-A164 573

INVESTIGATION OF T-2 MYCOTOXIN-INDUCED
CYTOTOXICITY *IN VITRO* AND PROTECTIVE
EFFECTS OF FLAVONOID COMPOUNDS (U)

by

DTIC
ELECTE
FEB 19 1986
S D

R.J.F. Markham, V.L. DiNinno, N.P. Erhardt,
D. Penman and A.R. Bhatti

Project No. 16A10

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

January 1986

DTIC FILE COPY



DEFENCE RESEARCH ESTABLISHMENT SUFFIELD, RALSTON, ALBERTA

WARNING
The use of this information is restricted without
recognition of proprietary and patent rights.

FEB - 6 1986

Canada

96 2 18 126

UNCLASSIFIED

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD
RALSTON, ALBERTA

SUFFIELD MEMORANDUM NO. 1150

INVESTIGATION OF T-2 MYCOTOXIN-INDUCED CYTOTOXICITY IN VITRO
AND PROTECTIVE EFFECTS OF FLAVONOID COMPOUNDS

by

R.J.F. Markham, V.L. DiNinno, N.P. Erhardt, D. Penman and A.R. Bhatti



16A10

WARNING
The use of this information is permitted only for recognition
of proprietary and patent rights.

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

UNCLASSIFIED

UNCLASSIFIED

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD
RALSTON, ALBERTA

SUFFIELD MEMORANDUM NO. 1150

INVESTIGATION OF T-2 MYCOTOXIN-INDUCED CYTOTOXICITY IN VITRO
AND PROTECTIVE EFFECTS OF FLAVONOID COMPOUNDS

by

R.J.F. Markham, V.L. DiNinno, N.P. Erhardt, D. Penman and A.R. Bhatti

ABSTRACT

Aspects of the in vitro cytotoxic effects of T-2 mycotoxin on murine thymocytes were investigated. Cytotoxicity was found to be consistent when tested bi-weekly for a four month period. Cytotoxicity reached maximal values over a narrow range of doses and was dependent on temperature (37°C optimum) and number of cells in the reaction mixtures. Quercetin, a flavonoid compound was able to decrease the effect of T-2 toxin when the drug was added within an hour of mixing the T-2 toxin with the thymocytes.

UNCLASSIFIED

UNCLASSIFIED

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD
RALSTON, ALBERTA

SUFFIELD MEMORANDUM NO. 1150

INVESTIGATION OF T-2 MYCOTOXIN-INDUCED CYTOTOXICITY IN VITRO
AND PROTECTIVE EFFECTS OF FLAVONOID COMPOUNDS

by

R.J.F. Markham, V.L. DiNinno, N.P. Erhardt, D. Penman and A.R. Bhatti

INTRODUCTION

Trichothecene mycotoxins are a chemically related group of secondary metabolites of *Fusarium* and some other fungal species which have been shown to be toxic in both man and animals. Mycotoxins have been implicated as the cause of inadvertent food intoxication following fungal contamination of foodstuffs (1-3) and recent studies have suggested their deliberate use as biological warfare agents (4, 5). Many organ systems in the body can be

UNCLASSIFIED

affected by this toxin through its radiomimetic effect i.e. its ability to inhibit protein, RNA and DNA synthesis in actively growing cells (6). Hence, a major target of its action is the haematopoietic system resulting in anemia, leukopenia and immunological aberrations (reviewed in 7). To date there are no effective drugs or treatment regimens for this type of poisoning.

Previous work from this laboratory has shown that mycotoxins, in particular T-2 toxin, have potent cytotoxic capabilities in vitro expressed as a loss of viability of thymocytes and other leukocytes (8). The intent of the present communication is to suggest possible mechanisms of this cytotoxicity and to describe compounds which offer some protection against T-2 induced cytotoxicity.

MATERIALS AND METHODS

Animals

C-57 black male mice, 3-6 weeks of age, were used in all but one experiment, in which case mice up to 24 weeks of age were used. They were housed and cared for in accordance with guidelines prescribed by the Canadian Council on Animal Care.

Collection of Cells and Culture

Isolated thymus cells were prepared as described by Mischell and Shiigi (9). Briefly, mice were sacrificed by CO₂ inhalation and the thorax was flooded with 70% alcohol prior to removal of the thymus. For each assay at least two thymuses were collected. After removal, the thymuses were placed in a petri dish containing Joklik's media (Flow Laboratories, McLean, Virginia) and minced with scissors to dissociate cells. The cell suspension was washed by centrifugation and the pellet was resuspended in the same medium to give a concentration of 2×10^7 cells/ml.

Cytotoxicity Assay

T-2 toxin and other mycotoxins (Sigma Chemical Company) were dissolved in dimethylsulphoxide (DMSO) to give a stock solution of 20 mg/ml. Further dilution in DMSO allowed addition of specified amounts of mycotoxin to 0.5 ml of cell suspension (1×10^7 cells). The amount of DMSO added did not exceed 25 μ l and controls, containing cells and DMSO alone, were tested to detect any inherent cytotoxicity of DMSO.

Cytotoxicity was assessed by determining viability of thymocytes using dye exclusion with 0.2% eosin (9). At specified times after addition of T-2 toxin to the cells, an aliquot of the mixture was removed and mixed with an equal volume of eosin solution. Cells were examined microscopically using a Neubauer hemocytometer and viability of at least 200 cells was determined.

Quercetin or other flavonoid compounds were dissolved in DMSO and added to the T-2/thymocyte mixture at specified times and concentrations. Controls were included to determine if quercetin and additional DMSO had a cytotoxic effect on the thymocytes.

RESULTS

Figure 1 shows the results of 8 separate experiments performed at 2 week intervals for 14 weeks. Despite possible variation between mice of the same and different ages and week to week variation there was a consistent pattern of cytotoxicity induced by T-2 toxin. For the controls, the mean percentages ($n = 8$) of viable cells at 0, 2 and 4 hours of incubation were 89.4 ± 5 , 93.0 ± 5 and 91.0 ± 6 , respectively. In the presence of T-2 toxin the mean percentage of viable cells at these time points were 84.9 ± 6 , 36.3 ± 10 and 6.6 ± 4 .

Previous work performed in this laboratory determined that 400 $\mu\text{g/ml}$ T-2 mixed with 1×10^7 cells could produce significant cytotoxicity in the thymocytes (8). To further define the dose response relationship of T-2 toxin cytotoxicity, thymocytes were incubated with a lower range of doses of T-2 toxin. Figure 2 indicates that T-2 toxin has a narrow dose response curve. Cytotoxicity is evident at 175 $\mu\text{g/ml}$ and reaches maximal levels at 200 $\mu\text{g/ml}$. Similarly, increasing the numbers of cells decreased the cytotoxicity induced by a constant amount of T-2 toxin (Figure 3) suggesting that a critical concentration is required before cytotoxicity can be expressed.

The temperature dependence of T-2 induced cytotoxicity is shown in Figure 4. Full expression of cytotoxicity was not apparent until the incubation temperature was raised to 37°C.

T-2 proved to be the most toxic of the mycotoxins tested in this system (Figure 5) and subsequently was used in the evaluation of protective compounds.

Addition of flavonoid compounds afforded protection against T-2 induced cytotoxicity. In Figure 6A, quercetin is shown to give a significant reduction of 44% in the cytotoxicity induced by 400 $\mu\text{g/ml}$ T-2 toxin. In separate experiments and with a different preparation of T-2 toxin, quercetin was able to decrease the cytotoxic effect of 200 $\mu\text{g/ml}$ T-2 by 37% (Figure 6B). Quercetin was most effective when added before or at the time of addition of the T-2 toxin to the thymocytes. Delaying the addition of quercetin by 2 hours significantly reduced its protective effect (Figure 7).

Quercetin was active over a range of concentrations (Table 1) with an apparent decrease in effect with increasing concentration. This may be due to an observed cytotoxic effect of the drug itself (Figure 8).

DISCUSSION

Although the toxic effects of T-2 toxin and other trichothecene mycotoxins have been known for some time, it is only recently that research has focused on the cellular and molecular basis of this toxicity. It is important that this knowledge be obtained as it will allow for a more logical and fruitful search for compounds which may be able to counteract the effects of T-2 toxin.

It has been suggested that inhibition of protein synthesis and, to a lesser degree, inhibition of DNA and RNA synthesis is responsible for the observed toxic effects of T-2 toxin (10, 11). More recent studies, however, have focused on possible membrane effects as mechanisms of cytotoxicity. Ethanol, detergents and other membrane modifying agents increase sensitivity of yeast to T-2 toxin as measured by inhibition of growth (12). Decreased temperatures, known to decrease membrane fluidity, also decrease the effect of T-2 toxin. It was also found that a yeast with reduced plasma membrane fluidity was resistant to T-2 toxin. Additional evidence for a major role of membranes in T-2 toxin induced cytotoxicity comes from studies of mycoplasma (13). Although T-2 toxin inhibited growth of the organism, no changes in gross protein DNA or RNA synthesis could be observed although precursors of these macromolecules were inhibited from crossing the cell membrane into the cytosol.

Membrane changes have also been implicated in T-2 toxin induced damage in eukaryotic cells. Hemolysis of red blood cells has been studied and an effect on the cell membrane has been described (14, 15). Hemolysis was evident over a narrow dose range and was temperature dependent. It was concluded that T-2 toxin induced membrane lesions of less than 5.5 Å which precipitated osmotic lysis.

Our studies suggest that interaction of T-2 toxin with thymocyte membranes is involved in the observed cytotoxicity. The narrow range of doses which distinguish between cytotoxicity and non-cytotoxicity (as observed in red cells) is similar to the effect of certain anesthetics (16, 17) and polyene antibiotics (18). These agents are incorporated into cell membranes up to a critical point after which damage quickly ensues. Their ability to intercollate into the membrane structure is likely responsible for their action. Likewise, due to the amphipathic nature of the T-2 toxin molecule (15), it too may have membrane reactivity and be able to bind to hydrophobic sites in that structure. T-2 toxin has been shown to bind to the membranes of thymocytes although the exact nature of the receptor was not identified (20). The dependence on temperature for cytotoxicity supports the concept that membranes could be involved as fluidity is dramatically altered by decreased temperature. The fact that T-2 toxin is the most active of the mycotoxins tested adds further evidence for the membrane being targets for its action. The HT-2 toxin has hydroxyl group at C4, whereas the T-2 toxin has a acetyl group at this position. It may be that increasing the hydrophobicity of the T-2 toxin molecule in this manner allowed better membrane interaction. A reported age difference in the susceptibility to T-2 toxin cytotoxicity in mice (19) may reside in membrane differences. Although we could not see any age effects between recently weaned and older mice, there may have been a difference if cells from newborn animals had been used.

Quercetin (3,3',4,5,7-pentahydroxyflavone) is among a large number of flavonoid compounds commonly found in a variety of fruits and vegetables. Flavonoid compounds have been reported to have substantial biological activity and have been used as antitumor drugs, heart stimulants, diuretics, antivirals and antihistaminics (21). Some of these compounds have been shown to protective against whole body irradiation which is significant with the knowledge of the radiomimetic effects of the mycotoxins (22). Quercetin itself has been investigated for its ability to reduce histamine release from mast cells as well as other anti-inflammatory properties (23). Quercetin, in

our experiments, clearly had a protective effect against T-2 toxin induced cytotoxicity. It was necessary for quercetin to be present during the early stages for full protection to be manifested.

There is no clear evidence of how these flavonoids, including quercetin, exert these effects. Quercetin has been shown to decrease calcium transport across membranes (24), inhibit NADPH oxidase in neutrophils (25) and decrease spontaneous lipid peroxidation following glutathione depletion in rats (26). Other flavonoids have similar antioxidant activity (27). From evidence implicating participation of cell membranes in the cytotoxicity, it may be possible that activities relating to inhibition of lipid peroxidation or cellular enzyme activation may be involved. This protection does not appear to be a general property of the flavonoids as related compounds do not possess the same protective effect as quercetin (data not shown). Rutin, in particular, a precursor molecule of quercetin, shows little activity. This molecule is far more hydrophilic than is quercetin and it may be that the interaction of quercetin with membranes in a competitive manner may be a mechanism of protection against membrane active T-2 toxin. Similar structure/function relationships have been found for other biological activities of flavonoid compounds (28).

It is imperative that the mechanisms of cytotoxicity of mycotoxins be elucidated so appropriate countermeasures can be devised.

REFERENCES

1. Hsu, I.C., Smalley, E.B., Strong, F.M. and Ribelin, W.E. Identification of T-2 toxin in moldy corn associated with lethal toxicosis in dairy cattle. *Appl. Microbiol.* 24: 684-690, 1972.
2. Greenway, J.A. and Puls, R. Fusariotoxicosis from barley in British Columbia I. Natural occurrence and diagnosis. *Can. J. Comp. Med.* 40: 12-15, 1976.
3. Ueno, Y., Ishii, K., Sakai, K., et al. Toxicological approaches to the metabolites of *Fusarium*. IV. Microbial survey on bean-hulls poisoning of horses with the isolation of toxic trichothecenes neosolaniol and T-2 toxin of *Fusarium solani*. *Jpn. J. Exp. Med.* 42: 187-203, 1972.
4. Rosen, R.T. and Rosen, J.D. Synthetic material in "yellow rain": Evidence for the use of chemical weapons in Laos. *Biomed. Mass. Spectrom.* 9: 443-450, 1982.
5. Mirocha, C.J., Pawlosky, R.A., Chatterjee, K., Watson, S. and Hayes, W. Analysis for *Fusarium* toxins in various samples implicated in biological warfare in Southeast Asia. *J. Assoc. Off. Anal. Chem.* 66: 1485-1499, 1983.
6. Saito, M., Enomoto, M. and Tatsuno, T. Radiomimetic biological properties of new scripine metabolites of *Fusarium nivale*. *Gann.* 60: 599-603, 1969.
7. Protection Against Trichothecene Mycotoxins. National Academy Press, Washington, D.C., 1983.

8. DiNinno, V.L., Penman, D., Bhatti, A.R., Erhardt, N.P. and Lockwood, P.A. An in vitro method for the screening of trichothecene mycotoxin cytotoxicity and the study of mycotoxin countermeasures. Suffield Memorandum No. 1147, 1985.
9. Mishell, B.B. and Shiigi, S.M. Selected Methods in Cellular Immunology, W.H. Freeman and Company, San Francisco, p. 3-27, 1980.
10. Ueno, Y., Nakajima, K., Sakai, K., Ishii, K., Sato, N., and Shimada, N. Comparative toxicology of trichothecene mycotoxins: inhibition of protein synthesis in animal cells. J. Biochem. 74: 285-296, 1973.
11. Oldham, J.W., Allred, L.E., Milo, G.E., Kindigo, O., and Capen, C.C. The toxicological evaluation of the mycotoxins T-2 and T-2 tetrol using normal human fibroblasts in vitro. Toxicol. Appl. Pharmacol. 52: 159-168, 1980.
12. Schappert, K.T. and Khachatourians, G.G. Influence of the membrane on T-2 toxicity in *Saccharomyces* spp. Appl. Environ. Microbiol. 47: 681-684, 1984.
13. Rottem, S., Yagen, B. and Katznel, A. Effect of trichothecenes on growth and intracellular pool size of *Mycoplasma gallisepticum*. FEBS Lett. 175: 189-192, 1984.
14. Segal, R., Milo-Goldzweig, I., Joffe, A.Z., and Yagen, B. Trichothecene-induced hemolysis. The hemolytic activity of T-2 toxin. Toxicol. Appl. Pharmacol. 70: 343-349, 1983.
15. Gyongyossy-Issa, M.I.C., Khanna, V., and Khachatourians, G.G.

- Characterization of hemolysis induced by T-2 toxin. *Biochim. Biophys. Acta* 838: 252-256, 1985.
16. Sheetz, M.P. and Singer, M.P. Biological membranes and bilayer couples. A molecular mechanism of drug erythrocyte interactions. *Proc. Natl. Acad. Sci.* 71: 4457-4461, 1974.
 17. Kwant, W.O. and Seeman, P. The membrane concentration of a local anesthetic (Chlorpromazine). *Biochim. Biophys. Acta* 183: 530-543, 1969.
 18. Kinsey, S.C., Avruch, J., Permutt, M. and Rogers, H.B. The lytic effect of polyene antifungal antibiotics on mammalian erythrocytes. *Biochem. Biophys. Res. Commun.* 9: 503-507, 1962.
 19. Ueno, Y. Toxicological features of T-2 toxin and related trichothecenes. *Fund. Appl. Toxicol.* 4: s124-s134, 1984.
 20. Gyongyossy-Issa, M.I.C. and Khachatourians, G.G. Interaction of T-2 toxin with murine lymphocytes. *Biochim. Biophys. Acta* 803: 197-202, 1984.
 21. Willaman, J.J. Some biological effects of the flavonoids. *J. Amer. Pharm. Assoc.* 44: 404-408, 1955.
 22. Agarwal, O.P. and Nagaratnam, A. Radioprotective property of flavonoids in mice. *Toxicol.* 19: 201-204, 1981.
 23. Middleton, E., Drzewiecki, G., and Krishnarao, D. Quercetin: An inhibitor of antigen induced human basophil release. *J. Immunol.* 127: 546-510, 1985.

24. Barzilai, A., and Rahamimoff, H. Inhibition of Ca^{++} transport ATPase from synaptosomal vesicles by flavonoids. *Biochim. Biophys. Acta* 730 245-254, 1983.
25. Tauber, A.I., Fay, J.R. and Marletta, M.A. Flavonoid inhibition of human neutrophil NADPH-oxidase. *Biochem. Pharmacol.* 33: 1367-1369, 1984.
26. Younes, M. and Siegers, C.P. Inhibitory action of some flavonoids on enhanced spontaneous lipid peroxidation following glutathione depletion. *Planta. Medica.* 43: 240-244, 1981.
27. Lambev, I., Belcheva, A., and Zhelyazkov, D. Flavonoids with anti-oxidant action (Naringin and Rutin) and the release of mastocytic and nonmastocytic histamine. *Acta Physiol. Pharmacol. (Bulgaria)* 6: 70-75, 1980.
28. Beretz, A., Cazenave, J.P. and Anton, R. Inhibition of aggregation and secretion of human platelets by quercetin and other flavonoids. Structure-activity relationships. *Agents and Actions* 12: 382-387, 1982.

UNCLASSIFIED

TABLE 1 Relation Between Concentration and Protective Effect of Quercetin
on Cytotoxicity of T-2 Toxin on Murine Thymocytes.

Concentration of Quercetin ($\mu\text{g/ml}$)	Protection* (%)
5	29 \pm 6**
10	37 \pm 4
30	33 \pm 4
50	23

* Present reduction in observed cytotoxicity of T-2 toxin

** Data represents mean \pm 1 standard deviation of at least three
(3) separate trials

UNCLASSIFIED




Figure 1

Longitudinal studies of the cytotoxicity of T-2 toxin on murine thymocytes. ● control cells incubated with DMSO; ○ cells incubated with 200 µg/mL T-2 toxin. Thymus cells from two mice were used in each experiment. Mean values ± 1 standard deviation for control cells and T-2 treated cells at 0, 2 and 4 hour are 89.4 ± 5, 93.0 ± 5, 91.0 ± 6 and 84.9 ± 36.3 ± 10 and 6.6 ± 4 respectively.




Figure 2

Effect of concentration on the cytotoxicity of T-2 toxin on murine thymocytes.
● cytotoxicity measured at 1 hr; ○ cytotoxicity measured at 2 hr; ▲ cytotoxicity measured at 5 hr. Cells representing a pool of 3 mice were treated with different amounts of T-2 toxin.




Figure 3

Effect of cell number on the cytotoxicity of T-2 toxin on murine thymocytes. ○ 2 x 10⁷ cells/mL; ● 4 x 10⁷ cells/mL; □ 6 x 10⁷ cells/mL. All cell preparations were incubated with 200 µg/mL T-2 toxin. Cells representing a pool of 3 mice were used.




Figure 4

Effect of temperature on the cytotoxicity of T-2 toxin on murine thymocytes.
● control cells incubated with DMSO; ○ cells incubated with 200 µg/mL T-2 toxin. Cells representing a pool of 3 mice were incubated at the different temperatures.




Figure 5

Comparative cytotoxicity of various mycotoxins on murine thymocytes. ● control cells incubated with DMSO; ○ cells incubated with T-2; ▼ HT-2; △ T-2 triol; □ T-2 tetrol. Cells representing a pool of 3 mice were incubated with 400 µg/mL of each of the compounds.




Figure 6




Figure 7

The effect of time of addition of quercetin on protection from cytotoxicity of T-2 toxin on murine thymocytes. Cells were mixed with 200 $\mu\text{g}/\text{mL}$ T-2 toxin at 0 hr. 10 $\mu\text{g}/\text{mL}$ quercetin was added at times indicated. Cytotoxicity Assay was performed 5 hours after addition of T-2 toxin. Except for -2 hour, which was a single trial, data represents the mean \pm 1 standard deviation of at least 3 separate trials.




Figure 8

Effect of quercetin on murine thymocytes. ○ control cells; cells incubated with □ 5 µg/mL; △ 10 µg/mL; ▼ 50 µg/mL quercetin. Data represents the mean \pm 1 standard deviation of at least 3 separate trials.

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall document is classified)

1. ORIGINATING ACTIVITY Defence Research Establishment Suffield		2a. DOCUMENT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP	
3. DOCUMENT TITLE Investigation of T-2 Mycotoxin-Induced Cytotoxicity <u>In Vitro</u> and Protective Effects of Flavonoid Compounds			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) SM-1150			
5. AUTHOR(S) (Last name, first name, middle initial) R.J.F. Markham, V.L. DiNinno, N.P. Erhardt, D. Penman and A.R. Bhatti			
6. DOCUMENT DATE January 1986		7a. TOTAL NO. OF PAGES 21	7b. NO. OF REFS 28
8a. PROJECT OR GRANT NO. 16A-10		9a. ORIGINATOR'S DOCUMENT NUMBER(S) SM1150	
8b. CONTRACT NO.		9b. OTHER DOCUMENT NO.(S) (Any other numbers that may be assigned this document)	
10. DISTRIBUTION STATEMENT Unlimited			
11. SUPPLEMENTARY NOTES N/A		12. SPONSORING ACTIVITY N/A	
13. ABSTRACT Aspects of the <u>in vitro</u> cytotoxic effects of T-2 mycotoxin on murine thymocytes were investigated. Cytotoxicity was found to be consistent when tested bi-weekly for a four month period. Cytotoxicity was found to be consistent when over a narrow range of doses and was dependent on temperature (37°C optimum) and number of cells in the reaction mixtures. Quercetin, a flavonoid compound was able to decrease the effect of T-2 toxin when the drug was added within an hour of mixing of T-2 toxin with the thymocytes.			

KEY WORDS

T-2 Mycotoxin
 Quecetin
 Flavonoids

INSTRUCTIONS

1. ORIGINATING ACTIVITY. Enter the name and address of the organization issuing the document.
2. DOCUMENT SECURITY CLASSIFICATION. Enter the overall security classification of the document including special warning terms whenever applicable.
3. GROUP. Enter security reclassification group number. The three groups are defined in Appendix 'M' of the DRB Security Regulations.
4. DOCUMENT TITLE. Enter the complete document title in all capital letters. Titles in all cases should be unclassified. If a sufficiently descriptive title cannot be selected without classification, show title classification with the usual one capital-letter abbreviation in parentheses immediately following the title.
5. DESCRIPTIVE NOTES. Enter the category of document, e.g. technical report, technical note or technical letter. If appropriate, enter the type of document, e.g. interim, progress, summary, annual or final. Give the inclusive dates when a specific reporting period is covered.
6. AUTHOR(S). Enter the name(s) of author(s) as shown on or in the document. Enter last name, first name, middle initial, if military, show rank. The name of the principal author is an absolute minimum requirement.
7. DOCUMENT DATE. Enter the date (month, year) of Exhibit Board approval for publication of the document.
8. TOTAL NUMBER OF PAGES. The total page count should follow normal pagination procedures, i.e. enter the number of pages containing information.
9. NUMBER OF REFERENCES. Enter the total number of references cited in the document.
10. PROJECT OR GRANT NUMBER. If appropriate, enter the applicable research and development project or grant number under which the document was written.
11. CONTRACT NUMBER. If appropriate, enter the applicable order number under which the document was written.
12. ORIGINATOR'S DOCUMENT NUMBER(S). Enter the official document number by which the document will be identified and controlled by the originating activity. This number must be unique to this document.
- 9b. OTHER DOCUMENT NUMBER(S). If the document has been assigned any other document numbers (either by the originator or by the sponsor), also enter this number(s).
10. DISTRIBUTION STATEMENT. Enter any limitations on further dissemination of the document, other than those imposed by security classification, using standard statements such as:
 - (1) "Qualified requesters may obtain copies of this document from their defence documentation center."
 - (2) "Announcement and dissemination of this document is not authorized without prior approval from originating activity."
11. SUPPLEMENTARY NOTES. Use for additional explanatory notes.
12. SPONSORING ACTIVITY. Enter the name of the departmental project office or laboratory sponsoring the research and development. Include address.
13. ABSTRACT. Enter an abstract giving a brief and factual summary of the document, even though it may also appear elsewhere in the body of the document itself. It is highly desirable that the abstract of classified documents be unclassified. Each paragraph of the abstract shall end with an indication of the security classification of the information in the paragraph (unless the document itself is unclassified) represented as (TS), (S), (C), (R), or (U).

The length of the abstract should be limited to 20 single-spaced standard typewritten lines 7 1/4 inches long.
14. KEY WORDS. Key words are technically meaningful terms or short phrases that characterize a document and could be helpful in cataloging the document. Key words should be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context.

END

FILMED

386

DTIC