

AD-A164 813

STRUCTURAL CHEMISTRY OF ENERGETIC COMPOUNDS(U) OFFICE
OF NAVAL RESEARCH ARLINGTON VA R GILARDI ET AL.


1/2


81 OCT 85

UNCLASSIFIED

F/G 7/4

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

(2)

AD-A164 813

Structural Chemistry of Energetic Compounds

ONR Annual Report

Contract No. N00014-85-WR-24060

1 October 1984 - 1 October 1985

Richard Gilardi, Clifford George and J. L. Flippin-Anderson

Laboratory for the Structure of Matter

Naval Research Laboratory, Washington, D.C. 20375

DTIC FILE COPY



86 2 26 025

This document has been approved
for public release and its
distribution is unlimited.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER N001485WR24060-1985A	2. GOVT ACCESSION NO. <i>AD-A164 813</i>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Structural Chemistry of Energetic Compounds	5. TYPE OF REPORT & PERIOD COVERED Annual Progress Report Oct. 1984-Oct. 1985	
7. AUTHOR(s) Richard Gilardi, Clifford George and Judith Flippen-Anderson	6. PERFORMING ORG. REPORT NUMBER N0001485WR24060	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Office of Naval Research Code 432 Arlington, VA 22217	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS <i>(16)</i> 61153N-RR024-02-01	
11. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE	
	13. NUMBER OF PAGES	
14. MONITORING AGENCY NAME & ADDRESS(if different from Controlling Office)	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
	15a. DECLASSIFICATION/ DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Reproduction in whole or in part is permitted for any purpose of the United States Government.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) X-ray diffraction, tetraazabicyclooctanes, oxetane, nitramines, nitro-adamantane, Nitric Radicals, Benzyl radicals, Tetraoxoles, Cyclic compounds		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Structural characterization of 16 energetic materials and precursors was performed by x-ray diffraction. The materials examined include: (CONTINUED REVERSE SIDE)		

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

- (a) 1-Nitroso-2,2-di(trifluoromethyl)-1,3-diazacyclopentan-4-one
 (b) 3,3,7,7,-Tetra(trifluoromethyl)-2,4,6,8-tetraazabicyclo[3.3.0]octane
 (c) 2,6-Dinitro-3,3,7,7-tetra(trifluoromethyl)-2,4,6,8-tetraazabicyclo[3.3.0]octane
 (d) 2,4,6-Trinitro-3,3,7,7-tetra(trifluoromethyl)-2,4,6,8-tetraazabicyclo[3.3.0]octane
 (e) 2,4,6,8-Tetrานитро-3,3,7,7-tetra(trifluoromethyl)-2,4,6,8-tetraazabicyclo[3.3.0]octane
 (f) 3-Benzamido-6,8-dibenzoyl-1,3,5,6,8-pentaazabicyclo[3.2.2]nonane
 (g) 1,3,5-Tribenzamido-1,3,5-hexahydrotriazine monohydrate
 (h) 1-Benzoyl-4-benzamido-4²-1,2,4-triazoline
 (i) 2,7-Dibenzoyl-4-benzamido-1,2,4,6,7-pentaazabicyclo[1.2.4]nonane
 (j) bis-Tetrazolo[1,5-a:1',5'-c]pyrazine
 (k) 1,4-Dinitro-2,3,5,6-tetrabromo-2,3,5,6-piperazine
 (l) 1,9-Diacetyl-3,5,7-trinitro-1,3,5,7-tetraaza-9-oxanonane
 (m) 1,9-Diacetyl-3,5,7-trinitro-1,3,5,7-9-pentaazanonane
 (n) 2-Oxa-6,9-diaza-6,9-dinitrospiro[3.6]decane
 (o) 1-Aza-3,5,7-trinitroadamantane
 (p) 7-Nitro-5,6-dihydro-7H-imidazolo[1,2-d]tetrazole












- Cont'd






→ The trinitro and tetrานитро tetraazabicyclo[3.3.0]octanes examined, compounds D and E, are significantly more dense than had been predicted. The crystal structure analysis showed that the increase in density is due to intramolecular crowding rather than extraordinary packing efficiencies which were at normal values of 0.73 and 0.75 respectively. Kly (W) 1/1

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unpublished	<input type="checkbox"/>
Journal Article	<input type="checkbox"/>
Event	<input type="checkbox"/>
Product	<input type="checkbox"/>
Process	<input type="checkbox"/>
Design	<input type="checkbox"/>
A-1	

Table of Contents

	Page
1. Structural Diagrams	1
2. Synopsis of Results	3
3. Individual Structure Reports	7
(a) 1-Nitroso-2,2-di(trifluoromethyl)-1,3-diazacyclopentan-4-one	7
(b) 3,3,7,7,-Tetra(trifluoromethyl)-2,4,6,8-tetraazabicyclo[3.3.0]octane	10
(c) 2,6-Dinitro-3,3,7,7-tetra(trifluoromethyl)-2,4,6,8-tetraazabicyclo[3.3.0]octane	12
(d) 2,4,6-Trinitro-3,3,7,7-tetra(trifluoromethyl)-2,4,6,8-tetraazabicyclo[3.3.0]octane	14
(e) 2,4,6,8-Tetranitro-3,3,7,7,-tetra(trifluoromethyl)-2,4,6,8-tetraazabicyclo[3.3.0]octane	16
(f) 3-Benzamido-6,8-dibenzoyl-1,3,5,6,8-pentaazabicyclo[3.2.2]nonane	18
(g) 1,3,5-Tribenzamido-1,3,5-hexahydrotriazine monohydrate	20
(h) 1-Benzoyl-4-benzamido- Δ^2 -1,2,4-triazoline	22
(i) 2,7-Dibenzoyl-4-benzamido-1,2,4,6,7-pentaazabicyclo[1.2.4]nonane	24
(j) bis-Tetrazolo[1,5-a:1',5'-c]pyrazine	26
(k) 1,4-Dinitro-2,3,5,6-tetrabromo-2,3,5,6-piperazine ...	29
(l) 1,9-Diacetyl-3,5,7-trinitro-1,3,5,7-tetraaza-9-oxanonane	32
(m) 1,9-Diacetyl-3,5,7-trinitro-1,3,5,7-9-pentaazanonane	34
(n) 2-Oxa-6,9-diaza-6,9-dinitrospiro[3.6]decane	36
(o) 1-Aza-3,5,7-trinitroadamantane	38
(p) 7-Nitro-5,6-dihydro-7H-imidazolo[1,2-d]tetrazole	40
4. Appendix: Bond distances, angles, and coordinates for FY85 crystal structure analyses	42





Synopsis of Results.

Molecules A through E and J and K were obtained from the Naval Surface Weapons Center (NSWC). Of these A, J, and K are either unexpected reaction products or synthetic intermediates which were produced in their energetic synthesis program. Although these three compounds are not themselves energetic materials, the identification of their structures and stereochemistry may aid in understanding the mechanisms of the reactions involved, and thus indirectly further the ONR synthesis program.

The series of tetraazabicyclooctanes, molecules B through E, are very interesting molecules closely related to bicyclo-HMX. In fact, compound E is bicyclo-HMX substituted with four trifluoromethyl groups; for certain applications, E may prove to be a useful explosive or fuel without further modification.

These four compounds are dense; each of the molecules has a density greater than the stable form of HMX. The crystallographic densities are 2.03, 1.98, 2.11, and 2.18 gm/cm³ for B through E respectively. The densities of D and E are anomalously high with respect to densities derived from Holdens (NSWC) density prediction program, 2.02 and 2.04 gm/cm³ respectively. Examination of the x-ray results show the source of the greater than expected density is intra-molecular crowding, not unusually close packing. All of the intermolecular contacts are greater than the usual close-packing limit (the sum of the van der Waals' radii of the atoms), except for a rather weak NH^{...}NO₂ hydrogen bond in D. The packing efficiency, which is the actual volume of the molecules in a cell divided by the volume of the cell, is near 0.75 for most crystals. For molecules

B through E, the packing efficiencies are 0.73, 0.68, 0.73, and 0.75, which are normal to low values.

The particular contacts which show severe crowding are the nitro-CF₃ contacts. This crowding also produces a number of distortions in the molecules (see table below). As one adds nitro groups to the parent compound, the bonds to the nitro and CF₃ groups lengthen and the N-C-N angles sharpen; in addition, the CF₃ group torsions deviate by almost 30° from a staggered configuration. All of these effects serve to lengthen the non-bonded CF₃-nitro distances; however, in compound E there are still 12 contacts which are extremely short (more than 0.1 Å below normal van der Waals separations).

Selected Bond Distances, Angles, and Torsions

	Zero-NO ₂	Di-NO ₂	Tri-NO ₂	Tetra-NO ₂
C-CF ₃ (Å)	1.529	1.544	1.575	1.582
N-NC ₂ (Å)	-	1.386	1.405	1.413
N-C-N	105.6°	99.5°	99.7°	98.5°
C-C-C-F	177.3°	173.2°	161.7°	152.0°
N-O(Å)	-	1.217	1.206	1.207
C-F(Å)	1.324	1.298	1.310	1.323
Ring Torsions	14.0°	13.9°	12.2°	20.7°

The strain caused by short internal contacts makes the synthetic chemistry of bicyclo-HMK and its derivatives very difficult and unpredictable; however, this internal strain adds to its total energy content, and makes it an even more desirable energetic target compound.

Molecules F, G, H and I were synthesized by Drs. J. Boyer and G. Kumar of the University of Illinois at Chicago, in a systematic study

of the heterocyclic compounds produced by hydrazine/formaldehyde/orthoformate condensation and rearrangement reactions. The specific aim of this program of study is the synthesis of a "tri-hydrazine" analog of adamantane [2,3,5,7,8,9-hexaazatricyclo(3.3.1.1^{3,7})decane]. This compound, if derivatized with nitro or nitramino groups, is expected to be a dense ($d > 2.1$) explosive. Two new hetero-N cage compounds (F and I) were produced in FY85 by Boyer and Kumar and their structures were characterized by our x-ray diffraction analyses; energetic substitution has not yet been accomplished. Compound G is a precursor heterocycle from which F and I were made; compound H is a minor, lower molecular weight product formed in the same reaction which produces I.

Crystals of both H & I were found in the crystalline reaction product.

Molecules L, M, and N were provided by C. Coon of Lawrence Livermore National Laboratories. Compounds L and M are energetic precursor molecules whose crystal structures were done primarily to corroborate their structural formulas. Compound N is a new energetic monomer, similar to 3,3-bis(methylnitroaminomethyl)oxetane, (BMNAMO). BMNAMO can be polymerized to form stable energetic homo- or co-polymers.

Molecule (O) is a novel monoazaadamantane synthesized by A. Nielsen of the Naval Weapons Center (China Lake). Of the four tertiary carbons in the adamantane cage, three are nitro-substituted and the other is replaced by a nitrogen atom. It is an energetic material, but its density (1.604 mg mm^{-3}) is much less than HMX; further nitro-substitution is theoretically possible but is considered to be unlikely.


Compound P was supplied by R. L. Willer of Morton Thiokol. This energetic molecule has the property of decomposing without going

through a melt phase.

The remainder of this report is a series of brief technical reports on each individual molecule and the x-ray experiment condidions.

Fractional coordinates are provided for those intending further computational study; these are 'crystallographic' coordinates and must be converted to Cartesian Angstrom coordinates for some programs. Contact the authors if this creates a problem. Bond distances and angles are provided for all molecules; if specific torsion angles or non-bonded distances are required, they are on file at the Naval Research Laboratory (NRL) and available upon request.

1-Nitroso-2,2-di(trifluoromethyl)-1,3-diazacyclopentan-4-one,



Abstract

$M_r = 251.088$, orthorhombic, $Pna2_1$, $a = 10.214(2)$, $b = 10.986(3)$,
 $c = 7.614(2)\text{\AA}$, $V = 854.4(3)\text{\AA}^3$, $Z = 4$, $D_x = 1.952 \text{ mg mm}^{-3}$, $\text{CuK}\alpha$,
 $\lambda = 1.54178\text{\AA}$, $\mu = 2.12 \text{ mm}^{-1}$, $F_{000} = 496$, $T = 295\text{K}$, $R = .0511$ and $R_w =$
.0609 for 524 observed reflections, $S = 1.295$. Crystals volatile, sublime
completely in a day; rapid data collection used to obtain unique set in
three hours.

Experimental

Clear, colorless needle-shaped crystals provided by W. Koppes of
the Naval Surface Weapons Center (Silver Spring, MD), grown by sublima-
tion at $90^\circ\text{C}/200 \text{ mm Hg}$. Data crystal ($0.10 \times 0.20 \times 0.70 \text{ mm}$) centered
on 25 reflections with $33^\circ < 2\theta < 67^\circ$ to obtain refined lattice parameters.
Nicolet R3m diffractometer used, incident-beam graphite monochromator,
 $\theta/2\theta$ scans, scan width = $(1.8 + \Delta_{\text{alc2}})^\circ$, $2\theta_{\text{max}} = 110^\circ$. In view
of volatility of crystals and scarcity of material, rapid data collection
used: scan rate = $30^\circ/\text{min}$ for all reflections (3 hours for 1 unique set
of data). First data set ($-10 \leq h \leq 0$, $0 \leq k \leq 11$, $0 \leq l \leq 8$) consisted of
696 observations, of which 36 were monitors, 524 were unique $F_o > 3\sigma(F_o)$
and 59 unique $F_o < 3\sigma(F_o)$. Monitor reflections (004, 620, and 040;
measured after every 60 reflections) decreased uniformly from 100 to 87%
of their original values (with individual variations from the mean of
 $\pm 4\%$) over the course of 3 hours collection. Smoothed curve of monitor
decrements used to correct all data. Second octant ($+h, +k, +l$) collected
subsequently, monitors decreasing from 87% to 80% of original values;



numerous peak profiles off-center in second octant; second set discarded to avoid introduction of systematic errors.

Lorentz and polarization, but not absorption, corrections made. Structure solved using direct methods in Pna₂₁ and Pnam (centric). Fifteen atoms in first Pna₂₁ E-map; one more appeared with tangent-formula recycling. Since this result was an unexpected reaction product, atoms were identified by assigning C scattering factors to all atoms and refining occupancies of atoms, holding isotropic U_{ij}'s constant. All nitrogen and oxygen occupancies refined to 1.20-1.40 range, while carbon atom occupancies remained in 1.0-1.15 range. Eventual appearance of all hydrogen atoms in difference map corroborated atomic assignment.

Refinement: 150 parameters refined, all non-H atoms anisotropic, amino hydrogen refined on position and isotropic T factor, methylene hydrogens fixed to ride on carbon atoms with isotropic U's and tetrahedral geometry. R = 0.0511, R_w = 0.0609, S = 1.295 for 524 Fc>3σ(Fc). Function minimized, $\Sigma w(F_o - F_c)^2$, where $w = [\sigma^2(F_o) + 0.0009F_o^2]^{-2}$. Maximum Δ/σ = 0.040, final difference map ripple extrema, 0.23 and -0.25e4-3.

1-Nitroso-2,2-dil(trifluoromethyl)-1,3-diazacyclopentan-4-one

(A)




3,3,7,7-Tetra(trifluoromethyl)-2,4,6,8-tetraazabicyclo[3.3.0]octane

Abstract



$C_8H_{14}N_4F_{12}$, $M_r = 336.14$, monoclinic, $P2/n$, $a = 10.431(4)$,
 $b = 7.050(2)$, $c = 25.773(14)\text{\AA}$, and $\beta = 90.80(4)$. M.P. = 94 $^{\circ}\text{C}$.
 $V = 1895.7(13)\text{\AA}^3$, $Z = 6$, $D_x = 2.029 \text{ mg mm}^{-3}$, $\lambda(\text{CuK}\alpha) = 1.54178\text{\AA}$,
 $\mu = 2.336 \text{ mm}^{-1}$, $F(000) = 1140$, $T = 295\text{K}$, Final $R = 0.053$, $wR = 0.056$ for
2294 independent observed reflections. There are one and a half molecules in the
asymmetric unit.

Experimental

Clear 0.35 x 0.22 x 0.15 mm. crystal crystallized from CH_2Cl_2 at 5 $^{\circ}\text{C}$,
Automated Nicolet R3m diffractometer with incident beam graphite mono-
chromator $\lambda = 1.54178\text{\AA}(\text{CuK}\alpha)$, 25 centered reflections within $30 < 2\theta < 60$
used for determining lattice parameters. $(\sin\theta/\lambda)_{\max} = 0.59\text{\AA}^{-1}$, range of
 hkl : $0 < h < 11$, $0 < k < 8$, $-30 < l < 30$. Standards 600, 020, 004, monitored
every 60 reflections with linear decrease of 8.9% during data collection, $\theta/2\theta$ mode,
scan width $(2.0 + \Delta_{\text{clsc2}})^{\circ}$, scan rate a function of count rate ($8^{\circ}/\text{min. minimum}$,
 $30^{\circ}/\text{min. maximum}$), 3601 reflections measured, 3112 unique, $R_{\text{int}} = 0.025$, 2294
observed $F_o > 3\sigma(F_o)$. Data corrected for Lorentz, polarization and absorption
effects, and for the linear decrease in monitored intensities. Structure solved by
direct methods. The least-squares refinement used program SHELXTL (Sheldrick 1980).
 $\Sigma w(|F_o| - |F_c|)^2$ minimized where $w = 1/[\sigma^2(|F_o|) + g \cdot (F_o)^2]$, $g = 0.00050$, 361
parameters refined: atom coordinates, anisotropic temperature factors for all
non-H atoms, isotropic temperature factors for H atoms. $(\Delta/\sigma)_{\max} = -.16$,
 $R = 0.053$, $wR = 0.056$, $S = 1.510$. Final difference Fourier excursions 0.26
and $-0.32e\text{\AA}^{-3}$. Atomic scattering factors from International Tables
for X-ray Crystallography (1974).



(B)



**3,3,7,7-Tetra(trifluoromethyl)-
2,4,6,8-tetraazabicyclo(3.3.0)octane**

2,5-Dinitro-3,3,7,7-tetra(trifluoromethyl)-2,4,5,8-tetraazabicyclo[3.3.0]octane



Abstract

$C_8H_4N_6O_4F_{12}$, $M_r = 476.14$, orthorhombic, $Pna2_1$, $a = 20.073(4)$,
 $b = 7.353(1)$, $c = 10.832(2)\text{\AA}$, $V = 1598.7(4)\text{\AA}^3$, $z = 4$, $D_x = 1.978 \text{ mg mm}^{-3}$,
 $\lambda(\text{CuK}\alpha) = 1.54178\text{\AA}$, $\mu = 2.22 \text{ mm}^{-1}$, $F(000) = 936$, $T = 295\text{K}$,
Final $R = 0.064$, $wR = 0.089$ for 1205 independent observed reflections.

Experimental

Clear 0.10x0.15x0.65 mm. crystal, crystallized from $\text{CH}_2\text{Cl}_2/\text{hexane}$ by W. Koppes of Naval Surface Weapons Center (Silver Springs, Md.). Automated Nicolet R3m diffractometer with incident beam graphite monochromator $\lambda = 1.54178\text{\AA}(\text{CuK}\alpha)$, 25 centered reflections within $40 < 2\theta < 60$ used for determining lattice parameters. Data corrected for Lorentz and polarization effects, but not for absorption. $(\sin\theta/\lambda)_{\text{max}} = 0.56\text{\AA}^{-1}$, range of hkl : $-22 < h < 14$, $-8 < k < 0$, $12 < l < 0$. Standards 004, 800, 020, monitored every 60 reflections with linear variation 4.1% over data collection, θ -2 θ mode, scan width $(2.0 + \Delta_{\text{slit}})^{\circ}$, scan rate a function of count rate (6°/min. minimum, 30°/min. maximum), 2393 reflections measured, 1263 unique, $R_{\text{int}} = 0.021$, 1205 observed $F_0 > 3\sigma(F_0)$. Structure solved by direct methods. The least-squares refinement used program SHELXTL (Sheldrick 1980). $\Sigma w(|F_0| - |F_c|)^2$ minimized where $w = 1/c^2(|F_0|) + g \cdot (F_0)^2$, $g = 0.00030$, 283 parameters refined: atom coordinates, anisotropic temperature factors for all non-H atoms, hydrogen isotropic temperature factors fixed at 1.2 $U_{\text{eq}}(\text{C})$. $(\Delta/\sigma)_{\text{max}} = -0.012$ $R = 0.064$, $wR = 0.089$, $S = 3.65$. Final difference Fourier excursions 0.34 and -0.28 e\AA^{-3} . Atomic scattering factors from International Tables for X-ray Crystallography (1974).

(C)



2,6-Dinitro-3,3,7,7-tetra(trifluoromethyl)-

2,4,6,8-tetraazabicyclo(3.3.0)octane

2,4,6-Trinitro-3,3,7,7-tetra(trifluoromethyl)-
2,4,6,8-tetraazabicyclo[3.3.0]octane



Abstract

$C_8H_3N_7O_6F_{12}$, $M_r = 521.13$, monoclinic, $P2_1/c$, $a = 6.390(6)$,
 $b = 19.090(2)$, $c = 12.347(1)\text{\AA}$, $\beta = 95.38(1)^\circ$, $V = 1640.1(3)\text{\AA}^3$,
 $Z = 4$, $D_x = 2.110 \text{ mg mm}^{-3}$, $\lambda(\text{CuKa}) = 1.54178\text{\AA}$, $\mu = 2.35 \text{ mm}^{-1}$,
 $F(000) = 1024$, $T = 295\text{K}$, Final $R = 0.049$, $wR = 0.074$ for independent 2153
observed reflections.

Experimental

Clear 0.62x0.23x0.35 mm. crystal, crystallized from CH_2Cl_2 /hexane by W. Koppes of Naval Surface Weapons Center (Silver Springs, Md) Automated Nicolet R3m diffractometer with incident beam graphite monochromator $\lambda = 1.54178\text{\AA}(\text{CuKa})$, 25 centered reflections within $30 < 2\theta < 60$ used for determining lattice parameters. Data corrected for Lorentz, polarization, and absorption effects, $(\sin\theta/\lambda)_{\max} = 0.55\text{\AA}^{-1}$, range of hkl : $-7 \leq h \leq 0$, $0 \leq k \leq 19$, $-13 \leq l \leq 12$. Standards 200,0 12 0, 006, monitored every 60 reflections with linear variation 8.3% over data collection, θ - 2θ mode, scan width $(2.0 + \Delta_{\alpha_1\alpha_2})^\circ$, scan rate a function of count rate, ($5^\circ/\text{min. minimum}$, $30^\circ/\text{min. maximum}$). 2568 reflections measured, 2259 unique, $R_{\text{int}} = 0.023$, 2163 observed $F_0 > 3\sigma(F_0)$. Structure solved by direct methods. The least-squares refinement used program SHELXL (Sheldrick 1980). $\sum w(|F_0| - |F_c|)^2$ minimized where $w = 1/[s^2(|F_0|) + g \cdot (F_0)^2]$, $s = 0.00023$, isotropic secondary extinction value, 0.0019, 367 parameters refined: atom coordinates, anisotropic temperature factors for all non-H atoms, isotropic temperature factors for H atoms. $(\Delta/c)_{\max} = 0.24$, $R = 0.049$, $wR = 0.074$, $S = 2.77$. Final difference Fourier excursions 0.27 and $-0.19\text{\textit{e}\AA}^{-3}$. Atomic scattering factors from International Tables for X-ray Crystallography (1974).

(D)



2,4,6-Trinitro-3,3,7,7-tetra(trifluoromethyl)-
2,4,6,8-tetraazabicyclo(3.3.0)octane

2,4,6,8-Tetranitro-3,3,7,7-tetra(trifluoromethyl)-2,4,6,8-tetraazabicyclo[3.3.0]octane

Abstract


$C_8H_2N_8O_8F_{12}$, $M_r = 566.13$, monoclinic, $C2/c$, $a = 34.074(5)$,
 $b = 7.456(1)$, $c = 13.877(2)\text{\AA}$, $\beta = 102.41(1)^\circ$, m.p. = 110°C ,
 $V = 3443.1(8)\text{\AA}^3$, $Z = 8$, $D_x = 2.184 \text{ mg mm}^{-3}$, $\lambda(\text{CuKa}) = 1.54178\text{\AA}$,
 $\mu = 2.413 \text{ mm}^{-1}$, $F(000) = 2224$, $T = 295\text{K}$, Final $R = 0.041$, $wR = 0.068$ for
2799 independent observed reflections.

Experimental

Clear $0.50 \times 0.35 \times 0.12$ mm. crystal, crystallized from CH_2Cl_2 /hexane,
Synthesized by W. Koppes of Naval Surface Weapons Center (Silver Springs, Md.)
Automated Nicolet R3m diffractometer with incident beam graphite mono-
chromator $\lambda = 1.54178\text{\AA}(\text{CuKa})$, 25 centered reflections within $45^\circ < 2\theta < 90^\circ$
used for determining lattice parameters. Data corrected for Lorentz,
polarization, and absorption effects, $(\sin\theta/\lambda)_{\max} = 0.59\text{\AA}^{-1}$, range of hkl :
 $-40 < h < 38$, $0 < k < 8$, $0 < l < 15$. Standards 15 1 1, 041, 0 0 10, monitored
every 60 reflections with linear variation 7.0% over data collection, θ -2 θ mode,
scan width $(2.0 + \Delta_{\text{cal}}/2)^\circ$, scan rate a function of count rate ($5^\circ/\text{min. minimum}$,
 $30^\circ/\text{min. maximum}$), 3503 reflections measured, 2830 unique, $R_{\text{int}} = 0.012$,
2799 observed $F_O > 3\sigma(F_O)$. Structure solved by direct methods. The
least-squares refinement used program SHELXL (Sheldrick 1980). $S \propto (\|F_O\| - |F_C\|)^2$
minimized where $w = 1/[o^2(\|F_O\|) + g \cdot (F_O)^2]$, $g = 0.00030$, 333 parameters
refined: atom coordinates, anisotropic temperature factors for all non-H
atoms, isotropic temperature factors for H atoms, $(\Delta/o)_{\max} = 0.13$,
 $R = 0.041$, $wR = 0.068$, $S = 3.077$. Final difference Fourier excursions 0.23
and -0.24 e\AA^{-3} . Atomic scattering factors from International Tables
for X-ray Crystallography (1974).

**2,4,6,8-Tetranitro-3,3,7,7-tetra(trifluoromethyl)-
2,4,6,8-tetraazabicyclo(3.3.0)octane**

(E)



3-Benzamido-6,8-dibenzoyl-1,3,5,6,8-pentaazabicyclo[3.2.2]nonane


Abstract

$C_{25}H_{23}N_6O_3$, $M_r = 455.50$, monoclinic, $P2_1/c$, $a = 11.325(3)$,
 $b = 19.161(4)$, $c = 11.952(5)\text{\AA}$, $\beta = 112.01(3)$, $V = 2280.3(10)\text{\AA}^3$.
 $Z = 4$, $D_x = 1.327 \text{ mg mm}^{-3}$, $\lambda(\text{CuK}\alpha) = 1.54178\text{\AA}$, $\mu = 0.70\text{mm}^{-1}$,
 $F(000) = 956$, $T = 295\text{K}^\circ$, Final $R = 0.072$, $wR = 0.061$ for 1269 independant
observed reflections.

Experimental

Clear 0.05x0.02x0.15mm. crystal, crystallized from methanol.
Synthesized by G. Kumar and J. Boyer of the University of Illinois at Chicago.
Automated Nicolet R3m diffractometer with incident beam graphite mono-
chromator $\lambda = 1.54178\text{\AA}(\text{CuK}\alpha)$, 20 centered reflections within $25^\circ < 2\theta < 50^\circ$
used for determining lattice parameters. Data corrected for Lorentz
and polarization effects, $(\sin\theta/\lambda)_{\max} = 0.53\text{\AA}^{-1}$, range of hkl :
 $0 \leq h \leq 10$, $0 \leq k \leq 19$, $-12 \leq l \leq 11$. Standards 202, 040, 002, monitored every
60 reflections with random variation 2.2% over data collection, θ -2 θ mode,
scan width $(2.0 + \Delta_{\text{calc}})^{\circ}$, scan rate a function of count rate (2°/min. minimum,
30°/min. maximum), 2541 reflections measured, 2268 unique, $R_{\text{int}} = 0.022$,
1269 observed $F_o > 3\sigma(F_o)$. Structure solved by direct methods. The
least-squares refinement used program SHELXML (Sheldrick 1980). $\Sigma w(|F_o| - |F_c|)^2$
minimized where $w = 1/[\sigma^2(|F_o|) + g \cdot (F_o)^2]$, $g = 0.00050$. 310 parameters
refined: atom coordinates, anisotropic temperature factors for all non-H
atoms, H atoms included using riding model, C-H = 0.96\AA, H-C-H = 109.5°,
 $U(H) = 1.2 U_{\text{eq}}(\text{C})$, amine hydrogen coordinates, $\bar{x}(10)$, refined. $(\Delta/\sigma)_{\max} = -0.19$,
 $R = 0.072$, $wR = 0.061$, $S = 1.293$. Final difference Fourier excursions 0.24 and
 -0.22eA^{-3} . Atomic scattering factors from International Tables for
X-ray Crystallography (1974).

(F)



3-Benzamido-6,8-pentaazabicyclo(3.2.2)nonane

1,3,5,6,8-pentaazabicyclo(3.2.2)nonane

1,3,5-Tribenzamido-1,3,5-hexahydrotriazine monohydrate


Abstract

$C_{24}H_{24}N_6O_3 \cdot H_2O$, $M_r = 462.49$, monoclinic, $P2_1/c$.
 $a = 12.655(1)$, $b = 14.721(2)$, $c = 13.339(2)\text{\AA}$, $\beta = 107.25(1)^\circ$,
 $V = 2373.0(5)\text{\AA}^3$, $Z = 4$, $D_x = 1.294 \text{ mg mm}^{-3}$, $\lambda(\text{CuK}\alpha) = 1.54178\text{\AA}$,
 $\mu = 0.709 \text{ mm}^{-1}$, $F(000) = 976$, $T = 295\text{K}$, Final $R = 0.042$, $wR = 0.045$ for
3054 independent observed reflections.

Experimental

Clear 0.08x0.25x0.30 mm. crystal, crystallized from methanol/water.
Synthesized by G.Kumar and J. Boyer of the Univ. of Ill. at Chicago.
Automated Nicolet R3m diffractometer with incident beam graphite monochromator
 $\lambda = 1.54178 \text{ \AA}(\text{CuK}\alpha)$, 25 centered reflections within $37^\circ < 2\theta < 90^\circ$ used
for determining lattice parameters. Data corrected for Lorentz and polarization,
but not absorption effects, $(\sin\theta/\lambda)_{\text{max}} = 0.57\text{\AA}^{-1}$, range of hkl :
 $-14 \leq h \leq 14$, $0 \leq k \leq 16$, $-15 \leq l \leq 0$. Standards $\bar{5}0\bar{2}$, $1\bar{0}\bar{6}$, $0\bar{6}0$, monitored every
60 reflections with random variation 1.3% over data collection, 9-29 mode,
scan width $(1.8 + \Delta_{\text{slip}})^{\circ}$, scan rate a function of count rate (6°/min. minimum,
30°/min. maximum), 4156 reflections measured, 3799 unique, $R_{\text{int}} = 0.011$,
3054 observed $F_o > 3\sigma(F_o)$. Structure solved by direct methods. The
least-squares refinement used program SHELXL (Sheldrick 1980). $\Sigma w(|F_o| - |F_c|)^2$
minimized where $w = 1/[s^2(|F_o|) + g \cdot (F_o)^2]$, $s = 0.00023$, isotropic secondary
extinction value, 0.00272, 412 parameters refined: atom coordinates,
anisotropic temperature factors for all non-H atoms, isotropic temperature
factors for H. $(\Delta/c)_{\text{max}} = -0.017$, $R = 0.042$, $wR = 0.045$, $S = 1.469$.
Final difference Fourier excursions 0.17 and $-0.15e\text{\AA}^{-3}$. Atomic
scattering factors from International Tables for X-ray Crystallography
(1974).

1,3,5-Tribenzamido-1,3,5-hexahydrotriazine monohydrate



1-Benzoyl-4-benzamido-5²-1,2,4-triazoline

Abstract


$C_{15}H_{14}N_4O_2$, $M_r = 294.31$, monoclinic, $P2_1/c$, $a = 14.604(2)$,
 $b = 10.339(1)$, $c = 9.714(1)\text{\AA}$, $\beta = 102.56(1)^\circ$, $V = 1431.7(3)\text{\AA}^3$,
 $Z = 4$, $D_x = 1.365 \text{ mg mm}^{-3}$, $\lambda(\text{Cu K}\alpha) = 1.54178\text{\AA}$, $\mu = 0.73 \text{ mm}^{-1}$,
 $F(000) = 616$, $T = 295\text{K}$. Final $R = 0.039$, $wR = 0.050$ for 2117 independent
observed reflections.

Experimental

Clear 0.15x0.25x0.40 mm. crystal, crystallized from benzene/chloroform.
Synthesized by G. Kumar and J. Boyer of the Univ. of Ill. at Chicago, m.p. =
216-218 $^\circ\text{C}$. Automated Nicolet R3m diffractometer with incident beam graphite
monochromator $\lambda = 1.54178\text{\AA}(\text{CuK}\alpha)$, 25 centered reflections within $30 < 2\theta < 60^\circ$
used for determining lattice parameters. Data corrected for Lorentz and
polarization effects. $(\sin\theta/\lambda)_{\text{max}} = 0.59\text{\AA}^{-1}$, range of hkl : $-17 \leq h \leq 4$, $0 \leq k \leq 11$,
 $-11 \leq l \leq 11$, Standards 800, 040, 004, monitored every 60 reflections with random
variation 2.4% over data collection, θ -2 θ mode, scan width ($2.0 + 4\sin\theta$)
scan rate a function of count rate (8°/min. minimum, 30°/min. maximum),
3408 reflections measured, 2437 unique, $R_{\text{int}} = 0.014$, 2117 observed $F_o > 3\sigma(F_o)$.

Structure solved by direct methods. The least-squares refinement
used program SHELXL (Sheldrick 1980). $\Sigma w(|F_o| - |F_c|)^2$ minimized where
 $w = 1/[s^2(|F_o|) + g \cdot (F_o)^2]$, $g = 0.0003$, isotropic secondary extinction value =
0.00206, 256 parameters refined: atom coordinates, anisotropic temperature
factors for all non-H atoms, isotropic temperature factors for H, $(\Delta/\sigma)_{\text{max.}} =$
0.016 ; $R = 0.039$, $wR = 0.0503$, $S = 1.741$. Final difference Fourier excursions
0.19 and -0.16e \AA^{-3} . Atomic scattering factors from International Tables
for X-ray Crystallography (1974).

1-Benzoyl-4-benzamido- Δ^2 -1,2,4-triazoline



2,7-Dibenzoyl-4-benzamido-1,2,4,6,7-pentaazabicyclo[1.2.4]nonane


Abstract

$C_{25}H_{24}N_6O_3$, $M_r = 456.51$, triclinic, $\bar{P}\bar{1}$, $a = 9.195(1)$.
 $b = 11.457(2)$, $c = 12.139(2)\text{\AA}$, $\alpha = 89.77(2)$, $\beta = 72.79(1)$, and $\gamma = 76.52(1)^\circ$.
 $v = 1134.9(4)\text{\AA}^3$, $Z = 2$, $D_x = 1.279 \text{ mg mm}^{-3}$, $\lambda(\text{CuK}\alpha) = 1.54178\text{\AA}$,
 $\mu = 0.674 \text{ mm}^{-1}$, $F(000) = 480$, $T = 295\text{K}$, Final $R = 0.056$, $wR = 0.057$ for
2421 independent observed reflections.

Experimental

Clear $0.32 \times 0.21 \times 0.08 \text{ mm}$. crystal, recrystallized from ethanol.
Synthesized by G. Kumar and J. Boyer of the Univ. of Ill. at Chicago.
Automated Nicolet R3m diffractometer with incident beam graphite mono-
chromator $\lambda = 1.54178\text{\AA}(\text{CuK}\alpha)$, 25 centered reflections within $39^\circ < 2\theta < 60^\circ$
used for determining lattice parameters. Data corrected for Lorentz, polarization,
but not for absorption effects, $(\sin\theta/\lambda)_{\max} = 0.573^{-1}$, range of hkl :
 $-10 \leq h \leq 6$, $-12 \leq k \leq 11$, $-13 \leq l \leq 8$. Standards $00\bar{5}$, 060 , $\bar{5}20$, monitored every
60 reflections with random variation 5.4% over data collection, θ -2 θ mode,
scan width $(1.8 + \Delta_{\text{cal}})_2$ °, scan rate a function of count rate (8°/min. minimum,
30°/min. maximum), 3448 reflections measured, 3131 unique, $R_{\text{int}} = 0.018$,
2421 observed $F_o > 3\sigma(F_o)$. Structure solved by direct methods. The
least-squares refinement used program SHELXTL (Sheldrick 1980). $\Sigma w(|F_o| - |F_c|)^2$
minimized where $w = 1/[c^2(|F_o|) + g \cdot (F_o)^2]$, $g = 0.00023$, isotropic secondary
extinction value, 0.0078, 344 parameters refined: atom coordinates,
anisotropic temperature factors for all non-H atoms, isotropic temperature
factors for H, benzene ring hydrogens included using riding model, C-H =
 0.96\AA , C-C-H = 120.0° , 1.1 U_{eq} (C). $(\Delta/\sigma)_{\max} = -0.003$, $R = 0.056$,
 $wR = 0.057$, $S = 1.512$. Final difference Fourier excursions 0.40 and -0.23 $e\text{\AA}^{-3}$.
Atomic scattering factors from International Tables for X-ray Crystallography (1974

(I)



2,7-Dibenzoyl-4-benzamido-
1,2,4,6,7-pentaazabicyclo(1.2.4)nonane

bis-Tetrazolo[1,5-a:1',5'-c]pyrazine

Abstract

$C_{14}H_2N_8$, $M_p = 162.11$, orthorhombic, $P2_12_12_1$, $a = 6.1155(5)$,
 $b = 8.2572(3)$, $c = 12.7401(11)\text{\AA}$, $V = 643.3(1)\text{\AA}^3$, $z = 4$, $D_x = 1.674$
 mg mm^{-3} , $\lambda(\text{CuKa}) = 1.54178\text{\AA}$, $\mu = 1.036 \text{ mm}^{-1}$, $F(000) = 328$, $T = 295\text{K}^\circ$
Final $R = 0.0326$, $wR = 0.0440$ for 1106 independent observed reflections.
All of the non-hydrogen atoms in this new heterocaromatic compound lie
within 0.014\AA of a common plane.

Experimental


Clear, colorless square prisms; data crystal $0.15 \times 0.15 \times 0.55\text{mm.}$, synthesized & crystallized by M. Chaykovsky of Naval Surface Weapons Center (Silver Spring, Md.). Automated Nicolet R3m diffractometer with incident beam graphite monochromator $\lambda = 1.54178\text{\AA}(\text{CuKa})$; 25 centered reflections within $43^\circ < 2\theta < 75^\circ$ used for determining lattice parameters. Data corrected for Lorentz & polarization, but not absorption effects, $(2\theta)_{\max} = 139^\circ$, range of hkl : $0 \leq h \leq 7$, $-9 \leq k \leq 10$, $-15 \leq l \leq 15$. Standards 400 , 040 , 0010 , monitored every 60 reflections with random variation $\pm 4\%$ over data collection, $\theta/2\theta$ mode, scan width $(1.4 + \Delta_{\text{calc}})^{\circ}$, scan rate a function of count rate ($4^\circ/\text{min. minimum, } 30^\circ/\text{min. maximum}$), 1976 reflections measured (incl. 97 monitors), 1106 unique (Friedel pairs not merged), $R_{\text{int}} = 0.0178$, no reflections considered unobserved.

Structure solved by direct methods. The least-squares refinement used program SHELXL (Sheldrick 1980). $\Sigma w(|F_0| - |F_c|)^2$ minimized where $w = 1/[c^2(|F_0|) + g \cdot (F_0)^2]$, $g = 0.000225$, isotropic secondary extinction correction applied, 116 parameters refined: atom coordinates (all atoms),

anisotropic temperature factors for all non-H atoms, H atom U_{ij} 's fixed at anisotropic values predicted by rigid-body vibration analysis of non-H atoms, $(\Delta/\sigma)_{\text{max}} = 0.005$, $R = 0.0326$, $wR = 0.0440$, $S = 2.25$. Final difference Fourier extrema 0.38 and $-0.28 \text{ e}\AA^{-3}$. Atomic scattering factors from International Tables for X-ray Crystallography (1974).

bis-Tetrazolo(1,5-a;1'5'-c)pyrazine

(J)



1,4-Dinitro-2,3,5,6-tetrabromo-2,3,5,6-piperazine

Abstract: $C_6H_6N_4O_4Br_4$, $M_r = 491.72$, triclinic, $\bar{P}1$, $a = 7.517(3)$, $b = 7.566(2)$, $c = 11.835(3)$ Å, $\alpha = 78.89(2)$, $\beta = 77.44(3)$, $\gamma = 63.84(2)^\circ$, $F(000) = 456$, $V = 586.1(3)$ Å 3 , $z = 2$, $D_x = 2.79$ mg mm $^{-3}$, λ (CuKα) = 1.54178 Å, $\mu = 170.96$ cm $^{-1}$, $T = 295K$, final $R = 0.069$ for 1891 unique reflections. Asymmetric unit consisted of two half-molecules. Results showed the bromine atoms are in trans-trans configuration, and are all axial with respect to the piperazine chair-shaped ring.


Experimental: Clear colorless 0.23x0.15x0.08 mm crystal provided by M. Chaykovsky of the Naval Surface Weapons Center (Silver Spring, MD). Nicolet R3M diffractometer, monochromator on incident beam, θ -2 θ scan technique with a constant scan speed of 60°/min. Unit cell parameters from least-squares analysis of 25 reflections with 2θ from 47 to 95°. $\bar{P}1$ checked for higher symmetry using program AIDS (Mighell, Hubbard and Stalick, 1980), 2038 independent reflections with $2\theta_{max} = 130^\circ$; three standard reflections measured after every 60 new reflections showed an average random variation of 3.0% in $|F_o|$. D_m not determined, crystals sank in CCl_4 ($d = 1.59$ mg mm $^{-3}$), Lorentz polarization corrections, empirical absorption correction applied (max. trans = 0.913, min trans = 0.363). Structure solved by direct methods (Karle and Karle, 1966), Br atoms found in E-map, all remaining non-hydrogen atoms in a difference map. Refined by full-matrix least-squares, function minimized $\Sigma w(|F_o - |F_c|)^2$, isotropic secondary extinction correction applied. Non-H atoms anisotropic, H atoms in calculated positions riding on bonded atoms, 1891 reflections having $|F_o| > 3\sigma|F_o|$ refined to a final R factor of 0.069 ($R_w = 0.083$, $S = 3.0$), $\Delta/\Delta_{max} = 0.20$, final difference map

was featureless except for ripples around Br atoms (max height = 1.76 Å).

Discussion

The asymmetric unit for this compound consisted of 2 half-molecules located on centers of symmetry rather than one full molecule in a general position. However, there are no significant differences in the conformations of the two molecules. In both molecules the six-membered ring has a slightly flattened chair conformation and the NO₂ group is coplanar with N1. The sum of angles around both N2 and N2' is 359.9. Pairs of bromine atoms substituted on adjacent carbon atoms of the ring are trans with respect to one another, and are axial with respect to the ring. Amino nitrogen atoms substituted with nitro groups are usually near planar in geometry. In this molecule, the amino nitrogens are slightly pyramidal, and the nitro groups both bend in the equatorial direction with respect to the ring. Even though this crystal has a very high density ($d_{\text{calc}} = 2.79$) there are only a few close intermolecular contacts: a Br1...Br2 approach at 3.69 Å and a Br2...O2' approach at 3.28 Å.

1,4-Dinitro-2,3,5,6-tetrabromopiperazine




1,9-Diacetyl-3,5,7-trinitro-1,3,5,7-tetraaza-9-oxanane

Abstract



$C_8H_{15}N_7O_9$, $M_r = 353.25$, orthorhombic Pbca, $a = 12.410(2)$,
 $b = 9.695(1)$, $c = 25.079(3)\text{\AA}$, $\alpha = \beta = \gamma = 90.00^\circ$, $\nu = 3017.5(6)\text{\AA}^3$,
 $Z = 8$, $D_x = 1.555 \text{ mg mm}^{-3}$, $\lambda(\text{CuK}\alpha) = 1.54178\text{\AA}$, $\mu = 1.189 \text{ mm}^{-1}$,
 $F(000) = 1472$, $T = 295\text{K}$. Final $R = 0.087$, $wR = 0.094$ for 1427 independent
observed reflections. The acetoxy group is disordered with an occupancy
of 57 and 43% respectively for the two orientations.

Clear 0.05 x 0.10 x 0.25 mm. crystal crystallized from methyl chloride.
Automated Nicolet R3m diffractometer with incident beam graphite mono-
chromator $\lambda = 1.54178\text{\AA}(\text{CuK}\alpha)$, 25 centered reflections within $30 < 2\theta < 50^\circ$
used for determining lattice parameters. $(\sin\theta/\lambda)_{\max} = 0.54\text{\AA}^{-1}$, range
of hkl : $-13 \leq h \leq 1$, $0 \leq k \leq 10$, $0 \leq l \leq 14$. Standards 400, 040, 006, monitored
every 60 reflections with random variation 4.6% over data collection, $\phi/2\theta$ mode,
scan width $(2.0 + \Delta_{\text{slit}})^*$, scan rate a function of count rate ($2^\circ/\text{min. minimum}$,
 $30^\circ/\text{min. maximum}$), 2782 reflections measured, 2082 unique, $R_{\text{int}} = 0.02$, 1427
observed $F_c > 3\sigma(F_o)$. Data corrected for Lorentz, polarization but not absorption
effects, Structure solved by direct methods. The least-squares refinement
used program SHELXTL (Sheldrick 1980). $\Sigma w(|F_o| - |F_c|)^2$ minimized where
 $w = 1/[s^2(|F_o|) + g \cdot (F_o)^2]$, $g = 0.00040$, Secondary extinction value, 0.0009.
256 parameters refined: atom coordinates, anisotropic temperature factors
for all non-H atoms, H atoms included using riding model, C-H = 0.96 \AA ,
H-C-H = 109.5° , $J(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$, $|(\Delta/c)_{\max} = 0.123$, $R = 0.066$,
 $wR = 0.068$, $S = 1.841$. Final difference Fourier excursions 0.38 and -0.34 $e\text{\AA}^{-3}$.
Atomic scattering factors from International Tables for X-ray Crystallography
(1974).

(L)



d - 1.555



1,9-DIACETYL-3,5,7-TRINITRO-1,3,5,7-TETRAAZA-9-OXANONANE

1,9-Diacetyl-3,5,7-trinitro-1,3,5,7,9-pentaazanonane

Abstract


$C_9H_{16}N_8O_8$, $M_r = 352.17$, monoclinic, $P2_1$, $a = 7.243(2)$, $b = 19.986(4)$, $c = 10.692(2)\text{\AA}$, $\beta = 96.18(2)^\circ$, $V = 1538.7(6)\text{\AA}^3$, $Z = 4$, $D_x = 1.520 \text{ mg mm}^{-3}$, $\lambda(\text{CuK}\alpha) = 1.54178\text{\AA}$, $\mu = 1.13 \text{ mm}^{-1}$, $F(000) = 736$, $T = 295\text{K}$. Final $R = 0.038$, $wR = 0.045$ for 2900 independent observed reflections. There are two independent molecules in the asymmetric unit.

Clear $0.25 \times 0.15 \times 0.45 \text{ mm}$. crystal crystallized from methyl chloride. Automated Nicolet R3m diffractometer with incident beam graphite monochromator $\lambda = 1.54178\text{\AA}(\text{CuK}\alpha)$, 25 centered reflections within $40 < 2\theta < 65^\circ$ used for determining lattice parameters. $(\sin\theta/\lambda)_{\max} = 0.61\text{\AA}^{-1}$, range of $hkl: -7 \leq h \leq 8, -24 \leq k \leq 0, -12 \leq l \leq 1$. Standards $\bar{2}0\bar{3}$, $0\bar{8}0$, $00\bar{5}$, monitored every 60 reflections with random variation 1.4% over data collection, $6/29$ mode, scan width $(2.0 + \Delta_{\text{alat}})^\circ$, scan rate a function of count rate ($4^\circ/\text{min. minimum}$, $30^\circ/\text{min. maximum}$), 4272 reflections measured, 2982 unique, $R_{\text{int}} = 0.027$, 2900 observed $F_o > 3\sigma(F_o)$. Data corrected for Lorentz, polarization and absorption effects, Structure solved by direct methods. The least-squares refinement used program SHELXL (Sheldrick 1980). $\Sigma w(|F_o| - |F_c|)^2$ minimized where $w = 1/[c^2(|F_c|) + g \cdot (F_o)^2]$, $g = 0.00030$.


458 parameters refined: atom coordinates, anisotropic temperature factors for all non-H atoms, amine hydrogen coordinates and isotropic temperature factors, other H atoms included using riding model, C-H = 0.96\AA , angle H-C-H = 109.5° , $U(H) = 1.2 U_{\text{eq}}(\text{C}) \cdot (\Delta/c)_{\max} = 0.40$, mean = 0.03, $R = 0.038$, $wR = 0.045$, $S = 1.737$. Final difference Fourier excursions 0.1^4 and $-0.31e4^{-3}$. Atomic scattering factors from International Tables for X-ray Crystallography (1974).

1,9-Diacetyl-3,5,7-trinitro-1,3,5,7,9-pentaazanonane

d - 1.52



(M)




2-Oxa-6,9-diaza-6,9-dinitrospiro[3.6]decane

Abstract

$C_7H_{12}N_4O_5$, $M_r = 232.20$, triclinic, $\bar{P}\bar{1}$, $a = 6.028(1)$,
 $b = 6.420(2)$, $c = 13.764(4)\text{\AA}$, $\alpha = 91.04(2)$, $\beta = 100.22(2)$, and $\gamma = 108.57(2)^\circ$,
 $V = 492.0(2)\text{\AA}^3$, $Z = 2$, $D_x = 1.567 \text{ mg mm}^{-3}$, $\lambda(\text{CuK}\alpha) = 1.54178\text{\AA}$,
 $\mu = 1.108 \text{ mm}^{-1}$, $F(000) = 244$, $T = 295\text{K}$, Final $R = 0.075$, $wR = 0.075$ for
877 independent observed reflections.

Clear 0.02 x 0.08 x 0.15 mm. crystal crystallized from ethyl acetate,
Automated Nicolet R3m diffractometer with incident beam graphite mono-
chromator $\lambda = 1.54178\text{\AA}(\text{CuK}\alpha)$, 25 centered reflections within $32 < 2\theta < 75^\circ$
used for determining lattice parameters. $(\sin\theta/\lambda)_{\max} = 0.58\text{\AA}^{-1}$, range
of hkl : $0 \leq h \leq 6$, $-7 \leq k \leq 7$, $-15 \leq l \leq 15$. Standards $0\bar{2}1$, 200 , 003 , monitored
every 60 reflections with random variation 2.5% over data collection, $\theta/2\theta$ mode,
scan width $(2.0 + \Delta_{\text{slit}}/2)^\circ$, scan rate a function of count rate ($3^\circ/\text{min. minimum}$,
 $30^\circ/\text{min. maximum}$), 1864 reflections measured, 1253 unique, $R_{\text{int}} = 0.026$, 877
observed $F_c > 3\sigma(F_o)$. Data corrected for Lorentz, polarization and absorption
effects, Structure solved by direct methods. The least-squares refinement
used program SHELXL (Sheldrick 1980). $\Sigma w(|F_o| - |F_c|)^2$ minimized where
 $w = 1/[c^2(|F_o|) + g \cdot (F_o)^2]$, $g = 0.00063$, Secondary extinction value 0.0022.
146 parameters refined: atom coordinates, anisotropic temperature factors
for all non-H atoms isotropic temperature factors for H atoms [H atoms
included using riding model, C-H = 0.96\AA, H-C-H = 109.5°, $U(H) = 1.1 U_{\text{eq}}(\text{C})$],
 $(\Delta/c)_{\max} = 0.01$, $R = 0.075$, $wR = 0.075$, $S = 1.64$. Final difference
Fourier excursions 0.28 and -0.29e4-3 . Atomic scattering factors from
International Tables for X-ray Crystallography (1974).



2-Oxa-6,9-diaza-6,9-dinitrosopropane (3,6-decane)

1-Aza-3,5,7-trinitroadamantane

Abstract

$C_9H_{12}N_4O_6$, $M_r = 272.22$, trigonal, $a = b = 12.821(3)$, $c = 5.937\text{\AA}$, $\alpha = 90.00$, $\beta = 90.00$, and $\gamma = 120.00^\circ$, $V = 845.2(4)\text{\AA}^3$, $Z = 3$, $D_x = 1.604$ mg mm^{-3} , $\lambda(\text{CuK}\alpha) = 1.54178\text{\AA}$, $\mu = 1.13 \text{ mm}^{-1}$, $F(000) = 426$, $T = 295\text{K}$.

The cage nitrogen lies on a three fold axis.



Clear $0.15 \times 0.08 \times 0.20 \text{ mm}$. crystal. Automated Nicolet R3m diffractometer with incident beam graphite monochromator $\lambda = 1.54178\text{\AA}(\text{CuK}\alpha$, 20 centered reflections within $20 < 2\theta < 50^\circ$ used for determining lattice parameters. $(\sin\theta/\lambda)_{\max} = 0.59 \text{ \AA}^{-1}$, range of $hkl: -15 \leq h \leq 12$, $-14 \leq k \leq 12$, $-6 \leq l \leq 0$. Standards $CO\bar{3}$, $3\bar{2}0$, $\bar{1}30$, monitored every 60 reflections with random variation 3.5% over data collection, $\theta/2\theta$ mode, scan width $(2.0 + \Delta_{\text{alc2}})^\circ$, scan rate a function of count rate ($2^\circ/\text{min. minimum}$, $30^\circ/\text{min. maximum}$), 3016 reflections measured, 1017 unique, $R_{\text{int}} = 0.035$. Data corrected for Lorentz, polarization but not absorption effects, Structure solved by direct methods. The least-squares refinement used program SHELXL (Sheldrick 1980). $\Sigma w(|F_O| - |F_C|)^2$ minimized where $w = 1/[c^2(|F_O|) + g \cdot (F_O)^2]$, $g = 0.00060$.

The structure analysis is not yet complete, however, atomic assignment is definitive.

1-Aza-3,5,7-trinitroadamantane

$d = 1.604$

(O)



7-Nitro-5,6-dihydro-7H-imidazo[1,2-d]tetrazole


Abstract

$C_3H_4N_6O_2$, $M_r = 156.11$, monoclinic, $P2_1/c$, $a = 6.214(2)$,
 $b = 8.692(3)$, $c = 11.664(3)\text{\AA}$, $\beta = 102.42(2)^\circ$. Decomposes without melting.
 $V = 615.2(3)\text{\AA}^3$, $Z = 4$, $D_x = 1.685 \text{ mg mm}^{-3}$, $\lambda(\text{CuKa}) = 1.54178\text{\AA}$,
 $\mu = 1.195 \text{ mm}^{-1}$, $F(000) = 320$, $T = 295\text{K}$, Final $R = 0.041$, $wR = 0.060$ for
986 independent observed reflections. The compound has a whole molecule
disorder with a 12% occupancy. The disordered molecule is related to the
primary molecule via a pseudo two fold axis passing midway through the
 $N(7)-C(8)$ and $C(5)-C(6)$ bonds.

Experimental

Clear $0.60 \times 0.12 \times 0.08 \text{ mm}$. crystal crystallized from acetone.
Automated Nicolet R3m diffractometer with incident beam graphite mono-
chromator $\lambda = 1.54178\text{\AA}(\text{CuKa})$, 25 centered reflections within $30^\circ < 2\theta < 73^\circ$
used for determining lattice parameters. Data corrected for Lorentz and
polarization effects, $(\sin\theta/\lambda)_{\text{max}} = 0.59\text{\AA}^{-1}$, range of hkl :
 $-7 \leq h \leq 0$, $0 \leq k \leq 9$, $-13 \leq l \leq 13$. Standards $\bar{3}00$, $0\bar{4}0$, 008 , monitored
every 60 reflections with random variation 2.6% over data collection, $\theta/2\theta$ mode,
scan width $(2.0 + \Delta_{\text{clat}})^\circ$, scan rate a function of count rate ($4^\circ/\text{min. minimum}$,
 $30^\circ/\text{min. maximum}$), 1243 reflections measured, 1049 unique, $R_{\text{int}} = 0.011$,
986 observed $F_o > 3\sigma(F_o)$. Structure solved by direct methods. The
least-squares refinement used program SHELXTL (Sheldrick 1980). $\Sigma w(|F_o| - |F_c|)^2$
minimized where $w = 1/[s^2(|F_o|) + g \cdot (F_o)^2]$, $s = 0.00023$, 130 parameters
refined: atom coordinates, anisotropic temperature factors for all non-H
atoms, isotropic temperature factors for H atoms. $(\Delta/c)_{\text{max}} = 0.10$,
 $R = 0.041$, $wR = 0.060$, $S = 2.657$. Final difference Fourier excursions 0.13
and -0.14 e\AA^{-3} . Atomic scattering factors from International Tables for
X-ray Crystallography (1974).

7-Nitro-5,6-dihydro-7H-1mildazolo(1,2-d)tetrazole



Appendix A

This appendix lists atom coordinates in a fractional or 'crystallographic' coordinate system. In addition tables of bond distances and angles as well as tables of anisotropic thermal parameters are provided. The alphabetic labels follow those given in the index.

TABLE 1a Atom coordinates ($\times 10^4$) and temperature factors ($\text{Å}^2 \times 10^3$)

atom	x/a	y/b	z/c	U_{eq}
N(1)	2878(6)	6381(6)	5825(10)	43(2)*
C(2)	2750(7)	6067(7)	7668(12)	38(3)*
N(3)	1436(6)	5548(8)	7676(10)	42(2)*
C(4)	837(7)	5543(7)	6118(12)	36(3)*
C(5)	1735(8)	6080(8)	4754(13)	44(3)*
N(6)	3908(7)	7050(7)	5336(12)	53(3)*
O(7)	3875(6)	7334(7)	3769(11)	66(3)*
C(8)	2802(8)	7150(9)	8882(14)	55(3)*
C(9)	3721(8)	5036(8)	8144(15)	49(3)*
O(10)	-244(5)	5108(6)	5841(8)	49(2)*
F(11)	2358(6)	6872(6)	10454(8)	81(2)*
F(12)	3998(5)	7593(5)	9110(10)	79(2)*
F(13)	2091(5)	8060(5)	8259(11)	85(2)*
F(14)	4933(5)	5409(5)	8169(10)	76(2)*
F(15)	3600(6)	4122(5)	7057(10)	83(3)*
F(16)	3442(5)	4610(6)	9745(10)	81(2)*
Hn(3)	1025(83)	5432(82)	8627(166)	61(29)

* Equivalent isotropic U defined as one third of the trace of the orthogonalised U_{ij} tensor

TABLE 2a Bond lengths (Å)

N(1)-C(2)	1.451(11)	N(1)-C(5)	1.461(11)
N(1)-N(6)	1.337(10)	C(2)-N(3)	1.457(10)
C(2)-C(8)	1.508(13)	C(2)-C(9)	1.549(12)
N(3)-C(4)	1.335(11)	N(3)-Hn(3)	0.847(116)
C(4)-C(5)	1.507(12)	C(4)-O(10)	1.221(9)
N(6)-O(7)	1.233(12)	C(8)-F(11)	1.315(12)
C(8)-F(12)	1.326(10)	C(8)-F(13)	1.324(11)
C(9)-F(14)	1.304(10)	C(9)-F(15)	1.307(12)
C(9)-F(16)	1.336(13)		

TABLE 3a Bond angles (deg.)

C(2)-N(1)-C(5)	114.4(6)	C(2)-N(1)-N(6)	118.1(7)
C(5)-N(1)-N(6)	126.7(7)	N(1)-C(2)-N(3)	100.4(6)
N(1)-C(2)-C(8)	113.7(7)	N(3)-C(2)-C(8)	109.8(7)
N(1)-C(2)-C(9)	110.1(7)	N(3)-C(2)-C(9)	107.6(7)
C(8)-C(2)-C(9)	114.3(7)	C(2)-N(3)-C(4)	114.8(7)
C(2)-N(3)-Hn(3)	121.3(69)	C(4)-N(3)-Hn(3)	122.1(69)
N(3)-C(4)-C(5)	109.4(7)	N(3)-C(4)-O(10)	124.7(8)
C(5)-C(4)-O(10)	125.8(8)	N(1)-C(5)-C(4)	101.0(7)
N(1)-N(6)-O(7)	112.8(7)	C(2)-C(8)-F(11)	111.2(8)
C(2)-C(8)-F(12)	113.7(7)	F(11)-C(8)-F(12)	106.5(8)
C(2)-C(8)-F(13)	110.9(8)	F(11)-C(8)-F(13)	108.2(8)
F(12)-C(8)-F(13)	106.0(7)	C(2)-C(9)-F(14)	112.4(7)
C(2)-C(9)-F(15)	110.7(8)	F(14)-C(9)-F(15)	109.9(8)
C(2)-C(9)-F(16)	109.5(7)	F(14)-C(9)-F(16)	107.4(8)
F(15)-C(9)-F(16)	106.8(7)		

TABLE 4a Anisotropic temperature factors ($\text{\AA}^2 \times 10^3$)

atom	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}
N(1)	31(3)	65(5)	32(4)	5(4)	2(3)	-17(3)
C(2)	26(4)	52(5)	36(5)	2(4)	-1(4)	1(4)
N(3)	31(4)	73(5)	23(4)	-3(4)	-7(3)	-12(4)
C(4)	34(4)	39(5)	34(5)	-3(4)	-2(3)	-1(4)
C(5)	37(4)	66(6)	28(4)	-9(5)	3(4)	-1(4)
N(6)	44(4)	69(5)	45(5)	10(4)	0(4)	-6(4)
O(7)	61(4)	89(5)	49(5)	12(4)	10(4)	-12(4)
C(8)	50(5)	55(6)	59(6)	-2(5)	-2(5)	-10(5)
C(9)	44(5)	51(5)	51(5)	-3(5)	-15(5)	-4(4)
O(10)	34(3)	78(4)	34(3)	-3(3)	-9(3)	-7(3)
F(11)	99(4)	112(5)	33(3)	-21(3)	20(3)	-26(4)
F(12)	59(3)	90(4)	87(4)	-28(3)	-7(3)	-31(3)
F(13)	98(4)	70(4)	88(4)	-13(4)	-8(4)	29(3)
F(14)	32(2)	89(4)	108(5)	23(4)	-12(3)	-8(3)
F(15)	88(4)	65(3)	96(5)	-16(4)	-18(4)	8(3)
F(16)	69(4)	109(4)	65(4)	38(4)	-4(3)	10(3)

The anisotropic temperature factor exponent takes the form:

$$-2\pi^2(a^2a^*{}^2U_{11} + k^2b^*{}^2U_{22} + \dots + 2hka^*b^*U_{12})$$

TABLE 5a Hydrogen coordinates ($\times 10^4$) and temperature factors ($\text{\AA}^2 \times 10^3$)

atom	x/a	y/b	z/c	U
H(5a)	1362	6793	4224	53
H(5b)	1947	5499	3857	53

TABLE 1b Atom coordinates ($\times 10^4$) and temperature factors ($\text{\AA}^2 \times 10^3$)

atom	x/a	y/b	z/c	U_{eq}
C(1x)	7105(3)	505(4)	7247(1)	44(1)*
N(2x)	5985(2)	1662(4)	7365(1)	48(1)*
C(3x)	5844(3)	1846(4)	7922(1)	42(1)*
N(4x)	7099(2)	1412(4)	8137(1)	52(1)*
C(3ax)	5532(3)	3917(5)	8044(1)	51(1)*
C(3bx)	4811(3)	518(5)	8128(1)	57(1)*
F(1x)	4429(2)	4464(3)	7839(1)	72(1)*
F(2x)	5481(2)	4252(3)	8550(1)	73(1)*
F(3x)	6436(2)	5065(3)	7865(1)	83(1)*
F(4x)	3682(2)	785(3)	7901(1)	80(1)*
F(5x)	4642(2)	690(4)	8633(1)	92(1)*
F(6x)	5134(2)	-1267(3)	8042(1)	97(1)*
C(1)	9188(3)	5485(4)	9432(1)	40(1)*
N(2)	10493(2)	6151(3)	9293(1)	40(1)*
C(3)	10620(3)	5871(4)	8738(1)	42(1)*
N(4)	9338(2)	5693(4)	8520(1)	49(1)*
C(5)	8412(3)	5586(4)	8929(1)	42(1)*
N(6)	7662(2)	7331(3)	8997(1)	44(1)*
C(7)	7545(3)	7738(4)	9544(1)	42(1)*
N(8)	8602(2)	6751(3)	9795(1)	44(1)*
C(3a)	11310(4)	7537(5)	8492(1)	67(1)*
C(3b)	11386(3)	4033(5)	8663(1)	56(1)*
C(7a)	7727(4)	9856(5)	9636(1)	59(1)*
C(7b)	6260(3)	7044(5)	9757(1)	57(1)*
F(1)	12385(3)	7972(4)	8721(1)	126(1)*
F(2)	11514(2)	7297(3)	7998(1)	97(1)*
F(3)	10568(3)	9089(3)	8526(1)	109(1)*
F(4)	12586(2)	4190(4)	8830(1)	97(1)*
F(5)	11440(2)	3496(3)	8167(1)	69(1)*
F(6)	10856(2)	2640(3)	8922(1)	95(1)*
F(7)	7720(2)	10333(3)	10131(1)	86(1)*
F(8)	8835(2)	10437(3)	9450(1)	94(1)*
F(9)	6817(2)	10886(3)	9397(1)	89(1)*
F(10)	6125(2)	7377(3)	10257(1)	82(1)*
F(11)	6171(2)	5173(3)	9692(1)	85(1)*
F(12)	5266(2)	7801(4)	9514(1)	96(1)*

* Equivalent isotropic U defined as one third of the trace of the orthogonalised U_{ij} tensor

TABLE 2b Bond lengths (Å)

C(1x)-N(2x)	1.452(4)	C(1x)-C(1xa)	1.531(6)
C(1x)-N(4xa)	1.460(4)	N(2x)-C(3x)	1.447(4)
C(3x)-N(4x)	1.452(4)	C(3x)-C(3ax)	1.532(5)
C(3x)-C(3bx)	1.528(5)	N(4x)-C(1xa)	1.460(4)
C(3ax)-F(1x)	1.325(4)	C(3ax)-F(2x)	1.317(4)
C(3ax)-F(3x)	1.329(4)	C(3bx)-F(4x)	1.321(4)
C(3bx)-F(5x)	1.323(4)	C(3bx)-F(6x)	1.321(4)
C(1)-N(2)	1.435(4)	C(1)-C(5)	1.524(4)
C(1)-N(8)	1.487(4)	N(2)-C(3)	1.447(4)
C(3)-N(4)	1.446(4)	C(3)-C(3a)	1.524(5)
C(3)-C(3b)	1.537(4)	N(4)-C(5)	1.470(4)
C(5)-N(6)	1.443(4)	N(6)-C(7)	1.448(4)
C(7)-N(3)	1.451(4)	C(7)-C(7a)	1.523(5)
C(7)-C(7b)	1.535(5)	C(3a)-F(1)	1.321(4)
C(3a)-F(2)	1.340(4)	C(3a)-F(3)	1.321(4)
C(3b)-F(4)	1.316(4)	C(3b)-F(5)	1.312(4)
C(3b)-F(6)	1.334(4)	C(7a)-F(7)	1.299(5)
C(7a)-F(8)	1.345(5)	C(7a)-F(9)	1.304(4)
C(7b)-F(10)	1.325(4)	C(7b)-F(11)	1.317(4)
C(7b)-F(12)	1.333(4)		

TABLE 3b Bond angles (deg.)

N(2x)-C(1x)-C(1xa)	106.0(3)	N(2x)-C(1x)-N(4xa)	111.3(2)
C(1xa)-C(1x)-N(4xa)	104.3(3)	C(1x)-N(2x)-C(3x)	110.7(2)
N(2x)-C(3x)-N(4x)	104.9(2)	N(2x)-C(3x)-C(3ax)	108.4(2)
N(4x)-C(3x)-C(3ax)	108.2(2)	N(2x)-C(3x)-C(3bx)	112.1(3)
N(4x)-C(3x)-C(3bx)	111.8(2)	C(3ax)-C(3x)-C(3bx)	111.1(3)
C(3x)-N(4x)-C(1xa)	110.3(2)	C(3x)-C(3ax)-F(1x)	112.5(3)
C(3x)-C(3ax)-F(2x)	112.5(3)	F(1x)-C(3ax)-F(2x)	107.3(3)
C(3x)-C(3ax)-F(3x)	110.8(3)	F(1x)-C(3ax)-F(3x)	105.8(3)
F(2x)-C(3ax)-F(3x)	107.6(3)	C(3x)-C(3bx)-F(4x)	113.0(3)
C(3x)-C(3bx)-F(5x)	112.7(3)	F(4x)-C(3bx)-F(5x)	107.1(3)
C(3x)-C(3bx)-F(6x)	110.0(3)	F(4x)-C(3bx)-F(6x)	106.9(3)
F(5x)-C(3bx)-F(6x)	106.7(3)	N(2)-C(1)-C(5)	107.3(2)
N(2)-C(1)-N(8)	111.1(2)	C(5)-C(1)-N(8)	104.8(2)
C(1)-N(2)-C(3)	104.8(2)	N(2)-C(3)-N(4)	105.5(2)
N(2)-C(3)-C(3a)	108.0(2)	N(4)-C(3)-C(3a)	109.5(2)
N(2)-C(3)-C(3b)	110.4(2)	N(4)-C(3)-C(3b)	111.4(2)
C(3a)-C(3)-C(3b)	111.3(3)	C(3)-N(4)-C(5)	109.4(2)
C(1)-C(5)-N(4)	102.6(2)	C(1)-C(5)-N(6)	105.7(2)
N(4)-C(5)-N(6)	114.1(2)	C(5)-N(6)-C(7)	110.1(2)
N(6)-C(7)-N(8)	107.4(2)	N(6)-C(7)-C(7a)	110.1(3)
N(8)-C(7)-C(7a)	110.9(3)	N(6)-C(7)-C(7b)	110.4(3)
N(8)-C(7)-C(7b)	107.2(2)	C(7a)-C(7)-C(7b)	110.2(3)
C(1)-N(8)-C(7)	107.0(2)	C(3)-C(3a)-F(1)	113.4(3)
C(3)-C(3a)-F(2)	112.0(3)	F(1)-C(3a)-F(2)	107.0(3)
C(3)-C(3a)-F(3)	110.8(3)	F(1)-C(3a)-F(3)	106.7(3)
F(2)-C(3a)-F(3)	106.5(3)	C(3)-C(3b)-F(4)	113.6(3)
C(3)-C(3b)-F(5)	112.6(3)	F(4)-C(3b)-F(5)	107.4(3)
C(3)-C(3b)-F(6)	104.2(3)	F(4)-C(3b)-F(6)	107.1(3)
F(5)-C(3b)-F(6)	106.5(3)	C(7)-C(7a)-F(7)	113.7(3)
C(7)-C(7a)-F(8)	108.9(3)	F(7)-C(7a)-F(8)	105.8(3)
C(7)-C(7a)-F(9)	113.1(3)	F(7)-C(7a)-F(9)	108.4(3)
F(8)-C(7a)-F(9)	105.8(3)	C(7)-C(7b)-F(10)	112.4(3)
C(7)-C(7b)-F(11)	104.9(3)	F(10)-C(7b)-F(11)	107.4(3)
C(7)-C(7b)-F(12)	113.1(3)	F(10)-C(7b)-F(12)	106.4(3)
F(11)-C(7b)-F(12)	107.4(3)		

TABLE 4b Anisotropic temperature factors ($\text{Å}^2 \times 10^3$)

atom	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}
C(1x)	46(2)	33(2)	52(2)	-8(1)	1(1)	-2(2)
N(2x)	41(1)	57(2)	46(1)	-7(1)	-2(1)	5(1)
C(3x)	37(2)	44(2)	44(2)	1(1)	0(1)	-5(1)
N(4x)	48(4)	65(2)	44(1)	8(1)	1(1)	-0(1)
C(3ax)	50(2)	50(2)	54(2)	-7(2)	4(2)	-4(2)
C(3bx)	50(2)	59(2)	64(2)	2(2)	3(2)	-8(2)
F(1x)	72(1)	62(1)	82(1)	-13(1)	-15(1)	21(1)
F(2x)	71(1)	87(1)	61(1)	-29(1)	-0(1)	-1(1)
F(3x)	90(2)	47(1)	114(2)	-4(1)	29(1)	-17(1)
F(4x)	45(1)	90(2)	104(2)	9(1)	-5(1)	-17(1)
F(5x)	89(2)	117(2)	70(1)	16(1)	18(1)	-33(1)
F(6x)	87(2)	43(1)	161(2)	8(1)	22(2)	-11(1)
C(1)	43(2)	34(2)	43(2)	3(1)	-0(1)	-6(1)
N(2)	40(1)	41(1)	38(1)	-3(1)	1(1)	-9(1)
C(3)	46(2)	40(2)	40(2)	-1(1)	3(1)	-5(2)
N(4)	46(1)	60(2)	40(1)	-11(1)	-2(1)	-0(1)
C(5)	40(2)	38(2)	47(2)	-5(1)	-5(1)	-8(1)
N(6)	45(1)	48(2)	40(1)	-0(1)	-3(1)	3(1)
C(7)	42(2)	41(2)	41(2)	1(1)	-1(1)	-2(1)
N(8)	45(1)	51(2)	37(1)	1(1)	1(1)	0(1)
C(3a)	89(3)	56(2)	57(2)	5(2)	9(2)	-22(2)
C(3b)	54(2)	63(2)	51(2)	-7(2)	4(2)	4(2)
C(7a)	73(2)	46(2)	58(2)	-3(2)	5(2)	-1(2)
C(7b)	50(2)	61(2)	61(2)	4(2)	2(2)	1(2)
F(1)	113(2)	159(3)	105(2)	29(2)	-13(2)	-97(2)
F(2)	142(2)	95(2)	56(1)	10(1)	32(1)	-34(2)
F(3)	174(3)	48(1)	106(2)	18(1)	28(2)	-13(2)
F(4)	62(1)	140(2)	87(1)	-31(1)	-20(1)	34(1)
F(5)	68(1)	82(1)	57(1)	-22(1)	1(1)	14(1)
F(6)	143(2)	43(1)	102(2)	7(1)	46(2)	16(1)
F(7)	126(2)	67(1)	64(1)	-26(1)	-0(1)	-5(1)
F(8)	100(2)	58(1)	123(2)	-14(1)	32(1)	-35(1)
F(9)	125(2)	49(1)	92(2)	-2(1)	-11(1)	26(1)
F(10)	70(1)	118(2)	60(1)	-3(1)	21(1)	-10(1)
F(11)	75(1)	71(1)	110(2)	-3(1)	27(1)	-30(1)
F(12)	46(1)	139(2)	103(2)	30(2)	-2(1)	8(1)

The anisotropic temperature factor exponent takes the form:

$$-2\pi^2 (h^2 a^2 U_{11} + k^2 b^2 U_{22} + \dots + 2hka^2 b^2 U_{12})$$

TABLE 5b Hydrogen coordinates ($\times 10^4$) and temperature factors ($\text{Å}^2 \times 10^3$)

atom	x/a	y/b	z/c	U_{eq}
H(1x)	6850(22)	-735(19)	7134(9)	36(7)
H(2x)	5308(29)	1202(45)	7254(11)	83(10)
H(4x)	7042(27)	783(41)	8389(10)	65(9)
H(1)	9247(22)	4213(17)	9569(8)	37(7)
H(2)	10568(27)	7460(41)	9393(10)	70(9)
H(4)	9216(27)	4758(41)	8293(10)	71(9)
H(5)	7889(19)	4498(23)	8862(9)	35(7)
H(6)	6919(24)	7275(37)	8853(9)	52(8)
H(8)	8435(25)	6175(39)	10078(9)	57(8)

TABLE 1C Atom coordinates ($\times 10^4$) and temperature factors ($\text{Å}^2 \times 10^3$)

atom	x/a	y/b	z/c	U_{eq}
C(1)	9012(3)	3079(8)	2432(6)	50(2)*
N(2)	8296(2)	3212(5)	2433(5)	49(2)*
C(3)	7956(3)	1515(7)	2074(6)	47(2)*
N(4)	4483(3)	303(7)	2293(7)	66(2)*
C(5)	9103(3)	1040(8)	2570(6)	51(2)*
N(6)	9304(3)	941(7)	3909(6)	52(2)*
C(7)	9565(3)	2632(8)	4379(7)	49(2)*
N(8)	9336(4)	3865(7)	3462(8)	82(3)*
N(2')	8025(3)	4888(6)	2107(7)	60(2)*
O(2a)	8426(3)	6151(6)	2076(7)	32(2)*
O(2b)	7433(3)	5022(7)	1931(8)	96(3)*
N(6')	9538(3)	-731(8)	4286(7)	70(2)*
O(6a)	9424(3)	-1990(5)	3589(6)	77(2)*
O(6b)	9811(4)	-816(8)	5286(7)	107(3)*
C(3a)	7354(4)	1042(11)	2881(9)	80(3)*
C(3b)	7766(4)	1384(11)	685(8)	70(3)*
C(7a)	9254(5)	3136(15)	5640(10)	100(4)*
C(7b)	10332(4)	2760(11)	4425(10)	78(3)*
F(1)	5830(4)	1977(8)	2696(13)	218(7)*
F(2)	7186(2)	-668(6)	2737(5)	38(2)*
F(3)	7545(5)	1186(11)	4068(7)	171(4)*
F(4)	7223(4)	2193(10)	379(8)	153(3)*
F(5)	7727(3)	-281(7)	336(5)	101(2)*
F(6)	8256(5)	2161(12)	49(6)	164(4)*
F(7)	9392(4)	4815(8)	5931(7)	136(3)*
F(8)	8546(4)	3019(12)	5546(10)	136(5)*
F(9)	9437(7)	2121(11)	6538(8)	202(6)*
F(10)	10529(3)	4399(7)	4413(8)	118(3)*
F(11)	10560(3)	1954(10)	3459(11)	173(4)*
F(12)	10601(3)	2047(8)	5397(9)	151(4)*

* Equivalent isotropic U defined as one third of the trace of the orthogonalised U_{ij} tensor

TABLE 2C Bond lengths (Å)

C(1)-N(2)	1.440(8)	C(1)-C(5)	1.517(5)
C(1)-N(8)	1.414(10)	N(2)-C(3)	1.474(7)
N(2)-N(2')	1.393(6)	C(3)-N(4)	1.403(3)
C(3)-C(3a)	1.532(11)	C(3)-C(3b)	1.555(11)
N(4)-C(5)	1.390(8)	C(5)-N(6)	1.507(9)
N(6)-C(7)	1.443(8)	N(6)-N(6')	1.378(3)
C(7)-N(8)	1.421(10)	C(7)-C(7a)	1.547(13)
C(7)-C(7b)	1.542(9)	N(2')-O(2a)	1.230(7)
N(2')-O(2b)	1.206(9)	N(6')-O(6a)	1.216(3)
N(6')-O(6b)	1.216(11)	C(3a)-F(1)	1.272(11)
C(3a)-F(2)	1.311(9)	C(3a)-F(3)	1.347(13)
C(3b)-F(4)	1.277(11)	C(3b)-F(5)	1.283(9)
C(3b)-F(6)	1.331(12)	C(7a)-F(7)	1.304(12)
C(7a)-F(8)	1.327(12)	C(7a)-F(9)	1.279(14)
C(7b)-F(10)	1.269(9)	C(7b)-F(11)	1.287(14)
C(7b)-F(12)	1.294(13)		

TABLE 3C Bond angles (deg.)

N(2)-C(1)-C(5)	100.8(4)	N(2)-C(1)-N(8)	115.4(6)
C(5)-C(1)-N(8)	105.7(5)	C(1)-N(2)-C(3)	113.8(4)
C(1)-N(2)-N(2')	116.8(4)	C(3)-N(2)-N(2')	120.0(5)
N(2)-C(3)-N(4)	98.3(5)	N(2)-C(3)-C(3a)	114.0(6)
N(4)-C(3)-C(3a)	110.7(6)	N(2)-C(3)-C(3b)	115.0(5)
N(4)-C(3)-C(3b)	108.0(6)	C(3a)-C(3)-C(3b)	110.1(6)
C(3)-N(4)-C(5)	117.6(5)	C(1)-C(5)-N(4)	104.9(5)
C(1)-C(5)-N(6)	100.1(5)	N(4)-C(5)-N(6)	115.4(6)
C(5)-N(6)-C(7)	113.3(5)	C(5)-N(6)-N(6')	114.8(5)
C(7)-N(6)-N(6')	122.7(6)	N(6)-C(7)-N(8)	100.7(6)
N(6)-C(7)-C(7a)	111.8(6)	N(8)-C(7)-C(7a)	109.5(6)
N(6)-C(7)-C(7b)	115.3(5)	N(8)-C(7)-C(7b)	107.9(6)
C(7a)-C(7)-C(7b)	111.0(7)	C(1)-N(8)-C(7)	116.1(5)
N(2)-N(2')-O(2a)	114.8(6)	N(2)-N(2')-O(2b)	119.9(5)
O(2a)-N(2')-O(2b)	125.3(5)	N(6)-N(6')-O(6a)	115.5(7)
N(6)-N(6')-O(6b)	117.6(6)	O(6a)-N(6')-O(6b)	126.8(6)
C(3)-C(3a)-F(1)	116.1(8)	C(3)-C(3a)-F(2)	110.7(7)
F(1)-C(3a)-F(2)	106.6(7)	C(3)-C(3a)-F(3)	107.6(7)
F(1)-C(3a)-F(3)	110.1(10)	F(2)-C(3a)-F(3)	105.2(8)
C(3)-C(3b)-F(4)	115.5(7)	C(3)-C(3b)-F(5)	111.1(6)
F(4)-C(3b)-F(5)	108.4(7)	C(3)-C(3b)-F(6)	107.0(6)
F(4)-C(3b)-F(6)	106.9(8)	F(5)-C(3b)-F(6)	107.5(7)
C(7)-C(7a)-F(7)	110.7(8)	C(7)-C(7a)-F(8)	108.5(9)
F(7)-C(7a)-F(8)	106.9(9)	C(7)-C(7a)-F(9)	114.6(9)
F(7)-C(7a)-F(9)	107.9(9)	F(8)-C(7a)-F(9)	107.9(11)
C(7)-C(7b)-F(10)	111.7(6)	C(7)-C(7b)-F(11)	107.5(7)
F(10)-C(7b)-F(11)	108.6(8)	C(7)-C(7b)-F(12)	114.7(8)
F(10)-C(7b)-F(12)	105.2(8)	F(11)-C(7b)-F(12)	109.0(8)

TABLE 4C Anisotropic temperature factors ($\text{\AA}^2 \times 10^3$)

atom	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}
C(1)	56(3)	38(3)	56(4)	11(3)	-7(3)	-4(3)
N(2)	54(3)	19(2)	74(3)	-1(2)	-10(3)	-2(2)
C(3)	58(4)	26(2)	57(4)	2(3)	-7(3)	-7(2)
N(4)	64(3)	39(3)	96(5)	-10(3)	-29(4)	3(2)
C(5)	55(4)	40(3)	59(4)	-5(3)	-10(3)	11(3)
N(6)	64(3)	42(3)	51(3)	6(2)	-17(3)	2(2)
C(7)	48(3)	37(3)	61(4)	-4(3)	-12(3)	3(3)
N(8)	115(6)	28(3)	104(5)	1(3)	-58(5)	-12(3)
N(2')	65(3)	29(2)	84(4)	3(3)	-22(3)	9(2)
O(2a)	91(4)	35(2)	121(5)	12(3)	-20(4)	-3(2)
O(2b)	96(4)	45(3)	153(6)	3(4)	-33(5)	22(3)
N(6')	62(4)	42(3)	85(5)	15(3)	-24(4)	-10(3)
O(6a)	104(4)	24(2)	103(4)	-1(3)	-35(3)	1(2)
O(6b)	151(6)	60(3)	112(5)	15(3)	-74(5)	-0(3)
C(3a)	74(5)	52(4)	112(7)	-4(4)	30(5)	-10(4)
C(3b)	74(5)	64(4)	72(5)	3(4)	-15(4)	-11(4)
C(7a)	92(6)	107(8)	102(7)	-48(6)	-8(6)	9(5)
C(7b)	64(4)	63(4)	108(7)	-1(5)	-9(5)	-11(4)
F(1)	116(5)	100(4)	438(19)	62(8)	146(9)	23(4)
F(2)	95(3)	69(3)	100(3)	4(3)	-3(3)	-36(2)
F(3)	265(10)	156(6)	93(4)	-43(5)	80(6)	-119(7)
F(4)	137(7)	127(5)	145(6)	-28(5)	-108(6)	42(5)
F(5)	157(5)	72(3)	75(3)	-24(3)	-33(3)	-12(3)
F(6)	240(9)	191(7)	60(3)	7(4)	-10(4)	-108(7)
F(7)	214(7)	82(4)	113(5)	-55(4)	-25(5)	13(4)
F(8)	110(5)	188(7)	260(11)	-133(8)	75(6)	-26(5)
F(9)	383(16)	126(6)	97(5)	20(5)	78(7)	90(8)
F(10)	98(3)	74(3)	181(6)	29(4)	-40(4)	-46(3)
F(11)	88(4)	179(7)	253(10)	-107(7)	66(5)	-57(4)
F(12)	97(4)	115(5)	239(10)	72(6)	-85(5)	-19(3)

The anisotropic temperature factor exponent takes the form:

$$-2\pi^2(h^2a^2U_{11} + k^2b^2U_{22} + \dots + 2hka^2c^2U_{12})$$

TABLE 5C Hydrogen coordinates ($\times 10^{-4}$) and temperature factors ($\text{\AA}^2 \times 10^3$)

atom	x/a	y/b	z/c	U
H(1)	9174(34)	3699(85)	1712(39)	59
H(4)	8390(37)	-754(31)	2230(87)	77
H(5)	9473(22)	579(57)	2102(57)	60
H(8)	9144(44)	4688(96)	3739(39)	98

TABLE 1d Atom coordinates ($\times 10^4$) and temperature factors ($\text{\AA}^2 \times 10^3$)

atom	x/a	y/b	z/c	U_{eq}
C(1)	-1858(4)	7515(2)	1347(3)	42(1)*
N(2)	-1761(4)	6931(1)	2120(2)	58(1)*
C(3)	-455(5)	6360(2)	1941(3)	50(1)*
N(4)	616(4)	6702(1)	1122(2)	48(1)*
C(5)	-22(4)	7403(2)	807(2)	40(1)*
N(6)	1210(3)	7943(1)	1353(2)	42(1)*
C(7)	116(4)	8509(2)	1793(3)	44(1)*
N(8)	-1695(4)	8175(1)	1867(3)	61(1)*
N(2')	-3083(5)	6918(2)	2898(3)	63(1)*
O(2a)	-4382(4)	7331(2)	2769(2)	74(1)*
O(2b)	-2800(5)	6509(2)	3636(3)	93(1)*
N(4')	1359(5)	6286(2)	280(3)	67(1)*
O(4a)	1985(5)	5722(2)	574(3)	98(1)*
O(4b)	1259(4)	6531(2)	-621(2)	75(1)*
N(6')	3004(4)	8035(2)	995(3)	56(1)*
O(6a)	3570(4)	7564(2)	452(2)	67(1)*
O(6b)	3897(4)	8549(2)	1328(3)	87(1)*
C(3a)	941(6)	6194(2)	2976(3)	65(1)*
C(3b)	-1644(6)	5698(2)	1487(3)	65(1)*
C(7a)	957(6)	8732(2)	2970(3)	62(1)*
C(7b)	-169(5)	9159(2)	1036(3)	55(1)*
F(1)	280(4)	5697(2)	3580(2)	94(1)*
F(2)	2613(4)	5968(2)	2703(2)	91(1)*
F(3)	1265(4)	6757(1)	3566(2)	92(1)*
F(4)	-3214(4)	5635(1)	1989(2)	92(1)*
F(5)	-684(4)	5120(1)	1629(2)	98(1)*
F(6)	-2166(4)	5798(1)	451(2)	85(1)*
F(7)	-336(6)	9140(2)	3372(3)	86(1)*
F(8)	1001(11)	8159(2)	3604(3)	127(3)*
F(9)	2595(6)	9029(4)	3044(3)	125(2)*
F(10)	-1690(12)	9520(5)	1251(10)	78(3)*
F(11)	-476(11)	8980(2)	35(3)	100(2)*
F(12)	1254(9)	9609(3)	1140(9)	111(3)*
F(7')	-277(12)	8956(6)	3556(7)	81(3)*
F(8')	2012(12)	8240(4)	3407(5)	71(3)*
F(9')	2110(11)	9298(4)	2908(6)	66(3)*
F(10')	-1182(34)	9633(13)	1482(25)	132(9)*
F(11')	-1175(14)	8898(4)	166(6)	89(3)*
F(12')	1421(22)	9387(7)	688(8)	77(4)*

* Equivalent isotropic U defined as one third of the trace of the orthogonalised U_{ij} tensor

TABLE 2D Bond lengths (Å)

C(1)-N(2)	1.464(4)	C(1)-C(5)	1.517(4)
C(1)-N(8)	1.414(4)	N(2)-C(3)	1.453(4)
N(2)-N(2')	1.395(5)	C(3)-N(4)	1.467(4)
C(3)-C(3a)	1.565(5)	C(3)-C(3b)	1.586(5)
N(4)-C(5)	1.451(4)	N(4)-N(4')	1.444(4)
C(5)-N(6)	1.467(4)	N(5)-C(7)	1.453(4)
N(6)-N(6')	1.381(4)	C(7)-N(8)	1.429(4)
C(7)-C(7a)	1.573(4)	C(7)-C(7b)	1.555(4)
N(2')-O(2a)	1.203(4)	N(2')-O(2b)	1.202(5)
N(4')-O(4a)	1.206(4)	N(4')-O(4b)	1.202(4)
N(6')-O(6a)	1.210(4)	N(6')-O(6b)	1.214(4)
C(3a)-F(1)	1.317(5)	C(3a)-F(2)	1.320(5)
C(3a)-F(3)	1.306(5)	C(3b)-F(4)	1.316(5)
C(3b)-F(5)	1.296(4)	C(3b)-F(6)	1.310(4)
C(7a)-F(7)	1.325(5)	C(7a)-F(8)	1.344(5)
C(7a)-F(9)	1.274(5)	C(7a)-F(7')	1.253(10)
C(7a)-F(8')	1.281(9)	C(7a)-F(10')	1.355(8)
C(7b)-F(10)	1.316(10)	C(7b)-F(11)	1.281(5)
C(7b)-F(12)	1.312(7)	C(7b)-F(10')	1.303(26)
C(7b)-F(11')	1.325(8)	C(7b)-F(12')	1.304(15)

Table 3d Bond angles (deg.)

π(2)-C(1)-C(5)	100.9(2)	π(2)-C(1)-N(8)	112.7(3)
C(5)-C(1)-N(8)	106.3(2)	C(1)-N(2)-C(3)	117.6(3)
C(1)-N(2)-N(2')	113.0(3)	C(3)-N(2)-N(2')	123.9(3)
N(2)-C(3)-N(4)	98.0(2)	N(2)-C(3)-C(3a)	112.4(3)
N(4)-C(3)-C(3a)	109.4(3)	N(2)-C(3)-C(3b)	109.6(3)
N(4)-C(3)-C(3b)	113.3(3)	C(3a)-C(3)-C(3b)	113.1(3)
C(3)-N(4)-C(5)	115.5(2)	C(3)-N(4)-N(4')	105.3(2)
C(1)-C(5)-N(6)	100.6(2)	N(4)-C(5)-N(6)	111.9(2)
C(5)-N(6)-C(7)	112.7(2)	C(5)-N(6)-N(6')	117.0(2)
C(7)-N(6)-N(6')	123.4(2)	N(6)-C(7)-N(8)	101.1(2)
N(6)-C(7)-C(7a)	112.3(2)	N(8)-C(7)-C(7a)	108.6(3)
N(6)-C(7)-C(7b)	114.2(3)	N(8)-C(8)-C(7b)	109.3(3)
C(7a)-C(7)-C(7b)	110.8(3)	C(1)-N(8)-C(7)	113.8(3)
π(2)-N(2')-C(2a)	116.0(3)	N(2)-N(2')-O(2b)	117.4(3)
O(2a)-N(2')-O(2b)	126.5(4)	N(4)-N(4')-O(4a)	114.8(3)
π(4)-N(4')-O(4b)	117.3(3)	O(4a)-N(4')-O(4b)	127.9(4)
N(6)-N(6')-C(6a)	115.9(3)	N(6)-N(6')-O(6b)	116.6(3)
O(6a)-N(6')-O(6b)	127.1(3)	C(3)-C(3a)-F(1)	112.5(3)
C(3)-C(3a)-F(2)	111.0(3)	F(1)-C(3a)-F(2)	106.0(3)
C(3)-C(3a)-F(3)	110.5(3)	F(1)-C(3a)-F(3)	109.3(3)
F(2)-C(3a)-F(3)	107.4(3)	C(3)-C(3b)-F(4)	109.3(3)
C(3)-C(3b)-F(5)	112.5(3)	F(4)-C(3b)-F(5)	107.7(3)
C(3)-C(3b)-F(6)	108.5(3)	F(4)-C(3b)-F(6)	107.7(3)
F(5)-C(3b)-F(6)	110.5(3)	C(7)-C(7a)-F(7)	107.2(3)
C(7)-C(7a)-F(8)	107.7(3)	F(7)-C(7a)-F(8)	103.9(4)
C(7)-C(7a)-F(9)	116.3(3)	F(7)-C(7a)-F(9)	110.6(4)
F(8)-C(7a)-F(9)	110.4(5)	C(7)-C(7a)-C(7')	114.0(5)
C(7)-C(7a)-F(9')	109.7(4)	F(3')-C(7a)-F(9')	106.6(6)
F(7')-C(7a)-F(9')	101.3(7)	C(7)-C(7b)-F(11)	111.5(3)
C(7)-C(7b)-F(10)	110.9(6)	C(7)-C(7b)-F(12)	114.3(5)
F(10)-C(7b)-F(11)	105.6(7)	F(11)-C(7b)-F(12)	109.0(5)
F(10)-C(7b)-F(12)	105.0(6)	C(7)-C(7b)-F(12')	113.6(5)
C(7)-C(7b)-F(10')	110.2(13)	F(10')-C(7b)-F(12')	114.9(12)
C(7)-C(7b)-C(11')	102.4(4)		
F(10')-C(7b)-F(11')	109.6(12)		
F(11')-C(7b)-F(12')	105.3(7)		

TABLE 4d Anisotropic temperature factors ($\text{Å}^2 \times 10^3$)

atom	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}
C(1)	40(2)	40(2)	46(2)	0(1)	3(1)	-4(1)
N(2)	69(2)	47(1)	61(2)	11(1)	30(1)	10(1)
C(3)	61(2)	36(2)	52(2)	3(1)	1(2)	1(1)
N(4)	58(2)	39(1)	49(1)	-2(1)	16(1)	2(1)
C(5)	39(1)	39(1)	41(2)	2(1)	2(1)	4(1)
N(6)	31(1)	40(1)	56(1)	0(1)	8(1)	1(1)
C(7)	45(2)	33(1)	53(2)	-4(1)	5(1)	2(1)
N(8)	47(2)	42(1)	99(2)	-14(2)	27(2)	-6(1)
N(2')	68(2)	68(2)	56(2)	3(2)	21(2)	-15(2)
O(2a)	66(2)	78(2)	81(2)	-4(2)	30(1)	-4(2)
O(2b)	104(2)	101(2)	77(2)	24(2)	34(2)	-13(2)
N(4')	72(2)	55(2)	74(2)	-18(2)	20(2)	7(2)
O(4a)	136(3)	61(2)	97(2)	-1(2)	23(2)	50(2)
O(4b)	98(2)	79(2)	52(1)	-8(1)	23(1)	11(2)
N(6')	41(1)	61(2)	68(2)	6(2)	8(1)	-4(1)
O(6a)	51(1)	81(2)	72(2)	-11(1)	21(1)	7(1)
O(6b)	58(2)	69(2)	136(3)	-18(2)	25(2)	-27(1)
C(3a)	81(3)	57(2)	54(2)	5(2)	-6(2)	-1(2)
C(3b)	83(3)	46(2)	64(2)	4(2)	-4(2)	-10(2)
C(7a)	80(3)	60(2)	45(2)	-4(2)	6(2)	3(2)
C(7b)	63(2)	43(2)	60(2)	7(2)	5(2)	4(2)
F(1)	112(2)	91(2)	75(1)	35(2)	-7(1)	-7(2)
F(2)	78(2)	95(2)	95(2)	9(2)	-16(1)	17(1)
F(3)	119(2)	82(2)	70(1)	-12(1)	-21(1)	-9(1)
F(4)	97(2)	81(2)	98(2)	11(2)	11(1)	-38(1)
F(5)	135(2)	42(1)	111(2)	-2(1)	-12(2)	-2(1)
F(6)	111(2)	78(2)	52(1)	-3(1)	-20(1)	-25(1)
F(7)	133(3)	66(2)	63(2)	-22(2)	30(2)	5(2)
F(8)	255(7)	72(2)	53(2)	9(2)	8(3)	38(3)
F(9)	67(2)	222(6)	82(2)	-59(3)	-11(2)	-22(3)
F(10)	84(3)	60(3)	93(6)	8(3)	22(3)	32(2)
F(11)	165(5)	80(2)	55(2)	12(2)	19(3)	54(3)
F(12)	85(3)	52(3)	191(8)	43(4)	-19(4)	-22(2)
F(7')	74(5)	109(7)	62(5)	-12(5)	11(4)	-27(5)
F(8')	85(6)	87(6)	38(4)	3(4)	-14(3)	-2(5)
F(9')	64(4)	52(4)	78(5)	-17(4)	-12(4)	-29(4)
F(10')	231(24)	77(11)	80(7)	-22(8)	-25(14)	106(14)
F(11')	115(7)	80(5)	60(4)	-4(4)	-51(4)	6(5)
F(12')	122(8)	57(7)	53(5)	-7(4)	15(5)	-31(6)

The anisotropic temperature factor exponent takes the form:

$$-2\pi^2(n^2a^*^2U_{11} + k^2b^*^2U_{22} + \dots + 2nka^*b^*U_{12})$$

TABLE 5d Hydrogen coordinates ($\times 10^4$) and temperature factors ($\text{Å}^2 \times 10^3$)

atom	x/a	y/b	z/c	U
H(1)	-3114(55)	7508(19)	960(29)	65(11)
H(5)	-177(38)	7451(13)	-32(21)	29(7)
H(8)	-3041(40)	8500(15)	1854(22)	47(8)

TABLE 1E Atom coordinates ($\times 10^4$) and temperature factors ($\text{Å}^2 \times 10^3$)

atom	x/a	y/b	z/c	U_{eq}
C(1)	3569(1)	6743(3)	3424(1)	34(1)*
N(2)	3744(1)	8556(2)	3387(1)	33(1)*
C(3)	4187(1)	8534(3)	3647(1)	33(1)*
N(4)	4248(1)	5627(2)	3549(1)	36(1)*
C(5)	3894(1)	5510(3)	3156(1)	34(1)*
N(6)	3756(1)	5666(3)	2060(1)	42(1)*
C(7)	3322(1)	5868(3)	1711(2)	40(1)*
N(8)	3225(1)	6511(3)	2632(1)	42(1)*
N(2')	3536(1)	9965(2)	3745(1)	44(1)*
O(2a)	3199(1)	9511(3)	3828(2)	55(1)*
O(2b)	3706(1)	11406(2)	3890(1)	57(1)*
N(4')	4634(1)	5804(3)	3709(2)	48(1)*
O(4a)	4911(1)	6794(3)	3656(2)	63(1)*
O(4b)	4641(1)	4236(2)	3896(2)	68(1)*
N(6')	4010(1)	5167(2)	1445(1)	46(1)*
O(6a)	4334(1)	4604(2)	1853(1)	59(1)*
O(6b)	3884(1)	5390(3)	560(1)	64(1)*
N(8')	2844(1)	6175(3)	2864(2)	53(1)*
O(8a)	2571(1)	5933(4)	2164(2)	80(1)*
O(8b)	2832(1)	6221(3)	3720(1)	64(1)*
C(3a)	4360(1)	9681(3)	2878(2)	43(1)*
C(3b)	4355(1)	9117(3)	4762(2)	40(1)*
C(7a)	3201(1)	7323(3)	891(2)	51(1)*
C(7b)	3131(1)	3946(3)	1385(2)	50(1)*
F(1)	4121(1)	11061(2)	2544(1)	62(1)*
F(2)	4720(1)	10299(2)	3243(1)	62(1)*
F(3)	4373(1)	8670(2)	2093(1)	57(1)*
F(4)	4447(1)	10829(2)	4861(1)	59(1)*
F(5)	4680(1)	8190(2)	5160(1)	61(1)*
F(6)	4079(1)	8745(2)	5277(1)	49(1)*
F(7)	2864(1)	8142(3)	980(1)	83(1)*
F(8)	3488(1)	8562(2)	1001(1)	68(1)*
F(9)	3141(1)	6670(2)	-6(1)	70(1)*
F(10)	2774(1)	4051(2)	786(1)	67(1)*
F(11)	3101(1)	3041(2)	2136(1)	71(1)*
F(12)	3367(1)	3027(2)	929(1)	69(1)*

* Equivalent isotropic U defined as one third of the trace of the orthogonalised U_{ij} tensor

TABLE 2E Bond lengths (Å)

C(1)-N(2)	1.482(3)	C(1)-C(5)	1.501(3)
C(1)-N(8)	1.435(2)	N(2)-C(3)	1.475(2)
N(2)-N(2')	1.417(2)	C(3)-N(4)	1.448(2)
C(3)-C(3a)	1.577(3)	C(3)-C(3b)	1.591(3)
N(4)-C(5)	1.431(2)	N(4)-N(4')	1.427(2)
C(5)-N(6)	1.493(2)	N(6)-C(7)	1.462(2)
N(6)-N(6')	1.393(3)	C(7)-N(8)	1.467(3)
C(7)-C(7a)	1.562(3)	C(7)-C(7b)	1.598(3)
N(8)-N(8')	1.425(3)	N(2')-O(2a)	1.207(3)
N(2')-O(2b)	1.217(2)	N(4')-O(4a)	1.211(3)
N(4')-O(4b)	1.197(3)	N(6')-O(6a)	1.203(2)
N(6')-O(6b)	1.222(2)	N(8')-O(8a)	1.205(2)
N(8')-O(8b)	1.197(3)	C(3a)-F(1)	1.331(2)
C(3a)-F(2)	1.306(2)	C(3a)-F(3)	1.333(3)
C(3b)-F(4)	1.314(2)	C(3b)-F(5)	1.322(2)
C(3b)-F(6)	1.325(3)	C(7a)-F(7)	1.328(3)
C(7a)-F(8)	1.330(3)	C(7a)-F(9)	1.312(3)
C(7b)-F(10)	1.321(3)	C(7b)-F(11)	1.322(3)
C(7b)-F(12)	1.319(3)		

TABLE 3E Bond angles (deg.)

N(2)-C(1)-C(5)	100.4(1)	N(2)-C(1)-N(8)	110.6(2)
C(5)-C(1)-N(8)	105.5(2)	C(1)-N(2)-C(3)	112.4(1)
C(1)-N(2)-N(2')	115.3(2)	C(3)-N(2)-N(2')	119.1(1)
N(2)-C(3)-N(4)	98.5(1)	N(2)-C(3)-C(3a)	109.9(1)
N(4)-C(3)-C(3a)	112.8(2)	N(2)-C(3)-C(3b)	111.7(2)
N(4)-C(3)-C(3b)	109.3(1)	C(3a)-C(3)-C(3b)	113.7(2)
C(3)-N(4)-C(5)	115.5(1)	C(3)-N(4)-N(4')	123.7(1)
C(5)-N(4)-N(4')	120.4(2)	C(1)-C(5)-N(4)	102.3(1)
C(1)-C(5)-N(6)	98.5(1)	N(4)-C(5)-N(6)	115.4(2)
C(5)-N(6)-C(7)	114.5(2)	C(5)-N(6)-N(6')	121.1(1)
C(7)-N(6)-N(6')	122.9(2)	N(6)-C(7)-N(8)	98.4(1)
N(6)-C(7)-C(7a)	113.7(2)	N(8)-C(7)-C(7a)	109.5(2)
N(6)-C(7)-C(7b)	108.8(2)	N(8)-C(7)-C(7b)	112.7(2)
C(7a)-C(7)-C(7b)	112.9(2)	C(1)-N(8)-C(7)	114.0(2)
C(1)-N(8)-N(8')	118.8(2)	C(7)-N(8)-N(8')	121.7(2)
N(2)-N(2')-O(2a)	115.5(2)	N(2)-N(2')-O(2b)	117.2(2)
O(2a)-N(2')-O(2b)	127.3(2)	N(4)-N(4')-O(4a)	115.6(2)
N(4)-N(4')-O(4b)	115.2(2)	O(4a)-N(4')-O(4b)	129.1(2)
N(6)-N(6')-O(6a)	115.7(2)	N(6)-N(6')-O(6b)	116.7(2)
O(6a)-N(6')-O(6b)	127.6(2)	N(8)-N(8')-O(8a)	115.2(2)
N(8)-N(8')-O(8b)	116.4(2)	O(8a)-N(8')-O(8b)	128.3(2)
C(3)-C(3a)-F(1)	111.4(2)	C(3)-C(3a)-F(2)	112.6(2)
F(1)-C(3a)-F(2)	108.3(2)	C(3)-C(3a)-F(3)	109.3(2)
F(1)-C(3a)-F(3)	106.1(2)	F(2)-C(3a)-F(3)	108.9(2)
C(3)-C(3b)-F(4)	113.2(2)	C(3)-C(3b)-F(5)	110.9(2)
F(4)-C(3b)-F(5)	107.6(2)	C(3)-C(3b)-F(6)	108.6(1)
F(4)-C(3b)-F(6)	109.2(2)	F(5)-C(3b)-F(6)	107.0(2)
C(7)-C(7a)-F(7)	110.4(2)	C(7)-C(7a)-F(8)	109.1(2)
F(7)-C(7a)-F(8)	107.4(2)	C(7)-C(7a)-F(9)	113.4(2)
F(7)-C(7a)-F(9)	107.3(2)	F(8)-C(7a)-F(9)	109.1(2)
C(7)-C(7b)-F(10)	112.9(2)	C(7)-C(7b)-F(11)	108.6(2)
F(10)-C(7b)-F(11)	109.5(2)	C(7)-C(7b)-F(12)	110.4(2)
F(10)-C(7b)-F(12)	107.7(2)	F(11)-C(7b)-F(12)	107.6(2)

TABLE 4B Anisotropic temperature factors ($\text{A}^2 \times 10^3$)

atom	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}
C(1)	31(1)	42(1)	28(1)	2(1)	4(1)	-3(1)
N(2)	29(1)	34(1)	34(1)	-2(1)	3(1)	3(1)
C(3)	30(1)	31(1)	36(1)	-1(1)	5(1)	0(1)
N(4)	27(1)	30(1)	47(1)	-1(1)	2(1)	-0(1)
C(5)	33(1)	31(1)	36(1)	2(1)	4(1)	-3(1)
N(6)	35(1)	56(1)	36(1)	-11(1)	9(1)	-6(1)
C(7)	37(1)	49(1)	33(1)	-4(1)	5(1)	-9(1)
N(8)	30(1)	63(1)	32(1)	-6(1)	5(1)	-8(1)
N(2')	42(1)	44(1)	42(1)	-3(1)	1(1)	13(1)
O(2a)	39(1)	73(1)	83(1)	-16(1)	12(1)	12(1)
O(2b)	71(1)	38(1)	63(1)	-10(1)	14(1)	4(1)
N(4')	34(1)	46(1)	59(1)	-4(1)	1(1)	8(1)
O(4a)	32(1)	68(1)	90(1)	-12(1)	15(1)	-4(1)
O(4b)	59(1)	42(1)	96(1)	5(1)	-0(1)	17(1)
N(6')	52(1)	42(1)	49(1)	-13(1)	21(1)	-11(1)
O(6a)	52(1)	56(1)	72(1)	-11(1)	24(1)	3(1)
O(6b)	81(1)	74(1)	44(1)	-11(1)	25(1)	-10(1)
N(8')	31(1)	78(1)	50(1)	-4(1)	8(1)	-5(1)
O(3a)	33(1)	138(2)	62(1)	-17(1)	-3(1)	-15(1)
O(3b)	43(1)	106(1)	48(1)	3(1)	17(1)	-9(1)
C(3a)	50(1)	36(1)	43(1)	-3(1)	11(1)	-8(1)
C(3b)	37(1)	41(1)	38(1)	-3(1)	3(1)	-5(1)
C(7a)	55(1)	56(1)	40(1)	1(1)	4(1)	-10(1)
C(7b)	53(1)	54(1)	40(1)	-4(1)	5(1)	-18(1)
F(1)	84(1)	43(1)	62(1)	18(1)	22(1)	5(1)
F(2)	58(1)	66(1)	67(1)	-6(1)	21(1)	-27(1)
F(3)	77(1)	55(1)	43(1)	-7(1)	26(1)	-13(1)
F(4)	74(1)	48(1)	51(1)	-15(1)	8(1)	-19(1)
F(5)	45(1)	76(1)	52(1)	-6(1)	-14(1)	9(1)
F(6)	52(1)	63(1)	33(1)	-0(1)	9(1)	-4(1)
F(7)	74(1)	97(1)	73(1)	21(1)	7(1)	30(1)
F(8)	84(1)	51(1)	61(1)	8(1)	1(1)	-21(1)
F(9)	88(1)	80(1)	35(1)	4(1)	-0(1)	-20(1)
F(10)	58(1)	89(1)	48(1)	-9(1)	-4(1)	-32(1)
F(11)	85(1)	67(1)	56(1)	12(1)	9(1)	-29(1)
F(12)	81(1)	55(1)	74(1)	-21(1)	21(1)	-19(1)

The anisotropic temperature factor exponent takes the form:

$$-2\pi^2(h^2a^*{}^2U_{11} + k^2b^*{}^2U_{22} + \dots + 2hka^*b^*U_{12})$$

TABLE 5B Hydrogen coordinates ($\times 10^4$) and temperature factors ($\text{A}^2 \times 10^3$)

atom	x	y	z	c
H(1)	3524(7)	6491(33)	3987(21)	53(7)
H(5)	3903(5)	4635(27)	3408(15)	29(5)

TABLE 1f Atom coordinates ($\times 10^4$) and temperature factors ($\text{\AA}^2 \times 10^3$)

atom	x/a	y/b	z/c	U_{eq}
N(1)	1814(5)	2572(3)	2623(5)	57(3)*
C(2)	1053(6)	3192(4)	2004(6)	64(4)*
N(3)	-304(5)	3061(3)	1682(5)	53(3)*
C(4)	-810(6)	2427(4)	890(6)	57(4)*
N(5)	-122(5)	1758(3)	1373(4)	51(3)*
N(6)	211(5)	1692(3)	2630(4)	57(3)*
C(7)	1284(7)	2166(4)	3366(6)	64(4)*
N(8)	2057(5)	2074(3)	1824(5)	60(3)*
C(9)	977(5)	1605(4)	1077(6)	60(3)*
N(10)	-1057(6)	3690(4)	1228(6)	64(3)*
C(11)	-1390(6)	4140(4)	1972(6)	59(3)*
O(11)	-1047(4)	4015(3)	3042(4)	75(2)*
C(12)	-2233(7)	4760(4)	1360(7)	60(4)*
C(13)	-2311(8)	5054(4)	260(8)	75(5)*
C(14)	-3198(11)	5603(5)	-281(9)	106(6)*
C(15)	-3976(9)	5869(5)	264(9)	109(6)*
C(16)	-3884(8)	5601(4)	1341(9)	104(6)*
C(17)	-3000(7)	5060(4)	1918(8)	83(5)*
C(18)	-517(7)	1307(4)	3111(6)	58(3)*
O(18)	-281(4)	1370(3)	4196(4)	77(2)*
C(19)	-1513(6)	822(3)	2300(5)	51(3)*
C(20)	-2672(9)	811(5)	2409(6)	70(4)*
C(21)	-3622(8)	326(6)	1744(8)	92(6)*
C(22)	-3373(8)	-166(5)	1004(8)	90(5)*
C(23)	-2231(7)	-167(4)	875(6)	78(4)*
C(24)	-1309(7)	324(4)	1520(6)	65(4)*
C(25)	3171(7)	2049(4)	1645(6)	62(4)*
O(25)	3220(5)	1619(3)	872(5)	80(3)*
C(26)	4305(7)	2499(4)	2382(6)	49(3)*
C(27)	5231(7)	2526(4)	1921(7)	86(5)*
C(28)	5331(8)	2931(5)	2490(8)	99(5)*
C(29)	6544(7)	3282(4)	3546(7)	71(4)*
C(30)	5535(7)	3243(4)	4013(7)	82(4)*
C(31)	4520(7)	2860(4)	3447(6)	75(4)*

* Equivalent isotropic U defined as one third of the trace of the orthogonalised U_{ij} tensor

TABLE 2^f Bond lengths (Å)

N(1)-C(2)	1.441(8)	N(1)-C(7)	1.447(11)
N(1)-N(8)	1.416(9)	C(2)-N(3)	1.457(9)
N(3)-C(4)	1.465(8)	N(3)-N(10)	1.406(9)
C(4)-N(5)	1.442(8)	N(5)-N(6)	1.411(7)
N(5)-C(9)	1.442(10)	N(6)-C(7)	1.480(3)
N(6)-C(18)	1.361(10)	N(8)-C(9)	1.483(5)
N(8)-C(25)	1.357(11)	N(10)-C(11)	1.360(11)
C(11)-O(11)	1.210(8)	C(11)-C(12)	1.481(9)
C(12)-C(13)	1.390(13)	C(12)-C(17)	1.388(13)
C(13)-C(14)	1.391(13)	C(14)-C(15)	1.365(18)
C(15)-C(16)	1.343(15)	C(16)-C(17)	1.387(11)
C(18)-O(18)	1.228(8)	C(18)-C(19)	1.472(3)
C(19)-C(20)	1.363(13)	C(19)-C(24)	1.379(11)
C(20)-C(21)	1.388(12)	C(21)-C(22)	1.358(15)
C(22)-C(23)	1.360(13)	C(23)-C(24)	1.369(9)
C(25)-O(25)	1.226(10)	C(25)-C(26)	1.499(9)
C(26)-C(27)	1.357(13)	C(26)-C(31)	1.369(10)
C(27)-C(28)	1.385(11)	C(28)-C(29)	1.352(13)
C(29)-C(30)	1.349(14)	C(30)-C(31)	1.378(10)

TABLE 3^f Bond angles (deg.)

C(2)-N(1)-C(7)	114.1(6)	C(2)-N(1)-N(8)	112.5(5)
C(7)-N(1)-N(8)	108.5(5)	N(1)-C(2)-N(3)	112.2(6)
C(2)-N(3)-C(4)	114.7(6)	C(2)-N(3)-N(10)	113.4(6)
C(4)-N(3)-N(10)	110.9(5)	N(3)-C(4)-N(5)	112.5(5)
C(4)-N(5)-N(6)	112.8(5)	C(4)-N(5)-C(9)	116.8(6)
N(6)-N(5)-C(9)	109.5(4)	N(5)-N(6)-C(7)	114.6(6)
N(5)-N(6)-C(18)	122.1(5)	C(7)-N(6)-C(18)	122.7(6)
N(1)-C(7)-N(5)	111.7(6)	N(1)-N(8)-C(9)	116.6(6)
N(1)-N(8)-C(25)	123.7(5)	C(9)-N(8)-C(25)	119.4(6)
N(5)-C(9)-C(8)	109.5(5)	N(3)-N(10)-C(11)	120.8(6)
N(10)-C(11)-O(11)	121.3(6)	N(10)-C(11)-C(12)	114.3(6)
O(11)-C(11)-C(12)	123.8(7)	C(11)-C(12)-C(13)	124.0(8)
C(11)-C(12)-C(17)	117.7(7)	C(13)-C(12)-C(17)	118.3(7)
C(12)-C(13)-C(14)	119.6(10)	C(13)-C(14)-C(15)	121.0(10)
C(14)-C(15)-C(16)	119.8(8)	C(15)-C(16)-C(17)	121.0(10)
C(12)-C(17)-C(16)	120.2(9)	N(6)-C(18)-O(18)	118.1(6)
N(6)-C(18)-C(19)	117.7(6)	O(18)-C(18)-C(19)	124.1(7)
C(18)-C(19)-C(20)	118.1(7)	C(18)-C(19)-C(24)	123.8(7)
C(20)-C(19)-C(24)	117.7(6)	C(19)-C(20)-C(21)	121.4(8)
C(20)-C(21)-C(22)	119.0(9)	C(21)-C(22)-C(23)	120.9(8)
C(22)-C(23)-C(24)	119.5(8)	C(19)-C(24)-C(23)	121.5(8)
N(8)-C(25)-O(25)	116.6(5)	N(8)-C(25)-C(26)	122.1(7)
O(25)-C(25)-C(26)	121.3(7)	C(25)-C(26)-C(27)	113.8(7)
C(25)-C(26)-C(31)	128.5(4)	C(27)-C(26)-C(31)	117.7(7)
C(26)-C(27)-C(28)	120.6(6)	C(27)-C(28)-C(29)	121.5(10)
C(28)-C(29)-C(30)	117.9(7)	C(29)-C(30)-C(31)	121.3(8)
C(26)-C(31)-C(30)	120.9(3)		

TABLE 4f Anisotropic temperature factors ($\text{\AA}^2 \times 10^3$)

atom	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}
N(1)	46(4)	70(5)	53(4)	-10(4)	15(3)	-1(4)
C(2)	48(5)	72(5)	67(5)	-21(4)	15(4)	-15(5)
N(3)	50(4)	57(4)	52(4)	-11(3)	18(3)	2(4)
C(4)	56(5)	70(6)	46(5)	-5(4)	20(4)	-12(5)
N(5)	51(4)	58(4)	44(3)	1(3)	17(3)	-2(3)
N(6)	54(4)	65(4)	45(4)	-3(3)	12(3)	-12(4)
C(7)	53(5)	80(6)	64(5)	-5(5)	29(5)	-2(5)
N(8)	54(4)	64(4)	58(4)	-22(4)	18(4)	-18(4)
C(9)	73(5)	50(5)	58(5)	-11(4)	37(4)	-11(5)
N(10)	60(4)	64(5)	63(5)	-7(4)	24(4)	3(4)
C(11)	43(5)	74(6)	59(5)	-16(4)	18(4)	-7(4)
O(11)	64(3)	106(4)	56(3)	-16(3)	24(3)	-10(3)
C(12)	67(5)	44(5)	70(5)	-19(4)	28(5)	-21(4)
C(13)	33(7)	54(6)	90(7)	-8(5)	35(6)	7(5)
C(14)	138(10)	59(7)	115(9)	-12(6)	41(8)	2(7)
C(15)	97(8)	42(6)	165(10)	-5(6)	21(7)	-9(6)
C(16)	92(7)	54(5)	183(10)	-17(6)	69(7)	8(6)
C(17)	90(7)	67(6)	106(7)	-31(5)	55(6)	-15(6)
C(18)	51(5)	69(6)	58(5)	10(5)	24(4)	7(5)
O(18)	87(4)	107(4)	46(3)	-0(3)	34(3)	-7(3)
C(19)	40(4)	61(5)	53(5)	16(4)	17(4)	1(4)
C(20)	54(6)	102(8)	57(6)	-9(5)	24(5)	3(6)
C(21)	49(7)	142(10)	91(8)	15(7)	30(6)	0(7)
C(22)	85(8)	78(7)	92(7)	19(6)	17(6)	-24(6)
C(23)	103(7)	48(5)	74(5)	-2(4)	25(5)	-17(5)
C(24)	73(6)	52(5)	79(6)	4(5)	39(5)	10(5)
C(25)	56(5)	71(6)	72(5)	3(5)	40(5)	-2(5)
O(25)	92(4)	79(4)	92(4)	-30(3)	61(4)	-10(3)
C(26)	52(5)	52(5)	54(5)	-3(4)	33(4)	2(4)
C(27)	90(7)	92(7)	91(7)	-21(5)	51(6)	-17(6)
C(28)	75(7)	144(9)	99(7)	-24(7)	56(6)	-10(6)
C(29)	51(6)	89(7)	72(6)	19(5)	21(5)	-5(5)
C(30)	67(6)	96(7)	79(6)	-15(5)	24(5)	-15(5)
C(31)	65(5)	87(6)	84(6)	-18(5)	42(5)	-2(5)

The anisotropic temperature factor exponent takes the form:

$$-2\pi^2 (n^2 a^2 U_{11} + k^2 b^2 U_{22} + \dots + 2hka^2 n U_{12})$$

TABLE Sf Hydrogen coordinates ($\times 10^4$) and temperature factors ($\text{Å}^2 \text{x}^2$)

atom	x/a	y/b	z/c	
H(2a)	1298	3515	2521	20
H(2b)	1209	3284	1280	60
H(4a)	-756	2525	122	20
H(4b)	-1686	2359	736	66
H(7a)	1933	1362	3913	76
H(7b)	984	2508	3812	76
H(8a)	1207	1096	1234	20
H(9b)	779	1709	239	66
H(10)	-1114(53)	3738(37)	494(54)	76
H(13)	-1757	4878	-123	83
H(14)	-3262	5798	-1048	115
H(15)	-4587	6246	-122	116
H(16)	-4439	5786	1716	119
H(17)	-2919	4893	2705	101
H(20)	-2834	1146	2957	90
H(21)	-4444	337	1804	95
H(22)	-4012	-518	567	90
H(23)	-3370	-508	335	91
H(24)	-503	322	1428	77
H(27)	5125	2262	1193	80
H(28)	5956	2963	2128	94
H(29)	7319	3550	3948	81
H(30)	5766	3488	4768	85
H(31)	3385	2845	3799	83

TABLE 1G Atom coordinates ($\times 10^4$) and temperature factors ($\text{\AA}^2 \times 10^3$)

atom	x/a	y/b	z/c	U_{eq}
N(1)	-42(1)	2495(1)	3183(1)	38(1)*
C(2)	-565(2)	3397(1)	2924(2)	44(1)*
N(3)	270(1)	4084(1)	2971(1)	42(1)*
C(4)	962(2)	3826(1)	2311(1)	43(1)*
N(5)	1388(1)	2894(1)	2432(1)	40(1)*
C(6)	515(2)	2250(1)	2400(1)	41(1)*
N(7)	-860(1)	1835(1)	3172(1)	44(1)*
C(8)	-946(2)	1519(1)	4090(1)	39(1)*
O(9)	-387(1)	1807(1)	4942(1)	58(1)*
C(10)	-1804(2)	798(1)	4002(2)	40(1)*
C(11)	-1666(2)	200(1)	4838(2)	52(1)*
C(12)	-2436(2)	-481(2)	4779(2)	64(1)*
C(13)	-3348(2)	-564(2)	3911(2)	67(1)*
C(14)	-3486(2)	29(2)	3094(2)	64(1)*
C(15)	-2723(2)	707(1)	3132(2)	50(1)*
N(16)	891(1)	4261(1)	4027(1)	40(1)*
C(17)	1366(2)	5083(1)	4281(1)	38(1)*
O(18)	1276(1)	5701(1)	3641(1)	60(1)*
C(19)	2046(2)	5187(1)	5398(1)	39(1)*
C(20)	2912(2)	5798(2)	5632(2)	64(1)*
C(21)	3588(2)	5899(2)	6658(2)	76(1)*
C(22)	3398(2)	5405(2)	7444(2)	69(1)*
C(23)	2524(2)	4815(2)	7226(2)	62(1)*
C(24)	1848(2)	4700(1)	6212(2)	49(1)*
N(25)	2305(1)	2777(1)	3329(1)	44(1)*
C(26)	3290(2)	2576(1)	3196(1)	40(1)*
O(27)	3422(1)	2459(1)	2331(1)	60(1)*
C(28)	4229(2)	2508(1)	4184(2)	42(1)*
C(29)	5107(2)	1944(2)	4195(2)	61(1)*
C(30)	6001(2)	1873(2)	5079(2)	73(1)*
C(31)	6044(2)	2373(2)	5957(2)	64(1)*
C(32)	5175(2)	2928(2)	5960(2)	62(1)*
C(33)	4270(2)	3006(2)	5076(2)	54(1)*
O(w)	1729(1)	2662(1)	5290(1)	45(1)*

* Equivalent isotropic U defined as one third of the trace of the orthogonalised U_{ij} tensor

TABLE 2G Bond lengths (Å)

N(1)-C(2)	1.478(2)	N(1)-C(6)	1.468(3)
N(1)-N(7)	1.418(2)	C(2)-N(3)	1.451(3)
N(3)-C(4)	1.465(3)	N(3)-N(16)	1.418(2)
C(4)-N(5)	1.465(3)	N(5)-C(6)	1.446(3)
N(5)-N(25)	1.408(2)	N(7)-C(8)	1.345(3)
C(8)-O(9)	1.223(2)	C(8)-C(10)	1.497(3)
C(10)-C(11)	1.390(3)	C(10)-C(15)	1.385(2)
C(11)-C(12)	1.383(3)	C(12)-C(13)	1.376(3)
C(13)-C(14)	1.366(4)	C(14)-C(15)	1.379(3)
N(16)-C(17)	1.349(2)	C(17)-O(18)	1.229(2)
C(17)-C(19)	1.490(2)	C(19)-C(20)	1.380(3)
C(19)-C(24)	1.385(3)	C(20)-C(21)	1.390(3)
C(21)-C(22)	1.354(4)	C(22)-C(23)	1.368(4)
C(23)-C(24)	1.379(3)	N(25)-C(26)	1.343(3)
C(26)-O(27)	1.226(3)	C(26)-C(28)	1.494(2)
C(28)-C(29)	1.382(3)	C(28)-C(33)	1.385(3)
C(29)-C(30)	1.375(3)	C(30)-C(31)	1.370(4)
C(31)-C(32)	1.370(4)	C(32)-C(33)	1.384(3)

TABLE 3G Bond angles (deg.)

C(2)-N(1)-C(5)	109.3(1)	C(2)-N(1)-N(7)	109.5(1)
C(5)-N(1)-N(7)	109.1(1)	N(1)-C(2)-N(3)	110.2(2)
C(2)-N(3)-C(4)	110.5(2)	C(2)-N(3)-N(16)	110.7(1)
C(4)-N(3)-N(16)	112.9(1)	N(3)-C(4)-N(5)	116.3(2)
C(4)-N(5)-C(6)	111.0(2)	C(4)-N(5)-N(25)	113.1(1)
C(6)-N(5)-N(25)	112.0(1)	N(1)-C(6)-N(5)	109.9(1)
N(1)-N(7)-C(8)	118.9(1)	N(7)-C(8)-O(9)	123.1(2)
N(7)-C(8)-C(10)	115.2(1)	O(9)-C(8)-C(10)	121.7(2)
C(8)-C(10)-C(11)	119.0(2)	C(8)-C(10)-C(15)	122.9(2)
C(11)-C(10)-C(15)	119.0(2)	C(10)-C(11)-C(12)	119.5(2)
C(11)-C(12)-C(13)	120.9(2)	C(12)-C(13)-C(14)	119.4(2)
C(13)-C(14)-C(15)	120.6(2)	C(10)-C(15)-C(14)	120.4(2)
N(3)-N(16)-C(17)	119.3(1)	N(16)-C(17)-O(18)	123.1(2)
N(16)-C(17)-C(19)	115.2(2)	O(18)-C(17)-C(19)	121.7(2)
C(17)-C(19)-C(20)	118.5(2)	C(17)-C(19)-C(24)	123.1(2)
C(20)-C(19)-C(24)	118.4(2)	C(19)-C(20)-C(21)	120.6(2)
C(20)-C(21)-C(22)	120.4(2)	C(21)-C(22)-C(23)	119.6(2)
C(22)-C(23)-C(24)	120.9(2)	C(19)-C(24)-C(23)	120.1(2)
N(5)-N(25)-C(26)	118.5(2)	N(25)-C(26)-O(27)	121.7(2)
N(25)-C(26)-C(28)	115.2(2)	O(27)-C(26)-C(28)	121.7(2)
C(26)-C(28)-C(29)	118.5(2)	C(26)-C(28)-C(33)	122.6(2)
C(29)-C(28)-C(33)	118.9(2)	C(28)-C(29)-C(30)	120.6(2)
C(29)-C(30)-C(31)	120.4(2)	C(30)-C(31)-C(32)	119.6(2)
C(31)-C(32)-C(33)	120.5(2)	C(28)-C(33)-C(32)	119.9(2)

TABLE 4g Anisotropic temperature factors ($\text{\AA}^2 \times 10^3$)

atom	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}
N(1)	41(1)	41(1)	32(1)	-5(1)	13(1)	-7(1)
C(2)	41(1)	48(1)	43(1)	-3(1)	12(1)	1(1)
N(3)	48(1)	44(1)	31(1)	-2(1)	6(1)	-2(1)
C(4)	53(1)	47(1)	29(1)	3(1)	11(1)	-3(1)
N(5)	40(1)	51(1)	29(1)	-1(1)	9(1)	-2(1)
C(6)	45(1)	47(1)	31(1)	-5(1)	11(1)	0(1)
N(7)	50(1)	51(1)	29(1)	-10(1)	11(1)	-16(1)
C(8)	44(1)	39(1)	35(1)	-3(1)	11(1)	-0(1)
O(9)	57(1)	72(1)	32(1)	-1(1)	8(1)	-23(1)
C(10)	43(1)	38(1)	42(1)	-3(1)	16(1)	2(1)
C(11)	63(1)	45(1)	51(1)	0(1)	21(1)	-2(1)
C(12)	85(2)	47(1)	72(2)	7(1)	38(1)	-4(1)
C(13)	67(2)	50(1)	92(2)	-8(1)	37(1)	-13(1)
C(14)	48(1)	64(2)	78(2)	-8(1)	16(1)	-10(1)
C(15)	46(1)	49(1)	55(1)	-0(1)	16(1)	-1(1)
N(16)	55(1)	34(1)	29(1)	4(1)	9(1)	1(1)
C(17)	48(1)	33(1)	36(1)	2(1)	14(1)	4(1)
O(18)	90(1)	41(1)	43(1)	11(1)	13(1)	-4(1)
C(19)	46(1)	34(1)	39(1)	-2(1)	14(1)	3(1)
C(20)	72(2)	68(2)	53(1)	-2(1)	20(1)	-24(1)
C(21)	63(2)	89(2)	68(2)	-13(1)	8(1)	-29(1)
C(22)	72(2)	72(2)	51(1)	-8(1)	1(1)	3(1)
C(23)	90(2)	53(1)	38(1)	-3(1)	12(1)	-5(1)
C(24)	68(1)	41(1)	38(1)	-3(1)	15(1)	-7(1)
N(25)	40(1)	65(1)	29(1)	-1(1)	12(1)	2(1)
C(26)	47(1)	40(1)	37(1)	1(1)	16(1)	1(1)
O(27)	59(1)	86(1)	39(1)	0(1)	22(1)	16(1)
C(28)	41(1)	46(1)	41(1)	5(1)	18(1)	2(1)
C(29)	63(1)	71(2)	51(1)	2(1)	17(1)	21(1)
C(30)	60(2)	86(2)	68(2)	8(1)	13(1)	27(1)
C(31)	49(1)	78(2)	58(1)	17(1)	4(1)	0(1)
C(32)	49(1)	85(2)	49(1)	-14(1)	10(1)	-10(1)
C(33)	43(1)	66(1)	52(1)	-12(1)	13(1)	2(1)
O(w)	54(1)	48(1)	29(1)	4(1)	10(1)	-2(1)

The anisotropic temperature factor exponent takes the form:

$$-2\pi^2(h^2 a^*{}^2 U_{11} + k^2 b^*{}^2 U_{22} + \dots + 2hka^*b^* U_{12})$$

TABLE 5G Hydrogen coordinates ($\times 10^4$) and temperature factors ($\text{A}^2 \times 10^3$)

atom	x/a	y/b	z/c	β
H(2a)	-1110(15)	3390(13)	2150(14)	54(5)
H(2b)	-973(14)	3521(12)	3446(14)	52(5)
H(4a)	434(14)	3857(12)	1526(14)	50(5)
H(4b)	1609(15)	4264(13)	2439(15)	59(5)
H(5a)	-7(14)	2237(11)	1683(13)	41(5)
H(5b)	831(15)	1620(13)	2559(14)	57(5)
H(7)	-1153(15)	1510(13)	2580(14)	53(5)
H(11)	-1027(16)	251(14)	5421(15)	65(5)
H(12)	-2340(17)	-853(15)	5312(16)	80(7)
H(13)	-3854(17)	-1064(15)	3883(16)	82(7)
H(14)	-4079(18)	-32(15)	2510(17)	86(7)
H(15)	-2806(14)	1119(12)	2570(14)	51(5)
H(16)	1090(14)	3811(13)	4435(14)	52(5)
H(20)	3071(16)	6101(15)	5069(15)	73(7)
H(21)	4201(19)	6317(19)	6800(19)	112(9)
H(22)	3912(20)	5455(18)	8167(19)	109(9)
H(23)	2346(20)	4429(18)	7800(19)	108(9)
H(24)	1209(16)	4303(14)	6071(15)	65(6)
H(25)	2205(15)	2740(13)	3974(14)	57(6)
H(29)	5049(18)	1618(15)	3559(16)	87(7)
H(30)	6642(20)	1504(18)	5056(19)	115(9)
H(31)	6643(17)	2304(15)	6556(16)	76(7)
H(32)	5174(18)	3271(15)	5562(16)	82(7)
H(33)	3685(14)	3418(12)	5088(14)	55(6)
H(1w)	1051(20)	2367(15)	5133(18)	99(8)
H(2w)	2092(16)	2584(14)	5904(16)	69(6)

TABLE 1h Atom coordinates ($\times 10^4$) and temperature factors ($\text{\AA}^2 \times 10^3$)

atom	x/a	y/b	z/c	U_{eq}
N(1)	6108(1)	2954(1)	4735(1)	51(1)*
N(2)	6239(1)	1619(1)	5002(1)	54(1)*
C(3)	5651(1)	1073(2)	4013(2)	51(1)*
N(4)	5148(1)	1875(1)	2981(1)	48(1)*
C(5)	5367(1)	3204(2)	3479(2)	52(1)*
C(6)	6550(1)	3930(2)	5525(2)	49(1)*
C(7)	7345(1)	3709(2)	6767(2)	48(1)*
C(8)	7600(1)	2536(2)	7426(2)	66(1)*
C(9)	8343(2)	2463(2)	8581(2)	78(1)*
C(10)	8846(1)	3549(2)	9088(2)	74(1)*
C(11)	8594(1)	4713(2)	8451(2)	73(1)*
C(12)	7848(1)	4800(2)	7310(2)	62(1)*
S(6)	6294(1)	5045(1)	5160(1)	63(1)*
N(13)	4207(1)	1595(1)	2424(1)	50(1)*
C(14)	3586(1)	1569(1)	3273(2)	44(1)*
O(14)	3839(1)	1708(1)	4556(1)	53(1)*
C(15)	2579(1)	1378(1)	2557(2)	46(1)*
C(16)	2213(1)	1772(2)	1187(2)	58(1)*
C(17)	1265(1)	1634(2)	614(2)	67(1)*
C(18)	681(1)	1097(2)	1396(2)	67(1)*
C(19)	1047(1)	683(2)	2752(2)	69(1)*
C(20)	1992(1)	822(2)	3334(2)	58(1)*

* Equivalent isotropic U defined as one third of the trace of the orthogonalised U_{ij} tensor

TABLE 2h Bond lengths (Å)

N(1)-N(2)	1.410(2)	N(1)-C(5)	1.467(2)
N(1)-C(6)	1.345(2)	N(2)-C(3)	1.273(2)
C(3)-N(4)	1.383(2)	N(4)-C(5)	1.468(2)
N(4)-N(13)	1.393(2)	C(6)-C(7)	1.499(2)
C(6)-O(6)	1.240(2)	C(7)-C(8)	1.384(2)
C(7)-C(12)	1.386(2)	C(8)-C(9)	1.383(3)
C(9)-C(10)	1.373(3)	C(10)-C(11)	1.367(3)
C(11)-C(12)	1.377(2)	N(13)-C(14)	1.353(2)
C(14)-O(14)	1.228(2)	C(14)-C(15)	1.497(2)
C(15)-C(16)	1.384(2)	C(15)-C(20)	1.384(2)
C(16)-C(17)	1.382(2)	C(17)-C(18)	1.377(3)
C(18)-C(19)	1.378(3)	C(19)-C(20)	1.380(2)

TABLE 3h Bond angles (deg.)

N(2)-N(1)-C(5)	111.8(1)	N(2)-N(1)-C(6)	126.9(1)
C(5)-N(1)-C(6)	121.2(1)	N(1)-N(2)-C(3)	104.7(1)
N(2)-C(3)-N(4)	116.5(1)	C(3)-N(4)-C(5)	106.2(1)
C(3)-N(4)-N(13)	118.6(1)	C(5)-N(4)-N(13)	116.6(1)
N(1)-C(5)-N(4)	100.2(1)	N(1)-C(5)-C(7)	122.4(1)
N(1)-C(6)-O(5)	117.2(1)	C(7)-C(6)-O(5)	120.3(1)
C(6)-C(7)-C(8)	126.1(1)	C(6)-C(7)-C(12)	115.9(1)
C(5)-C(7)-C(12)	118.0(1)	C(7)-C(8)-C(9)	120.5(2)
C(5)-C(9)-C(10)	120.7(2)	C(9)-C(10)-C(11)	119.2(2)
C(10)-C(11)-C(12)	120.6(2)	C(7)-C(12)-C(11)	121.0(2)
N(4)-N(13)-C(14)	120.2(1)	N(13)-C(14)-O(14)	121.5(1)
C(13)-C(14)-C(15)	115.1(1)	O(14)-C(14)-C(15)	122.4(2)
C(14)-C(15)-C(16)	122.9(2)	C(14)-C(15)-C(20)	117.7(1)
C(15)-C(16)-C(20)	119.4(1)	C(15)-C(16)-C(17)	120.0(2)
C(16)-C(17)-C(18)	120.4(2)	C(17)-C(18)-C(19)	119.6(2)
C(18)-C(19)-C(20)	120.3(2)	C(15)-C(20)-C(19)	120.2(2)

TABLE 4h Anisotropic temperature factors ($\text{A}^2 \times 10^3$)

atom	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}
N(1)	54(1)	41(1)	49(1)	4(1)	-6(1)	-1(1)
N(2)	58(1)	42(1)	53(1)	0(1)	-5(1)	2(1)
C(3)	53(1)	45(1)	49(1)	-2(1)	-1(1)	3(1)
N(4)	43(1)	54(1)	43(1)	-1(1)	0(1)	-1(1)
C(5)	51(1)	51(1)	49(1)	8(1)	-1(1)	-2(1)
C(6)	51(1)	43(1)	50(1)	2(1)	7(1)	-1(1)
C(7)	47(1)	49(1)	47(1)	-5(1)	8(1)	-0(1)
C(8)	70(1)	54(1)	61(1)	-4(1)	-12(1)	1(1)
C(9)	82(1)	69(1)	68(1)	-7(1)	-18(1)	15(1)
C(10)	59(1)	89(1)	64(1)	-20(1)	-9(1)	11(1)
C(11)	62(1)	76(1)	73(1)	-23(1)	-0(1)	-13(1)
C(12)	63(1)	56(1)	63(1)	-7(1)	6(1)	-7(1)
O(5)	58(1)	45(1)	69(1)	7(1)	-8(1)	1(1)
N(13)	45(1)	65(1)	38(1)	-6(1)	2(1)	-4(1)
C(14)	51(1)	38(1)	43(1)	-0(1)	5(1)	2(1)
O(14)	50(1)	56(1)	40(1)	-3(1)	6(1)	1(1)
C(15)	47(1)	40(1)	49(1)	-2(1)	7(1)	2(1)
C(16)	52(1)	64(1)	55(1)	10(1)	7(1)	4(1)
C(17)	57(1)	77(1)	61(1)	11(1)	-2(1)	5(1)
C(18)	47(1)	69(1)	80(1)	-2(1)	2(1)	-2(1)
C(19)	56(1)	61(1)	73(1)	5(1)	16(1)	-8(1)
C(20)	56(1)	61(1)	55(1)	2(1)	11(1)	-4(1)

The anisotropic temperature factor exponent takes the form:

$$-2\pi^2 (h^2 a^*{}^2 U_{11} + k^2 b^*{}^2 U_{22} + \dots + 2hk a^* b^* U_{12})$$

TABLE 5h Hydrogen coordinates ($\times 10^4$) and temperature factors ($\text{Å}^2 \times 10^3$)

atom	x/a	y/b	z/c	U
H(3)	5579(11)	113(17)	3949(16)	64(5)
H(5a)	4799(12)	3668(16)	3724(17)	58(5)
H(20)	2268(12)	489(17)	4292(17)	73(5)
H(5b)	5625(12)	3706(16)	2834(18)	58(5)
H(18)	24(12)	983(17)	1027(18)	73(5)
H(13)	4059(13)	1388(17)	1554(19)	75(5)
H(10)	9377(15)	3482(18)	9933(21)	88(6)
H(15)	2610(12)	2125(17)	644(17)	71(5)
H(17)	1015(14)	1933(18)	-337(19)	83(6)
H(19)	618(14)	349(19)	3291(19)	93(6)
H(8)	7230(15)	1786(18)	7050(20)	77(6)
H(9)	8551(17)	1626(22)	8974(25)	110(8)
H(12)	7638(12)	5615(18)	6849(17)	75(5)
H(11)	8912(15)	5474(22)	8806(21)	101(7)

TABLE II Atom coordinates ($\times 10^4$) and temperature factors ($\text{Å}^2 \times 10^3$)

atom	x/a	y/c	z/c	U_{eq}
N(1)	11097(3)	2273(2)	2142(2)	52(1)*
N(2)	9636(3)	2549(2)	3018(2)	54(1)*
C(3)	9000(4)	1595(3)	3681(3)	53(1)*
N(4)	9356(3)	545(2)	2903(2)	48(1)*
C(5)	10910(4)	-267(3)	2758(3)	53(1)*
N(6)	12201(3)	174(2)	1998(2)	51(1)*
N(7)	11863(3)	509(2)	950(2)	51(1)*
C(8)	11032(4)	1795(3)	1051(3)	59(1)*
C(9)	12304(4)	1346(3)	2420(2)	54(1)*
C(10)	9200(4)	3669(3)	3526(3)	68(2)*
O(11)	8270(3)	3899(2)	4509(2)	98(1)*
C(12)	9840(4)	4637(3)	2861(3)	63(2)*
C(13)	9466(6)	4974(4)	1895(3)	104(2)*
C(14)	9976(7)	5923(4)	1303(4)	131(3)*
C(15)	10846(6)	6522(3)	1698(4)	110(3)*
C(16)	11237(5)	6174(4)	2661(4)	96(2)*
C(17)	10718(4)	5242(3)	3259(3)	31(2)*
N(18)	8160(3)	-67(2)	3119(2)	54(1)*
C(19)	7832(4)	-724(3)	4048(3)	53(1)*
O(20)	8428(3)	-703(2)	4820(2)	71(1)*
C(21)	6729(4)	-1494(3)	4052(2)	50(1)*
C(22)	6882(4)	-2530(3)	4620(3)	72(2)*
C(23)	5929(5)	-3304(3)	4617(4)	85(2)*
C(24)	4812(4)	-3040(4)	4066(4)	83(2)*
C(25)	4633(4)	-2002(4)	3524(4)	90(2)*
C(26)	5593(4)	-1230(3)	3504(3)	73(2)*
C(27)	12488(4)	-112(3)	-78(3)	54(1)*
O(28)	12137(3)	373(2)	-909(2)	77(1)*
C(29)	13515(4)	-1349(3)	-223(3)	61(1)*
C(30)	14510(4)	-1776(3)	422(3)	75(2)*
C(31)	15473(5)	-2936(4)	176(4)	107(2)*
C(32)	15453(5)	-3660(4)	-716(4)	124(3)*
C(33)	14485(5)	-3232(3)	-1362(4)	117(2)*
C(34)	13526(4)	-2088(3)	-1134(3)	83(2)*

* Equivalent isotropic U defined as one third of the trace of the orthogonalised U_{ij} tensor

TABLE 2i Bond lengths (Å)

N(1)-N(2)	1.412(3)	N(1)-C(8)	1.458(4)
N(1)-C(9)	1.464(4)	N(2)-C(3)	1.482(4)
N(2)-C(10)	1.345(4)	C(3)-N(4)	1.445(4)
N(4)-C(5)	1.466(4)	N(4)-N(18)	1.406(4)
C(5)-N(6)	1.464(4)	N(6)-N(7)	1.426(4)
N(6)-C(9)	1.473(4)	N(7)-C(8)	1.481(4)
N(7)-C(27)	1.337(3)	C(10)-O(11)	1.232(4)
C(10)-C(12)	1.498(5)	C(12)-C(13)	1.347(6)
C(12)-C(17)	1.364(6)	C(13)-C(14)	1.396(6)
C(14)-C(15)	1.352(8)	C(15)-C(16)	1.354(7)
C(16)-C(17)	1.386(6)	N(18)-C(19)	1.351(4)
C(19)-O(20)	1.219(4)	C(19)-C(21)	1.490(5)
C(21)-C(22)	1.370(5)	C(21)-C(26)	1.374(5)
C(22)-C(23)	1.386(6)	C(23)-C(24)	1.361(7)
C(24)-C(25)	1.357(6)	C(25)-C(26)	1.383(6)
C(27)-O(28)	1.239(4)	C(27)-C(29)	1.487(4)
C(29)-C(30)	1.379(5)	C(29)-C(34)	1.391(5)
C(30)-C(31)	1.392(5)	C(31)-C(32)	1.373(8)
C(32)-C(33)	1.361(8)	C(33)-C(34)	1.375(5)

TABLE 3i Bond angles (deg.)

N(2)-N(1)-C(8)	113.8(3)	N(2)-N(1)-C(9)	114.0(2)
C(8)-N(1)-C(9)	103.4(2)	N(1)-N(2)-C(3)	121.3(2)
N(1)-N(2)-C(10)	115.0(3)	C(3)-N(2)-C(10)	118.3(2)
N(2)-C(3)-N(4)	108.5(2)	C(3)-N(4)-C(5)	114.6(3)
C(3)-N(4)-N(18)	113.7(2)	C(5)-N(4)-N(18)	112.6(2)
N(4)-C(5)-N(6)	113.1(2)	C(5)-N(6)-N(7)	109.7(3)
C(5)-N(6)-C(9)	113.2(2)	N(7)-N(6)-C(9)	99.4(2)
N(6)-N(7)-C(8)	110.6(2)	N(6)-N(7)-C(27)	126.7(2)
C(8)-N(7)-C(27)	121.0(2)	N(1)-C(8)-N(7)	104.9(2)
N(1)-C(9)-N(6)	107.4(3)	N(2)-C(10)-O(11)	120.7(3)
N(2)-C(10)-C(12)	118.6(3)	O(11)-C(10)-C(12)	120.7(3)
C(10)-C(12)-C(13)	120.0(4)	C(10)-C(12)-C(17)	120.9(3)
C(13)-C(12)-C(17)	119.0(4)	C(12)-C(13)-C(14)	120.7(5)
C(13)-C(14)-C(15)	120.2(5)	C(14)-C(15)-C(16)	119.0(4)
C(15)-C(16)-C(17)	120.9(5)	C(12)-C(17)-C(16)	120.1(4)
N(4)-N(18)-C(19)	121.9(3)	N(18)-C(19)-O(20)	122.0(3)
N(18)-C(19)-C(21)	115.6(3)	O(20)-C(19)-C(21)	122.4(3)
C(19)-C(21)-C(22)	117.7(3)	C(19)-C(21)-C(26)	123.5(3)
C(22)-C(21)-C(26)	118.6(4)	C(21)-C(22)-C(23)	120.3(4)
C(22)-C(23)-C(24)	120.6(4)	C(23)-C(24)-C(25)	119.2(4)
C(24)-C(25)-C(26)	120.8(4)	C(21)-C(26)-C(25)	120.4(3)
N(7)-C(27)-O(28)	117.1(3)	N(7)-C(27)-C(29)	122.0(3)
O(28)-C(27)-C(29)	120.9(3)	C(27)-C(29)-C(30)	125.0(3)
C(27)-C(29)-C(34)	116.2(3)	C(30)-C(29)-C(34)	118.7(3)
C(29)-C(30)-C(31)	120.1(4)	C(30)-C(31)-C(32)	120.5(5)
C(31)-C(32)-C(33)	119.4(4)	C(32)-C(33)-C(34)	121.0(4)
C(29)-C(34)-C(33)	120.3(4)		

TABLE 41 Anisotropic temperature factors ($\text{\AA}^2 \times 10^3$)

atom	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}
N(1)	64(2)	49(2)	39(1)	8(1)	-10(1)	-16(1)
N(2)	66(2)	43(1)	45(2)	2(1)	-2(1)	-17(1)
C(3)	67(2)	50(2)	43(2)	4(1)	-14(2)	-19(2)
N(4)	63(2)	44(1)	42(1)	4(1)	-15(1)	-22(1)
C(5)	73(2)	46(2)	43(2)	9(1)	-19(2)	-17(2)
N(6)	66(2)	54(2)	36(1)	12(1)	-16(1)	-18(1)
N(7)	66(2)	51(2)	32(1)	4(1)	-14(1)	-5(1)
C(8)	70(2)	62(2)	44(2)	13(2)	-18(2)	-15(2)
C(9)	66(2)	63(2)	40(2)	8(2)	-16(2)	-24(2)
C(10)	82(2)	55(2)	59(2)	-0(2)	-5(2)	-22(2)
O(11)	126(2)	69(2)	74(2)	-13(1)	18(2)	-38(2)
C(12)	86(3)	43(2)	54(2)	-1(2)	-8(2)	-19(2)
C(13)	165(4)	88(3)	92(3)	30(2)	-61(3)	-64(3)
C(14)	222(6)	92(3)	104(4)	44(3)	-68(4)	-65(4)
C(15)	182(5)	58(3)	80(3)	9(2)	-7(3)	-53(3)
C(16)	113(3)	81(3)	98(3)	0(3)	-12(3)	-56(3)
C(17)	95(3)	76(3)	75(3)	8(2)	-20(2)	-36(2)
N(18)	77(2)	60(2)	38(1)	15(1)	-23(1)	-37(1)
C(19)	68(2)	56(2)	41(2)	12(2)	-17(2)	-23(2)
O(20)	95(2)	95(2)	45(1)	26(1)	-34(1)	-47(1)
C(21)	58(2)	53(2)	39(2)	8(1)	-10(2)	-21(2)
C(22)	81(2)	71(2)	74(2)	28(2)	-29(2)	-31(2)
C(23)	89(3)	61(2)	105(3)	25(2)	-20(3)	-31(2)
C(24)	73(3)	88(3)	94(3)	11(2)	-13(2)	-45(2)
C(25)	73(3)	117(3)	99(3)	30(3)	-39(2)	-46(3)
C(26)	70(2)	79(3)	82(3)	31(2)	-30(2)	-32(2)
C(27)	61(2)	64(2)	42(2)	6(2)	-15(2)	-27(2)
O(28)	103(2)	90(2)	39(1)	5(1)	-26(1)	-20(2)
C(29)	63(2)	61(2)	50(2)	3(2)	1(2)	-21(2)
C(30)	71(2)	72(2)	67(2)	7(2)	-3(2)	-9(2)
C(31)	77(3)	89(3)	120(4)	26(3)	6(3)	2(3)
C(32)	119(4)	60(3)	141(5)	4(3)	22(4)	-1(3)
C(33)	126(4)	72(3)	125(4)	-29(3)	8(3)	-26(3)
C(34)	87(3)	75(3)	78(3)	-16(2)	-2(2)	-31(2)

The anisotropic temperature factor exponent takes the form:

$$-2\pi^2 (h^2 a^*{}^2 U_{11} + k^2 b^*{}^2 U_{22} + \dots + 2hka*b^* U_{12})$$

TABLE 5i Hydrogen coordinates ($\times 10^4$) and temperature factors ($\text{A}^2 \times 10^3$)

atom	x/a	y/b	z/c	U
H(13)	8840	4559	1612	114
H(14)	9708	6150	612	143
H(15)	11184	7184	1301	120
H(16)	11880	6579	2935	105
H(17)	10977	5021	3954	88
H(22)	7654	-2720	5022	78
H(23)	6057	-4034	5008	93
H(24)	4159	-3581	4061	91
H(25)	3831	-1801	3149	98
H(26)	5465	-506	3105	79
H(30)	14537	-1273	1043	82
H(31)	16158	-3231	631	116
H(32)	16114	-4461	-882	135
H(33)	14472	-3737	-1987	130
H(34)	12862	-1798	-1605	90
H(3a)	9438(30)	1414(21)	4335(22)	58(8)
H(3b)	7883(31)	1900(22)	3991(22)	66(8)
H(5a)	11154(26)	-417(19)	3482(19)	43(7)
H(5b)	10914(29)	-1065(21)	2397(22)	61(8)
H(8a)	10010(29)	1886(21)	1026(21)	55(8)
H(8b)	11657(33)	2148(24)	381(24)	77(9)
H(9a)	13409(30)	1445(23)	1956(23)	56(8)
H(9b)	12158(25)	1381(18)	3212(18)	37(6)
H(18)	7992(27)	-225(20)	2498(21)	52(7)

TABLE 1j Atom coordinates ($\times 10^4$) and temperature factors ($\text{\AA}^2 \times 10^3$)

atom	x/a	y/b	z/c	U_{eq}
N(1)	2240(2)	2848(1)	6836(1)	33(1)*
C(2)	406(2)	2943(2)	6248(1)	33(1)*
C(3)	350(2)	4180(2)	5468(1)	34(1)*
N(4)	2116(2)	5164(2)	5411(1)	37(1)*
C(5)	3981(3)	5024(2)	6030(1)	42(1)*
C(6)	4042(3)	3867(2)	6747(1)	39(1)*
N(7)	1999(3)	1598(2)	7508(1)	45(1)*
N(8)	81(3)	995(2)	7315(1)	52(1)*
N(9)	-971(3)	1806(2)	6336(1)	48(1)*
N(10)	-1098(3)	4629(2)	4763(1)	46(1)*
N(11)	-164(3)	5900(2)	4260(1)	52(1)*
N(12)	1757(3)	6242(2)	4635(1)	50(1)*

* Equivalent isotropic U defined as one third of the trace of the orthogonalised U_{ij} tensor

TABLE 2j Bond lengths (Å)

N(1)-C(2)	1.351(2)	N(1)-C(6)	1.391(2)
N(1)-N(7)	1.349(2)	C(2)-C(3)	1.426(2)
C(2)-N(9)	1.313(2)	C(3)-N(4)	1.352(2)
C(3)-N(10)	1.315(2)	N(4)-C(5)	1.392(2)
N(4)-N(12)	1.348(2)	C(5)-C(6)	1.323(2)
N(7)-N(8)	1.298(2)	N(8)-N(9)	1.359(2)
N(10)-N(11)	1.356(2)	N(11)-N(12)	1.299(2)

TABLE 3j Bond angles (deg.)

C(2)-N(1)-C(6)	125.2(1)	C(2)-N(1)-N(7)	107.8(1)
C(6)-N(1)-N(7)	126.9(1)	N(1)-C(2)-C(3)	116.6(1)
N(1)-C(2)-N(9)	109.6(1)	C(3)-C(2)-N(9)	133.8(1)
C(2)-C(3)-N(4)	116.7(1)	C(2)-C(3)-N(10)	133.9(1)
N(4)-C(3)-N(10)	109.4(1)	C(3)-N(4)-C(5)	125.0(1)
C(3)-N(4)-N(12)	107.8(1)	C(5)-N(4)-N(12)	127.2(1)
N(4)-C(5)-C(6)	118.3(1)	N(1)-C(6)-C(5)	113.1(1)
N(1)-N(7)-N(8)	105.3(1)	N(7)-N(8)-N(9)	112.1(1)
C(2)-N(9)-N(8)	104.6(1)	C(3)-N(10)-N(11)	104.8(1)
N(10)-N(11)-N(12)	112.1(1)	N(4)-N(12)-N(11)	105.8(1)

TABLE 4j Anisotropic temperature factors ($\text{Å}^2 \times 10^3$)

atom	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}
N(1)	35(1)	33(1)	32(1)	0(1)	-1(1)	1(1)
C(2)	32(1)	32(1)	34(1)	-4(1)	-0(1)	-0(1)
C(3)	37(1)	34(1)	32(1)	-6(1)	-0(1)	1(1)
N(4)	44(1)	32(1)	33(1)	0(1)	2(1)	-3(1)
C(5)	35(1)	43(1)	47(1)	-3(1)	1(1)	-7(1)
C(6)	32(1)	43(1)	43(1)	-3(1)	-2(1)	1(1)
N(7)	52(1)	41(1)	42(1)	7(1)	-0(1)	2(1)
N(8)	56(1)	46(1)	54(1)	14(1)	3(1)	-9(1)
N(9)	44(1)	45(1)	54(1)	5(1)	0(1)	-12(1)
N(10)	53(1)	46(1)	39(1)	-2(1)	-11(1)	7(1)
N(11)	71(1)	45(1)	40(1)	3(1)	-7(1)	7(1)
N(12)	69(1)	38(1)	41(1)	5(1)	0(1)	1(1)
H(5)	47	51	62	2	1	-17
H(6)	37	57	53	-1	-10	0

The anisotropic temperature factor exponent takes the form:

$$-2\pi^2(h^2 a^*{}^2 U_{11} + k^2 b^*{}^2 U_{22} + \dots + 2hka^*b^* U_{12})$$

TABLE 5j Hydrogen coordinates ($\times 10^4$) and temperature factors ($\text{Å}^2 \times 10^3$)

atom	x/a	y/b	z/c	U
H(5)	4984(31)	5828(27)	5964(14)	53*
H(6)	5194(29)	3690(21)	7215(14)	49*

Table 1k Atomic coordinates and temperature factors (\AA^2)*

Atom	x/a	y/b	z/c	U_{eq}
Br1	0.1041(2)	0.0615(2)	1.1230(1)	4.3(1)
Br2	0.3816(2)	0.3325(2)	1.1286(1)	4.7(1)
N1	0.3786(12)	0.1538(11)	0.9362(7)	2.7(3)
N2	0.2374(15)	0.3193(14)	0.8724(8)	4.2(4)
O1	0.1171(13)	0.2899(14)	0.8356(8)	5.9(4)
O2	0.2631(13)	0.1725(12)	0.8598(7)	5.4(4)
C1	0.3116(15)	-0.0003(15)	0.9822(8)	2.9(4)
C6	0.5093(15)	0.1914(14)	0.9875(8)	3.0(4)
Br1'	0.3341(2)	0.8776(2)	0.6236(1)	4.9(1)
Br2'	0.0599(2)	0.6119(2)	0.6114(1)	4.6(1)
N1'	0.1563(12)	0.8795(12)	0.4304(7)	3.0(3)
N2'	0.3232(15)	0.7538(14)	0.3587(8)	4.6(4)
O1'	0.4685(13)	0.7889(15)	0.3337(8)	5.9(4)
O2'	0.2944(15)	0.6252(14)	0.3252(8)	6.3(5)
C2'	0.1942(16)	1.0104(16)	0.4863(8)	3.2(4)
C6'	-0.0015(16)	0.8172(15)	0.4772(9)	3.2(4)

Hydrogen Coordinates

H2	0.251	-0.014	0.923	3.5
H6	0.552	0.278	0.929	3.6
H2'	0.282	1.056	0.429	4.0
H6'	-0.015	0.761	0.415	3.8

Coordinates are given for both half-molecules in the asymmetric unit. The remaining half of each molecule is generated by crystallographic symmetry ($1-x$, $-y$, $2-z$ for the unprimed half-molecule; $-x$, $2-y$, $1-z$ for the primed one).

*

Equivalent isotropic U defined as one third of the trace of the orthogonalised U_{ij} tensor.

Table 2k

Bond lengths (Å)

	Molecule 1	Molecule 2(')
Brl-C2	1.980(9)	1.984(10)
Br2-C6	1.984(10)	1.971(10)
N1-C2	1.437(16)	1.443(18)
N1-C6	1.410(17)	1.429(16)
N1-N2	1.441(1)	1.412(11)
N2-O1	1.201(17)	1.198(17)
N2-O2	1.233(16)	1.234(16)
C2-C6A	1.525(12)	1.505(13)

Table 3k

Bond angles (deg.)

	Molecule 1	Molecule 2(')
C2-N1-C6	123.6(8)	122.4(8)
C2-N1-N2	113.1(9)	115.3(9)
C6-N1-N2	117.3(9)	117.4(10)
N1-N2-O1	117.2(10)	117.7(11)
N1-N2-O2	113.9(11)	113.9(10)
O1-N2-O2	128.8(9)	128.3(9)
Brl-C2-N1	113.9(7)	113.6(7)
Brl-C2-C6A	108.9(6)	109.0(7)
N1-C2-C6A	109.2(9)	110.6(10)
Br2-C6-N1	114.6(6)	113.3(7)
Br2-C6-C2A	108.7(7)	108.4(6)
N1-C6-C2A	111.2(10)	111.2(10)

Table 4K Torsion angles (deg.)

	Molecule 1	Molecule 2
N1-C2-C6A-N1A	40.3	40.5
C2-C6A-N1A-C2A	-17.1	-46.2
C6A-N1A-C2A-C6	46.3	45.9
Br1-N1A-C2A-Br2A	148.4	149.3
C2-N1-N2-O1	-21.3	-12.9
C2-N1-N2-O2	161.2	170.4
C6-N1-N2-O1	-175.0	-168.9
C6-N1-N2-O2	7.5	14.1

TABLE 11 Atom coordinates ($\times 10^4$) and temperature factors ($\text{\AA}^2 \times 10^3$)

atom	x/a	y/b	z/c	U_{eq}
N(1)	-328(3)	629(4)	2225(1)	48(1)*
C(2)	-1052(4)	874(5)	1789(2)	47(1)*
N(3)	-526(3)	1064(4)	1274(1)	47(1)*
C(4)	-390(4)	2381(4)	1019(2)	46(1)*
N(5)	-1267(3)	2767(4)	662(1)	44(1)*
C(6)	-1350(4)	2312(3)	114(2)	45(1)*
N(7)	-1120(3)	3356(4)	-283(1)	59(1)*
C(10)	186(4)	1667(5)	2468(2)	46(1)*
O(11)	65(3)	2865(3)	2321(1)	57(1)*
C(12)	877(5)	1271(6)	2923(2)	72(2)*
N(13)	-80(3)	-76(4)	1043(2)	55(1)*
O(14)	356(3)	67(4)	608(1)	71(1)*
O(15)	-161(3)	-1172(4)	1286(2)	71(1)*
N(16)	-2058(3)	3585(4)	855(2)	51(1)*
O(17)	-2714(3)	4034(4)	538(1)	69(1)*
O(18)	-2063(3)	3821(4)	1337(1)	70(1)*
N(19)	-57(4)	3642(4)	-382(2)	70(2)*
O(20)	609(3)	3127(4)	-84(2)	82(1)*
O(21)	144(4)	4396(4)	-756(2)	112(2)*
C(8)	-1922(5)	3880(6)	-627(2)	104(2)*
O(9)	-1702(5)	3723(6)	-1177(2)	84(2)*
C(22)	-2246(6)	2481(7)	-1280(3)	53(2)*
O(23)	-2774(6)	1916(7)	-929(2)	78(2)*
C(24)	-2295(8)	2064(10)	-1857(3)	89(3)*
O(9a)	-2113(9)	2692(11)	-1005(3)	103(3)*
C(22a)	-2590(8)	2593(3)	-1518(3)	106(3)*
O(23a)	-2521(16)	3792(4)	-1674(4)	265(3)*
C(24a)	-2199(12)	1383(4)	-1840(5)	75(3)*

* Equivalent isotropic U defined as one third of the trace of the orthogonalised U_{ij} tensor

TABLE 21 Bond lengths (Å)

N(1)-C(2)	1.434(6)	N(1)-C(10)	1.338(6)
C(2)-N(3)	1.460(6)	N(3)-C(4)	1.438(6)
N(3)-N(13)	1.364(6)	C(4)-N(5)	1.458(6)
N(5)-C(6)	1.447(5)	N(5)-N(16)	1.351(5)
C(6)-N(7)	1.448(6)	N(7)-N(19)	1.370(7)
N(7)-C(8)	1.413(7)	C(10)-O(11)	1.228(5)
C(10)-C(12)	1.479(7)	N(13)-O(14)	1.225(6)
N(13)-O(15)	1.228(6)	N(16)-O(17)	1.217(5)
N(15)-O(18)	1.231(5)	N(19)-O(20)	1.221(6)
N(19)-O(21)	1.216(6)	C(8)-O(9)	1.413(7)
C(8)-O(9a)	1.509(11)	O(9)-C(22)	1.404(9)
C(22)-O(23)	1.226(10)	C(22)-C(24)	1.504(10)
O(9a)-C(22a)	1.421(12)	C(22a)-O(23a)	1.229(6)
C(22a)-O(24a)	1.504(10)		

TABLE 3 | Bond angles (deg.)

C(2)-N(1)-C(10)	121.5(4)	N(1)-C(2)-N(3)	114.5(4)
C(2)-N(3)-C(4)	123.9(4)	C(2)-N(3)-N(13)	117.0(4)
C(4)-N(3)-H(13)	118.9(4)	N(3)-C(4)-N(5)	114.4(4)
C(4)-N(5)-C(6)	123.9(3)	C(4)-N(5)-H(16)	118.3(3)
C(6)-N(5)-H(16)	117.8(3)	N(5)-C(6)-N(7)	115.2(4)
C(6)-N(7)-H(19)	117.2(4)	C(6)-N(7)-C(8)	122.2(4)
H(19)-N(7)-C(8)	119.6(4)	N(1)-C(10)-O(11)	121.1(4)
N(1)-C(10)-C(12)	115.7(4)	O(11)-C(10)-C(12)	123.3(4)
N(3)-N(13)-O(14)	117.7(4)	N(3)-N(13)-O(15)	117.2(4)
O(14)-N(13)-O(15)	125.1(4)	N(5)-N(16)-O(17)	117.5(4)
N(5)-N(16)-O(18)	117.7(4)	O(17)-N(16)-O(18)	124.8(4)
N(7)-N(19)-O(20)	117.3(4)	N(7)-N(19)-O(21)	117.4(5)
O(20)-N(19)-O(21)	125.3(5)	N(7)-C(8)-O(9)	115.0(5)
H(7)-C(8)-O(3a)	102.7(6)	O(9)-C(22)-O(23)	120.6(7)
C(3)-C(9)-C(22)	100.3(5)	O(23)-C(22)-C(24)	123.2(7)
O(9)-C(22)-C(24)	115.3(7)	C(3)-O(9a)-C(22a)	133.2(3)
O(9a)-C(22a)-O(23a)	101.3(8)	O(9a)-C(22a)-C(24a)	113.9(3)
O(23a)-C(22a)-C(24a)	123.0(9)		

TABLE 4 | Anisotropic temperature factors ($\text{\AA}^2 \times 10^3$)

atom	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}
N(1)	66(2)	34(2)	44(2)	1(2)	-9(2)	-2(2)
C(2)	47(3)	44(3)	51(2)	-10(2)	5(2)	-4(2)
N(3)	65(2)	40(2)	37(2)	-11(2)	-1(2)	16(2)
C(4)	48(2)	44(3)	44(2)	-10(2)	-1(2)	9(2)
N(5)	51(2)	44(2)	36(2)	-6(2)	-3(2)	16(2)
C(6)	54(3)	42(2)	40(2)	-4(2)	-5(2)	3(2)
N(7)	80(3)	53(2)	43(2)	6(2)	-3(2)	13(2)
C(10)	62(3)	42(2)	36(2)	-5(2)	1(2)	6(2)
O(11)	81(2)	37(2)	54(2)	-1(2)	-13(2)	-5(2)
C(12)	94(4)	61(3)	62(3)	-2(3)	-20(3)	7(3)
N(13)	65(2)	30(2)	50(2)	-11(2)	-10(2)	17(2)
O(14)	98(3)	66(2)	50(2)	-12(2)	11(2)	29(2)
O(15)	93(3)	43(2)	71(2)	-2(2)	-15(2)	11(2)
N(16)	50(2)	53(2)	48(2)	-6(2)	3(2)	16(2)
O(17)	50(2)	32(2)	64(2)	-6(2)	-10(2)	34(2)
O(18)	73(2)	37(2)	50(2)	-18(2)	5(2)	28(2)
N(19)	109(3)	50(3)	51(2)	-11(2)	22(3)	-1(3)
O(20)	74(2)	100(3)	71(2)	-12(2)	7(2)	-16(2)
O(21)	173(4)	63(2)	91(3)	15(2)	57(3)	-6(3)
C(8)	133(4)	115(4)	44(3)	7(3)	1(2)	73(4)
O(9)	105(4)	72(4)	74(4)	41(3)	-3(4)	-13(3)
C(22)	23(2)	31(3)	104(5)	-15(1)	-24(4)	-14(3)
O(23)	39(4)	72(4)	72(4)	-17(3)	-11(4)	-9(3)
C(24)	73(3)	98(5)	98(5)	37(5)	21(5)	74(5)
O(9a)	107(5)	160(6)	41(4)	-46(4)	-34(4)	49(5)
C(22a)	40(5)	49(5)	223(6)	-11(6)	-28(6)	15(5)
O(23a)	643(6)	63(5)	84(5)	4(4)	-147(6)	27(6)
C(24a)	83(5)	37(6)	54(5)	4(5)	-17(5)	26(5)

The anisotropic temperature factor exponent takes the form:

$$-2^{-2}(h^2a^2U_{11} + k^2b^2U_{22} + \dots + 2hka^2b^2U_{12})$$

TABLE 5 | Hydrogen coordinates ($\times 10^4$) and temperature factors ($\text{\AA}^2 \times 10^3$)

atom	x/a	y/b	z/c	U_{eq}
H(1)	-208	-299	2344	45
H(2a)	-1530	98	1762	45
H(2b)	-1461	1690	1867	45
H(4a)	-334	3072	1292	39
H(4b)	265	2356	816	39
H(6a)	-853	1565	64	48
H(6b)	-2072	1989	57	48
H(12a)	1240	421	2843	70
H(12b)	440	1152	3235	70
H(12c)	1401	1981	2987	70
H(8a)	-2611	3489	-541	112
H(8b)	-1936	4855	-562	112
H(8a')	-2536	4182	-426	112
H(8b')	-1635	4641	-827	112
H(24a)	-2681	1212	-1898	91
H(24b)	-1567	1947	-1979	91
H(24c)	-2641	2771	-2063	91
H(24x)	-1967	625	-1622	90
H(24y)	-1681	1541	-2118	90
H(24z)	-2886	1171	-1996	90

TABLE 1M Atom coordinates ($\times 10^4$) and temperature factors ($\text{Å}^2 \times 10^3$)

atom	x/a	y/b	z/c	U_{eq}
N(1)	2367(3)	1369(1)	2528(2)	47(1)*
C(2)	3136(4)	1823(1)	3493(2)	46(1)*
N(3)	3250(3)	2513(1)	3113(2)	40(1)*
C(4)	4877(3)	2793(1)	2614(2)	38(1)*
H(5)	5857(2)	3312(1)	3369(2)	35(1)*
C(6)	5753(3)	4011(1)	3020(2)	40(1)*
H(7)	7499(3)	4282(1)	2677(2)	44(1)*
C(8)	8577(4)	4790(1)	3417(3)	49(1)*
N(9)	8070(3)	5462(1)	3053(2)	49(1)*
C(10)	3090(3)	1118(1)	1665(2)	45(1)*
O(11)	4990(3)	1321(1)	1586(2)	61(1)*
C(12)	2499(4)	587(2)	822(3)	65(1)*
N(13)	1660(3)	2893(1)	3006(2)	46(1)*
O(14)	213(2)	2597(1)	3165(2)	62(1)*
O(15)	1801(3)	3485(1)	2774(2)	57(1)*
N(16)	6806(3)	3149(1)	4433(2)	39(1)*
O(17)	6733(3)	2556(1)	4831(2)	51(1)*
O(18)	7665(2)	3599(1)	5062(2)	51(1)*
N(19)	7955(3)	4141(1)	1500(2)	47(1)*
O(20)	7046(3)	3710(1)	911(2)	65(1)*
O(21)	9234(3)	4467(1)	1129(2)	61(1)*
C(22)	6701(4)	5795(1)	3540(3)	49(1)*
O(23)	5761(3)	5529(1)	4291(2)	66(1)*
C(24)	5387(5)	6502(2)	3095(3)	64(1)*
N(1')	7453(3)	1470(1)	-289(2)	51(1)*
C(2')	8334(4)	2090(2)	1(2)	49(1)*
N(3')	8202(3)	2572(1)	-1034(2)	44(1)*
C(4')	9675(3)	2727(1)	-1802(2)	40(1)*
N(5')	10597(3)	3359(1)	-1514(2)	42(1)*
C(6')	10515(3)	3922(1)	-2365(2)	40(1)*
N(7')	12199(3)	4029(1)	-2985(2)	44(1)*
C(8')	13446(3)	4592(1)	-2757(3)	49(1)*
N(9')	13180(3)	5117(1)	-3687(3)	50(1)*
C(10')	8070(4)	1027(1)	-1090(3)	48(1)*
O(11')	9411(3)	1170(1)	-1669(2)	59(1)*
C(12')	7072(4)	378(2)	-1253(4)	65(1)*
N(13')	6530(3)	2871(1)	-1346(2)	48(1)*
O(14')	5241(2)	2674(1)	-782(2)	66(1)*
O(15')	6439(3)	3299(1)	-2162(2)	63(1)*
N(16')	11732(3)	3404(1)	-429(2)	44(1)*
O(17')	11814(3)	2915(1)	272(2)	57(1)*
O(18')	12641(3)	3920(1)	-223(2)	56(1)*
N(19')	12561(3)	3591(1)	-3902(2)	46(1)*
O(20')	11627(3)	3085(1)	-4030(2)	58(1)*
O(21')	13819(3)	3750(1)	-4527(2)	63(1)*
C(22')	11825(4)	5566(1)	-3632(3)	55(1)*
O(23')	10748(3)	5536(1)	-2826(3)	31(1)*
C(24')	11690(4)	6101(2)	-4640(4)	65(1)*

* Equivalent isotropic U defined as one third of the trace of the orthogonalised U_{ij} tensor

TABLE 2M Bond lengths (Å)

N(1)-C(2)	1.441(3)	N(1)-C(10)	1.341(4)
C(2)-N(3)	1.451(3)	N(3)-C(4)	1.454(3)
N(3)-N(13)	1.369(3)	C(4)-N(5)	1.452(3)
N(5)-C(6)	1.445(3)	N(5)-N(16)	1.354(2)
C(6)-N(7)	1.458(3)	N(7)-C(8)	1.461(3)
N(7)-N(19)	1.364(3)	C(8)-N(9)	1.436(4)
N(9)-C(22)	1.345(4)	C(10)-O(11)	1.239(3)
C(10)-C(12)	1.494(4)	N(13)-O(14)	1.230(3)
N(13)-O(15)	1.216(3)	N(16)-O(17)	1.225(3)
N(16)-O(18)	1.220(3)	N(19)-O(20)	1.218(3)
N(19)-O(21)	1.232(3)	C(22)-O(23)	1.229(4)
C(22)-C(24)	1.500(4)	N(1')-C(2')	1.429(4)
N(1')-C(10')	1.341(4)	C(2')-N(3')	1.463(4)
N(3')-C(4')	1.448(3)	N(3')-N(13')	1.358(3)
C(4')-N(5')	1.447(3)	N(5')-C(6')	1.444(3)
N(5')-N(16')	1.358(3)	C(6')-N(7')	1.465(3)
N(7')-C(8')	1.446(3)	N(7')-N(19')	1.361(3)
C(8')-N(9')	1.444(4)	N(9')-C(22')	1.334(4)
C(10')-O(11')	1.240(4)	C(10')-C(12')	1.487(4)
N(13')-O(14')	1.229(3)	N(13')-O(15')	1.220(3)
N(16')-O(17')	1.230(3)	N(16')-O(18')	1.222(3)
N(19')-O(20')	1.216(3)	N(19')-O(21')	1.229(3)
C(22')-O(23')	1.241(4)	C(22')-C(24')	1.499(5)

TABLE 3M Bond angles (deg.)

C(2)-N(1)-C(10)	121.9(2)	N(1)-C(2)-N(3)	115.5(2)
C(2)-N(3)-C(4)	122.6(2)	C(2)-N(3)-N(13)	118.3(2)
C(4)-N(3)-N(13)	118.1(2)	N(3)-C(4)-N(5)	115.7(2)
C(4)-N(5)-C(6)	122.5(2)	C(4)-N(5)-N(16)	119.4(2)
C(6)-N(5)-N(16)	117.9(2)	N(5)-C(6)-N(7)	113.7(2)
C(6)-N(7)-C(8)	123.3(2)	C(6)-N(7)-N(19)	117.3(2)
C(8)-N(7)-N(19)	118.3(2)	N(7)-C(8)-N(9)	113.4(2)
C(8)-N(9)-C(22)	122.4(2)	N(1)-C(10)-O(11)	121.0(2)
N(1)-C(10)-C(12)	116.5(2)	O(11)-C(10)-C(12)	122.5(3)
N(3)-N(13)-O(14)	116.6(2)	N(3)-N(13)-O(15)	117.5(2)
O(14)-N(13)-O(15)	125.9(2)	N(5)-N(16)-O(17)	117.4(2)
N(5)-N(16)-O(18)	116.8(2)	O(17)-N(16)-O(18)	125.9(2)
N(7)-N(19)-O(20)	116.8(2)	N(7)-N(19)-O(21)	117.0(2)
O(20)-N(19)-O(21)	126.2(2)	N(9)-C(22)-O(23)	121.4(3)
N(9)-C(22)-C(24)	115.9(3)	O(23)-C(22)-C(24)	122.7(3)
C(2')-N(1')-C(10')	121.9(2)	N(1')-C(2')-N(3')	113.9(2)
C(2')-N(3')-C(4')	123.9(2)	C(2')-N(3')-N(13')	118.7(2)
C(4')-N(3')-N(13')	117.3(2)	N(3')-C(4')-N(5')	114.6(2)
C(4')-N(5')-C(6')	124.0(2)	C(4')-N(5')-N(16')	117.9(2)
C(6')-N(5')-N(16')	117.8(2)	N(5')-C(6')-N(7')	114.7(2)
C(6')-N(7')-C(8')	124.9(2)	C(6')-N(7')-N(19')	117.8(2)
C(8')-N(7')-N(19')	117.2(2)	N(7')-C(8')-N(9')	114.5(2)
C(8')-N(9')-C(22')	121.0(3)	N(1')-C(10')-O(11')	119.9(3)
N(1')-C(10')-C(12')	117.4(3)	O(11')-C(10')-C(12')	122.7(3)
N(3')-N(13')-O(14')	116.3(2)	N(3')-N(13')-O(15')	117.5(2)
O(14')-N(13')-O(15')	126.2(2)	N(5')-N(16')-O(17')	116.6(2)
N(5')-N(16')-O(18')	118.1(2)	O(17')-N(16')-O(18')	125.3(2)
N(7')-N(19')-O(20')	117.9(2)	N(7')-N(19')-O(21')	115.9(2)
O(20')-N(19')-O(21')	126.2(2)	N(9')-C(22')-C(23')	120.4(3)
N(9')-C(22')-C(24')	117.0(3)	O(23')-C(22')-C(24')	122.6(3)

TABLE 4M Anisotropic temperature factors ($\text{\AA}^2 \times 10^3$)

atom	U_{11}	U_{22}	U_{33}	U_{12}	U_{13}	U_{23}
N(1)	39(1)	47(1)	57(1)	-7(1)	16(1)	-14(1)
C(2)	48(1)	48(1)	42(1)	3(1)	10(1)	-8(1)
N(3)	32(1)	41(1)	49(1)	-1(1)	9(1)	-5(1)
C(4)	36(1)	40(1)	39(1)	-3(1)	10(1)	-6(1)
N(5)	35(1)	34(1)	36(1)	1(1)	3(1)	-3(1)
C(6)	38(1)	37(1)	45(1)	1(1)	11(1)	-3(1)
N(7)	50(1)	40(1)	44(1)	-2(1)	15(1)	-11(1)
C(8)	47(1)	45(1)	55(1)	-2(1)	9(1)	-9(1)
N(9)	56(1)	39(1)	57(1)	-0(1)	22(1)	-12(1)
C(10)	47(1)	42(1)	48(1)	1(1)	10(1)	-5(1)
O(11)	50(1)	66(1)	70(1)	-10(1)	27(1)	-14(1)
C(12)	59(2)	63(2)	73(2)	-21(2)	11(1)	-5(1)
N(13)	37(1)	55(1)	47(1)	-9(1)	5(1)	-2(1)
O(14)	36(1)	78(1)	74(1)	-9(1)	15(1)	-10(1)
O(15)	52(1)	55(1)	64(1)	-2(1)	7(1)	10(1)
N(16)	34(1)	42(1)	41(1)	-1(1)	8(1)	2(1)
O(17)	56(1)	44(1)	52(1)	11(1)	4(1)	6(1)
C(18)	49(1)	56(1)	46(1)	-4(1)	-1(1)	-10(1)
N(19)	51(1)	45(1)	49(1)	3(1)	17(1)	-2(1)
O(20)	74(1)	72(1)	51(1)	-15(1)	19(1)	-18(1)
O(21)	58(1)	64(1)	67(1)	12(1)	29(1)	-7(1)
C(22)	59(2)	42(1)	48(1)	-3(1)	12(1)	-8(1)
O(23)	75(1)	55(1)	74(1)	5(1)	38(1)	-4(1)
C(24)	36(2)	47(2)	63(2)	-0(1)	16(2)	-1(2)
N(1')	52(1)	47(1)	60(1)	4(1)	29(1)	-2(1)
C(2')	56(1)	51(2)	43(1)	-1(1)	15(1)	2(1)
N(3')	37(1)	45(1)	51(1)	2(1)	14(1)	4(1)
C(4')	39(1)	44(1)	38(1)	-4(1)	11(1)	-1(1)
N(5')	41(1)	51(1)	33(1)	0(1)	4(1)	-3(1)
C(6')	36(1)	42(1)	44(1)	-3(1)	11(1)	4(1)
N(7')	46(1)	41(1)	48(1)	-3(1)	19(1)	2(1)
C(8')	42(1)	49(1)	56(1)	0(1)	11(1)	2(1)
N(9')	46(1)	41(1)	69(1)	1(1)	24(1)	3(1)
C(10')	43(1)	43(1)	60(2)	8(1)	14(1)	8(1)
O(11')	59(1)	47(1)	77(1)	-2(1)	36(1)	5(1)
C(12')	54(2)	44(2)	99(2)	2(2)	18(2)	3(1)
N(13')	41(1)	48(1)	57(1)	-9(1)	10(1)	3(1)
C(14')	44(1)	58(1)	90(1)	-15(1)	27(1)	-3(1)
O(15')	61(1)	76(2)	67(1)	10(1)	5(1)	22(1)
N(16')	35(1)	62(1)	37(1)	-8(1)	8(1)	-2(1)
O(17')	49(1)	80(1)	40(1)	14(1)	3(1)	-3(1)
C(18')	49(1)	67(1)	51(1)	-15(1)	8(1)	-11(1)
N(19')	50(1)	47(1)	42(1)	1(1)	13(1)	10(1)
O(20')	57(1)	52(1)	56(1)	-11(1)	15(1)	1(1)
O(21')	66(1)	58(1)	60(1)	2(1)	34(1)	7(1)
C(22')	41(1)	39(1)	85(2)	1(1)	16(1)	-1(1)
O(23')	63(1)	68(1)	122(2)	21(1)	52(1)	23(1)
C(24')	54(2)	46(2)	97(2)	10(2)	13(2)	-1(1)

The anisotropic temperature factor exponent takes the form:

$$-2\pi^2(h^2a^2U_{11}+k^2b^2U_{22}+\dots+2hka^2b^2U_{12})$$

TABLE 5M Hydrogen coordinates ($\times 10^4$) and temperature factors ($\text{Å}^2 \times 10^3$)

atom	x	y	z	u
H(1)	1381(33)	1185(17)	2566(30)	56
H(2a)	2377	1803	4175	54
H(2b)	4369	1674	3781	54
H(4a)	4486	2979	1800	45
H(4b)	5732	2433	2528	45
H(6a)	4825	4061	2315	47
H(6b)	5396	4262	3720	47
H(8a)	9866	4726	3316	58
H(8b)	8389	4732	4285	58
H(9)	8678(38)	5703(16)	2664(26)	54
H(12a)	1284	484	1049	77
H(12b)	3246	189	873	77
H(12c)	2395	757	-22	77
H(24a)	6289	6510	2192	76
H(24b)	7456	6755	3427	76
H(24c)	5291	6696	3376	76
H(1')	6635(33)	1357(17)	75(28)	61
H(2'a)	7877	2287	708	59
H(2'b)	9681	1999	219	59
H(4'a)	9153	2737	-2666	48
H(4'b)	10589	2378	-1687	48
H(6'a)	9498	3852	-3004	48
H(6'b)	10293	4318	-1896	48
H(8'a)	14697	4430	-2741	58
H(8'b)	13267	4779	-1952	58
H(9')	13903(36)	5150(18)	-4187(25)	60
H(12d)	5855	394	-977	78
H(12e)	7798	41	-787	78
H(12f)	6970	271	-2133	78
H(24d)	11745	6534	-4249	78
H(24e)	10539	6056	-5170	78
H(24f)	12708	6053	-5139	78

TABLE 1n Atom coordinates ($\times 10^4$) and temperature factors ($\text{Å}^2 \times 10^3$)

atom	x/a	y/b	z/c	U_{eq}
C(1)	164(10)	2834(11)	851(4)	60(3)*
O(2)	1709(8)	3174(8)	142(3)	77(2)*
C(3)	3659(11)	2927(11)	878(4)	61(3)*
C(4)	2085(9)	2447(9)	1688(4)	44(2)*
C(5)	1568(9)	49(9)	1973(4)	46(2)*
N(6)	510(7)	-422(7)	2862(3)	45(2)*
C(?)	1964(11)	-64(9)	3850(4)	57(3)*
C(8)	2906(11)	2350(10)	4210(4)	58(3)*
N(9)	4132(7)	3663(7)	3484(3)	46(2)*
C(10)	2788(9)	4127(8)	2570(4)	46(2)*
N(6a)	-1878(8)	-740(7)	2763(3)	53(2)*
O(6a)	-3023(7)	-790(7)	1931(3)	66(2)*
O(6b)	-2689(8)	-1014(7)	3542(3)	73(2)*
N(9a)	6474(8)	3855(7)	3526(4)	52(2)*
O(9a)	7321(7)	4492(7)	2790(3)	64(2)*
O(9b)	7564(8)	3444(7)	4303(3)	74(2)*

* Equivalent isotropic U defined as one third of the trace of the orthogonalised U_{ij} tensor

TABLE 2n Bond lengths (Å)

C(1)-O(2)	1.433(8)	C(1)-C(4)	1.561(8)
O(2)-C(3)	1.457(8)	C(3)-C(4)	1.556(8)
C(4)-C(5)	1.542(8)	C(4)-C(10)	1.509(7)
C(5)-N(6)	1.463(7)	N(6)-C(7)	1.445(6)
N(6)-N(6a)	1.369(7)	C(7)-C(8)	1.511(8)
C(8)-N(9)	1.461(7)	N(9)-C(10)	1.454(7)
N(9)-N(9a)	1.367(7)	N(6a)-O(6a)	1.215(6)
N(6a)-O(6b)	1.242(7)	N(9a)-O(9a)	1.227(7)
N(9a)-O(9b)	1.226(7)		

TABLE 3n Bond angles (deg.)

O(2)-C(1)-C(4)	92.3(5)	C(1)-O(2)-C(3)	92.2(4)
O(2)-C(3)-C(4)	91.6(5)	C(1)-O(4)-C(3)	85.8(4)
C(1)-C(4)-C(5)	115.2(4)	C(3)-C(4)-C(5)	111.1(5)
C(1)-C(4)-C(10)	112.2(5)	C(3)-C(4)-C(10)	117.2(4)
C(5)-C(4)-C(10)	114.0(4)	C(4)-C(5)-N(6)	114.7(5)
C(5)-N(6)-C(7)	121.7(4)	C(5)-N(6)-N(6a)	117.8(4)
C(7)-N(6)-N(6a)	119.0(5)	N(6)-C(7)-C(8)	111.3(5)
C(7)-C(8)-N(9)	110.6(5)	C(8)-N(9)-C(10)	120.7(4)
C(8)-N(9)-N(9a)	117.5(5)	C(10)-N(9)-N(9a)	116.8(4)
C(4)-C(10)-N(9)	117.3(5)	N(6)-N(6a)-C(6a)	116.1(5)
N(6)-N(6a)-O(6b)	116.0(4)	O(6a)-N(6a)-O(6b)	125.8(5)
N(9)-N(9a)-C(9a)	116.8(5)	N(9)-N(9a)-O(9b)	117.5(5)
C(9a)-N(9a)-O(9b)	125.6(5)		

TABLE 4n Anisotropic temperature factors ($\text{\AA}^2 \times 10^3$)

atom	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}
C(1)	46(4)	77(4)	61(4)	27(3)	6(3)	25(4)
O(2)	81(3)	111(4)	54(2)	36(2)	22(2)	46(3)
C(3)	48(4)	81(4)	61(4)	27(3)	24(3)	21(4)
C(4)	31(3)	55(3)	48(3)	22(3)	8(3)	14(3)
C(5)	37(3)	54(3)	51(3)	9(3)	14(3)	17(3)
N(6)	32(2)	47(3)	51(3)	10(2)	9(2)	7(2)
C(7)	56(4)	59(4)	60(4)	22(3)	20(3)	19(3)
C(8)	50(4)	75(4)	49(3)	11(3)	9(3)	19(3)
N(9)	27(2)	50(3)	59(3)	7(2)	8(2)	9(2)
C(10)	29(3)	43(3)	68(3)	16(3)	10(3)	12(3)
N(6a)	44(3)	42(3)	71(3)	5(2)	18(3)	9(2)
O(6a)	36(2)	78(3)	83(3)	15(2)	10(2)	18(2)
O(6b)	66(3)	67(3)	92(3)	9(2)	51(3)	12(2)
N(9a)	36(3)	45(3)	73(3)	-4(3)	2(3)	15(2)
O(9a)	40(2)	69(3)	91(3)	14(2)	25(2)	22(2)
O(9b)	54(3)	81(3)	84(3)	2(2)	-14(2)	32(3)

The anisotropic temperature factor exponent takes the form:

$$-2\pi^2(h^2a^2U_{11} + k^2b^2U_{22} + \dots + 2hka^2b^2U_{12})$$

TABLE 5n Hydrogen coordinates ($\times 10^4$) and temperature factors ($\text{\AA}^2 \times 10^3$)

atom	x/a	y/b	z/c	U_{eq}
H(1a)	-279	4099	998	65
H(1b)	-1245	1569	671	65
H(3a)	4168	1717	707	66
H(3b)	5017	4243	1034	66
H(5a)	3042	-267	2085	49
H(5b)	483	-902	1423	49
H(7a)	1010	-867	4299	61
H(7b)	3282	-593	3839	61
H(8a)	4009	2550	4831	62
H(8b)	1603	2836	4306	62
H(10a)	1351	4285	2721	49
H(10b)	3745	5498	2372	49

TABLE 1P Atom coordinates ($\times 10^4$) and temperature factors ($\text{\AA}^2 \times 10^3$)

atom	x/a	y/b	z/c	U_{eq}
N(1)	-2263(3)	1336(2)	278(2)	58(1)*
N(2)	-2609(3)	2348(3)	-639(2)	66(1)*
N(3)	-851(4)	3165(3)	-682(3)	64(1)*
N(4)	690(2)	2549(2)	242(1)	47(1)*
C(5)	2961(5)	2972(3)	729(2)	55(1)*
C(6)	3541(5)	1810(4)	1747(2)	53(1)*
N(7)	1359(3)	1089(2)	1729(2)	53(1)*
C(8)	-211(3)	1579(2)	794(2)	42(1)*
N(9)	1066(3)	-64(2)	2464(2)	57(1)*
O(10)	-769(3)	-645(2)	2258(2)	74(1)*
O(11)	2622(3)	-374(2)	3224(2)	79(1)*
N(1x)	-1045(11)	-62(9)	1682(7)	57(4)
N(2x)	231(11)	-745(9)	2648(7)	70(4)
N(3x)	2252(11)	-203(9)	2903(7)	115(12)
N(4x)	2267(11)	878(9)	2066(7)	50(3)
C(5x)	3843(11)	1939(9)	1794(7)	83(16)
C(6x)	2505(11)	2834(9)	737(7)	122(17)
N(7x)	358(11)	1998(9)	515(7)	32(4)
C(8x)	296(11)	918(9)	1356(7)	39(3)
N(9x)	-1403(11)	2419(9)	-332(7)	52(4)
O(10x)	-3111(11)	1723(9)	-323(7)	96(7)
O(11x)	-1085(11)	3406(9)	-985(7)	105(10)

* Equivalent isotropic U defined as one third of the trace of the orthogonalised U_{ij} tensor

Table 2P Bond Length \AA

N(1)-N(2)	1.366(3)	C(5)-C(6)	1.541(4)
N(1)-C(8)	1.305(2)	C(6)-N(7)	1.490(4)
N(2)-N(3)	1.313(3)	N(7)-C(8)	1.365(2)
N(3)-N(4)	1.356(3)	N(7)-N(9)	1.357(3)
N(1)-C(5)	1.431(3)	N(9)-O(10)	1.223(3)
N(4)-C(8)	1.322(3)	N(9)-O(11)	1.194(3)

Table 3P Bond Angles (deg.)

N(2)-N(1)-C(5)	103.3(2)	C(6)-N(7)-N(9)	123.2(2)
N(1)-N(2)-N(3)	112.7(2)	C(8)-N(7)-N(9)	124.4(2)
N(2)-N(3)-N(4)	104.3(2)	N(1)-C(8)-N(5)	111.3(2)
N(3)-N(4)-C(5)	108.6(2)	N(1)-C(8)-N(7)	141.0(2)
N(3)-N(4)-C(6)	135.9(2)	N(4)-C(8)-N(7)	107.6(2)
C(5)-N(4)-C(6)	115.1(2)	N(7)-N(9)-C(10)	115.3(2)
N(1)-C(5)-C(6)	103.0(2)	N(7)-N(9)-O(11)	116.0(2)
C(5)-C(6)-N(7)	101.7(2)	O(10)-N(9)-C(11)	128.7(2)
C(6)-N(7)-C(8)	111.9(2)		

*Bond distances and angles for the minor disorder form do not differ significantly from the major form.

TABLE 4p Anisotropic temperature factors ($\text{Å}^2 \times 10^3$)

atom	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}
N(1)	40(1)	65(1)	66(1)	-9(1)	2(1)	-1(1)
N(2)	46(1)	86(1)	56(1)	-11(1)	-10(1)	14(1)
N(3)	56(1)	80(2)	50(1)	8(1)	-0(1)	16(1)
N(4)	42(1)	51(1)	45(1)	3(1)	7(1)	2(1)
C(5)	36(1)	68(2)	61(2)	10(1)	11(1)	1(1)
C(6)	38(1)	62(2)	54(2)	2(1)	1(1)	0(1)
N(7)	44(1)	56(1)	57(1)	14(1)	4(1)	-4(1)
C(8)	40(1)	42(1)	43(1)	-2(1)	8(1)	1(1)
N(9)	70(1)	50(1)	53(1)	2(1)	18(1)	4(1)
O(10)	83(1)	67(1)	75(1)	5(1)	26(1)	-22(1)
O(11)	86(1)	83(1)	65(1)	24(1)	7(1)	18(1)

The anisotropic temperature factor exponent takes the form:

$$-2\pi^2(h^2a^{*2}U_{11} + k^2b^{*2}U_{22} + \dots + 2hka^{*}b^{*}U_{12})$$

TABLE 5p Hydrogen coordinates ($\times 10^4$) and temperature factors ($\text{Å}^2 \times 10^3$)

atom	x	y	z	U
H(5a)	3743(32)	2745(27)	117(15)	70
H(5b)	3222(36)	4004(15)	1036(19)	70
H(6a)	4508(30)	1037(20)	1563(20)	62
H(6b)	4158(32)	2237(24)	2502(12)	62
H(5ax)	4291(11)	2591(9)	2486(7)	70
H(5bx)	5104(11)	1444(9)	1560(7)	70
H(6ax)	2260(11)	3951(9)	969(7)	70
H(6bx)	3103(11)	2837(9)	49(7)	70

(SYN)

DISTRIBUTION LIST

Dr. R.S. Miller
Office of Naval Research
Code 432P
Arlington, VA 22217
(10 copies)

Dr. A.L. Slafkosky
Scientific Advisor
Commandant of the Marine Corps
Code RD-1
Washington, DC 20380

JHU Applied Physics Laboratory
ATTN: CPLA (Mr. T.W. Christian)
Johns Hopkins Rd.
Laurel, MD 20707

Dr. Kenneth D. Hartman
Hercules Aerospace Division
Hercules Incorporated
Alleghany Ballistic Lab
P.O. Box 210
Washington, DC 21502

Mr. Otto K. Heiney
AFATL-DLJG
Elgin AFB, FL 32542

Dr. Merrill K. King
Atlantic Research Corp.
5390 Cherokee Avenue
Alexandria, VA 22312

Dr. R.L. Lou
Aerojet Strategic Propulsion Co.
Bldg. 05025 - Dept 5400 - MS 167
P.O. Box 15699C
Sacramento, CA 95813

Dr. R. Olsen
Aerojet Strategic Propulsion Co.
Bldg. 05025 - Dept 5400 - MS 167
P.O. Box 15699C
Sacramento, CA 95813

Dr. J. Pastine
Naval Sea Systems Command
Code 06R
Washington, DC 20362

Dr. Henry P. Marshall
Dept. 93-50, Bldg 204
Lockheed Missile & Space Co.
3251 Hanover St.
Palo Alto, CA 94304

Dr. Ingo W. May
Army Ballistic Research Lab.
ARRADCOM
Code DRXBR - 1BD
Aberdeen Proving Ground, MD 21005

Dr. R. McGuire
Lawrence Livermore Laboratory
University of California
Code L-324
Livermore, CA 94550

P.A. Miller
736 Leavenworth Street, #6
San Francisco, CA 94109

Dr. W. Moniz
Naval Research Lab.
Code 6120
Washington, DC 20375

Dr. K.F. Mueller
Naval Surface Weapons Center
Code R11
White Oak
Silver Spring, MD 20910

Prof. M. Nicol
Dept. of Chemistry & Biochemistry
University of California
Los Angeles, CA 90024

(SYN)

DISTRIBUTION LIST

Dr. Randy Peters
Aerojet Strategic Propulsion Co.
Bldg. 05025 - Dept 5400 - MS 167
P.O. Box 15699C
Sacramento, CA 95813

Dr. D. Mann
U.S. Army Research Office
Engineering Division
Box 12211
Research Triangle Park, NC 27709-2211

Mr. R. Geisler
ATTN: DY/MS-24
AFRPL
Edwards AFB, CA 93523

Naval Air Systems Command
ATTN: Mr. Bertram P. Sobers
NAVAIR-320G
Jefferson Plaza 1, RM 472
Washington, DC 20361

R.B. Steele
Aerojet Strategic Propulsion Co.
P.O. Box 15699C
Sacramento, CA 95813

Mr. M. Stossz
Naval Surface Weapons Center
Code R103
White Oak
Silver Spring, MD 20910

Mr. E.S. Sutton
 Thiokol Corporation
 Elkton Division
 P.O. Box 241
 Elkton, MD 21921

Dr. Grant Thompson
Morton Thiokol, Inc.
Wasatch Division
MS 240 P.O. Box 524
Bingham City, UT 84302

Mr. L. Roslund
Naval Surface Weapons Center
Code R10C
White Oak, Silver Spring, MD 20910

Dr. David C. Sayles
Ballistic Missile Defense
Advanced Technology Center
P.O. Box 1500
Huntsville, AL 35807

Director
US Army Ballistic Research Lab.
ATTN: DRXER-TED
Aberdeen Proving Ground, MD 21005

Commander
US Army Missile Command
ATTN: DRSMI-RKL
Walter W. Wharton
Redstone Arsenal, AL 35898

T. Yee
Naval Weapons Center
Code 3265
China Lake, CA 93555

Dr. E. Zimet
Office of Naval Technology
Code 071
Arlington, VA 22217

Dr. Ronald L. Derr
Naval Weapons Center
Code 389
China Lake, CA 93555

T. Boggs
Naval Weapons Center
Code 389
China Lake, CA 93555

Lee C. Estabrook, P.E.
Morton Thiokol, Inc.
P.O. Box 30058
Shreveport, LA 71130

(SYN)

DISTRIBUTION LIST

Dr. G. Neece
Office of Naval Research
Code 413
Arlington, VA 22217

Mr. C.M. Havlik
0/83-10, B/157-3W
Lockheed Missiles & Space Co., Inc.
P.O. Box 504
Sunnyvale, CA 94086

Dr. Philip Howe
Ballistic Research Laboratory
Code DRXBR-TBD
Aberdeen Proving Ground, MD 21005

Prof. C. Sue Kim
Department of Chemistry
California State University, Sacramento
Sacramento, California 95819

Mr. J. Moniz
Naval Ordnance Station
Code 5253L
Indian Head, MD 20640

Dr. R. Reed Jr.
Naval Weapons Center
Code 38904
China Lake, CA 93555

L.H. Sperling
Materials Research Center #32
Lehigh University
Bethlehem, PA 18015

Dr. Kurt Baum
Fluorocchem, Inc.
680 South Ayon Ave.
Azusa, CA 91702

Dr. Andrew C. Victor
Naval Weapons Center
Code 3208
China Lake, CA 93555

Dr. J.C. Hinshaw
Morton Thiokol Inc.
P.O. Box 524
Mail Stop 240
Brigham City, Utah 84302

Dr. V.J. Keenan
Anal-Syn Lab. Inc.
P.O. Box 547
Paoli, PA 19301

G.E. Manser
Morton Thiokol
Wasatch Division
P.O. Box 524
Brigham City, Utah 84302

P. Politzer
Chemistry Department
University of New Orleans
New Orleans, Louisiana 70148

Mr. David Siegel
Office of Naval Research
Code 253
Arlington, VA 22217

Dr. Rodney L. Willer
Morton Thiokol, Inc.
P.O. Box 241
Elkton, MD 21921

Dr. R. Atkins
Naval Weapons Center
Code 3852
China Lake, CA 93555

(CONT)

DISTRIBUTION LIST

Prof. J.H. Boyer
University of Illinois
Department of Chemistry
Box 4348
Chicago, Illinois 60680

Prof. J.C. Chien
University of Massachusetts
Department of Chemistry
Amherst, MA 01003

Dr. B. David Halpern
Polysciences, Inc.
Paul Valley Industrial Park
Warrington, PA 18976

Dr. M.B. Frankel
Rockwell International
Rockwood Division
6631 Canoga Avenue
Canoga Park, CA 91304

Dr. R.A. Earl
Hercules, Inc.
Magna, Utah 84109

Dr. C. Bedford
SRI International
333 Ravenswood Avenue
Menlo Park, CA 94025

Dr. Robert R. Ryan
INC-4, MS C346
Los Alamos National Laboratory
Los Alamos, New Mexico 87545

Dr. Robert D. Chapman
AFPL/LMR
Edwards AFB, CA 93523

Dr. L. Erwin
MIT
Room 36-306
Cambridge, MA 02139

Dr. W. Farber
Space Sciences, Inc.
105 W. Maple Avenue
Monrovia, CA 91016

Dr. W.H. Graham
Morton Thiokol, Inc.
Hunstville Division
Hunstville, AL 35807-7501

Dr. C. Coon
Lawrence Livermore Lab.
University of California
P.O. Box 308
Livermore, CA 94550

Dr. R. Gilardi
Naval Research Laboratory
Code 6030
Washington, DC 20373

Dr. Alan Marchand
Dept. of Chemistry
North Texas State University
NTSU Station, Box 5068
Denton, Texas 76203

T.B. Brill
Department of Chemistry
University of Delaware
Newark, Delaware 19716

Dr. A.A. Defusco
Code 3858
Naval Weapons Center
China Lake, CA 93553

Dr. Richard A. Hollins
Naval Weapons Center
Code 3853
China Lake, CA 93553

Dr. R. Armstrong
MIT
Room 46-505
Cambridge, MA 02139

Professor Philip E. Eaton
Department of Chemistry
University of Chicago
5735 South Ellis Avenue
Chicago, IL 60637

(SYN)

DISTRIBUTION LIST

Mr. J. Consaga
Naval Surface Weapons Center
Code R-16
Indian Head, MD 20640

Naval Sea Systems Command
ATTN: Mr. Charles M. Christensen
NAVSEA62R2
Crystal Plaza, Bldg. 6, Rm 806
Washington, DC 20362

Mr. R. Beauregard
Naval Sea Systems Command
SEA 64E
Washington, DC 20362

Dr. Anthony J. Matuszko
Air Force Office of Scientific Research
Directorate of Chemical & Atmospheric
Sciences
Bolling Air Force Base
Washington, DC 20332

Dr. Michael Chaykovsky
Naval Surface Weapons Center
Code R11
White Oak
Silver Spring, MD 20910

J.J. Rocchio
USA Ballistic Research Lab.
Aberdeen Proving Ground, MD 21005-5066

G.A. Zimmerman
Aerojet Tactical Systems
P.O. Box 13400
Sacramento, CA 95813

B. Swanson
INC-4 MS C-346
Los Alamos National Laboratory
Los Alamos, NM 87545

Dr. L.H. Caveny
Air Force Office of Scientific
Research
Directorate of Aerospace Sciences
Bolling Air Force Base
Washington, DC 20332

W.G. Roger
Code 5253
Naval Ordnance Station
Indian Head, MD 20640

Dr. Donald L. Bell
Air Force Office of Scientific
Research
Directorate of Chemical &
Atmospheric Sciences
Bolling Air Force Base
Washington, DC 20332

Dr. H.G. Adolph
Naval Surface Weapons Center
Code R11
White Oak
Silver Spring, MD 20910

U.S. Army Research Office
Chemical & Biological Sciences
Division
P.O. Box 12211
Research Triangle Park, NC 27709

G. Butcher
Hercules, Inc.
MS K2H
P.O. Box 98
Magna, Utah 84044

W. Waesche
Atlantic Research Corp.
7511 Wellington Road
Gainesville, VA 22065

RD-A164 813

STRUCTURAL CHEMISTRY OF ENERGETIC COMPOUNDS(U) OFFICE
OF NAVAL RESEARCH ARLINGTON VA R GILARDI ET AL.

01 OCT 85


2/2

UNCLASSIFIED

F/G 7/4

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

(SYN)

DISTRIBUTION LIST

Dr. James T. Bryant,
Naval Weapons Center
Code 32053
China Lake, CA 93555

Dr. L. Rothstein
Assistant Director
Naval Explosives Dev. Engineering Dept.
Naval Weapons Station
Yorktown, VA 23691

Dr. M.J. Kamlet
Naval Surface Weapons Center
Code R11
White Oak, Silver Spring, MD 20910

Dr. Henry Webster, III
Manager, Chemical Sciences Branch
ATTN: Code 5063
Crane, IN 47522

Dr. R.S. Valentini
United Technologies Chemical Systems
P.O. Box 50015
San Jose, CA 95150-0015

Administrative Contracting
Officer (see contract for
address)
(1 copy)

Defense Technical Information Center
Bldg. 5, Cameron Station
Alexandria, VA 22314
(12 copies)

Arpad Junasz
Code DRDAR-IBD
Ballistic Research Lab
Aberdeen, MD 21005

Mr. C. Gotzmer
Naval Surface Weapons Center
Code R-11
White Oak
Silver Spring, MD 20910

Dr. John S. Wilkes, Jr.
FJSRL/NC
USAF Academy, CO 80840

Dr. H. Roserwasser
Naval Air Systems Command
AIR-320R
Washington, DC 20361

Dr. A. Nielsen
Naval Weapons Center
Code 385
China Lake, CA 93555

Dr. Joyce J. Kaufman
The Johns Hopkins University
Department of Chemistry
Baltimore, MD 21218

Dr. J.R. West
Morton Thiokol, Inc.
P.O. Box 30058
Shreveport, LA 71130

Director
Naval Research Laboratory
Attn: Code 2627
Washington, DC 20375
(6 copies)

Dr. Robert J. Schmidt
SRI International
333 Ravenswood Avenue
Menlo Park, CA 94025

Dr. Michael D. Coburn
Los Alamos National Lab
M-1, Explosives Technology
Mail Stop, C920
Los Alamos, NM 87545

(SYN)

DISTRIBUTION LIST

Dr. R.F. Walker
Chief, Energetic Materials Division
DRSMC-LCE (D), B-3022
USA ARDC
Dover, NJ 07801

Dr. Janet Wall
Code 012
Director, Research Administration
Naval Postgraduate School
Monterey, CA 93943

R.E. Shenton
Atlantic Research Corp.
7511 Wellington Road
Gainesville, VA 22065

Mike Barnes
Atlantic Research Corp.
7511 Wellington Road
Gainesville, VA 22065

Dr. Lionel Dickinson
Naval Explosive Ordnance
Disposal Tech. Center
Code D
Indian Head, MD 20340

Prof. J.T. Dickinson
Washington State University
Dept. of Physics 4
Pullman, WA 99164-2814

M.H. Miles
Dept. of Physics
Washington State University
Pullman, WA 99164-2814

Dr. T.F. Davidson
Vice President, Technical
Morton Thiokol, Inc.
Aerospace Group
110 North Wacker Drive
Chicago, IL 60606

Dr. D.D. Dillehay
Morton Thiokol, Inc.
Longhorn Division
Marshall, TX 75670

G.T. Bowman
Atlantic Research Corp.
7511 Wellington Road
Gainesville, VA 22065

Brian Wheatley
Atlantic Research Corp.
7511 Wellington Road
Gainesville, VA 22065

Mr. G. Edwards
Naval Sea Systems Command
Code 62R32
Washington, DC 20362

C. Dickinson
Naval Surface Weapons Center
White Oak, Code R-13
Silver Spring, MD 20910

Prof. John Deutch
MIT
Department of Chemistry
Cambridge, MA 02139

Dr. E.H. deButts
Hercules Aerospace Co.
P.O. Box 27408
Salt Lake City, UT 84127

David A. Flanigan
Director, Advanced Technology
Morton Thiokol, Inc.
Aerospace Group
110 North Wacker Drive
Chicago, IL 60606

DTIC

END

4-86