

AD-A164 930

HOST CELL SURFACE EXPRESSION OF RICKETTSIA TYPHI
ANTIGENS DURING INFECTION(U) NAVAL MEDICAL RESEARCH
INST BETHESDA MD F M ROLLWAGEN ET AL. 1985 NMRI-85-100


1/1

UNCLASSIFIED

F/G 6/5

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A164 930 PAGE

| | | | |
|--|--|---|--------------------------------------|
| 1a. REPORT SECURITY CLASSIFICATION | | MARKINGS | |
| 2a. SECURITY CLASSIFICATION AUTHORITY | | 3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution is unlimited | |
| 2b. DECLASSIFICATION / DOWNGRADING SCHEDULE | | | |
| 4. PERFORMING ORGANIZATION REPORT NUMBER(S) NMRI 85-100 | | 5. MONITORING ORGANIZATION REPORT NUMBER(S) | |
| 6a. NAME OF PERFORMING ORGANIZATION Naval Medical Research Institute | 6b. OFFICE SYMBOL (if applicable) | 7a. NAME OF MONITORING ORGANIZATION Naval Medical Command | |
| 6c. ADDRESS (City, State, and ZIP Code) Bethesda, Maryland 20814-5055 | | 7b. ADDRESS (City, State, and ZIP Code) Department of the Navy Washington, D. C. 20372-5120 | |
| 9a. NAME OF FUNDING / SPONSORING ORGANIZATION Naval Medical Research & Development Command | 8b. OFFICE SYMBOL (if applicable) | 9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER | |
| 8c. ADDRESS (City, State, and ZIP Code) Bethesda, Maryland 20814-5044 | | 10. SOURCE OF FUNDING NUMBERS | |
| | | PROGRAM ELEMENT NO 61102A | PROJECT NO 3M161102BS10 AC |
| | | TASK NO. | WORK UNIT ACCESSION NO. DA301604 |
| 11. TITLE (Include Security Classification) Host Cell Surface Expression of Rickettsia typhi Antigens During Infection. | | | |
| 12. PERSONAL AUTHOR(S) F. M. Rollwagen and G. A. Dasch | | | |
| 13a. TYPE OF REPORT | 13b. TIME COVERED FROM _____ TO _____ | 14. DATE OF REPORT (Year, Month, Day) 1985 | 15. PAGE COUNT |
| 16. SUPPLEMENTARY NOTATION IN: RICKETTSIAE AND RICKETTSIAL DISEASES. Proceedings of the 3d International Symposium. (Cont | | | |
| 17. COSATI CODES | | 18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) | |
| FIELD | GROUP | Endemic Typhus Rickettsial Diseases | |
| | | Rickettsia typhi Rickettsial Immunity | |
| | | Rickettsial antigens | |
| 19. ABSTRACT (Continue on reverse if necessary and identify by block number) | | | |
| 20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS | | 21. ABSTRACT SECURITY CLASSIFICATION | |
| | | | |
| 22a. NAME OF RESPONSIBLE INDIVIDUAL Rosemary Spitzen, Information Services Branch | | 22b. TELEPHONE (Include Area Code) 202-295-2188 | 22c. OFFICE SYMBOL ISB/ADMIN/NMRI |

DTIC FILE COPY

S DTIC ELECTED D

MAR 04 1988

D

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

16. (Continued from front)

Edited by J. Kazar. Bratislava, Publishing House of the Slovak Academy of Sciences,
1985. pp.203-210

DTIC

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

RICKETTSIAE AND RICKETTSIAL DISEASES

Proceedings of the IIIrd International Symposium

HELD AT SMOLENICE CASTLE, SEPTEMBER 10-14, 1984

Edited by J. KAZÁR

**PUBLISHING HOUSE
OF THE SLOVAK ACADEMY OF SCIENCES
BRATISLAVA 1985**


Host Cell Surface Expression of Rickettsia typhi
Antigens During Infection

F. M. Rollwagen and G. A. Dasch

Naval Medical Research Institute
Bethesda, Maryland

This investigation was supported by the Naval Medical Research and Development Command, Department of the Navy, Research Task No. 3M161102BS10.AC.423.

The opinions and assertions contained herein are the private ones of the writer and are not to be construed as official or reflecting the views of the Navy Department or the naval service at large.

| | |
|---------------------|---|
| Accession For | |
| NTIS CRA&I | <input checked="" type="checkbox"/> |
| DTIC TAB | <input type="checkbox"/> |
| Unannounced | <input type="checkbox"/> |
| Justification | |
| By | |
| Distribution / | |
| Availability Codes | |
| Dist | Avail and/or Special |
| A-1 |  |

HOST CELL SURFACE EXPRESSION OF RICKETTSIA TYPHI
ANTIGENS DURING INFECTION

F. M. ROLLWAGEN and G. A. DASCH

Naval Medical Research Institute,
Bethesda, Maryland 20814 U.S.A.

Immune responses to rickettsial infection have been attributed to both humoral and cellular mechanisms. Although it has been shown that transfer of immune cells (1) or serum (2) can mediate resistance to challenge, the exact immune mechanisms involved remain poorly understood. Since rickettsiae are obligate intracellular bacteria, an immune attack upon rickettsiae while they are still inside the host cell would represent a powerful immunologic defense. In the present report, we show that tissue culture cells infected with Rickettsia typhi express rickettsia-specific antigens on their surfaces. Such antigens may provide triggering mechanisms for both the afferent and efferent arms of the immune system.

Keywords: Endemic typhus

MATERIALS AND METHODS

1. Cell line: The LS-929 cell line is a suspension variant of the fibroblast-like cell line derived from a C3H/AnN mouse. It is maintained in Eagle's MEM supplemented with L-glutamine and fetal calf serum, without added antibiotics.

2. Rickettsia: Rickettsia typhi strain Wilmington was grown in the yolk sac of embryonated chicken eggs, purified by isopycnic banding in Renografin 76 density gradients, and controlled-rate frozen to -100°C at 0.5 mg protein/0.5 ml in Bovarnick's sucrose phosphate glutamate supplemented with 5 mM MgCl_2 and 1% Renografin 76.

3. Fluoresceinated reagents: Rabbit antiserum to R. typhi (RbRt) was raised by multiple injections of intact and disrupted purified rickettsia in Freund's incomplete adjuvant. Antibodies were purified by salt precipitation and fluoresceinated by standard methods. As a negative control, rabbit anti-yolk sac (RbYS) antibodies were raised, purified and fluoresceinated in the same way as RbRt.

4. Flow cytometry: Fluorescence analysis was carried out on a FACSII (Becton, Dickinson, Sunnyvale, CA) equipped as described elsewhere (3,4). For surface analysis, the cells to be tested were washed twice in HBSS containing 0.01% NaN_3 . The cell pellet was resuspended and stained with fluoresceinated reagents prepared as described above. After 45 min on ice, the cells were washed again and fixed in 1% paraformaldehyde.

RESULTS

In Fig. 1 it can be seen that a subpopulation of LS-929 cells infected with R. typhi 72 hours earlier expresses surface antigens that are detectable with RbRt antiserum. That this reaction is not caused by nonspecific stickiness of the infected cells was shown by the negative staining profile of RbYS tested on R. typhi infected LS cells which completely overlapped the background fluorescence profile given by unstained infected or uninfected LS-929 (not shown). Dead cells are excluded from the analysis by forward light scatter. When the kinetics of acquisition of this surface antigen were examined it was found that as early as 24 h after infection, a shoulder of positively staining cells appeared. These positive cells were increased in number on day 2, reaching a maximum on day 3.

As noted in Fig. 1, only a subpopulation of infected cells bears the antigen on its surface. Even though the entire population was exposed to R. typhi organisms, it is possible that only some LS-929 cells became infected, thereby giving the biphasic pattern seen. However, by light

microscopy on day 3 all the cells infected with R. typhi contained numerous rickettsiae, although only 42% expressed surface antigens, as shown in Fig. 1. The possibility that expression of rickettsial antigens on the cell surface is dependent on cell cycle is under investigation.

Since R. typhi infected cells may liberate intact organisms, or soluble antigens into the culture medium, it is possible that this antigen may passively adsorb on the surfaces of cells. Four different approaches were used in this study.

In the first approach, LS929 cells were infected at a high multiplicity, and stained one hour after infection. This experiment (not shown) examined the possibility that rickettsiae were leaving antigen on the cell surface upon penetration. No staining was observed.

In the second approach, uninfected LS-929 cells were incubated with sonicated whole rickettsiae for three days at 37°C. The cells were then washed and stained with antiserum as described above. The results of a representative experiment are shown in Fig. 2A. At the highest doses of sonicate used (4 ug/flask) the FACS pattern virtually overlaps that of cells incubated with no antigen.

The third approach used was to measure the amount of rickettsial antigen in supernatant from infected cells. Using a sensitive enzyme-linked immunosorbent assay (ELISA) and an antigen-capture assay, titrated amounts of supernatant from three-day infected LS-929 cells were analyzed for the amounts of antigen they contained. The results of these experiments, suggested that only small amounts of antigen were present: about 100-300 ng/ml of supernatant. This range is marked with an asterisk on Fig. 2A.

The fourth approach involved transfer of culture supernatant from three day infected LS-929 to uninfected cells, incubating an additional three days, and then examining the cells for their ability to bind RbxR.t. antiserum. Figure 2B shows the results of such an

experiment. It can be seen that LS-929 cells incubated for three days with culture supernatant from infected cells (containing 100-300 ng/ml antigen) did not exhibit the binding pattern seen after a three day infection (as shown in Fig. 1). These experiments strongly suggest that the appearance of rickettsial antigens on the surface of infected cells is dependent upon an ongoing infection, and is not due to external adsorption of antigen from the culture supernatant.

DISCUSSION

In theory organisms which replicate intracellularly and migrate from cell to cell might be relatively sequestered from the systemic effects of antibodies, or the local effects of cellular immune mechanisms. Further, intracellular organisms may not be presented as immunogens in an efficient manner. In the present report, we show that L-cells infected with the obligate intracellular bacterium, Rickettsia typhi express antigens on their surface which are recognized by an antibody directed to rickettsiae.

It is known that cells infected with other intracellular organisms such as viruses (5) protozoa (6) or bacteria (7) express antigen on their surface. In contrast to viral antigens, which are expressed on the infected cell surface within hours (7), the rickettsial antigens required a few days for maximum expression. This may reflect the slower replication time of rickettsiae relative to viruses. Since cycloheximide inhibits expression of rickettsial antigen under these conditions (Rollwagen and Dasch, paper in preparation), it might be inferred that a eukaryotic host processing (glycosylation?) step is involved. These mechanisms are currently under study in our laboratory.

ACKNOWLEDGEMENTS

The authors wish to acknowledge the capable technical assistance of A.J. Bakun and C. Campo; Neil Hardegan for the FACS analysis; D. Boyle for editorial assistance; M. Dobson for the antigen-capture ELISA; and L. Yaffe and R. Wistar for critical review of the manuscript.

REFERENCES

1. CRIST, JR., A.E., WISSEMAN, JR., C.L., and MURPHY, J.R. (1984): *Infect. Immun.*, 43:38.
2. ANACKER, R.L., PHILIP, R.N., CASPER, E., TODD, W.J., MANN, R.E., JOHNSTON, M.R., and NAUCK, C.J. (1983): *Infect. Immun.*, 40:292.
3. LOKEN, M.R., and HERZENBERG, L.A. (1975): *Ann. N.Y. Acad. Sci.*, 245:163.
4. MILLER, H.M., POWELL, J.I., SHARROW, S.O., and SCHULTZE, A.R. (1978): *Rev. Sci. Instrum.*, 49:1137.
5. PLATA, F., KALIL, G., ZILBER, M-T., FELLOWS, M., and LEVY, D. (1983): *J. Immunol.*, 131:2551.
6. EMERY, D.L., and KAR, S.K. (1983): *Immunol.*, 48:723.
7. ZINKERNAGEL, R.M., ALTHAGE, A., ADLER, B., BLANDEN, F.V., DAVIDSON, W.F., KEES, V., DUNLOP, M.B.C., and SCHREFFLER, D.C. (1977): *J. Exp. Med.*, 145:1353.




Figure 1. FACS profile of LS-929 cells uninfected or infected with R. typhi stained with fluoresceinated antibody to R. typhi or unstained.





Figure 2. A. Uninfected LS-929 cells incubated with sonicated whole rickettsiae for 3 days in vitro, washed and stained with Rb α Rt
 B. Supernatant from 72-hour infected or uninfected LS-929 was transferred (1:1) to uninfected LS-929 and incubated for 3 days. The cells were washed and stained with Rb α Rt

END

FILMED

3 - 86

DTIC