

MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

①

AD-A 165 024

DTIC
ELECTE
MAR 1 1 1986
S D

A REVIEW
OF
THE AQUATIC BIOLOGICAL RESOURCES OF
THE ATLANTIC COASTAL AREA
OFF VIRGINIA BEACH, VIRGINIA

Planning Aid Report by the
U.S. Fish and Wildlife Service
Under Provisions of the
Fish and Wildlife Coordination Act
For a Proposed Ocean Dredged Material Disposal Site
by the Norfolk District,
U.S. Army Corps of Engineers

February, 1984

DTIC FILE COPY

Prepared by: George Ruddy
Karen Mayne

Feb 28, 1984
Date

Approved:

Glenn Kinser
Glenn Kinser
Supervisor
Annapolis Field Office

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited



US Army Corps
Of Engineers
Norfolk District

Report B-10

86 3 1 8

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release, distribution unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		4. PERFORMING ORGANIZATION REPORT NUMBER(S)	
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		5. MONITORING ORGANIZATION REPORT NUMBER(S) B-10	
6a. NAME OF PERFORMING ORGANIZATION U.S. Fish and Wildlife Service	6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION U.S. Army Corps of Engineers, Norfolk District	
6c. ADDRESS (City, State, and ZIP Code) Annapolis, MD 21401		7b. ADDRESS (City, State, and ZIP Code) Norfolk, Virginia 23510-1096	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION U.S. Army Corps of Engineers, Norfolk District	8b. OFFICE SYMBOL (if applicable) NAOPL; NAOEN	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code) Norfolk, Virginia 23510-1096		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO.	PROJECT NO.
		TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (through Security Classification) A review of the Aquatic Biological Resources of the Atlantic Coastal Area Off Virginia Beach, Virginia			
12. PERSONAL AUTHOR(S) Ruddy, G. and K. Mayne			
13a. TYPE OF REPORT Final	13b. TIME COVERED FROM TO	14. DATE OF REPORT (Year, Month, Day) 1984, February	15. PAGE COUNT 37
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
		Dam Neck Disposal Site, Virginia, biological resources, phyto plankton, zooplankton, benthic fauna, finfish, commercial fisheries, recreational fisheries, endangered species, spawn-	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Concludes that DNDS and vicinity is one of the most biologically unique and important areas along the Atlantic coast.			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Craig L. Seltzer		22b. TELEPHONE (Include Area Code) (804) 441-3767/827-3767	22c. OFFICE SYMBOL NAOPL-R

18. ing, nursery areas, Chesapeake Bay plume

This report will provide planning aid information for the Norfolk Harbor Deepening Study. Consideration is currently being given to establishing a new dredged material disposal site further inshore from the previously proposed site, located approximately 17 miles east of the Bay mouth. We understand that although no firm alternative site has been selected for study, one alternative being considered is deposition of disposal material north of the previously utilized Dam Neck disposal site in a manner to create an off-shore bar.

This report will attempt to describe the baseline biological conditions in the study area. Because of the short time frame, absence of project specifics, and broad scope of this report, the level of detail of our treatment of this ecologically complex area will necessarily be limited. As the project requirements become better defined, we will be able to provide more detailed information on particular areas of concern.

For the purposes of this report we are considering the study area to be the area bounded by 37 degrees 10' latitude on the north, 36 degrees 40' latitude on the south, 75 degrees 40' longitude on the east and the shoreline and Bay mouth on the west (Figure 1). The ecology of this area is extremely dynamic. As is typical of mid-Atlantic coastal areas, the biological make-up changes dramatically on a seasonal basis. However, it is the proximity of the Chesapeake Bay, the largest estuary on the east coast, which primarily adds to the complexity and importance of this area.

Phytoplankton

Phytoplankton in the study area are dominated by lower Chesapeake Bay species which are transported eastward from the mouth of the Bay and then southward along the Virginia coast by the Chesapeake Bay plume. Species composition fluctuates seasonally. Diatoms are typically dominant in the fall, winter and spring, while in the summer there is a mixture of diatoms, phytoflagellates and nannoplankters. Moving away from the Bay mouth and out over the continental shelf, the phytoplankton composition changes to reflect an increase in shelf species and a decrease in concentration. Marshall (1980) investigated the phytoplankton within the Bay plume and adjacent shelf waters. Based on his sampling (Figure 2) he identified phytoplankton assemblages within the Bay plume and adjacent shelf water for March, June and October (Table 1). His findings also show that phytoplankton cell concentrations in the Bay plume water are significantly higher than in shelf waters outside the plume (Figure 3). Similarly, an examination of respiration rates in the Bay plume (Robertson and Thomas, 1981) indicated that the plume supports a higher level of biological activity than adjacent non-affected shelf waters.

Zooplankton

The zooplankton forms a vital part of the ecological picture in the study area. Not only does it provide an important trophic link between the primary production of the phytoplankton and higher animal forms, but it

<input checked="" type="checkbox"/>
<input type="checkbox"/>
<input type="checkbox"/>
<input type="checkbox"/>
<input type="checkbox"/>
<input type="checkbox"/>
<input type="checkbox"/>
<input type="checkbox"/>
Codes



Dist	Special
A-1	

also includes the planktonic larval stages of many marine and estuarine invertebrates and fish. Unfortunately, detailed comprehensive zooplankton studies in the study area are lacking. Researchers from Old Dominion University have been sampling zooplankton in the area regularly for two years, but their collections have not yet been analyzed. There are, however, a few studies available from which information on the zooplankton composition in the study area can be inferred. Focusing only on molluscs, decapods and fish, the U.S. Fish and Wildlife Service (1981) made six meroplankton collections in and around Rudee Inlet. Species and abundance from two of their stations at the mouth of Rudee Inlet are shown in Table 2. Grant and Olney (1979 and 1983) studied zooplankton in the lower Chesapeake Bay. Their sampling stations closest to the study area were in the vicinity of the Bay Bridge Tunnel. A list of species which they found at these stations is shown in Table 3. Grant (1979) surveyed zooplankton on a seasonal basis in the Middle Atlantic Bight. A list of species from his LI station, located approximately 20 miles offshore of Wachapreague is shown in Table 4. The locations of the zooplankton collections made in the above mentioned studies are shown in Figure 4. The zooplankton composition in the study area is likely to include a mixture of the species identified in these studies; that is to say, there will be a mixture of estuarine species from the Bay and continental shelf species. The relative composition of these species assemblages will vary with changes in the Bay discharge and meteorological conditions. Other complicating factors include tremendous seasonal shifts in species composition, diurnal vertical migrations which can bias sampling efforts, and substantial natural year to year variation. For these reasons it seems likely that the study area would include a large number of species which do not appear in any of the surveys we have reported.

The zooplankton includes many important species from higher trophic levels which are present during their developmental stages, mainly eggs and larvae. Of the species which were identified in the three studies mentioned above, there were a total of 32 species of molluscs, 38 species of decapods and 41 species of fish present in the zooplankton. The eggs and larvae of many decapods and fish were particularly prevalent in the late spring and summer, and were frequently concentrated in the surface layers.

Although there are several commercially and recreationally important fish species which will occur in the zooplankton in the study area, the blue crab (*Callinectes sapidus*) may be the most important species to regularly occur in high concentrations. Chesapeake Bay blue crabs spawn in the lower Bay during the summer months. Recent studies indicate that after hatching the larvae move into the neuston (surface layer) and are transported out of the Bay (Provenzano et al., 1982). Larval development occurs in offshore waters in the neuston (Smyth, 1980; McConaughy et al., 1983). Larval development to the megalopa stage has been estimated to take 30 to 40 days (McConaughy et al., 1983). Since the Chesapeake Bay plume moves southward toward Cape Hatteras, it is not clear how the crab larvae avoid being trapped in this flow and lost from the Chesapeake Bay system. However, since the larvae occupy the surface layer, wind driven surface currents may play a major role in determining their distribution. Johnson et al. (1983) have postulated that the southerly winds common in this area in the summer could retain the

larvae in the offshore waters near the Bay mouth, thereby facilitating their recruitment back into Chesapeake Bay. The dimensions of this nearshore "nursery area" have not as yet been determined.

Benthic Fauna

There have been several studies on benthic fauna at various locations within the study area. Our office conducted a one time survey of benthic infauna at 8 stations and epifauna at 3 additional stations approximately 3 miles offshore from Virginia Beach (Figure 5) in June of 1982. The results from this study were submitted to your office in a planning aid report for the Virginia Beach Hurricane Protection Study (USFWS, 1982). In this report we also reviewed other benthic studies conducted on the mid-Atlantic continental shelf. Species lists from this study are shown in Tables 5 and 6. The Hampton Roads Sanitation District has been conducting benthic surveys in the vicinity of their Virginia Beach sewage treatment plant discharge pipe for several years. Although these data are not presently available, a species inventory from an earlier study done for them by Hydrosience, Inc. (1974) is shown in Table 7 and Figure 6. Based on these studies it appears that the study area supports a productive benthic faunal population which in general character is typical of the mid-Atlantic inner continental shelf.

Finfish

The fishery resources in the study area are rich and varied. A species list compiled from National Marine Fisheries Service groundfish trawl surveys between 1977 and 1980 is given in Table 8. The surveys identified 76 species, but this is far from a complete list of the species that use the area. As is typical for the mid-Atlantic region, most species undergo seasonal migrations. The species composition in the study area is also strongly influenced by the large number of species which migrate in and out of the Chesapeake Bay. Of the 209 marine and estuarine fish species known to occur in Chesapeake Bay, 180 species migrate in and out of the Bay (Musick, 1972). Spring and fall are the peak migration periods. It is of particular interest that many species migrate through the area during their more vulnerable larval or early juvenile stages on their way to nursery areas within Chesapeake Bay. These would include species of commercial and recreational importance such as menhaden (*Brevoortia tyrannus*), weakfish (*Cynoscion regalis*), croaker (*Micropogon undulatus*), spot (*Leiostomus xanthurus*) and summer flounder (*Paralichthys dentatus*) (Gosselein and Azarovitz, 1982). Weakfish spawn in the lower Bay and coastal waters in the summer. Their larvae are planktonic until the later stages of development when they become demersal (Mercer, 1983). In a survey of lower Chesapeake Bay ichthyoplankton, Olney (1983) found early stage weakfish larvae at densities up to 966/100 m² near the Bay mouth. Menhaden, croaker, spot and summer flounder spawn in offshore waters. Their larval and postlarval stages appear to follow inshore bottom currents into Chesapeake Bay (U.S. Dept of the Interior, 1970).

Commercial Fisheries

The commercial fisheries in the Atlantic Ocean off Virginia Beach are found in two general areas. The "offshore" fishery occurs in deep water from 20 to 100 miles or more off the coast and consists of oceanic species such as the surf clam, scallop, and lobster, and deep water finfish such as tuna, dolphin and bonito. The "inshore" fishery occurs in the nearshore area from just outside the breaker zone to about 3 to 4 miles offshore. This fishery mainly harvests the finfish that migrate along the Atlantic coast and into and out of the Chesapeake Bay. The following is a brief discussion of the major inshore fisheries, which is accompanied by a map that shows the locations of these fisheries. It is based on discussions with the Virginia Institute of Marine Science, the Virginia Marine Resources Commission, and local watermen and seafood dealers.

There is a small fishery for blue crabs in the area, which is incidental to the major blue crab fishery within the Chesapeake Bay. A summer crab pot fishery exists from Cape Henry south to below Rudee Inlet, immediately east of the breaker zone and in Smith Inlet on the Eastern Shore. The crab pot fishery has been restricted within the mouth of the Chesapeake Bay by the State of Virginia because the lower Chesapeake Bay is the blue crab spawning and nursery area. Although the Virginia Beach pot fishery is outside of the restricted zone, most of the crabs that are caught in this area are also the spawning females. A winter crab dredge fishery exists along the flanks of the Chesapeake Bay entrance channels. This area is only fished during good weather conditions. The majority of the winter fishery occurs inside the mouth of the Chesapeake Bay.

A small conch fishery (*Busycon* sp. and *Strombus* sp.) occurs along Virginia Beach from about 1.5 to 3 miles offshore. This fishery takes place year round but is usually incidental to the more important finfisheries. The abundance of conch appears to vary from year to year, which may be related to fishing pressure.

A purse seine fishery for menhaden is located at the mouth of the Chesapeake Bay from late May through October. Purse seining may also occur in the nearshore area of the Atlantic Ocean during this period if spotter planes locate large concentrations of fish.

A gill net fishery is located from just outside the breaker zone to about the 20-foot contour south of the mouth of the Chesapeake Bay down to the North Carolina state line. Restrictions by the Navy limit fishing off the Dam Neck area. A shad fishery (*Alosa* sp.), utilizing floating gill nets, occurs during the spring run of shad into the Chesapeake Bay from March through April. Anchored gill nets are fished in the area between April and September, with the largest catches during the spring and fall migrational periods. The primary species which are caught are bluefish (*Pomatomus saltatrix*), croaker, spot and sea trout (*Cynoscion regalis* and *C. nebulosus*).

The most important fishery in the study area is the trawl fishery, which extends from 1/4 to 4 miles offshore, both north and south of the mouth of the Chesapeake Bay. The primary fishery, however, is south of Cape Henry.

There is an intense flounder fishery (primarily summer flounder, (Paralichthys dentatus) in the fall between about October and December. Other major species which are caught during the spring and fall migrations into and out of the Chesapeake Bay include spot, croaker, sea trout and bluefish. Striped bass (Morone saxatilis) were previously caught during the winter; however, the decline of the striped bass stocks in recent years has eliminated this fishery.

It is difficult to determine the size and economic value of the commercial fisheries in the study area. The methods that have been used by the National Marine Fisheries Service and the Virginia Marine Resources Commission to compile the statistics on the annual landings of commercial species from the Atlantic Ocean will not allow a breakdown of the catches on a small scale. Therefore, catch data for the study area includes a large area that extends from the shoreline to 75 degrees longitude and from 36 degrees to 37 degrees latitude. Catch data from the period of 1972 to 1981 have previously been compiled for the Norfolk Harbor Deepening Study and can be provided in draft form if necessary.

One of the unanswered questions regarding the commercial fisheries is whether the location of the fisheries adequately reflects the actual limits of the resources. For example, it is known that blue crabs extend much further offshore than what the extent of the crab pot fishery would indicate. However, fishermen find that it is both uneconomical and impractical to set their pots in offshore areas. This may also be the case with the finfisheries. Several watermen have indicated that they do not trawl further offshore than 4 miles because the bottom is bad, there are many obstructions, and because of the shipping lanes. Therefore, while it is speculated that the major concentrations of finfish occur within the 3 mile zone, it is possible that these fish are also found further offshore, but in lesser numbers. Of course, the location of migrating fish along the Virginia coast can vary greatly from year to year, depending on such variables as weather conditions, water temperatures, current patterns and food sources.

Recreational Fisheries

There is a substantial amount of recreational fishing in and around the study area. It is derived from commercial charter and head boats, private boats, pier fishing and beach fishing. The accompanying map displays some particularly popular fishing spots and target species. The map is based on information from the publication Saltwater Sport Fishing and Boating in Virginia (1981), and discussions with VIMS and local charter boat captains. It should be realized, however, that substantial year to year variation is expected depending on the condition of the stocks, climatic conditions, etc. The sport fishing season usually begins in March and early April with the arrival of Atlantic (Boston) mackerel (Scomber scombrus) in water 4 to 12 miles off the Virginia coast. These fish remain in the area only a short time before migrating northward. Black sea bass (Centropristis striata) and tautog (Tautoga onitis) are sought after on wrecks and artificial reefs which exist in the area. Black sea bass are caught from

spring to fall, while tautog are more restricted to just the spring and fall months (Marshall and Lucy, 1981). Bluefish are caught throughout the area, especially around the Chesapeake Light Tower, from spring to fall. Weakfish, croaker, spot and summer flounder are also caught from late spring to fall, although these fish are more frequently sought after inside the Bay mouth. It should be noted that fishing even occurs within the restricted area around Cape Henry despite the regulations prohibiting "anchoring, trawling, fishing and dragging." Fishing for bluefin tuna (Thunnus thynnus) occurs 10 to 30 miles offshore in June and July. King mackerel (Scomberomorus cavella) are also caught in this vicinity from late August to October. Other pelagic game species such as white marlin (Tottrapturus albidus), wahoo (Acanthocybium solardii), false albacore (Euthynnus alletteratus), skipjack tuna (Katsuwonus pelamis), dolphin (Coryphaena hippurus), Atlantic bonita (Sarda sarda), blue marlin (Makaira nigricans), and yellowfin tuna (Thunnus albacares) are caught in deeper waters, generally 30 to 50 miles offshore.

Endangered Species

The only endangered or threatened species which would occur in the study area on more than an occasional or transient basis are marine turtles. The loggerhead turtle (Caretta caretta), which is Federally listed as threatened, and the leatherback (Dermochelys coriacea) and Atlantic ridley (Lepidochelys kempis) turtles, which are Federally listed as endangered, are known to frequently occur in the study area from May through October (Richard Byles, VIMS, personal communication). Leatherbacks occur around the Bay mouth where they often feed on jellyfish. Loggerheads and ridleys, especially the juveniles, have more of a tendency to penetrate further up the Bay. Both these species spend a good deal of time on the bottom feeding on epifaunal invertebrates. All three species are under the jurisdiction of the National Marine Fisheries Service (except when nesting) and further information can be obtained by contacting Doug Beach, NMFS, Gloucester, Massachusetts, at (617) 281-3600.

Miscellaneous Resource Considerations

The area extending from Assateague to Fisherman's Island and offshore for 10 miles is on the NOAA site evaluation list for consideration as a marine sanctuary. This does not confer any specific statutory protection on the area, but merely indicates that the site is included within a pool of 39 nationwide sites from which individual sites can be chosen for study when funds become available. Most of Fisherman's Island, located on the north side of the Bay mouth, is a National Wildlife Refuge known for having a wide variety of bird life which are particularly abundant during seasonal migratory periods. The Back Bay National Wildlife Refuge and False Cape State Park are located to the south of the study area.

Summary

The dominant factor influencing the biological make-up of the study area is its position adjacent to the mouth of the Chesapeake Bay. The Bay, which is the largest estuary on the east coast, exerts a tremendous influence. As the Bay plume sweeps over this portion of the inner continental shelf, it contributes nutrients and other materials, resulting in enhanced primary production compared to adjacent shelf waters. A critical period in the life cycle of the Chesapeake Bay blue crab occurs in this area when, after spawning in the lower Bay, the crab larvae enter the surface waters and are transported out of the Bay. After undergoing larval development in shelf waters, the post-larval crabs must pass through the area on their way back into the Bay.

The proximity of the Bay mouth also dictates that there will be extensive fish migrations through the study area. This would also include the vulnerable larval and juvenile stages of many ocean spawning species passing through the area on their way to estuarine nursery grounds.

Detailed information on the nearshore spawning and nursery areas and migratory routes of most of the finfish species and blue crabs is not available. Since the study area comprises one of the most biologically unique and important areas along the Atlantic Coast, and is vital to the maintenance of the Chesapeake Bay resources, it is imperative that any proposal to modify this area must accurately define the impacts of such a project. Unless it can be reasonably determined that there will be no adverse impacts associated with dredged material disposal, this area should not be used for a permanent, long-term disposal site.

References

- Byles, R. VIMS, personal communication, Jan. 1984.
- Grant, G. C. 1979. Middle Atlantic Bight zooplankton: second year results and a discussion of the two-year BLM-VIMS survey. Special Report in Applied Marine Science and Ocean Engineering No. 192, Virginia Institute of Marine Science. Gloucester Point, VA.
- Grant, G. C. and J. E. Olney, 1979. Lower Bay zooplankton monitoring program: an introduction to the program and results of the initial survey of March 1978. Special Scientific Report No. 93, Virginia Institute of Marine Science. Gloucester Point, VA.
- Grant, G. C. and J. E. Olney, 1983. Lower Bay zooplankton monitoring program: the August 1978 survey. Special Scientific Report No. 115, Virginia Institute of Marine Science. Gloucester Point, VA.
- Grosslein, D. M. and T. R. Azarowitz, 1982. Fish distribution. MESA New York Bight Atlas Monograph 15. New York Sea Grant Institute.
- Hydroscience, Inc. 1974. Water quality studies for marine disposal systems, Atlantic treatment plant, Virginia Beach, Virginia.
- Johnson, D. R., Hester, B. S. and J. R. McConaugha, 1983. Studies of wind mechanisms influencing the recruitment of blue crabs in the Middle Atlantic Bight. Technical report No. 83-5, Old Dominion University. Norfolk, VA.
- Marshall, A. R. and J. A. Lucy, 1981. Virginia's charter and head boat fishery, analysis of catch and socioeconomic impacts. Special Report In Applied Science and Ocean Engineering No. 253. Virginia Sea Grant Program, Virginia Institute of Marine Science. Gloucester Point, VA.
- Marshall, H. G. 1981. Phytoplankton assemblages within the Chesapeake Bay plume and adjacent waters of the continental shelf. In: Chesapeake Bay Plume Study, Proceedings of a symposium held in Williamsburg, VA, Jan. 21-23, 1981. NASA Conf. Publ. 2188.
- McConaugha, J. R.; Johnson, D. F.; Provenzano, A. J. and R. C. Maris, 1983. Seasonal distribution of larvae of Callinectes sapidus (Crustacea : Decapoda) in the waters adjacent to Chesapeake Bay. J. of Crustacean Biology, Vol. 3(4):582-591.
- Mercer, L. P. 1983. A biological and fisheries profile of weakfish, Cynoscion regalis. Special Scientific Report No. 39, North Carolina Dept. of Natural Resources and Community Development.
- Musick, J. A. 1972. "Fishes of Chesapeake Bay and the adjacent coastal plain," In: A Check List of the Biota of Lower Chesapeake Bay. Marvin L. Wass (ed.), Virginia Inst. Mar. Sci., Spec. Sci. Rept. No. 65, pp. 175-212. Gloucester Point, VA.

National Marine Fisheries Service, Northeast Fisheries Center, unpublished data.

Olney, J. E. 1983. Eggs and early larvae of the Bay anchovy, Anchoa mitchilli, and the weakfish, Cynoscion regalis, in the lower Chesapeake Bay with notes on associated ichthyoplankton. *Estuaries*, Vol. 6(1):20-35.

Provenzano, A. J.; McConaughy, J. R.; Philips, K. B.; Johnson, D. F. and J. Clark. 1983. Vertical distribution of first stage larvae of the blue crab, Callinectes sapidus, at the mouth of Chesapeake Bay. *Estuarine, Coastal and Shelf Science*, 16:489-499.

Robertson, C. N. and J. P. Thomas. 1981. Total plankton respiration in the Chesapeake Bay plume. In: Chesapeake Bay Plume Study, Proceedings of a symposium held in Williamsburg, Virginia Jan. 21-23, 1981. NASA Conf. Publ. 2188.

Smyth, P. O. 1980. Callinectes (Decapoda : Portunidae) larvae in the Middle Atlantic Bight, 1975-77. *Fishery Bulletin* 78(2):251-265.

U.S. Dept. of Interior, 1970. National Estuary Study. Vol. 3.

U.S. Fish and Wildlife Service, 1981. Pelagic resources of Rudee Inlet and vicinity. Report submitted to the Norfolk District, Corps of Engineers. Gloucester Point, VA.

U.S. Fish and Wildlife Service, 1982. Resources and impacts associated with potential sand sources in the Atlantic Ocean. A planning aid report submitted to the Norfolk District, Corps of Engineers, for the Virginia Beach Erosion and Hurricane Protection Study. Gloucester Point, VA.




Fig. 1. Location of study area.

Fig. 2. Stations locations for the phytoplankton sampling of Marshall (1981).




Fig. 3. Phytoplankton cell concentrations from Marshall (1981) within the Chesapeake Bay plum and adjacent shelf waters for March, June, and October 1980.





Fig. 4. Station locations for zooplankton collections.



sampled by USEWS (1982).

CHESAPEAKE BAY

78° 00'

SHOAL
DEEP

CHESAPEAKE BAY ENTRANCE

CAPE HENRY

CHESAPEAKE BAY ENTRANCE CHANNEL

STUDY
AREA

30'

36'

VIRGINIA BEACH

RUDEE INLET




Fig. 6. Locations of benthic stations sampled by Hydrosience, Inc. (1974).




Table 1. Phytoplankton assemblages found by Marshall (1981) within the Chesapeake Bay plume and adjacent shelf waters for March, June, and October 1980. Numerical dominance is indicated for each collection period.

	<u>Bay Entrance - Plume</u>	<u>Shelf</u>
<u>March</u>	<u>Asterionella glacialis</u> <u>Cyclotella sp.</u> <u>Guinardia flaccida</u> <u>Leptocylindrus danicus</u> <u>Leptocylindrus minimus</u> <u>Nitzschia pungens</u> <u>Paralia sulcata</u> <u>Rhizosolenia delicatula</u> <u>Rhizosolenia fragilissima</u> * <u>Skeletonema costatum</u> <u>Thalassiosira nordenskioldi</u> <u>Gomphosphaeria aponina</u> <u>Nostoc commune</u> * <u>Prorocentrum minimum</u> *Green cells <3 microns *Green cells 3-5 microns	<u>Bacteriastrium hyalinum</u> <u>Chaetoceros costatum</u> <u>Nitzschia longissima</u> <u>Rhizosolenia delicatula</u> <u>Thalassiosira nordenskioldii</u> <u>Thalassiosira rotula</u> * <u>Emiliana huxleyi</u> Green cells 5-10 microns
<u>June</u>	<u>Chaetoceros spp.</u> <u>Cylindrotheca closterium</u> <u>Leptocylindrus danicus</u> <u>Nitzschia pungens</u> <u>Rhizosolenia delicatula</u> * <u>Skeletonema costatum</u> Green cells 3-5 microns	<u>Rhizosolenia alata</u> * <u>Emiliana huxleyi</u> <u>Pontosphaera sp.</u> <u>Rhabdosphaera sp.</u> <u>Syracosphaera pulchra</u>
<u>October</u>	* <u>Asterionella glacialis</u> <u>Cerataulina pelagica</u> <u>Cylindrotheca closterium</u> <u>Lauderia borealis</u> <u>Leptocylindrus danicus</u> <u>Nitzschia pungens</u> <u>Rhizosolenia delicatula</u> * <u>Skeletonema costatum</u> * <u>Emiliana huxleyi</u> *Green cells <3 microns *Green cells 3-5 microns <u>Anacystis sp.</u> <u>Cryptomonas sp.</u>	<u>Nitzschia pungens</u> <u>Rhizosolenia delicatula</u> <u>Rhizosolenia fragilissima</u> <u>Skeletonema costatum</u> * <u>Emiliana huxleyi</u> *Green cells <3 microns *Green cells 3-5 microns Mixed phytoflagellates

*Dominant phytoplankters

Table 2. Species and abundance of fish, decapods, and molluscs collected by U.S. Fish and Wildlife Service (1981) using oblique daylight plankton tows at two stations near the mouth of Rudee Inlet (Fig. 4).

	Number/100m ²					
	<u>May</u>	<u>June</u>	<u>July</u>	<u>Aug</u>	<u>Oct</u>	<u>Jan</u>
FISH						
<i>Ammodytes</i> sp.						2
<i>Anchoa hepsetus</i> eggs				42		
<i>Anchoa mitchilli</i> eggs	11	1549	199	23		
<i>Anchoa mitchilli</i>		12	593			
<i>Anchoa</i> sp.				3261		
<i>Cynoscion regalis</i>				204		
<i>Etropus microstomus</i>			3			
<i>Gobiesox strumosus</i>				2		
<i>Gobiosoma</i> sp.				4		
<i>Leiostomus xanthurus</i>	2					
<i>Membras martinica</i>		3				
<i>Morone americanus</i>		2		31		
<i>Peprilus</i> sp. eggs			77	106		
<i>Pogonias cromis</i> (?)	30					
<i>Tautoga omitis</i> eggs	11					
<i>Trinectes maculatus</i> eggs		16		68		
<i>Trinectes maculatus</i>			8			
unid. Sciaenidae			10			
unid. Sciaenidae eggs	6	199	63	142		
DECAPODS						
<i>Callinassa atlantica</i>			55	837	2	
<i>Callinectes sapidus</i>		289	15	1973		
<i>Cancer irrorata</i>	4	4				
<i>Crangon septemspinosa</i>	15					10
<i>Emerita talpoida</i>		5	6	5	1	
<i>Lucifer faxoni</i>				20	52	
<i>Naushonia crangonoides</i>			5			
<i>Neopanope sayi</i>		20	35	46		
<i>Ogyrides limicola</i>					20	
<i>Ovalipes ocellatus</i>					2	
<i>Pagurus longicarpus</i>			3	33		
<i>Palaemonetes</i> sp.	1	3				
<i>Pinnixa</i> sp.		3	185	134	2	
<i>Pinnotheres maculatus</i>			6			
<i>P. ostreum</i>				5	38	
<i>Rhithropanopeus harrissi</i>				284		
<i>Sesarma</i> sp.			18	155		
<i>Uca</i> sp.		15	27	10		
<i>Upogebia affinis</i>		4	130	3476	5	

Table 2. (Cont.)

	Number/100m ²					
	<u>May</u>	<u>June</u>	<u>July</u>	<u>Aug</u>	<u>Oct</u>	<u>Jan</u>
MOLLUSCS						
Acteon punctostriatus		28	1602	2		
Cerithiopsis (?)					6	
Littorina sp.	6					
Mulinia lateralis				7		
Mytilus edulis	1					
Nassarius vibex			3	2		
Petricola pholadiformis				5		
Solen viridis				7		
Spiratella trochiformis				2		
unid. Turridae			2			

Table 3. List of zooplankton identified by Grant and Olney (1979, 1983) from bongo (oblique) and neuston (surface) tows conducted during day and night periods at stations in the vicinity of the Bay Bridge Tunnel (Fig. 4).

	<u>March '79</u>	<u>August '79</u>
COELENTERATA		
Bougainvillea sp.		x
Chrysaoura quinquecirrha		x
Cunina octonoria		x
Cyanea capillata	x	
Hydrocodon prolifer	x	
Liriope tetraphylla		x
Muggiaea kochei		x
Nemopsis bachei		x
Obelia sp.	x	x
Ruthkea octopunctata	x	
CTENOPHORA		
Pleurobrachia pileus	x	
NEMATODA		
unid. nemotodes	x	
POLYCHAETA		
Autolytus sp.	x	
Glycera dibranchiata	x	
Gyptis vittata		x
Harmathoe sp.	x	x
Hesionidae		x
Nereis succinea		x
Paranaitis speciosa	x	
Paraprionospio pinnata		x
Phyllodocidae		x
Polydora sp.	x	x
Polynoidae		x
Pseudeurythoe ambigua		x
Spionidae larvae	x	x
Syllidae		x
Terebellidae		x
Tomopteris helgolandica	x	x
MOLLUSCA		
Cerithiopsis sp.		x
Crassostrea virginica		x
Crepidula sp.		x
Ensis directus	x	
Epitonium sp.		x
Laevicardium mortoni	x	
Lilliguncula brevis		x

Table 3. (Cont.)

	<u>March '79</u>	<u>August '79</u>
MOLLUSCA (cont.)		
<i>Limacina retroversa</i>	x	
<i>Littorina irrorata</i>	x	x
<i>Loligo pealeii</i>		x
<i>Lyonsia hyalina</i>	x	
<i>Macoma balthica</i>	x	
<i>Mulina lateralis</i>	x	x
<i>Mytilus edulis</i>	x	
<i>Nassarius obsoletus</i>		x
<i>N. vibex</i>		x
Naticidae		x
Pholadidae		x
<i>Tellina agilis</i>	x	x
MEROSTOMATA		
<i>Limulus polyphemus</i>		x
MYSIPACEA		
<i>Metamysidopsis mexicana</i>		x
<i>Mysidopsis bigelowi</i>	x	x
<i>Neomysis americana</i>	x	x
CUMACEA		
<i>Leptocuma minor</i>	x	
<i>Mancocuma</i> sp.	x	x
<i>Oxyurostylis smithi</i>	x	x
ISAPODA		
<i>Aegathea medialis</i>		x
<i>A. oculata</i>		x
<i>Edotea triloba</i>	x	
AMPHIPODA		
<i>Ampelisca abdita</i>	x	x
<i>A. agassizi</i>		x
<i>A. vadorum</i>		x
<i>Ampithoe longimana</i>		x
<i>Atylus minidoi</i>		x
<i>Batea catharinensis</i>		x
<i>Caprella equilibra</i>	x	
<i>Corophium insidiosum</i>	x	
<i>C. lacustre</i>		x
<i>Gammarus mucronatus</i>	x	x
<i>Idunella</i> sp.		x
<i>Jassa falacata</i>	x	x
<i>Lestrigonus bengalensis</i>		x
<i>Melita</i> sp.		x
<i>Microprotopus raneyi</i>	x	x
<i>Monoculodes edwardsi</i>	x	

Table 3. (Cont.)

	<u>March '79</u>	<u>August '79</u>
AMPHIPODA (cont.)		
Paraplenes aestuarius		x
Phoxocephalus spinosus		x
Unciola irrorata	x	
CLADOCERA		
Evadne nordmani	x	x
Penilia avirostris		x
Podon polyphemoides	x	
COPEPODA		
Acartia clausi	x	
A. tonsa	x	x
Caligus chelifer		x
Centropages hamatus	x	x
C. typicus	x	x
C. velificatus		x
Corycaeus speciosus		x
Eucalanus crassus		x
E. pileatus	x	x
Eurytemora affinis	x	
Hemicyclops sp.	x	
Labidocera aestiva		x
Oithona sp.	x	x
Paracalanus crassirostris	x	x
P. indicus		x
P. quosimoda		x
Pseudocalanus sp.	x	
Pseudodiaptomus coronatus		x
Temora longicornis	x	
CIRRIPEDIA		
Balanus balanoides (?)	x	
Lepus sp.	x	
unid. barnacle larvae		x
STOMATOPODA		
Squilla empusa		x
DECAPODA		
Alpheus normanni		x
Callinassa atlantica		x
C. biformis		x
Callinectes sp.		x
Crangon septemspinosa	x	x
Euceramus praelongus		x
Hexapanopeus angustifons		x
Libinia emarginata		x
L. dubia		x

Table 3. (Cont.)

	<u>March '79</u>	<u>August '79</u>
DECAPODA (cont.)		
Lucifer faxoni		x
Naushonia crangonoides		x
Neopanope sayi		x
Ogyrides limicola		x
Ovalipes ocellatus		x
O. stephensoni		x
Pagurus longicarpus		x
Palaemonetes sp.		x
Panopeus herbstii		x
Pinnixa chaetoptera		x
P. cylindrica		x
P. sayana		x
Pinnotheres maculatus		x
P. ostreum		x
Portunus sp.		
Uca sp.		x
Upogebia affinis		x
PHORONIDA		
unid. phoronid larvae	x	
CHAETOGNATHA		
Sagitta elegans	x	
S. enflata		x
S. hispida		x
S. tenuis		x
TUNICATA		
Oikopleura sp.		x
PISCES		
Ammodytes hexapterus	x	
Anchoa hepsetus		x
A. mitchilli	x	
Anguilla rostrata	x	
Atherinidae	x	x
Centropristis striata		x
Cynoscion nebulosus		x
C. regalis		x
Etropis microstomus		x
Gadus morhua	x	
Gobiosoma sp.		x
Hypsoblennius hentzi		x
Menticirrhus sp.		x
Peprilus paru		x
P. trincanthus		x
Prionotus sp.		x

Table 3. (Cont.)

	<u>March '79</u>	<u>August '79</u>
PISCES (cont.)		
Pseudopleuronectes americanus	x	
Rissola marginata		x
Symphurus plagiusa		x
Syngnathus fuscus	x	
Trinectes maculatus		x
unid. ophidiid		x
unid. pleuronectiform		x

Table 4. List of zooplankton identified by Grant et al (1979) from oblique and surface tows conducted during day and night periods at a coastal station off the eastern shore of Virginia (Fig. 4).

	<u>Fall '76</u>	<u>Winter '77</u>	<u>Spring '77</u>	<u>Summer '77</u>
CNIDARIA				
Abylopsis eschscholtzii	x			
Abylopsis tetragona	x			
Aequorea sp.				x
Agalma elegans	x			
Bassia bassensis	x			
Catablema vesicarium				x
Diphyes chamissoni	x			
Diphyes dispar	x			
Diphyopsis campanulifer	x			
Liriope sp.	x			x
Liriope tetraphylla	x			
Muggiaea kochei	x			
Obelia sp.				x
Pelagia noctiluca	x			
Rhacostoma atlanticum				x
unid. anthozoans	x			
unid. hydrozoans				x
TURBELLARIA				
unid. flatworms	x			x
ANNELIDA				
Tomopteris helgolandica				x
unid. polychaetes	x	x	x	x
MOLLUSCA				
Atlanta peroni	x			x
Cavolina longirostris	x			
Cerastoderma pinnulatum		x		x
Creseis virgula	x			x
Ensis directus		x	x	
Hyalocyclis striata				x
Lima tenera	x			
Limacina retroversa			x	
Limacina trochiformis	x			x
Loligo pealii	x		x	x
Lunatia triseriata			x	
Macoma balthica		x	x	
Paedoclione doliiformis			x	x
Rossia tenera				x
Spisula solidissima	x	x	x	
unid. gastropods				x

Table 4. (Cont.)

	<u>Fall '76</u>	<u>Winter '77</u>	<u>Spring '77</u>	<u>Summer '77</u>
CLADOCERA				
Evadne nordmanni			x	x
E. spinifera				x
E. tergestina	x			x
Penilia avirostris	x			x
Podon sp.				x
Podon leuckartii			x	
OSTRACODA				
Conchoecia bispinosa		x		
Euconchoecia chierchine	x			
COPEPODA				
Acartia sp.				x
A. clausi				x
A. danae	x			
A. tonsa				x
Anomalocera sp.			x	
A. ornata	x		x	
A. patersonii			x	
Calanus finmarchicus		x	x	x
Candacia armata	x			x
Caligus chelifer				x
Centropages furcatus	x			x
C. hamatus		x	x	x
C. typicus	x	x	x	x
C. violacens	x			
Clausocalanus arevicornis		x		
Copilia mirabilis				x
Corycaeus sp.	x	x	x	x
C. lautus				x
C. speciosus	x			x
Eucalanus sp.	x	x		
E. crassus	x			x
E. pileatus	x			x
Euchaeta marina		x		
Eurytemora sp.				x
E. americana	x			
Labidocera sp.	x			x
L. aestiva	x			x
Mecynocera clausi	x			
Metridia leicens	x	x		
Nannocalanus minor	x			x
Oithona sp.	x		x	x
Oncaea conifera	x			
O. mediterranea	x			
O. venusta	x			
Paracalanus sp.	x	x	x	

Table 4. (Cont.)

	<u>Fall '76</u>	<u>Winter '77</u>	<u>Spring '77</u>	<u>Summer '77</u>
COPEPODA (cont.)				
<i>P. parvus</i>	x			
<i>Pleuromamma gracilis</i>	x	x		
<i>Pontella</i> sp.	x			x
<i>P. meadii</i>	x			x
<i>Pontellopsis villosa</i>	x			
<i>Pseudocalanus</i> sp.			x	x
<i>Rhincalanus nasutus</i>	x	x		
<i>Sapphirina</i> sp.	x		x	
<i>S. nigromaculata</i>	x			x
<i>Scolecithrix danae</i>	x			
<i>Temora</i> sp.			x	
<i>T. longicornis</i>			x	x
<i>T. stylifera</i>	x			x
<i>T. turbinata</i>	x		x	
<i>Tortanus discaudatus</i>			x	x
<i>Undinula vulgaris</i>	x			
CIRRIPIEDIA				
<i>Chthamalus fragilis</i>		x		x
unid. barnacle larvae			x	x
STOMATOPODA				
unid. stomatopod larvae	x		x	x
MYSIDACEA				
<i>Bowmaniella</i> sp.	x			x
<i>Heteromysis formosa</i>		x	x	
<i>Mysidopsis bigelowi</i>	x	x	x	
<i>Neomysis americana</i>		x		
<i>Promysis atlantica</i>	x			
CUMACEA				
<i>Cyclaspis</i> sp.			x	x
<i>Leptocuma minor</i>		x	x	x
<i>Leucon americanus</i>	x		x	x
<i>Oxyurostylis smithi</i>	x		x	x
ISOPODA				
<i>Chiridotea tuftsi</i>			x	x
<i>Edotea triloba</i>				x
<i>Idotea balthica</i>				x
<i>I. metallica</i>	x		x	

Table 4. (Cont.)

	<u>Fall '76</u>	<u>Winter '77</u>	<u>Spring '77</u>	<u>Summer '77</u>
AMPHIPODA				
Acanthostepheia sp.			x	
Ampelisca agassizi			x	
A. verrilli				x
Ampithoe longimana				x
Argissa hamatipes			x	x
Byblis serrata			x	
Corophium acheruscium			x	
Gammarus sp.			x	
Lestrigonus sp.				x
L. bengalensis	x			x
Lycaea sp.				x
Lycaeopsis neglecta	x			
Microprotopus raneyi				x
Monoculodes sp.		x	x	
Parametopella cypris			x	
Parathemisto gandichandii	x	x		
Protohaustorius wigleyi			x	
Syncholidium americanum				x
Themistella fusca	x			x
Trichophoxus epistomus				x
Unciola irrorata				x
unid. gammarids		x		
unid. caprellids	x			
unid. hyperiids				x
EUPHAUSIACEA				
Euphausia sp.	x			
Thysanoessa sp.	x	x		
unid. euphausiids	x	x		
DECAPODA				
Arenaens sp.				x
Callianassa sp.	x			
Callinectes sp.	x			x
Cancer sp.	x		x	
Crangon septemspinosa	x	x	x	x
Dichelopandalus leptocerus			x	
Dromida antillensis	x			
Emerita sp.				x
Hexapanopeus augustifrons				x
Homarus americanus			x	
Leptochela sp.	x			
Libinia sp.	x			
Lucifer faxoni	x			x
Naushonia crangonoides				x
Ocypode quadrata				x
Ovalipes sp.	x			

Table 4. (Cont.)

	<u>Fall '76</u>	<u>Winter '77</u>	<u>Spring '77</u>	<u>Summer '77</u>
DECAPODA (cont.)				
Portunus sp.	x			x
Processa sp.	x			
Sergestes sp.	x			
Uca sp.				x
Upogebia affinis				x
unid. calappids			x	
unid. hippolytids				x
unid. leucosiids	x			
unid. pagurids	x			x
unid. palaemonids				x
unid. penaeids				x
unid. xanthids	x			
ECHINODERMATA				
unid. asteroids	x			
unid. ophinroids	x			
CHAETOGNATHA				
Sagitta elegans		x	x	x
S. enflata	x			x
S. helenae	x			
S. hispida	x			x
S. minima	x			
S. serratodentata	x			
S. tasnanica	x	x		
S. tenuis	x			x
TUNICATA				
Doliolum sp.				x
D. nationalis	x			x
Oikopleura sp.	x			x
PISCES				
Ammodytes sp.		x	x	
Anchoa sp.	x			x
Astroscopus guttatus	x			x
Bothus sp.	x			
Citharichthys arctifrons				x
Conger oceanicus			x	
Cynoscion regalis	x			
Enchelyopus cimbrius			x	
Etropus microstomus				x
Gasterosteus aculeatus			x	
Gobiosoma ginsburgi	x		x	x
Hypsoblennius hentzi				x
Limanda ferruginea			x	
Lophius americanus			x	

Table 4. (Cont.)

	<u>Fall '76</u>	<u>Winter '77</u>	<u>Spring '77</u>	<u>Summer '77</u>
PISCES (cont.)				
Menidia minidia		x		
Merluccius sp.	x			
Paralichthys dentatus	x			x
Peprilus triacanthus				x
Pomatomus saltatrix			x	x
Scophthalmus aquosus	x		x	x
Symphurus sp.				x
Tylosurus acus				x
Urophycis sp.	x		x	
unid. engraulids	x			
unid. ophidiids				x
unid. scorpaenids				x

Table 5. Macrobenthic species sampled with Shipek grab by FWS (1982) and the stations (see Fig. 4) at which they occurred.

<u>Taxa</u>	<u>Common Name</u>	<u>Stations Where Collected</u>							
CNIDARIA									
Anthozoa (unidentified)	anemone	1	2				6		
PLATYHELMINTHES									
Turbellaria (unidentified)	flatworm								8
NEMERTEA									
unidentified Nemertean	nemertean worm	1	2	3	4	5	6	7	8
PHORONIDA									
<u>Phoronis</u> sp.	phoronid worm		2	3	4			7	8
MOLLUSCA									
Gastropoda									
<u>Crepidula fornicata</u>	slipper shell					5			8
<u>Mitrella lunata</u>	lunar mitrella	1	2		4				
<u>Nassarius trivittatus</u>	mud snail	1		3					8
<u>Natica pusilla</u>	moon shell			3					
<u>Retusa candea</u>	lathe shell		2						
<u>Turbonilla interrupta</u>	pyramid shell		2		4	5		7	
Pelecypoda									
<u>Anadara transversa</u>	transverse ark shell	1							
<u>Ensis directus</u>	razor clam	1	2	3	4	5	6	7	8
<u>Gemma gemma</u>	gem shell								8
<u>Lyonsia hyalina</u>	transparent lyonsia							7	8
<u>Mulinia lateralis</u>	little surf clam		2						
<u>Mytilus edulis</u>	blue mussel	1	2	3	4	5	6		
<u>Nucula proxima</u>	nut shell		2			5	6		8
<u>Siliqua costata</u>	ribbed pod shell								7
<u>Spisula solidissima</u>	surf clam	1			4	5	6	7	8
<u>Tellina agilis</u>	tellin shell	1	2	3	4	5	6	7	8
<u>Yoldia limatula</u>			2	3	4				
ANNELIDA									
Polychaeta									
<u>Ampharete acutifrons</u>		1	2	3	4	5	6	7	8
<u>Amparete americana</u>			2	3		5	6		8
Ampharetidae sp. (juvenile)			1	2	3	4	5		
<u>Amastigos caperatus</u>		1	2	3	4	5	6	7	8
<u>Ancistrosyllis hartmanae</u>		1				5	6		8
<u>Aproprionospio pygmaea</u>									7 8
<u>Aricidia catherinae</u>		1				5	6	7	8
<u>Aricidia wassi</u>						5		7	
<u>Asabellides oculata</u>							6		
<u>Capitella capitata</u>		1	2	3	4				
Capitella sp. (juvenile)									
Cirratulidae spp.		1	2	3	4	5	6	7	8

Table 5. (Cont.)

<u>Taxa</u>	<u>Common Name</u>	<u>Stations Where Collected</u>							
ANNELIDA, cont'd.									
Polychaeta									
<u>Cirrophorus furcatus</u>		1	2	4	5	6		8	
<u>Clymenella torquata</u>	bamboo worm			3	4			7	8
<u>Diopatra cuprea</u>		1	2						
<u>Dorvilleidae sp. (juvenile)</u>							6	7	
<u>Eteone heteropoda</u>		1	2				6		
<u>Eteone lactea</u>		1				5	6	7	8
<u>Eumida sanguinea</u>				4					
<u>Glycera americana</u>	bloodworm	1	2	3	4	5	6	7	8
<u>Glycera dibranchiata</u>	bloodworm	1		3	4	5	6	7	8
<u>Glycera sp. (juvenile)</u>	bloodworm	1	2	3	4		6	7	
<u>Goniadella gracilis</u>			2	3					
<u>Harmothoe extenuata</u>		1	2	3	4				8
<u>Hemipodus roseus</u>									8
<u>Linopherus ambigua</u>							6		
<u>Lumbrineris tenuis</u>						5	6		8
<u>Macroclymene zonalis</u>				4	5	6			8
<u>Magelona sp.</u>		1	2	3	4	5	6	7	8
<u>Maldanidae sp. (juvenile)</u>		1		4				7	8
<u>Mediomastus ambiseta</u>		1	2	3	4	5	6	7	8
<u>Microphthalmus sczelkowi</u>							6		
<u>Nephtys picta</u>			2	3	4	5	6	7	
<u>Nereidae sp. (juvenile)</u>	clam worm	1							
<u>Nereis succinea</u>	clam worm	1							
<u>Notomastus hemipodus</u>				4				7	8
<u>Ophelia sp.</u>						6			
<u>Owenia fusiformis</u>									8
<u>Paleonotus heteroseta</u>				3					
<u>Paranaitis speciosa</u>				3	4				8
<u>Phyllodoce arenae</u>	paddle worm		2	3	4	5	6	7	8
<u>Phyllodoce sp. (juvenile)</u>	paddle worm							7	
<u>Phyllodocidae sp. (juvenile)</u>	paddle worm			3					
<u>Pista cristata</u>			2						
<u>Polydora websteri</u>			2		4				8
<u>Polydora sp. (juvenile)</u>		1	2	3	4	5		7	
<u>Polygordius sp.</u>		1	2	3	4	5	6	7	8
<u>Protodorvillae kefersteini</u>			1			4	5	6	
<u>Sabellaria vulgaris</u>	sand builder worm	1							
<u>Scalibregma inflatum</u>							6		8
<u>Schistomeringos caeca</u>		1		4	5	6	7	8	
<u>Schistomeringos rudolfi</u>		1			5	6			
<u>Scolecopsis sp.</u>				4				7	
<u>Scoloplos robustus</u>		1		4				7	8
<u>Scoloplos sp. (juvenile)</u>		1				6			
<u>Sigambra tentaculata</u>			2	3			6		
<u>Spio setosa</u>		1	2	3	4	5	6	7	8

Table 5. (Cont.)

<u>Taxa</u>	<u>Common Name</u>	<u>Stations Where Collected</u>							
ANNELIDA, Cont'd.									
Polychaeta									
<u>Spiochaetopterus oculatus</u>			2						
<u>Spiophanes bombyx</u>		1	2	3	4	5	6	7	8
<u>Streblospio benedicti</u>			2	3	4	5		7	8
Syllidae spp.		1	2		4	5	6		8
Oligochaeta									
Oligochaeta spp.	oligochaete worms	1	2	3	4	5	6	7	8
ARTHROPODA									
Crustacea									
Cumacea									
<u>Oxyurostylis smithi</u>						5	6	7	8
Isopoda									
<u>Edotea</u> sp.		1			4		6	7	8
Amphipoda									
<u>Ampelisca verrilli</u>								7	
<u>Corophium</u> spp.		1	2	3					
<u>Synchelidium americanum</u>		1							
<u>Unciola irrorata</u>			2	3	4	5	6	7	8
<u>Unciola serrata</u>							6		
<u>Unciola</u> sp.				3		5			
Decapoda									
<u>Pagurus</u> sp.	hermit crab					5			
<u>Pinnixa</u> sp.	commensal crab		2			5		7	
ECHINODERMATA									
Echinoidea sp.	sand dollar	1	2			5	6	7	8
CEPHALOCHORDATA									
<u>Branchiostoma virginiae</u>	lancelet	1			4	5	6	7	8

Table 6. Epibenthic species sampled with otter trawl by FWS (1982) and the stations (see Fig. 4) at which they occurred.

<u>Taxa</u>	<u>Common Name</u>	<u>Stations Where Collected</u>		
MOLLUSCA				
Gastropoda				
<u>Busycon canaliculatum</u>	channeled whelk		T2	
<u>Crepidula fornicata</u>	slipper shell	T1		
<u>Nassarius trivittatus</u>	mud snail		T2	
Pelecypoda				
<u>Mytilus edulis</u>	blue mussel		T2	T3
Cephalopoda				
<u>Loligo pealei</u>	common squid	T1		T3
<u>Lolliguncula brevis</u>	brief squid		T2	
ARTHROPODA				
Merostomata				
<u>Limulus polyphemus</u>	horseshoe crab		T2	
Crustacea				
<u>Callinectes sapidus</u>	blue crab			T3
<u>Cancer irroratus</u>	rock crab	T1	T2	T3
<u>Libinia emarginata</u>	spider crab		T2	
<u>Pagurus longicarpus</u>	hermit crab	T1	T2	T3
<u>Pagurus pollicaris</u>	hermit crab	T1		
<u>Panopeus herbstii</u>	mud crab			T3
ECHINODERMATA				
Echinoidea				
<u>Asterias forbesi</u>	common starfish		T2	
CHORDATA				
Pisces				
Carangidae sp.	scad			T3
<u>Centropristes striata</u>	black sea bass		T2	T3
<u>Cottus sp.</u>	sculpin	T1	T2	T3
<u>Etropus microstomus</u>	smallmouth flounder	T1		T3
<u>Prionotus carolinus</u>	searobin	T1		T3
<u>Raja eglanteria</u>	clearnose skate	T1	T2	T3
<u>Scophthalmus aquosus</u>	windowpane		T2	
<u>Stenotomus chrysops</u>	southern scup	T1	T2	
<u>Urophycis regia</u>	spotted hake	T1	T2	T3

Table 7. List of benthic species reported by Hydrosience, Inc. (1974) and the stations at which they occurred (see Fig. 5).

TAXA	Stations					
	<u>T3-1</u>	<u>T3-3</u>	<u>T5-1</u>	<u>T5-3</u>	<u>T7-1</u>	<u>T7-3</u>
CNIDARIA						
Halcampoides sp.	x					
Hydractinia sp.				x		
Lovenella gracilis		x				
NEMERTEA						
unid. nemertean			x	x		x
NEMATODA						
unid. nematode		x	x	x		
BRYZOA						
Electra sp.				x		
POLYCHAETA						
Ampharete acutifrons				x	x	
Amphicteis gunneri						x
Aricidea cerruti	x	x			x	x
Branchiostoma caribeum					x	x
Brania dispar						x
Ceratonereis irritabilis				x		
Diopatra cuprea				x		
Drilonereis filum	x	x	x	x		
D. magna					x	
Exogone setosa						x
Glycera dibranchiata	x	x	x	x	x	
Heteromastus filiformis				x		
Lepidonotus sublevis				x		
Loimia medusa				x		
Magelona rosea	x		x			
Nephtys magellanica	x					
N. picta		x			x	
Nereis succinea				x		
Notomostus sp.	x	x	x		x	x
Odontosyllis fulgurans					x	x
Owenia fusiformis	x		x			x
Paraonis fulgons		x				
Phyllodoce mucosa				x		
Phylo felix					x	
Polycirrus eximius				x	x	
Polydora sp.					x	
P. socialis				x		
Polygordius sp.		x		x	x	x
Prionospio cirrifera				x		
P. dayi			x			

Table 7. (Cont.)

TAXA	Stations					
	<u>T3-1</u>	<u>T3-3</u>	<u>T5-1</u>	<u>T5-3</u>	<u>T7-1</u>	<u>T7-3</u>
POLYCHAETA (cont.)						
<i>Sabellaria vulgaris</i>		x		x	x	
<i>Scolecopides viridis</i>						x
<i>Scoloplos fragilis</i>	x					
<i>Sigalion arenicola</i>					x	x
<i>Spio setosa</i>					x	x
<i>Spiochaetopterus oculatus</i>				x		
<i>Spiophanes bombyx</i>	x	x	x			
<i>Tharyx marioni</i>			x	x	x	x
<i>Tharyx setigera</i>	x					
<i>Travisia parva</i>					x	x
OLIGOCHAETA						
unid. oligochaetes	x	x		x	x	
GASTROPODA						
<i>Acteocina canaliculata</i>			x			
<i>Clione</i> sp.		x			x	
<i>Crepidula convexa</i>		x		x	x	
<i>Cylichna alba</i>			x			
<i>Doridella obscura</i>				x		
<i>Mitrella lunata</i>				x		
<i>Polinices duplicatus</i>					x	
<i>Turbonilla interrupta</i>			x			
BIVALVIA						
<i>Abra aequalis</i>	x	x	x	x	x	x
<i>Amygdalum papyria</i>			x			
<i>Mercenaria</i> sp.	x					
<i>Mercenaria mercenaria</i>			x			
<i>Modiolus</i> sp.				x		
<i>Nucula proxima</i>		x		x	x	
<i>Pandora trilineata</i>		x			x	
<i>Spisula solidissima</i>		x			x	
<i>Tellina virsicolor</i>	x		x	x	x	x
CIRRIPEDIA						
unid. barnacles	x					
STOMATOPODA						
<i>Squilla empusa</i>					x	
CUMACEA						
<i>Oxyurostylis smithi</i>	x		x		x	
ISOPODA						
<i>Edotea triloba</i>			x			

Table 7. (Cont.)

TAXA	Stations					
	<u>T3-1</u>	<u>T3-3</u>	<u>T5-1</u>	<u>T5-3</u>	<u>T7-1</u>	<u>T7-3</u>
AMPHIPODA						
<i>Ampelisca verrilli</i>				x		
<i>Corophium acherusicum</i>				x		
<i>Erichthonius brasiliensis</i>				x		
<i>Photis</i> sp.				x		
<i>Protohaustorius wigleyi</i>						x
<i>Stenothoe minuta</i>				x		
<i>Trichophoxus epistomus</i>	x	x		x		
<i>Unciola irrorata</i>				x	x	
MYSIDACEA						
<i>Neomysis americanus</i>	x	x	x	x	x	
DECAPODA						
<i>Callinassa atlantica</i>					x	
<i>Crangon septemspinosus</i>	x			x		
<i>Euceramus praelongus</i>				x	x	
<i>Pagurus annulipes</i>	x		x	x		
<i>P. pollicaris</i>				x		
<i>Pinnixa chaetoptera</i>	x	x		x	x	
<i>Portunus sayi</i>				x		
<i>Pelia mutica</i>				x		
unid. xanthid		x		x		
ECHINODERMATA						
<i>Mellita quinquesperforata</i>						x

Table 3. List of species collected during NMFS groundfish surveys conducted along three depth transects between the mouth of Chesapeake Bay and the Virginia/North Carolina border. The list is based on four annual surveys from 1977 through 1980 with 3 cruises conducted each year, usually in late March, early August and late September.

<u>Species</u>	<u>Fathoms</u>		
	<u>0-5</u>	<u>5-10</u>	<u>10-15</u>
<i>Odontaspis taurus</i>			x
<i>Mustelus canis</i>	x	x	x
<i>Carcharhinus obscurus</i>		x	
<i>Rhizoprionodon terraenovae</i>	x	x	x
<i>Sphyrna lewini</i>	x		
<i>Squalus acanthias</i>	x	x	x
<i>Squatina dumerili</i>	x	x	x
<i>Raja eglanteria</i>	x		x
<i>R. erinacea</i>		x	
<i>Aetobatus narinari</i>	x		
<i>Dasyatis centroura</i>		x	
<i>D. sabina</i>	x		
<i>Gymnura altavela</i>	x	x	x
<i>G. micrura</i>	x		
<i>Myliobatis freminvilli</i>	x	x	x
<i>M. goodei</i>	x		
<i>Rhinoptera bonasus</i>	x		
<i>Accipenser oxyrhynchus</i>		x	
<i>Clupea harengus</i>	x	x	x
<i>Alosa aestivalis</i>	x	x	x
<i>A. pseudoharengus</i>	x	x	x
<i>A. sapidissima</i>	x	x	
<i>Opisthonema oglinum</i>	x	x	
<i>Brevoortia tyrannus</i>	x	x	
<i>Etrumens salina</i>		x	x
<i>Anchoa mitchilli</i>	x	x	x
<i>A. hepsetus</i>	x	x	
<i>A. lyolepis</i>		x	x
<i>Conger oceanicus</i>		x	
<i>Synodus foetens</i>	x	x	x
<i>Urophycis chuss</i>	x	x	x
<i>U. regia</i>	x	x	x
<i>Merluccius bilinearis</i>	x	x	x
<i>Paralichthys dentatus</i>	x	x	x
<i>Pseudopleuronectes americanus</i>	x		x
<i>Etropus microstomus</i>	x	x	x
<i>Citharichthys arctifrons</i>		x	x
<i>Scophthalmus aquosus</i>	x	x	x
<i>Trinectes maculatus</i>	x		
<i>Symphurus plagiusa</i>		x	x
<i>Syngnathus fuscus</i>		x	
<i>Menidia menidia</i>	x	x	x

Table 8. (Cont.)

<u>Species</u>	<u>Fathoms</u>		
	<u>0-5</u>	<u>5-10</u>	<u>10-15</u>
<i>Ammodytes americanus</i>	x	x	x
<i>Scomber scombrus</i>		x	x
<i>Trichiurus lepturus</i>		x	
<i>Peprilus alepidotus</i>	x	x	
<i>P. triacanthos</i>	x	x	x
<i>Selar crumenophthalmus</i>	x	x	
<i>Seriola zonata</i>	x		
<i>Caranx fusus</i>	x	x	
<i>C. bartholomaei</i>	x		
<i>Selene setapinnis</i>		x	
<i>Trachurus lathami</i>		x	x
<i>Pomatomus saltatrix</i>	x	x	x
<i>Centropristis striata</i>		x	x
<i>Stenotomus chrysops</i>	x	x	x
<i>S. caprinus</i>	x	x	x
<i>Leiostomus xanthurus</i>	x	x	x
<i>Micropogonias undulatus</i>	x	x	x
<i>Menticirrhus saxatilis</i>	x	x	
<i>M. americanus</i>	x	x	
<i>Cynoscion regalis</i>	x	x	x
<i>Prionotus evolans</i>	x	x	x
<i>P. tribulus</i>			x
<i>Tautoga onitis</i>			x
<i>Astroscopus guttatus</i>		x	
<i>Ophidium marginatum</i>	x	x	x
<i>Monacanthus hispidus</i>	x	x	
<i>Sphoeroides maculatus</i>	x	x	x
<i>S. trichocephalus</i>	x		
<i>Lophius americanus</i>	x		
<i>Decapterus punctatus</i>		x	x
<i>Macrozoarces americanus</i>			x
Total	53	58	45
All total	76 species		

DTIC

FILMED

4-86

END