

AD-A167 669


GENETIC AND MOLECULAR STUDIES OF THE PHLEBOTOMUS FEVER 1/1
GROUP OF VIRUSES(U) ALABAMA UNIV IN BIRMINGHAM
D H BISHOP MAR 85 DAMD17-84-C-4116

UNCLASSIFIED

F/G 6/13

NL





MICROCOPY

CHART

AD-A167 669

12

GENETIC AND MOLECULAR STUDIES OF THE PHLEBOTOMUS FEVER
GROUP OF VIRUSES

FINAL REPORT

DR. DAVID H.L. BISHOP

MARCH 1985

DTIC
ELECTE
MAY 16 1985
S D

Supported by

U.S. ARMY MEDICAL RESEARCH AND
DEVELOPMENT COMMAND
Fort Detrick, Frederick, Maryland 21701-5012

Contract No. DAMD17-84-C-4116

University of Alabama in Birmingham
Birmingham, Alabama 35294

DOD DISTRIBUTION STATEMENT

Approved for public release;
distribution unlimited

Findings in this report are not to be construed as an
official Department of the Army position unless so designated
by other authorized documents

DTIC FILE COPY

86 5 16 004

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS			
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited			
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			4. PERFORMING ORGANIZATION REPORT NUMBER(S)			
6a. NAME OF PERFORMING ORGANIZATION University of Alabama in Birmingham			6b. OFFICE SYMBOL (if applicable)	5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6c. ADDRESS (City, State, and ZIP Code) Birmingham, Alabama 35294			7a. NAME OF MONITORING ORGANIZATION			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION U.S. Army Medical Research and Development Command			8b. OFFICE SYMBOL (if applicable)	7b. ADDRESS (City, State, and ZIP Code)		
8c. ADDRESS (City, State, and ZIP Code) Fort Detrick Frederick, Maryland 21701-5012			9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DAMD17-84-C-4116			
11. TITLE (Include Security Classification) Genetic and Molecular Studies of the Phlebotomus Fever Group of Viruses			10. SOURCE OF FUNDING NUMBERS			
			PROGRAM ELEMENT NO. 61102A	PROJECT NO. 3M1- 61102BS12	TASK NO. AC	WORK UNIT ACCESSION NO. 104
12. PERSONAL AUTHOR(S) David H. L. Bishop, Ph.D.						
13a. TYPE OF REPORT Final Report		13b. TIME COVERED FROM 84/3/1 TO 84/7/31		14. DATE OF REPORT (Year, Month, Day) 1985 March	15. PAGE COUNT 12	
16. SUPPLEMENTARY NOTATION						
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)			
FIELD	GROUP	SUB-GROUP	Virus vaccines, Phlebotomus fever group viruses			
06	13					
06	03					
19. ABSTRACT (Continue on reverse if necessary and identify by block number) See reverse side.						
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> OTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified			
22a. NAME OF RESPONSIBLE INDIVIDUAL Mrs. Virginia Miller			22b. TELEPHONE (Include Area Code) 391/663-7325	22c. OFFICE SYMBOL SGRD-RMS		

Item 19:

10 to the + power 6, 10 to the 6th power

Studies supported by this contract have reported the characterization of the structural components of several Phlebotomus fever serogroup viruses (viral polypeptides and RNA species). The viruses studied include Punta Toro (PT), Karimabad (KAR), Chagres (CHG), Sandfly fever Sicilian (SFS Tesh and Sabin isolates), Sandfly fever Naples (SFN), Rift Valley fever (RVF), Phlebotomus 3 virus (Phl 3), Icoraci (ICO) and Buenaventura (BUE) viruses. In summary, the evidence has shown that the viruses have 3 virion RNA species (large, L, $2.6-2.8 \times 10^6$, medium, M, $1.85-2.2 \times 10^6$ and small, S, $0.7-0.8 \times 10^6$), which are comparable in number to those of other members of the family, but different in size for the S RNA (which is significantly larger than the S RNA of Bunyavirus genus members i.e., $0.4-0.5 \times 10^6$). The studies have also shown that, although the nucleocapsid, N, polypeptides of Phlebotomus fever group viruses ($20-24 \times 10^3$ daltons) are similar in size to those of bunyaviruses, the glycoproteins (G1, G2, $57-69 \times 10^3$ daltons) are quite different (e.g., bunyaviruses: G1 115×10^3 , and G2 38×10^3 daltons).

With the demonstration of a tripartite RNA genome, genetic analyses were also undertaken in order to determine the coding strategies of the 3 RNA species and determine if recombinant viruses could be obtained and used for vaccine purposes. From analyses of intertypic reassortant PT viruses we showed that the viral S RNA species codes for the viral N protein. Although not yet proven, based on the precedent of the bunyavirus coding strategy, it is probable that the phlebovirus M RNA codes for the 2 viral glycoproteins leaving the L RNA species to code for the 200×10^3 dalton protein (putative transcriptase component) found in viruses. The genetic studies, including interference assays, indicated that although intertypic reassortant viruses could be obtained, heterotypic virus interactions are not demonstrable (i.e., heterotypic viruses do not interfere with each other and do not reassort their genomes in dual virus infections).

The coding strategy of the 3 RNA species have been investigated with the goal of characterizing the epitopes on viral antigens that elicit and interact with neutralizing antigens. Such goals were realized by cloning and sequencing studies. The S RNA of PT phlebovirus has been cloned and sequenced. From the data obtained an open reading frame in the viral complementary RNA species has been identified (coding for N). However another open reading frame has been found in the viral sense strand. The mRNA species corresponding to the 2 open reading frames have been identified and characterized. Correlations of these predicted products to viral gene products remain to be completed as does cloning and sequencing the PT M and L RNA species.

TABLE OF CONTENTS

I. SUMMARY	2
II. REPORT	5
A. Introduction	5
B. Results from this Reporting Period	6
C. Summary of Progress Report	7
D. Literature Cited	8
E. Publications emanating from this contract	9

QUALITY

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution /	
Availability Codes	
Dist	Avail and / or Special
A-1	

I. SUMMARY

We have undertaken the biochemical and genetic characterization of members of the Phlebovirus genus of the negative sense RNA virus family Bunyaviridae (Bishop *et al.*, 1980). This genus of viruses include members that are known human pathogens and that are of consequence to military and civilian personnel in particular regions of the world (Bishop and Shope, 1979). Rift valley fever (RVF) virus has caused epidemics of infection including several hundred deaths in a single outbreak in Egypt in 1977-1978. The sandfly fever virus isolates, Sicilian and Naples (SFN, SFS), originally were recovered from American troops in 1943-1944 during epidemics of sandfly fever among military personnel. Isolates representing New World phleboviruses have been obtained from military personnel stationed in Panama (Punta Toro, PT). Other New World isolates some of which are the etiologic agents of human infection include Aguagate (AGU), Anhangá (ANH), Bujaru (BUJ), Icoaraci (ICO), Chagres (CHG), Buenaventura (BUE), Candiru (CDU) and Itaporanga (ITP) viruses. Other Old World isolates include Karimabad (KAR) virus.

Our analyses have been directed towards developing a strategy for phlebovirus vaccine development. Initial studies were therefore aimed at delineating the structural components of member viruses. With the demonstration of a tripartite RNA genome, genetic analyses were also undertaken in order to determine the coding strategies of the 3 RNA species and determine if recombinant viruses could be obtained and used for vaccine purposes. From analyses of intertypic reassortant PT viruses we showed that the viral 7×10^5 dalton small (S) RNA species codes for the viral $20-25 \times 10^3$ dalton nucleocapsid (N) protein. Although not proven, based on the precedent of the bunyavirus coding strategy, it is probable that the middle (M) size phlebovirus RNA (2×10^6 daltons) codes for the 2 viral glycoproteins ($50-70 \times 10^3$ daltons) leaving the 3×10^6 dalton large (L) RNA species to code for the 200×10^3 dalton protein (putative transcriptase component) found in viruses. The genetic studies, including interference assays we established, indicated that although intertypic reassortant viruses can be obtained, heterotypic virus interactions are not demonstrable (i.e., heterotypic viruses do not interfere with each other and do not reassort their genomes in dual virus infections). Although not all phleboviruses have been tested for genetic interactions (see below), the results do not hold out much hope for using such an approach for vaccine development. Consequently, we have taken an alternate approach, that of developing subunit vaccines. Central to this approach is a determination of the coding strategy of the 3 RNA species and the goal of characterizing the epitopes on viral antigens that elicit and interact with neutralizing antigens. Such goals will be realized by cloning and sequencing studies here and antigenic epitope analyses being undertaken at the U.S. Army Medical Research Institute of Infectious Diseases U.S. Army Medical Research Institute of Infectious Diseases (USAMRIID) (Drs. J. Smith and J. Dalrymple). In the previous studies the results of cloning and sequencing the S RNA of PT phlebovirus were given. From the data obtained an open reading frame in the viral complementary RNA species (mRNA) was identified (coding for N). An open reading frame was also found in the viral sense strand. Since the renewal of the contract (March 1, 1984) until the transfer of the Principal Investigator to the NERC Institute of Virology in Oxford (U.K.) and termination of the contract on July 31, 1984, the PT S induced mRNA species have been characterized.

This final report summarizes the results obtained through the entire contract, (listed below) including the results obtained in the final period of the contract.

- (1). Analyses of the major structural components of PHL group viruses have established that KAR, PT, CHG, Candiru (CDU), ICO, PHL 3, Itaporanga (ITP), Buenaventura (BUE), and the Sicilian and Naples sandfly fever (SFS, SFN) viruses, each has a tripartite RNA genome and three major structural polypeptides (two external glycoproteins, G1 and G2, $57-69 \times 10^3$ daltons, and an internal, nucleocapsid

associated, protein N, $20-24 \times 10^3$ daltons). Both the mol. wt. of the major structural polypeptides and the virion RNA segments of the different PHL group viruses can be easily distinguished from those of bunyaviruses (e.g., the California serogroup, CAL, viruses) and, to various extents (depending on the virus), from each other. Tryptic peptide analyses of ^{35}S and ^3H methionine labeled G1 and G2 polypeptides of KAR have established that these two polypeptides have distinguishable sequences (Robeson *et al.*, 1979). The behaviour of reduced KAR G1 and G2 polypeptides on polyacrylamide gel electrophoresis was found to be aberrant by comparison with unreduced preparations.

(2). Twenty four ts mutants of PT virus were isolated following mutagenesis of the wild-type virus by growth in the presence of 5-fluorouracil. Recombination assays with these mutants have allowed them to be categorized into 3 non-overlapping recombination groups (Group I has 8 ts mutants; Group II has 5 ts mutants; Group III has 1 ts mutant - so strictly is not a group - and 1 ts mutant is probably a double mutant).

(3). Oligonucleotide fingerprint analyses of prototype DEN 1, 2, 3 and 4 viruses involving both single and mixed coelectropherograms of ribonuclease T1 digests of ^{32}P labeled 40S viral RNA samples showed that each has a unique fingerprint that is easily distinguished from that of another prototype DEN virus (Veza *et al.*, 1980). These procedures have now been adopted by Walter Reed Army Institute of Research (WRAIR) personnel for their Dengue virus studies). Evidence was obtained indicating that the 5' sequence of DEN 2 RNA is m7GpppAmpXp... Oligonucleotide fingerprint analyses of alternate RVF virus isolates obtained from different countries of Africa have similarly been used to categorize and relate/distinguish RVF isolates (Cash *et al.*, 1981).

(4). Electron microscopic analyses of the surface structure arrangements of PHL group viruses have been undertaken and indicate that unlike bunyaviruses, but like uukuviruses (2 of the other Bunyaviridae genera) PHL group viruses have a particular "chimney-pot" arrangements of their surface glycoproteins as evidenced by glutaraldehyde fixation prior to staining (Robeson *et al.*, 1979).

(5). Competition RIA assays using iodinated nucleocapsid and glycoprotein preparations of KAR virus, KAR antisera and the competing antigens of KAR, CHG and SFS indicate that the KAR and SFS N polypeptides have more antigenic determinants in common than have the N polypeptides of KAR and CHG. Also the KAR and SFS G polypeptides share more antigenic determinants than the G polypeptides of KAR and CHG (Klimas *et al.*, 1981). No shared antigenic determinants were detected between KAR and vesicular stomatitis virus (VSV), or the bunyaviruses La Crosse (LAC), Oriboca (ORI), or Bunyamwera (BUN) viruses.

(6). ICO virus has been adapted to produce plaques in Vero cell monolayers at 39.8°C . The original virus stock, which gave 10^5 plaques at 35°C , gave none at 39.8°C . By high temperature passaging of the virus stock and cloning at 39.8°C , a stock of ICO virus has been derived which gives 8×10^4 PFU at 35°C and 2.3×10^3 PFU at 39.8°C .

(7). The viral induced polypeptides, (immune precipitated from infected cell extracts by their homologous antisera), have been characterized for KAR, AGU, BUE, CDU, SFS and SFN viruses. Several alternate isolates of PT virus obtained from the eastern, central and western regions of Panama have been cloned and their L, M and S RNA species fingerprinted and shown to be distinct/related to each other (depending on the virus isolate).

(8). By dual wild-type virus crosses, reassortant viruses have been obtained between prototype PT and an alternate PT isolate (PT-ada). The genotypes of 2 such reassortants were shown by fingerprint analyses to be the L/M/S combinations of ada/PT/ada and PT/PT/ada.

(9). The N polypeptides of PT-ada and prototype PT viruses can be distinguished by tryptic peptide analyses allowing the N polypeptides of the reassortants to be analyzed. By such analyses it has been shown that their S RNA codes for the virion N polypeptide. This conclusion has been confirmed by S mRNA in vitro translation analyses.

(10). Dual wild-type virus infections have failed to detect reassortant virus formation between PT and BUE viruses, or BUE and ICO viruses.

(11). An assay of the homologous and heterologous virus interference capabilities by stocks of PT, or ICO, viruses that contain their respective defective interfering (DI) virus has been developed. Using such assays, homologous virus interference has been demonstrated. No heterologous virus interference has been detected from ICO DI and PT coinfections, or ICO DI and BUE, or ICO DI and snowshoe hare (SSH) bunyavirus coinfections.

(12). The stocks of PT and ICO DI virus have been shown to contain new RNA species not found in virus stocks lacking DI virus. Fingerprint analyses have shown that for both viruses the new RNA species represent deletion derivatives of their respective L RNA species.

(13). The 3' end sequence of the 3 RNA species of both PT and BUE viruses have been shown to be like those of the uukuvirus, Uukuniemi, i.e., HO^UUC.

(14). Heterologous recombination attempts involving dual infections with wild-type PT and SFS viruses, PT and CHG viruses, PT and ICO viruses, or PT and KAR viruses have failed to yield recombinant viruses. Similar studies with different PT virus varieties yielded recombinant viruses.

(15). Further analyses of the homologous and heterologous virus interference capabilities of PT (or ICO) virus preparations that contain DI particles have failed to detect heterologous virus interference with KAR, SFS, SFN, CHG, CDU or ITP phleboviruses under conditions in which homologous PT (or ICO) interference was demonstrated. We conclude that these heterologous viruses are distantly related to PT (or ICO) viruses, and (by this test) not genetically compatible.

(16). Current data indicate that at the structural level the tick-borne uukuviruses (Uukuvirus genus, Bunyaviridae family) closely resemble phleboviruses. The 3' end sequences of their RNA species, and the virion polypeptides and RNA sizes are similar, as are their viral-induced non-structural polypeptides. The structural components of several non-assigned tick-borne members of the family have been characterized and found to conform to neither the phleboviruses/uukuviruses, nor to the mosquito-borne bunyaviruses (Clerx and Bishop, 1981; Clerx et al., 1981). Because of their unique characteristics these other viruses have been constituted into a new genus, the Nairovirus genus. Member viruses have 3' AGAGUUUC.. end sequences (Clerx van-Haaster et al., 1982), in contrast to bunyaviruses (3' UCAUCAUGA...), or uukuviruses (3' UGUGUUUUCUGG...), or phleboviruses (3' UGUGUUUCG..). The nairovirus viral RNA species and virion polypeptide sizes also differ significantly from those of uukuviruses, phleboviruses, or bunyaviruses.

(17). Characterization of the unassigned tick-borne Dhori (DHO) virus, a virus that heretofore has been considered as a possible member of the family, has been undertaken in order to determine if this virus has the structural attributes of phleboviruses (or members of other defined genera of the family). The data obtained indicate that DHO virus does not conform to any other member of the family, rather it resembles an orthomyxovirus (Clerx et al., 1983).

(18). In order to clone, sequence and thereby determine the complete coding content of the RNA species of PHL group viruses, the Principal Investigator spent a 10 month sabbatical in the laboratory of Dr. G.G. Brownlee in Oxford, England. During this time he cloned and sequenced complete copies of influenza A/PR/8/34 RNA segments 2 and 3, as well as bunyavirus SSH S RNA species (Bishop et al., 1982a,b,c).

(19). Using a PT S RNA 3' end specific viral complementary oligodeoxynucleotide, clones of DNA copies of the S RNA have been obtained and sequenced. The data have been used to produce other viral complementary oligodeoxynucleotides and further, overlapping, clones obtained and sequenced allowing a DNA sequence of the S RNA to be obtained. Open reading frames have been identified in the viral and viral complementary RNA sequences. The results indicate a novel coding strategy for phleboviruses, that of an ambisense viral RNA (Ihara et al., 1984).

(20). The S mRNA species induced by PT virus in permissive cells have been characterized, confirming that they are subgenomic and correspond to different viral sequences.

II. REPORT

A. Introduction.

The objectives of this contract are to determine the infection strategy of members of the PHL group viruses (Phlebovirus genus, Bunyaviridae) and develop protocols for vaccine development. Since this group of exotic viruses includes agents that cause illnesses in epidemic proportions (e.g., RVF) in different parts of the world, and therefore are of military importance, our objectives relate to the question of deriving vaccines that will be useful in protecting military personnel against virus infections.

To realize these objectives, initial studies involved determining if reassortant PHL viruses could be used for preparing vaccines. This was studied by determining which PHL group viruses are genetically compatible (capable of producing reassortant viruses with other members of the genus, or members of other genera of the family). Studies conducted in our laboratories have demonstrated that certain virus serotypes of the CAL serogroup of the Bunyavirus genus (another genus of the Bunyaviridae) are capable of producing reassortant viruses with other CAL group members (Gentsch & Bishop, 1976; Gentsch, et al., 1977; Gentsch et al., 1979; Rozhon et al., 1981). However, not all CAL group bunyavirus crosses yield reassortants, indicating that their gene pool is limited. Also no crosses involving viruses representing different bunyavirus serogroups (e.g., CAL and the Group C, or Bunyamwera, serogroups) have yet yielded reassortant viruses. Analyses of reassortant bunyaviruses have shown that the S RNA codes for the nucleocapsid (N) protein and a non-structural protein (NS_G) (Gentsch and Bishop, 1978; Cash et al., 1979; Fuller and Bishop, 1982). The M RNA codes for the viral glycoproteins.

Results reported previously from this contract have documented that PHL group viruses have a tripartite RNA genome with RNA segments that are designated L, M and S. From dual PT ts mutant, or wild-type, virus coinfections recombinant intertypic PT

viruses have been obtained. However, no heterotypic reassortants have been obtained using different PHL group viruses. Although not all PHL virus combinations have been tested, this negative result suggests (but does not prove) that there are restrictions at the genetic level between different PHL group viruses. DI interference studies also suggest that there are restrictions on genetic interactions between many of the PHL group viruses.

An alternate strategy for vaccine development is to identify and prepare adequate quantities of the viral antigen(s) that elicit protective antibodies. Central to this issue is knowledge of the RNA segment coding assignments and characterization of the viral antigen(s) that interact with such antibodies (see Gentsch & Bishop, 1978, 1979). Our current studies have been directed towards defining the RNA segment coding assignments of phleboviruses. We have already demonstrated that the S RNA segment codes for N protein. Proof that the M RNA codes for the glycoproteins (as shown for bunyaviruses) has not yet been obtained, and remains a goal of our continuing efforts. Studies by USAMRIID personnel with monoclonal antibodies indicate that the glycoproteins interact with neutralizing sera. Therefore, it is important to determine which PHL viral RNA species codes for the glycoproteins. Sequencing the phlebovirus S RNA species has been reported (Ihara et al., 1984).

B. Results from this Reporting Period.

The presence of open reading frames in the PT vRNA as well as the PT vcRNA sequences contrasts with the S RNA sequence data of SSH bunyavirus where only the vcRNA has a significant length of open reading (Bishop et al., 1981).

In the PT S viral complementary (vc) RNA sequence there is a large contiguous open reading frame initiated with a methionine codon that extends from vcRNA 5' nucleotide no. 35 through 763 (243 amino acid residues). Shorter open reading frames in the viral complementary sequences correspond to amino acid stretches of no more than 60-70 amino acids in length (Ihara et al., 1984). These data indicate that at least 1200 nucleotides of the vc RNA sequence are not involved in coding for a gene product. In the PT S viral sequence there is also a large contiguous open reading frame extending from the end of the clone i.e., corresponding to vRNA 5' nucleotide 29 through 778 (258 amino acid residues). It is possible that the existence of the large open reading frame in the viral sense strand is fortuitous. If so, then it would be unlikely that the viral RNA (or viral sense RNA made in infected cells) could be translated in vitro. Also it would be unlikely that other phleboviruses would have a similar open reading frame sequence in their viral sense S RNA sequence. This latter question remains to be investigated. However the observation of open reading frames in the viral-complementary and viral sense S RNA sequences has been followed up by the demonstration of the requisite subgenomic mRNA species in PT virus infected cells (Ihara et al., 1984). Also the subgenomic S viral-complementary mRNA species that codes for N (but not the PT S viral mRNA species that codes for NS protein) has been observed in virus infected cells grown in the presence of protein synthesis inhibitors (puromycin or cycloheximide). The coding assignment of the S vc mRNA species has been confirmed by limited peptide sequence analyses of viral N protein. The question of the relevance of coding information in the viral RNA needs to be addressed by examining other phleboviruses and translating their cellular viral sense and viral complementary RNA species.

C. Summary of Progress Report.

The research supported by this contract has shown that representative PHL group viruses (Phlebovirus genus) have a tripartite RNA genome consisting of 3 unique RNA species (L, M and S). The L and M RNA species have mol. wts. like those of Bunyavirus and Uukuvirus genus members. Like uukuviruses, the S RNA species of phleboviruses is significantly larger than that of bunyaviruses even though the N proteins coded by the S RNA species of both virus types are similar in size ($20-24 \times 10^3$ daltons).

The sequencing data from PT S RNA postulate that the viral-complementing RNA codes for a 22×10^3 dalton protein. This protein is N. In the viral sense strand there is an open reading frame that codes for a non-structural protein, designated NS_G.

Our analyses have shown that the size ranges of phlebovirus glycoproteins and their surface arrangement are similar to those of uukuviruses, and unlike those of Bunyavirus genus viruses. Whether the viral M RNA codes for the glycoprotein has yet to be established.

Genetic studies initiated with Punta Toro (PT) virus have shown that high frequency intertypic genetic recombination can result from certain PT mixed virus infections (ts, or dual wild-type coinfections using alternate isolates of PT virus). From analyses of the reassortants generated from dual infections involving prototype PT and an alternate PT strain, PT-ada, it has been demonstrated that the viral S RNA codes for the N polypeptide. In vitro translation by subgenomic polysome mRNA indicates that N is coded by a small mRNA.

So far no recombination has been detected between PT and BUE, or PT and SFS, or PT and CHG, or PT and ICO, or PT and KAR, or BUE and ICO viruses, either by analyses of the progeny of dual wild-type virus infections, or from dual infections involving PT ts mutants and the wild-type alternate virus (neither of which give plaques at 39.8°) and screening for reassortant progeny at the non-permissive temperature. These results suggest that these phleboviruses do not belong to the same gene pool.

In addition to the heterologous phlebovirus recombination assays, we have screened for genetic interaction between phleboviruses based on heterologous virus interference assays. We have failed to detect heterologous virus interference using PT (or ICO) dI virus preparations and BUE, KAR, SFS, SFN, CHG, CDU or ITP phleboviruses under conditions in which interference of homologous PT, or alternate PT varieties, (or ICO) viruses could be demonstrated.

We conclude from these results that the phleboviruses are genetically diverse. From the vaccine development point of view it now appears unlikely that heterologous phleboviruses could be used to produce reassortants for vaccine development. Evidently, however, intertypic reassortants could be obtained and may be of use for vaccine development.

As an alternate strategy for vaccine development we have initiated a program to characterize the proteins coded by the 3 viral RNA species with the aim of identifying the exact sites on viral antigens that induce and interact with neutralizing antibodies. This objective is being realized by DNA cloning and sequencing the viral RNA species. The goal of vaccine development will then be addressed by expressing cloned viral gene products and, or preparing subunit vaccines of cloned products (or peptides) that protect against virulent virus challenge.

D. Literature Cited.

- Bishop, D.H.L., Calisher, C., Casals, J., Chumakov, M.P., Gaidomovich, S.Ya., Hannoun, C., Lvov, D.K., Marshall, I.D., Oker-Blom, N., Petterson, R.F., Porterfield, J.S., Russell, P.K., Shope, R.E., and Westaway, E.G., 1980, Bunyaviridae. Intervirology 14:125-143.
- Bishop, D.H.L., Gould, K.G., Akashi, H. and Clerx-van Haaster, C.M., 1982a, The complete sequence and coding content of snowshoe hare bunyavirus small (S) viral RNA species. Nucl. Acids Res., 10: 3703-3713.
- Bishop, D.H.L., Huddleston, J.A. and Brownlee, G.G., 1982b, The complete sequence of RNA segment 2 of influenza A/NT/60/68 and its encoded P1 protein. Nucl. Acids Res., 10:1335-1343.
- Bishop, D.H.L., Jones, K.L., Huddleston, J.A. and Brownlee, G.G., 1982c, Influenza A virus evolution. Complete sequences of influenza A/NT/60/68 RNA segment 3 and its predicted acidic P polypeptide compared with those of influenza A/PR/8/34. Virology, 120:481-489.
- Bishop, D.H.L., and Shope, R. E., Bunyaviridae, 1979, In Comprehensive Virology, Volume 14, (ed. H. Fraenkel-Conrat and R. R. Wagner), pp.1-156. Plenum Press, New York.
- Cash, P., Vezza, A.C., Gentsch, J.R. and Bishop, D.H.L, 1979, Genome complexities of the three mRNA species of snowshoe hare bunyavirus and in vitro translation of S mRNA to viral N polypeptide. J. Virol. 31:685-694.
- Cash, P., Robeson, G., Erlich, B.J., and Bishop, D.H.L., 1981, Biochemical characterization of Rift Valley Fever and other phlebotomus fever group viruses. Contr. Epidem. Biostatist. 3:1-20.
- Clerx, J.P.M. and Bishop, D.H.L., 1981, Qalyub Virus, a member of the newly proposed Nairovirus genus (Bunyaviridae), Virology, 108:361-372.
- Clerx, J.P.M., Casals, J., and Bishop, D.H.L., 1981, Structural characteristics of nairoviruses (Genus Nairovirus Bunyaviridae). J. Gen. Virol., 35:1-13.
- Clerx, J., Fuller, F. and Bishop, D.H.L. Tick-borne viruses structurally similar to influenza viruses. Virology, 127:205-219, 1983.
- Clerx-van Haaster, C.M., Clerx, J.P.M., Ushijima, H., Akashi, H., Fuller, F., and Bishop, D.H.L., 1982, The 3' terminal RNA sequences of bunyaviruses and nairoviruses (Bunyaviridae). Evidence of end sequence generic differences within the virus family. J. Gen. Virol., 61:289-292, 1872.
- Fuller, F. and Bishop, D.H.L., 1982, Identification of viral coded non-structural polypeptides in bunyavirus infected cells. J. Virol., 41:643-648.
- Gentsch, J., and Bishop, D.H.L., 1976, Recombination and complementation between temperature sensitive mutants of the bunyavirus, snowshoe hare virus. J. Virol. 20:351-354.
- Gentsch, J., and Bishop, D.H.L., 1978, Small RNA segment of bunyaviruses codes for the viral nucleocapsid (N) protein. J. Virol. 28:417-419.

Gentsch, J.R., and Bishop, D.H.L., 1979, M viral RNA segment of bunyavirus codes for two unique glycoproteins: G1 and G2. J. Virol. 30:767-776.

Gentsch, J.R., Robeson, G., and Bishop, D.H.L., 1979, Recombination between snowshoe hare and La Crosse bunyaviruses. J. Virol. 31:707-717.

Gentsch, J., Wynne, L.R., Clewley, J.P., Shope, R.E., and Bishop, D.H.L., 1977, Formation of recombinants between snowshoe hare and La Crosse bunyaviruses. J. Virol. 24:893-902.

Ihara, T., Akashi, H., and Bishop, 1984, D.H.L. Novel coding strategy (ambisense genomic RNA) revealed by sequence analyses of Punta Toro phlebovirus S RNA. Virology, 136:293-306.

Klimas, R.A., Ushijima, H., Clerx-van Haaster, C.M., and Bishop, D.H.L., 1981, Radioimmune assays and molecular studies that place Anopheles B and Turlock serogroup viruses in the Bunyavirus genus (Bunyaviridae). Amer. J. Trop. Med. & Hyg. 30:876-887.

Robeson, G., El Said, L.H., Brandt, W., Dalrymple, J., and Bishop, D.H.L., 1979, Biochemical studies on the Phlebotomus fever group viruses (Bunyaviridae family). J. Virol. 30: 339-350.

Rozhon, E.J., Gensemer, P., Shope, R.E., and Bishop, D.H.L., 1981, Attenuation of virulence of a bunyavirus involving an L RNA defect and isolation of LAC/SSH/LAC and LAC/SSH/SSH reassortants, Virology 111:125-138.

Veza, A.C., Rosen, L., Repik, P., Dalrymple, J., and Bishop, D.H.L., 1980, Characterization of the viral RNA species of prototype Dengue viruses. Amer. J. Trop. Med. & Hyg. 29:643-652.

E. Publications emanating from this contract.

Bishop, D.H.L., 1979, Genetic potential of bunyaviruses. Curr. Top. Micro. Immun. 86:1-33.

Bishop, D.H.L., 1982, The molecular properties of bunyaviruses including group C isolates, and the genetic capabilities of bunyaviruses to form recombinants. Proc. Intern. Symp. Trop. Arboviruses and Hemorrhagic Fevers, edited by F. de P. Pinheiro, Imprensa Nat. Fund. Sci. Dev. Tech., Belem, Brazil, pp. 117-133.

Bishop, D.H.L., 1983, The application of RNA fingerprinting and sequencing to viral diagnosis. Current Topics in Microbiology and Immunology, 104:259-271.

Bishop, D.H.L., Beaty, B.J., and Shope, R.E., 1980, Recombination and gene coding assignments of bunyaviruses and arenaviruses. Annals New York Acad. Sci. 354:84-106.

Bishop, D.H.L., Clerx, J.P.M., Clerx-van Haaster, C.M., Robeson, G., Rozhon, E.J., Ushijima, H., and Veerisetty, V., 1981, Molecular and genetic properties of members of the Bunyaviridae, In: Replication of Negative Strand Viruses (D.H.L. Bishop and R.W. Compans, eds.), pp. 135-145, Elsevier, New York.

- Bishop, D.H.L., Gentsch, J.R., and El-Said, L.H., 1978, Genetic capabilities of bunyaviruses. J. Egypt Pub. Health Assoc. 53:217-234.
- Bishop, D.H.L., Gould, K.G., Akashi, H. and Clerx-van Haaster, C.M., 1982a, The complete sequence and coding content of snowshoe hare bunyavirus small (S) viral RNA species. Nucl. Acids Res., 10: 3703-3713.
- Bishop, D.H.L., and Shope, R.E., 1979, Bunyaviridae, In Comprehensive Virology, Volume 14, (ed. H. Frankel-Conrat and R.R. Wagner), pp. 1-156, Plenum Press, New York.
- Cash, P., Robeson, G., Erlich, B.J., and Bishop, D.H.L., 1981, Biochemical characterization of Rift Valley Fever and other phlebotomus fever group viruses. Contr. Epidem. Biostatist. 3:1-20.
- Clerx, J.P.M., and Bishop, D.H.L., 1980, Qalyub Virus, a member of the newly proposed Nairovirus genus (Bunyaviridae), Virology 108:361-372.
- Clerx, J.P.M., Casals, J., and Bishop, D.H.L., 1981, Structural characteristics of nairoviruses (Genus Nairovirus Bunyaviridae). J. Gen. Virol., 35:1-13.
- Clerx, J., Fuller, F. and Bishop, D.H.L. Tick-borne viruses structurally similar to influenza viruses. Virology, 127:205-219, 1983.
- Clerx-van Haaster, C.M., Clerx, J.P.M., Ushijima, H., Akashi, H., Fuller, F., and Bishop, D.H.L., 1982, The 3' terminal RNA sequences of bunyaviruses and nairoviruses (Bunyaviridae). Evidence of end sequence generic differences within the virus family. J. Gen. Virol., 61:289-292.
- Clewley, J.P., and Bishop, D.H.L., 1982, Oligonucleotide fingerprinting of viral genomes. In: New Techniques in Practical Virology (ed. C. Howard), pp. 231-277, Alan Liss Inc., N.Y.
- Dalrymple, J.M., Smith, J.F., Harrison, S.A., Bishop, D.H.L. and Ussery, M.A., 1982, Immunological and biochemical characterization of phlebotomus fever bunyaviruses. In: Proc. Intern. Symp. Trop. Arboviruses and Hemorrhagic Fevers, edited by F. de P. Pinheiro, Impresso Nat. Fund. Sci. Dev. Tech., Belem, Brazil, pp. 67-78.
- Gentsch, J.R., and Bishop, D.H.L., 1980, The genetic recombination potential of bunyaviruses, J. Vesenjak-Hirjan et al. (eds.): Arboviruses in the Mediterranean countries, pp. 81-96, Zbl. Bakt. Suppl. 9. Gustav Fischer Verlag, Stuttgart.
- Klimas, R.A., Ushijima, H., Clerx-van Haaster, C.M., and Bishop, D.H.L., 1981, Radioimmune assays and molecular studies that place Anopheles B and Turlock serogroup viruses in the Bunyavirus genus (Bunyaviridae). Amer. J. Trop. Med. & Hyg. 30:876-887.
- Ihara, T., Akashi, H., and Bishop, 1984, D.H.L. Novel coding strategy (ambisense genomic RNA) revealed by sequence analyses of Punta Toro phlebovirus S RNA. Virology, 136:293-306.
- Robeson, G., El-Said, L.H., Brandt, W., Dalrymple, J., and Bishop, D.H.L., 1979, Biochemical studies on the Phlebotomus fever group viruses (Bunyaviridae family), J. Virol. 30:339-350.

Ushijima, H., Klimas R., Kim, S., Cash, P., and Bishop, D.H.L., 1980, Characterization of the viral ribonucleic acids and structural polypeptides of Anopheles A, Bunyamwera, Group C, California, Capim, Guama, Patois, and Simbu bunyaviruses. Amer. J. Trop. Med. & Hyg. 29:1441-1452.

Veza, A.C., Rosen, L., Repik, P., Dalrymple, J., and Bishop, D.H.L., 1980, Characterization of the viral RNA species of prototype Dengue viruses. Amer. J. Trop. Med. & Hyg. 29:643-652.

DISTRIBUTION LIST

5 copies **Commander**
US Army Medical Research Institute of
Infectious Diseases
ATTN: SGRD-UIZ-E
Fort Detrick, Frederick, MD 21701-5011

4 copies **Commander**
US Army Medical Research and Development Command
ATTN: SGRD-RMS
Fort Detrick, Frederick, Maryland 21701-5012

12 copies **Defense Technical Information Center (DTIC)**
ATTN: DTIC-DDAC
Cameron Station
Alexandria, VA 22304-6145

1 copy **Dean**
School of Medicine
Uniformed Services University of the
Health Sciences
4301 Jones Bridge Road
Bethesda, MD 20814-4799

1 copy **Commandant**
Academy of Health Sciences, US Army
ATTN: AHS-CDM
Fort Sam Houston, TX 78234-6100

END