

NO-A167 022

A LONGITUDINAL STUDY OF DISEASE INCIDENCE AMONG
ANTARCTIC WINTER-OVER PERSONNEL(U) NAVAL HEALTH
RESEARCH CENTER SAN DIEGO CA L A PALINKAS FEB 86

1/1

UNCLASSIFIED

NAVHLTHRSCHC-86-3

F/G 6/19

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A167 822

2

A LONGITUDINAL STUDY OF DISEASE INCIDENCE AMONG ANTARCTIC WINTER-OVER PERSONNEL

L. A. PALINKAS

REPORT NO. 86-3

DTIC
ELECTE
MAY 19 1986
S D

DTIC FILE COPY



DISTRIBUTION STATEMENT A

Approved for public release
Distribution Unlimited

NAVAL HEALTH RESEARCH CENTER

P.O. BOX 85122
SAN DIEGO, CALIFORNIA 92138-9174

NAVAL MEDICAL RESEARCH AND DEVELOPMENT COMMAND
BETHESDA, MARYLAND

A LONGITUDINAL STUDY OF DISEASE INCIDENCE AMONG ANTARCTIC WINTER-OVER PERSONNEL

Lawrence A. Palinkas, Ph.D.
Manager, Psychiatric Effectiveness Program
Environmental Medicine Department

Naval Health Research Center
P.O. Box 85122
San Diego, California 92138-9174

To expedite communication of our research, this is a preprint of a paper submitted to Aviation, Space, and Environmental Medicine, and should be cited as a personal communication.

Report No. 86-3

supported by the Naval Medical Research and Development Command, Department of the Navy, under Research Work Unit MR000.01.01-6035. The views represented in this paper are those of the author and do not reflect the official policy or position of the Department of the Navy, Department of Defense, nor the U.S. Government.

SUMMARY

Problem

Previous research has indicated that Antarctic winter-over personnel are at increased risk for disease and psychosocial stress upon their return to the outside world. However, an earlier follow-up study of enlisted Navy personnel found no long-term risk for first hospital admissions.

Objective

The objective of this study was to test the hypothesis that even though there may be no long-term risk, winter-over personnel are at increased risk during the first year subsequent to the winter-over duty.

Approach

Subjects were 327 enlisted Navy men who wintered over between 1963 and 1974 and a control group of 2,396 enlisted men who volunteered and were accepted for winter-over duty but who did not winter over. A fifteen-year period from 1965 to 1979 was established for follow-up. Follow-up of subjects subsequent to screening for Operation Deep Freeze was conducted in six-month intervals.

Results

Results indicated that the total rates of first hospitalization during the six months prior to Antarctic duty and the first six months in Antarctica among winter-over personnel were significantly lower ($p < .05$) than the rates for the control group. No significant difference in the rates of the two groups was observed for the twelve months subsequent to winter-over duty.

Conclusions

The results of this study lend further support to the hypothesis that the winter-over experience is not associated with any subsequent risk for disease incidence. This may be due either to the quality of the candidates who actually winter-over or the process of psychosocial adjustment to prolonged isolation in this extreme environment which may promote health on a long-term basis.

Recommendations

Further research is required to identify which of the two alternative explanations best accounts for the lowered long-term risk of first hospitalizations among winter-over personnel.



Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Dist. Info.	
Availability Codes	
Dist	Avail and/or Special
A-1	

A Longitudinal Study of Disease Incidence among Antarctic Winter-Over Personnel

INTRODUCTION

A previous study of long-term disease incidence among U.S. Navy personnel who have wintered-over in small, isolated Antarctic research stations found that over a fifteen-year study period, these personnel have significantly lower total first hospitalization rates than a control group of enlisted personnel who were positively evaluated for Antarctic duty but who did not winter-over (12). When individual diagnostic categories were examined, the winter-over group had significantly fewer first hospitalizations for neoplasms; endocrine, nutritional and metabolic diseases; and diseases of the musculoskeletal system. Physiological changes which occur during the Antarctic winter and demographic, personality, and socioenvironmental factors which mediate the physical and psychological stress associated with prolonged isolation in an extreme environment were suggested as partial explanations for the lowered risk.

However, it has been observed that winter-over personnel are at risk for disease and psychological stress upon their return to the outside world. Outbreaks of common colds have been noted among personnel at many isolated antarctic stations immediately following the end of the isolation period and the resumption of contact with the outside world (1, 7-9). Although the evidence for an increased risk is not conclusive (6), the relative absence of viruses in the Antarctic and the lower resistance associated with neutropenia (14) may result in a reduction in immunocompetence, leaving the individual exposed to the risk of diseases of the respiratory system and infective and parasitic diseases. A study by Oliver (11) also found that adjustment problems and "culture shock" to home and new surroundings were common for returning winter-over personnel, and Natani and Shurley (10) observed that the process of readjustment to the larger society was potentially very stressful for winter-over personnel.

While the earlier follow-up study indicated that the winter-over experience has no long-term effects on the health and performance of enlisted Navy personnel, the possibility of a short-term risk immediately following the end of winter-over duty was not adequately addressed. Incidence rates for the follow-up period were averaged for the entire fifteen-year period and included time spent in the study immediately prior to and during the winter-over period. This study attempted to expand on the results of the first study by examining disease incidence of enlisted Navy winter-over personnel across time. A longitudinal perspective was employed to test the hypothesis that even though there may be no long-term health risk, winter-over personnel are at increased risk for hospitalization during the first year subsequent to the winter-over duty.

METHODS

Military and civilian applicants are evaluated for Antarctic winter-over duty by teams of psychiatrists and clinical psychologists through the Operation Deep Freeze Program. Records of screening evaluations conducted between 1963 and 1974 were compiled into a computerized file at the Naval Health Research Center. This file was then matched with the Inpatient Medical Data File at NHRC and service history data files obtained from the

Manpower and Personnel Management Information System (NMPC 15642) to obtain follow-up data on all enlisted personnel who participated in the Operation Deep Freeze Program during these years. Civilian applicants were not included because of the lack of available follow-up data. Follow-up information was obtained on 3,076 individuals on the basis of matching social security or service numbers.

For this study, cases were defined as those who wintered over at one of a number of small scientific stations between 1963 and 1974. A control group was comprised of individuals who applied for winter-over duty during the same period and were given a favorable combined evaluation by the screening team but who did not winter-over. Some of the program participants who were given unfavorable evaluations but who subsequently did winter-over were excluded from the study in order to reduce the potential confounding effects of personality variables. Of the 3,076 enlisted Navy personnel on whom follow-up information was available, 327 were identified as having wintered over and not having a negative combined evaluation by the psychiatrist/psychologist screening team. The control group was comprised of 2,396 enlisted Navy men who were screened during the study period and were considered to be acceptable by the screening team but who nevertheless did not winter over. The demographic characteristics and service histories of these two groups have been described elsewhere (12).

A fifteen-year period from 1965 to 1979 was established for follow-up. This was based on the period of time for which medical and service history information was available for both groups at the time the study was conducted. The start date for participation in the study was established as 1 January 1965 or the year an individual was evaluated for the Operation Deep Freeze Program if after this date. Withdrawal was defined as the date of last discharge from the Navy or 31 December 1979, whichever came first. Follow-up of subjects subsequent to entry into the study (i.e., subsequent to screening for Operation Deep Freeze) was conducted in six-month intervals for the first four years, allowing us to focus specifically on the period immediately prior to, during, and immediately following winter-over duty. Screening of enlisted personnel usually occurs in April of each year. Antarctic duty begins in September at the earliest and usually lasts for one year. The winter-over period usually begins in February and lasts until October.

Inpatient medical data included all first hospitalizations for all diagnoses which occurred during the follow-up period. Diagnoses were in accordance with the Eighth Revision, International Classification of Disease Adapted for Use in the United States (ICDA-8). Population at risk was defined as those subjects who remained in the study at the end of each interval.

Age-adjusted rates of total first hospitalizations were calculated for each time interval using the direct method of adjustment (5). The standard population was comprised of all study participants at risk during the time interval. The rates for the winter-over and control groups were compared to obtain estimates of relative risk by taking the ratio of rates for the winter-over group to rates for the control group. Levels of significance of

these associations were obtained using 95 percent confidence intervals (2).

RESULTS

The average rate of total first hospitalizations (incidence) for the fifteen-year period in the earlier study was significantly lower for the winter-over group, yielding a relative risk of 0.79 (12). Because of the reduced populations in each time interval in the current study, statistical significance was evident only in a few instances. Therefore, only total first hospitalizations are reported here. While some trends were evident in specific disease categories, such as diseases of the respiratory, circulatory, and digestive systems, the number of cases among the winter-over group were too few to achieve statistical significance and hence will not be reported.

The incidence of total first hospitalizations for all diagnoses among cases and controls by six-month intervals for the first four years of study participation is represented in Table 1. The rate for the winter-over group during the first two six-month periods of study participation was significantly lower than the rate for the control group. The rate

Table 1

Age-Adjusted Rates of Total First Hospitalizations for Winter-Over and Control Groups,
Operation Deep Freeze Participants, 1965-1979 by Six-Month Intervals
(per 10,000 person years)

Interval Number	Period (Months)	Mid-point (Months)	Winter Over		Controls		Relative Risk	95% Confidence Limits
			N	Rate	N	Rate		
1	0-6	3	3	104.7	79	359.0	0.29	-0.04 - 0.62*
2	6-12	9	4	137.3	65	345.2	0.40	0.00 - 0.80*
3	12-18	15	9	309.4	81	478.2	0.65	0.20 - 1.10
4	18-24	21	12	444.5	82	518.5	0.86	0.34 - 1.38
5	24-30	27	14	577.3	59	403.0	1.43	0.60 - 2.26
6	30-36	33	7	306.1	55	395.9	0.77	0.16 - 1.38
7	36-42	39	12	546.3	72	548.0	1.00	0.39 - 1.61
8	42-48	45	9	420.0	59	610.5	0.69	0.21 - 1.17
6-30	30-180	90	114	577.3	824	703.7	0.82	0.67 - 0.97*

* p. < .05

for the winter-over group displayed a steady increase, however, reaching a peak during the fifth interval which was about 27 months after entry into the study and nine months after the end of winter-over duty. With the increase in the rate of total first hospitalizations, the relative risk also increased, reaching a peak at the same time (i.e. nine months after the winter-over period). This is clearly evident from Figure 1. However, this relative risk of 1.43 was not statistically significant. Moreover, the rate of total first hospitalizations among the winter-over group for the remainder of the study period (months 30 to 180) was significantly less than the rate for the control group.




Figure 1. Relative risk of total first hospitalizations for winter-over group, 1965-1979. First four years of study participation by six-month intervals.

DISCUSSION

Although both groups may be similarly qualified and given favorable evaluations by the screening team, the winter-over group demonstrated a significantly smaller disease incidence during the six months immediately prior to winter-over duty and during the first six months of assignment in the Antarctic. The question remains as to why this is the case. One possible explanation is that the significantly lower rates of total first hospitalizations among the winter-over group, particularly during the period immediately prior to Antarctic duty, when compared with the control group, may be indicative of the success of the Operation Deep Freeze Program in selecting the healthiest candidates for winter-over duty. However, both groups were evaluated as acceptable for winter-over duty by the

screening team. No significant differences between the two groups were observed on any of the screening personality measures, and differences in age and, indirectly, pay grade and length of service were taken into consideration by the age-adjusted rates. With the exception of differences in education and occupation, the two groups appear to be adequately matched with respect to known health risk factors.

An alternative explanation is that the harshness of the environment combined with the processes of psychosocial adjustment to prolonged isolation minimize the risk of disease both in the short-term and, in the case of adaptation, in the long-term as well. Previous research has attributed the low incidence of infectious diseases and diseases of the respiratory system among Antarctic personnel during the austral winter as the result of prolonged isolation in a relatively sterile and disease-free environment (1,6,8,14). This would help to account for the smaller rates of first hospitalizations during the intervals of Antarctic duty. In addition, despite the presence of the psychological symptoms of the "winter-over syndrome" which appear to affect most winter-over personnel in varying degrees, the microcultures of Antarctic research stations support and promote values and personality traits in individual members which enable them to adapt to isolation and which also may promote health on a long-term basis as well (13, 15). The significantly lower rate of total first hospitalizations among the winter-over group after the fifth interval (27 months after entry into the study and nine months after returning from the Antarctic) would appear to suggest such a long-term benefit from this process of adaptation. This process may also begin immediately upon selection for winter-over duty, which would account for the significantly lower rates during the six months prior to Antarctic duty. Hence, the process of psychosocial adjustment may have long-term as well as short-term benefits for one's health and well-being.

During the last six months of Antarctic duty and the first six months subsequent to returning to the outside world, the relative risk of disease incidence among the winter-over group was no greater than that of the control group (i.e., it remained below 1.0). However, during the second six months upon returning to the outside world, the relative risk for disease incidence among the winter-over group showed a marked increase, followed by a decline and a second, but smaller increase. These variations were not statistically significant, however, indicating that although winter-over personnel may have a greater rate of total first hospitalizations upon return from Antarctic duty than either prior to or during such duty, this increase does not place them at greater risk for disease incidence relative to other Navy personnel who do not winter-over.

Because of the small number of cases in each interval, especially among the winter-over group, it is difficult to generalize the results of this study. Statistical significance was achieved only in the first two of the six-month intervals during the first four years of study participation. Moreover, while differences in age were taken into consideration when comparing rates of the two groups, other characteristics such as education and occupation which are known to influence hospitalization rates in the Navy (3-4) were not controlled because of the small number of cases.

Nevertheless, the results of this study lend further support to the hypothesis that the winter-over experience is not associated with any long-term risk for disease incidence. While some increase in disease incidence was observed upon return from winter-over duty, this increase failed to reach statistical significance and, hence, suggested no threat to the health and well-being of winter-over personnel either in the short-term or the long-term. In order to account for these results, two alternative explanations were discussed. However, further investigation is necessary to identify the specific causal factors.

ACKNOWLEDGMENTS

Patricia Coben provided invaluable assistance in the analysis of data.

REFERENCES

1. Allen JT. Common colds in Antarctica. *J. Hygiene*, 1973; 71:49-56.
2. Daniel WW. *Biostatistics: A foundation for analysis in the health sciences*. New York: John Wiley and Sons, 1983.
3. Gunderson EKE, Colcord C. Health risks in naval occupations. Report No. 82-1. San Diego: Naval Health Research Center, 1982.
4. Gunderson EKE, Rahe RH, Arthur RJ. The epidemiology of illness in naval environments. II. demographic, social background, and occupational factors. *Milit. Med.*, 1970; 135:453-8.
5. Lilienfeld AM, Lilienfeld DE. *Foundations of epidemiology*. 2nd ed. New York: Oxford University Press, 1980.
6. Meschievitz CK, Raynor WJ, Dick EC. Cold severity, duration, and epidemiology in persons emerging from isolation compared to newly arrived persons at McMurdo Station. *Antarctic J. U.S.*, 1983; 18:232-4.
7. Muchmore HG, Blackburn AB, Shurley JT, Pierce CM, McKown BA. Neutropenia in healthy men at the south polar plateau. *Arch. Intern. Med.*, 1970; 125:646-8.
8. Muchmore HG, Parkinson AJ, Scott EN. Respiratory virus infections during the winter at the south pole. *Antarctic J. U.S.*, 1983; 17:229-30.
9. Muchmore HG, Tatem, BA, Worley RA, Shurley JT, Scott N. Immunoglobins during south polar isolation. In: Edholm OG, Gunderson EKE, eds. *Polar human biology: proceedings of the SCAR/IUPS/IUBS symposium on human biology and medicine in the Antarctic*. Chicago: William Heinemann, 1974:135-140.
10. Natani K, Shurley JT. Sociopsychological aspects of a winter vigil at South Pole Station. In: Gunderson EKE, ed. *Human adaptability to antarctic conditions*. Antarctic Research Series, Vol. 22. Washington, D.C.: American Geophysical Union, 1974: 89-114.
11. Oliver DM. Some psychological effects of isolation and confinement in an antarctic winter-over group. Unpublished Ph.D. dissertation, School of Human Behavior, United States International University, San Diego, 1979.
12. Palinkas LA. Health and performance among Antarctic winter-over personnel: a follow-up study. Report No. 85-18. San Diego: Naval Health Research Center, 1985.
13. Palinkas LA. Sociocultural influences on psychosocial adjustment in Antarctica. Report No. 85-49. San Diego: Naval Health Research Center, 1985.

14. Parkinson AJ, Muchmore HG, Scott LV. Parainfluenzavirus upper respiratory infections at McMurdo Station during the austral summer 1975-76. *Antarctic J. U.S.*, 1979; 14:186-7.
15. Taylor AJ. The adaptation of New Zealand research personnel in the Antarctic. In: Edholm OG, Gunderson EKE, eds. *Polar human biology: proceedings of the SCAR/IUPS/IUBS symposium on human biology and medicine in the Antarctic*. Chicago: William Heinemann, 1974:417-429.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 86- 3	2. GOVT ACCESSION NO. ADA 167822	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A LONGITUDINAL STUDY OF DISEASE INCIDENCE AMONG ANTARCTIC WINTER-OVER PERSONNEL		5. TYPE OF REPORT & PERIOD COVERED Final
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Lawrence A. Palinkas, Ph.D.		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Health Research Center P.O. Box 85122 San Diego, CA 92138-9174		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS MR000.01.01-6035
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Medical Research & Development Command Naval Medical Command, National Capitol Region Berhesda, MD 20814		12. REPORT DATE February 1986
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Commander, Navy Medical Command Department of the Navy Washington, D.C.		13. NUMBER OF PAGES
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Approved for public release; distribution unlimited		
18. SUPPLEMENTARY NOTES To be submitted to <u>Aviation, Space and Environmental Medicine</u>		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Antarctica Extreme Environments Morbidity Hospitalizations Operation Deep Freeze Isolation		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A longitudinal perspective was employed to test the hypothesis that there is an increased risk of hospitalization among personnel who winter-over in Antarctica during the first year subsequent to the winter-over duty. Subjects were 327 enlisted Navy men who wintered over between 1963 and 1974 and a control group of 2,396 enlisted men who volunteered and were accepted for winter-over duty but who did not winter-over. A fifteen year period from 1965 to 1979 was established for follow-up. Follow-up of subjects subsequent to screening for		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102 LF 014 6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

Operation Deep Freeze was conducted in six month intervals for the first four years. Results indicated that the total rate of first hospitalization during the six months prior to Antarctic duty and the first six months in Antarctica was significantly lower than the rate for the control group. The rate for the winter-over group displayed a steady increase, peaking at nine months subsequent to returning from Antarctic duty. This peak rate was not significantly different from the rate of the control group, however. These results lend further support to the hypothesis that the winter-over experience is not associated with any long-term risk of disease incidence.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

END

DTIC

6-86