AD-A170 109	STATIONARY MARKOV SETS(U) FLORIDA STATE UNIV TALLAMASSEE DEPT OF STATISTICS M I TAKSAR APR 86					1/1				
UNCLASSIFIED	FSÜ-STATI	STICS-N76	95 AFOSR-	-TR-86-	-0366	F49628	-85-C- F/G 1	-8887 12/1	NL	
	> U									
			··							



MICROCOPY RESOLUTION TEST CHART NATIONAL BRIEFAL STAN AND LODGE A

1				RT DOCU	ENTATION PAG	E			
1. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			16. RESTRICTIVE MARKINGS						
28. SECURI NA	28. SECURITY CLASSIFICATION AUTHORITY NA			3. DISTRIBUTION/AVAILABILITY OF REPORT					
26. DECLASSIFICATION/DOWNGRADING SCHEDULE					- Approved for Public Release; Distribution Unlimited				
A PERFORMING ORGANIZATION REPORT NUMBER(S) FSU Technical Report No. M705					S. MONITORING ORGANIZATION REPORT NUMBER(S)				
Flori	de State	Univers	IZATION	6b, OFFICE SYMBOL (If applicable)	AFOSR/NM			<u></u>	
6c. ADDRE Depart Tallal	ss (City, State tment of hassee, F	and ZIP Con Statist L 3230	ics 6-3033	1	76. ADDRESS (City, Bldg. 410 Bolling AFB	State and ZIP Code	2-6448		
B. NAME ( ORGAN	F FUNDING	SPONSORIN	NG	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMB			UMBER	
AFC	DSR			NM_	F49020-85-0	-0007			
BC. ADDRESS (City. State and ZIP Code) Bldg. 410 Bolling AUR DC 20332 6448			10. SOURCE OF FU PROGRAM ELEMENT NO.	NDING NOS. PROJECT NO.	TASK NO.	WORK UN			
11. TITLE (Include Security Classification) Stationary Markov Sets				6.1102F	2304				
12. PERSON	AL AUTHOR	Mich	ael I. T	aksar			- 1		
Tech	nical		136. TIME C	TO	April, 1986		15. PAGE COUNT		
10. SUPPLE	COSATI	CODES		18. SUBJECT TERMS	Continue on reverse if n	eccuary a identify			
FIELD	GROUP	SUE	3. G A.					U	
19. A85TR/	ACT (Continue	on reverse i	necemary and	d identify by block numb	er)		D		
	A Marko nendert	v set i: of the '	s a rande "past" sl	om set on a rea hape given "pro	al line whose " esent". Such s	'future" sha sets appear	pe is cond in the stu	itionally dy of	
indc visi proc dist that	ribution In this cach sta	es of sp tationan does no paper v ationary merativo	pecial Ma ry then ot depend we will d w Markov c sets, h	arkov (but not the correspond: d on the choice describe all cl set which is r ov either takir	strong Markov) ing set is also of the origin osed stationar not regenerativ up a mixture of	y Markov se can be co	that is, that is, l line. ts. We wi nstructed nerative s	its Il show from two	
indc visi proc dist that spec taki loos piec	In this each sta ial regen ng a "sup ely as en es of iic	es of sp tationar does no paper w ationary merative perposit utting t d length	ve will of ve will of ve will of ve sets, h tion" of two real n and the	arkov (but not the correspond: d on the choice describe all cl set which is r by either takir two regenerati lines $\mathbb{R}_1$ and en combine them	strong Markov) ing set is also of the origin osed stationar not regenerativ ig a mixture of ve sets. Supe $\mathbb{R}_2$ with two s into one line	y Markov se cy Markov se ce can be co these rege prosition c ets M <sub>1</sub> and alternatin	ts. We winstructed nerative so an be desc M <sub>2</sub> in them	its 11 show from two ets or ribed , into rom	

20. DISTRIBUTION/AVAILABILITY OF A	BSTRACT .	21. ABSTRACT SECURITY CLASSIFICATION			
UNCLASSIFIED/UNLIMITED 🗷 SAME	AS RPT. C DTIC USERS				
228. NAME OF RESPONSIBLE INDIVIDUA	AL	225. TELEPHONE NUMBER	22c. OFFICE SYMBOL		
Frank Proschan/Myles Holl	ander	(904)644-3218	AFOSR/NM		
DD FORM 1473, 83 APR EDITION OF 1 JAN		IS OBSOLETE.	UNIX A CONTEN		
			THE REPORT OF A DATA OF A DATA OF A DATA OF A DATA		

E



SECURITY CLASSIFICATION OF THIS PAGE

5

Ļ

é

 $\mathbb{R}_1$  and  $\mathbb{R}_2.$  The union of the cut offs from  $M_1$  and  $M_2$  will be the superposition of the sets  $M_1$  and  $M_2$  .

. .

SEC

# AFOSR-TR- 86-0366

にたたたちをとうたたとうという

シー・・・ うちゅうし

Ľ

e

## STATIONARY MARKOV SETS

by

Michael I. Taksar

FSU Statistics Report M705 AFOSR Technical Report No. 86-187

April, 1986

The Florida State University Department of Statistics Tallahassee, Florida 32306-3033

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC) NOTICE OF TRANSMITTAL TO DTIC This technical report has been reviewed and is approved for public balense LAW AFR 190-18. Distribution is unlikited. MATTHEW J. KERPER Chief, Technical Information Division

Research supported by the United States Air Force Office of Scientific Research, Grant Number AFOSR F49620-85-C-0007.

> Approved for public release; distribution unlimited.

### STATIONARY MARKOV SETS

M. I. Taksar\*

Department of Statistics Florida State University Tallahassee, Florida United States

#### 1. Introduction

If one looks at the set of times when a strong Markov process visits a point in the state space, then this set is a regenerative set. It forms a replica of itself after each stopping time whose graph lies in this set. Closed regenerative sets have been studied for a long time (see Hoffman-Jørgensen [4], Maisonneuve [8], Meyer [9] and others).

Since the studies of regenerative sets were motivated by the theory of Markov processes, such sets were originally called (strong) Markov. In addition it was always supposed that any regenerative set M is a subset of the positive half-line and  $P\{0 \in M\} = 1$ .

However, if one considers visiting times of a stationary strong Markov process, then the corresponding set M is stationary, that is the probability law of the set M+t is the same as the one of M. The "natural" state space for stationary sets would be the set of closed subsets of a real line and the condition  $0 \in M$  a.s. should be dropped. The first study of such sets was done in Taksar [11]. It was shown that all such sets are in one-to-one correspondence with the\_\_\_\_\_\_ ranges (closures of the images) of the processes with independent increments having finite expectation.

\*) This research was supported by the AFOSR, Grant No. AFOSR F49620-85-C-0007.

wy Codes

Avail and for

Succial

Dit

The paper of Maisonneuve [8] gives a simple and comprehensive approach to the regenerative sets on a real line. It also gives an easy proof of the main results of [11]. Further development of the theory of regenerative sets on a real line is done in the recent work of Fitzsimmons, Frisdedt and Maisonneuve [3].

All regenerative sets have a (weak) Markov property. The "future" after time t of such set and its "past" are conditionally independent given "present". A Markov set is the set for which conditional independence of the "future" and the "past" holds, but stronger regenerative property might not be true.

Apparently, Markov sets form a larger class than regenerative sets. In a stationary case, however, the difference is not as big as one could expect. It was shown in [11] that stationary Markov sets are "almost" regenerative. There are two types of regeneration after each point t; one occurs if the point t belongs to the set and the other type of regeneration takes place if t does not belong to the set. In particular, every stationary Markov set which almost surely has Lebesgue measure zero, is regenerative, (see [11] Theorem 2).

In this paper we will describe all closed stationary Markov sets. We will show that each stationary Markov set which is not regenerative can be constructed from two special regenerative sets, by either taking a mixture of these regenerative sets or taking a "superposition" of two regenerative sets. Superposition can be described loosely as cutting two real lines  $\mathbb{R}_1$  and  $\mathbb{R}_2$  with two sets  $\mathbb{M}_1$  and  $\mathbb{M}_2$  in them, into pieces of iid length and then combine them into one line alternating pieces from  $\mathbb{R}_1$  and  $\mathbb{R}_2$ . The union of the cut offs from  $\mathbb{M}_1$  and  $\mathbb{M}_2$  will be the superposition of the sets  $\mathbb{M}_1$  and  $\mathbb{M}_2$ .

The paper is structured as follows. In section 2 we give definitions and formulate the main results. In section 3 we establish the main properties of stationary Markov sets. Section 4 studies the operation which transforms a stationary Markov set into a stationary regenerative set. Section 5 analyses those stationary Markov sets which are neither regenerative nor are mixtures of

2

regenerative sets. In section 6 we study the "residual life" process associated with the stationary Markov set, and find its stationary distribution. The last section is devoted to reversability properties. We outline a necessary and sufficient condition for the set -M to have the same distribution as M.

# 2. Basic definition. Formulation of the main result.

In our definition and notations we follow Maisonneuve [8] and Fitzimmons, Fristedt, and Maisonneuve [3]. Let  $\Omega^{\circ}$  be the set of all closed sets in R. For each  $\omega^{\circ} \in \Omega^{\circ}$  and t  $\in \mathbb{R}$  put (assuming inf  $\emptyset = -$ , sup  $\emptyset = --$ )

 $\begin{aligned} d_{t}(\omega^{\circ}) &\stackrel{\Delta}{=} \inf\{s > t: s \in \omega^{\circ}\}, & \ell_{t}(\omega^{\circ}) \stackrel{\Delta}{=} \sup\{u < t: u \in \omega^{\circ}\} \\ r_{t}(\omega^{\circ}) &\stackrel{\Delta}{=} d_{t}(\omega^{\circ}) - t, & n_{t}(\omega^{\circ}) \stackrel{\Delta}{=} t - \ell_{t}(\omega^{\circ}) \\ \tau_{t}(\omega^{\circ}) \stackrel{\Delta}{=} \{s - t: s \ge t, s \in \omega^{\circ}\} \\ c_{t}(\omega^{\circ}) \stackrel{\Delta}{=} \{s - t: s \le t s \in \omega^{\circ}\} \end{aligned}$ 

Let  $G^{\circ}$  ( $G_{t}^{\circ}$  respectively) be the  $\sigma$ -field generated by all functions  $d_{s}^{\circ}$ ,  $s \in \mathbb{R}$ ( $s \leq t$  respectively). Let  $J^{\circ}$  ( $J_{t}^{\circ}$  respectively) be the  $\sigma$ -field generated by all functions  $\ell_{u}^{\circ}$ ,  $u \in \mathbb{R}$  ( $u \geq t$  respectively). It is easy to see that  $G_{t}^{\circ}$  is an increasing and  $J_{t}^{\circ}$  is a decreasing filtration and  $J^{\circ} = G^{\circ}$ .

A closed random set M on a space  $(\Omega, F)$  is a measurable mapping of  $(\Omega, F)$  into  $(\Omega^{\circ}, G^{\circ})$ .

In this paper we will deal only with closed random sets, so in the sequel we will not write "closed" each time. Put

$$\begin{split} & \overset{\mathsf{D}}{}_{\mathsf{t}} \stackrel{\Delta}{=} \overset{\mathsf{d}}{}_{\mathsf{t}} \circ \mathsf{M}, \quad \mathsf{R}_{\mathsf{t}} \stackrel{\Delta}{=} \overset{\mathsf{r}}{}_{\mathsf{t}} \circ \mathsf{M} \\ & \overset{\mathsf{L}}{}_{\mathsf{t}} \stackrel{\Delta}{=} \overset{\mathsf{L}}{}_{\mathsf{t}} \circ \mathsf{M}, \quad \mathsf{N}_{\mathsf{t}} \stackrel{\Delta}{=} \overset{\mathsf{n}}{}_{\mathsf{t}} \circ \mathsf{M} \\ & \mathsf{M}^{\mathsf{t}} \stackrel{\Delta}{=} \overset{\mathsf{T}}{}_{\mathsf{D}} \circ \mathsf{M}, \quad \mathsf{M}_{\mathsf{t}} \stackrel{\Delta}{=} \overset{\mathsf{r}}{}_{\mathsf{L}} \circ \mathsf{M} \end{split}$$

It is obvious that all the mappings  $D_t$ ,  $R_t$ ,  $L_t$  and  $N_t$  are measurable and so are  $M^t$  and  $M_t$ .

3

Let  $(\Omega, F, P)$  be a complete probability space and M be a random set on this space. Let G,  $G_t$  and  $J_t$  be the preimages in F of the  $\sigma$ -fields  $G^\circ$ ,  $G_t^\circ$  and  $J_t^\circ$  under the mapping M.

(2.1) A set M is called right Markov (r.M.) if for any two bounded measurable functions f and g on  $(\Omega^0, G^0)$ 

 $P\{f(Mn[t,\infty[)g(Mn]-\infty,t]|D_t\}=P\{f(Mn[t,\infty[)|D_t\}P\{g(Mn]-\infty,t])|D_t\}.$ 

(2.2) A set M is called left Markov (1.M.) if for any two bounded measurable functions f and g on  $(\Omega^{\circ}, G^{\circ})$ 

$$P\{f(M\cap[t,\infty[)g(M\cap]-\infty,t)|L_{t}\} = P\{f(M\cap[t,\infty[)|L_{t}\} P\{g(M\cap]-\infty,t])|L_{t}\}.$$

For brevity here and in sequel we write equations with conditional expectations without adding a.s. after equalities. Given a random set M, we denote by M + s the set {t+s: t  $\in M$ }.

(2.3) A set M is called stationary if for any bounded measurable function f on  $(\Omega^{\circ}, G^{\circ})$  and any  $s \in \mathbb{R}$ 

$$P{f(M+s)} = P{f(M)}.$$

Our aim is to describe all stationary r.M. sets. We will need results from the theory of regenerative sets. The precise notion of regenerative set used in this paper is due to Maisonneuve [8].

(2.4) A random set M is right regenerative (r.r.) if there exists a measure P<sub>0</sub> on  $(\Omega^{\circ}, \mathbb{C})$  such that for each  $f \in b\mathbb{C}^{\circ}$  (set of bounded  $\mathbb{C}^{\circ}$ -measureable functions)

$$P\{f \in M^{\mathsf{L}} | G_{\mathsf{t}}\} = P_0\{f\} \text{ on } \{D_{\mathsf{t}} < \infty\}.$$

Following [8], the measure  $P_0$  is called the law of (right) regeneration of M.

(2.5) A set M is left regenerative (1.r.) if there exists a measure  $P^0$  on  $(\Omega^{\circ}, G^{\circ})$  such that for each  $f \in bG^{\circ}$ 

 $P\{f \circ M_t | J_t\} = P^0\{f\} \text{ on } \{L_t > -\infty\}.$ 

In the sequel for brevity, we will use the term regenerative (r.) and Markov (M) instead of right regenerative and right Markov respectively.

Increasing processes with independent increments (subordinators) play an important role in the description of regenerative sets and, as we will see in the sequel, stationary Markov sets as well. Each subordinator z is characterized by a constant  $\alpha < 0$  and a measure  $\Pi$  on ] 0,  $\infty$ [. We call such a subordinator an  $(\alpha, \Pi)$ -process. Let  $z_t(\omega)$ ,  $t \ge 0$ , be a stochastic process on a probability space  $(\Omega, F, P)$ . The image M of this process is defined as

$$M(\omega) = \overline{z_{\mathbb{R}_{+}}(\omega)}$$

where bar above the set stands for closure. If z is a subordinator, then the image of z is a right regenerative set. If z is a decreasing process with independent increments then the image of z is a left regenerative set.

Let us recall the main results of [8] and [11] regarding stationary regenerative sets. There is one-to-one correspondence between all stationary r.r. sets M and all pairs  $(\alpha, \Pi)$  defined up to proportionality, where  $\alpha$  and  $\Pi$  are characteristics of a subordinator subject to

 $\int_0^\infty x^{JI} (dx) < \infty .$ 

The stationary set M which corresponds to the pair  $(\alpha, \Pi)$  is called  $(\alpha, \Pi)$ -generated. Any stationary r.r. set M is also l.r. Moreover the set -M has the same distribution in  $(\Omega^{\circ}, G^{\circ})$  as M.

Since the definition of r.M. set is weaker than that of r.r. set, any r.r. set is r.M., however the opposite is not true.

An example of a stationary r. M. set which is not r.r. was constructed in [11]. Any mixture a  $(0, \Pi)$ -generated set and a real line  $\mathbb{R}$  with "weights" 0 and <math>q = 1 - p is a r. M. set but not a r.r. set.

**DEFINITION.** Right Markov sets of the first type are right regenerative sets. Right Markov sets which can be represented as a mixture of a  $(0, \Pi)$ -generated and a real line are called r.M. sets of the second type. Right Markov sets which are neither of the first or the second type are called right Markov sets of the third type.

Markov processes provide good examples of different types of stationary Markov sets. If  $x_t$  is a strong Markov process and b is a point in the state space then the "visiting set"

$$M = \{t: x_t = b\}$$

is regenerative and if in addition  $x_t$  is stationary, then M is stationary.

To obtain a Markov set of the second type, consider a strong Markov process  $x_t^1$ , for which  $P\{x_t^1 = b\} = 0$  for each t, but point b is not a polar set and a process  $x_t^2$  which stays deterministically at the point b. The mixture  $x_t$  of the processes  $x_t^1$  and  $x_t^2$  will be a Markov (but not a strong Markov) process. The visiting times of b by  $x_t$  is a Markov set of the second type, and if  $x_t^1$  is stationary then so is the visiting times set.

To give an example of a Markov set of the third type, consider a particle moving on the positive half line according to a diffusion law. An infinitely thin elastic screen is placed at the origin. The particle is reflected from this screen until time

 $\tau = \{ \inf t : :_{t} \ge S \}$ 

where  $f_t$  is the local time at zero of the reflected diffusion and S is a random variable with exponential distribution independent of the process  $x_t$ . At the moment  $\tau$  the particle moves to the other side of the screen where it stays for time X, where X is another exponential random variable independent of x. and S. At the time  $X + \tau$  the partical is placed back to a random point on the positive half line and the whole process starts anew. The closure of the set of times when this particle visits the origin is a Markov set of the third type. If this Markov

process is stationary (which can be easily achieved, provided that there exists a constant downward drift, or there exists a reflecting upper barrier)then this Markov set is stationary.

In the remainder of this section we define rigorously the superposition of two regenerative sets and formulate the main result.

Let  $\Pi$  be a measure on  $]0,\infty]$  and  $\mu$  be a probability measure on  $[0,\infty[$  and  $\lambda$  and  $\alpha$  be two positive constants. Let  $y_t$  be a  $(0, \Pi)$ -process and  $\{S_k\}$ ,  $k = 1, 2, \ldots$ ,  $\{X_k\}$  and  $\{Y_k\}$ ,  $k = 0, 1, 2, \ldots$  be three sequences of iid random variables, independent of  $y_t$  and independent of each other. The distributions of  $S_i$  and  $X_k$  are exponential with parameters  $\alpha$  and  $\lambda$  respectively. The distribution of  $Y_j$  is given by  $\mu$ . Consider a subordinator  $x_t$  of a pure jump type constructed in the following manner (we assume below  $\sigma_0 = 0$ )

$$\sigma_{k} - \sigma_{k-1} = S_{k}, \quad k = 1, 2, ...$$

$$x_{\sigma_{k}} - x_{\sigma_{k}} = Y_{k} + X_{k}, \quad k = 1, 2, ...$$

$$x_{s} = x_{u} \quad \text{if} \quad \sigma_{k} \le s \le u < \sigma_{k+1}, \quad k = 0, 1, ...$$
(2.6)

Put

$$Z_{t} = y_{t} + x_{t}$$

$$L = \bigcup_{k=1}^{\infty} \{x : \mathbb{Z}_{\sigma_{k}} \le x \le \mathbb{Z}_{\sigma_{k}} + x_{k}\}$$

$$M = \overline{\mathbb{Z}_{\mathbb{R}_{+}} \cup L}$$

$$(2.8)$$

The set M defined by (2.8) is called  $(\Pi, \alpha, \lambda, \mu)$ -set. (Note that there are many  $(\Pi, \alpha, \lambda, \mu)$ -sets corresponding to different initial distributions of the process  $y_t$ ).

Let  $\mu'$  be the restriction of  $\mu$  on  $]0, \infty[$ . We say that quadruple  $(\Pi, \alpha, \lambda, \mu)$  is equivalent to  $(\Pi_{1}, \alpha_{1}, \lambda_{1}, \mu_{1})$  if there exists a constant c such that

7

$$(\Pi, \alpha) = c(\Pi_1, \alpha_1)$$
(2.9)

$$\nu' - \nu_{1}' = \frac{\nu\{0\} - \nu_{1}\{0\}}{\pi (\mathbb{R}_{+})} \pi$$
(2.10)

$$\lambda(1 - \alpha \mu\{0\} / (\alpha + \Pi(\mathbf{R}_{+})) = \lambda(1 - \alpha_{1} \mu_{1}\{0\} / (\alpha_{1} + \Pi_{1}(\mathbf{R}_{+}))$$
(2.11)

In particular, when  $\Pi$  is an infinite measure, equivalency of  $(\Pi, \alpha, \lambda, \mu)$  and  $(\Pi_1, \alpha_1, \lambda_1, \mu_1)$  means proportionality of  $(\Pi, \alpha)$  and  $(\Pi_1, \alpha_1)$  and equality of  $(\lambda, \mu)$  and  $(\lambda_1, \mu_1)$ .

It is easy to see that if  $\Pi(\mathbb{R}_{+}) = \infty$  and quadruples  $(\Pi, \alpha, \lambda, \mu)$  and  $(\Pi_{1}, \alpha_{1}, \lambda_{1}, \mu_{1})$ are equivalent then every  $(\Pi, \alpha, \lambda, \mu)$ -set is a  $(\Pi_{1}, \alpha_{1}, \lambda_{1}, \mu_{1})$ -set as well. In fact, if we construct processes x, y and Z by (2,6) and (2,7), then processes  $x_{t}' = x_{ct}$ ,  $y_{t}' = y_{ct}$  and  $Z_{t}' = Z_{ct}$  generate the same set M given by (2.8). However, the Levi's measure of the process  $x_{ct}$  is c $\Pi$  and the rate of jumps of the process  $y_{ct}$  is c $\alpha$ , which shows that  $(\Pi, \alpha, \lambda, \mu)$ -set is  $(c\Pi, c \alpha, \lambda, \mu)$ -set as well.

If  $\Pi$  is a finite measure then both processes  $x_t$  and  $y_t$  have jumps governed by Poisson processes with rates a and  $\Pi(\mathbb{R}_+)$  respectively. In particular

$$p \stackrel{\Delta}{=} P\{y_{\sigma_1} = y_0\} = \alpha / (\alpha + \Pi(\mathbb{R}_+))$$

(see (2.6) for definition of  $\sigma_1$ ). The set M given by (2.7) consists of the intervals of L and discrete points of the image of Z. The length of the first interval  $I_1$  of  $\downarrow$  is equal to  $X_1 + X_2 + \ldots + X_N$  where N has geometric distribution with parameter  $p_{-1}(0)$ . Thus the distribution of  $I_1$  is exponential with parameter  $\lambda(1 - p_{-1}(0))$ . The distribution of the interval  $J_1$  which is contingent to  $I_1$  in M

from the right (i.e.,  $\inf\{t:t\in J_1\} = \sup\{s:s\in I_1\}$ ) has distribution  $\mu' + (\mu\{0\}/\Pi(\mathbb{R}_+))\Pi$ (note that  $(\Pi(\mathbb{R}_t))^{-1}\Pi$  is the distribution of the jumps of the process y). Likewise for any other interval  $I_k$  in L and contingent to  $I_k$  interval  $J_k$ . The distribution of any interval contingent to M which does not coincide with any of  $J_k$  is equal to the distribution of jumps of y, i.e. to  $(\Pi(\mathbb{R}_t))^{-1}\Pi$ . From the above it is easy to show that if M is a  $(\Pi, \alpha, \lambda, \mu)$ -set and  $(\Pi, \alpha, \lambda, \mu)$  is equivalent to  $(\Pi_1, \alpha_1, \lambda_1, \mu_1)$  then there exists a  $(\Pi_1, \alpha_1, \lambda_1, \mu_1)$ -set whose distribution is the same as that of M.

8

**DEFINITION.** A random set M is called  $(\Pi, \alpha, \lambda, \mu)$ -generated if for each t there exists a random variable  $\phi_t$ , such that  $\phi_t \ge t$  a.s. and  $M \cap [\phi_t, \infty]$  has the same distribution as a  $(\Pi, \alpha, \lambda, \mu)$ -set. In this case the quadruple  $(\Pi, \alpha, \lambda, \mu)$  is called the generator of the set M.

The next two theorems give the main result of this paper.

(2.12) **THEOREM.** Every stationary r.M. set M of the third type is  $(\Pi, \alpha, \lambda, \mu)$ -generated. The generator of M is unique up to equivalency and is subject to

 $\int_{0}^{\infty} x \Pi (dx) < \infty$  (2.13)

$$\int_{0}^{\infty} x \mu(dx) < \infty$$
 (2.14)

Each quadruple  $(\Pi, \alpha, \lambda, \mu)$  subject to (2.13) and (2.14) is a generator of a unique stationary right Markov set.

Let  $\delta_a$  denote a unit measure concentrated at point a.

(2.15) THEOREM <u>A stationary r.M. set M of the third type is left Markov iff</u> its generator  $(\Pi, \alpha, \lambda, u)$  is equivalent to  $(\Pi, \alpha, \lambda, \delta_0)$ . In this case the set -M has the same distribution as M.

In the diffusion example presented above the set of visiting times of 0 becomes a left Markov set when the diffusion process is made continuous. That can be done if at the time  $\pi + \lambda$  the particle is moved on the other side of the elastic screen and starts again moving according to the original reflected diffusion law.

3. General properties of stationary Markov sets.

Here and in the sequel we will deal only with those stationary Markov sets which are a.s. nonempty. This is equivalent to

$$P\{D_{+} < \infty\} = 1 \quad \text{for all } t \in \mathbb{R}. \tag{3.1}$$

The following proposition was proved in [11] (see Lemma 7.3).

(3.2) **PROPOSITION.** If M is stationary Markov set then for each function  $f \in bG^{\circ}$ there exist two constants a and b such that for each t

$$P\{f \circ M^{t} | G_{t}\} = a 1_{D_{t} > t} + b 1_{D_{t} = t}$$

For brevity we will denote indicator functions of  $]-\infty,t[, ]-\infty,t]$ ,  $[t,\infty[, ]t,\infty[$  by  $l_{<t}, l_{\leq t}, l_{\geq t}$  and  $l_{>t}$  respectively.

The following corollary is a simple consequence of Propostion (3.2).

(3.3) COROLLARY. If M is a stationary Markov set then there exist two measures  $P_{0} \text{ and } P_{1} \text{ on } (\Omega^{\circ}, G^{\circ}) \text{ such that for each } f \in bG^{\circ}$   $P\{f \circ M^{t} | G_{t}\} = 1_{<t} (D_{t}) P_{0}\{f\} + 1_{t} (D_{t}) P_{1}\{f\}.$ (3.4)

Let  $\hat{M}$  denote the set of all points of M which belong to M with its right neighborhood.

(3.5) **PROPOSITION.** For each  $f \in bG^{O}$  and any stopping time T with respect to the filtration  $G_{t+}$ 

$$P\{f \circ M^{T} | G_{T+}\} = 1_{T \in \widehat{M}} P_{0}\{f\} + 1_{T}(\widehat{M}) P_{1}\{f\}$$
(3.6)

 $P_{teed}$ . Usual arguments show that Proposition (3.3) remains true if t in (3.4) is replaced by any stopping time with respect to  $G_t$ , taking finite or countable number of values.

It is sufficient to prove (3.6) for f of the form

$$f = g(r_{s_1}, r_{s_2}, ..., r_{s_k})$$

where g is a bounded continuous function of k variables. For such f the function  $f \circ M^{\tt t}$  is continuous in t and

$$P\{f \circ M^{T} | G_{T_{+}}\} = \lim_{n \to \infty} P\{f \circ M^{T_{n}} | G_{T_{n}}\} =$$

$$\lim_{n \to \infty} [1_{>T_{n}} (D_{T_{n}}) P_{0}\{f\} + 1_{T_{n}} (D_{T_{n}}) P_{1}\{f\}]$$
(3.7)

where  $T_n$  is any sequence of stopping times, taking on finite or countable number of values and such that  $T_n \neq T$ .

Put

$$ae_n(x) \stackrel{\Delta}{=} k 2^{-n}$$
, if  $(k-1)2^{-n} \le x \le k 2^{-n}$  (3.8)

and let (assuming  $\inf \emptyset = +\infty$ )

 $T_n^+ = \inf \{ a_n^-(s) : s \ge T, u \in M \text{ for all } s \le u \le a_n^-(s) \} .$ 

The random variable  $T_n^+$  is a stopping time (see [2], Ch VI) and so is

$$T_{n} = I_{T}(\hat{M}) a e_{n}(T) + I_{TZM} T_{n}^{\dagger}$$
, (5.9)

Each  $T_n$  given by (3.9) takes at most a countable number of values and  $T_n + T$ . By the construction  $P_{T_n} > T_n$  on the set  $\{T \in \hat{M}\}$  and  $\{T_n = P_{T_n}\}$  converges to the set  $\{T \in \hat{M}\}$ . Hence we can pass to a limit in (3.7) and obtain (3.6). (3.10) **PROPOSITION.** For each  $f \in bG^{\circ}$  and each stopping time T with respect to  $G_{t+}^{\circ}$  and each i = 0, 1,

$${}^{P}{}_{i} \{ f \circ \tau_{d_{T}} | G_{T+} \} = {}^{1}{}_{T \bar{\epsilon} \hat{\omega}^{\circ}} {}^{P}{}_{0} \{ f \} + {}^{1}{}_{T} (\hat{\omega}^{\circ}) {}^{P}{}_{1} \{ f \}$$
(3.11)

The proof is similar to the proof of previous proposition.

From now on we will consider only stationary sets of the third type, for which

$$P\{D_{+} = t\} > 0. \tag{3.12}$$

(Theorem 2 of [11] shows that failure of (3.12) implies that M is regenerative.)

(3.13) **PROPOSITION.** For each t

$$P\{D_{+} = t, t \in \tilde{M}\} = 0.$$
 (3.14)

*Proof.* Suppose the left hand side of (3.14) is equal to  $\varepsilon > 0$ . By virtue of Proposition (3.5)

$$P\{f \circ M^{t} | G_{t+}\} = P_{0}\{f\} \text{ on } \{D_{t} = t, t \in \hat{M}\}.$$
(3.15)

On the other hand, using sequentially (3.4) and (3.15)

(3.16)

$$P\{f | M^{t} | G_{t}\} = P_{1}\{f\} = [\epsilon P_{0}\{f\} + P\{D_{t}=t, t \in \hat{W}\}P_{1}\{f\} ]/P\{D_{t}=t\} \text{ on } \{D_{t}=t\},$$

Equality (3.16), which is true for each f, shows  $P_0 = P_1$ , which contradicts the assumption that M is the set of the third type.

(3.17) **COROLLARY.**  $P_1 \{ 0 \in \hat{\omega}_0 \} = 1.$ 

Precé. By proposition (3.13) the sets  $\{D_t = t\}$  and  $\{t \in \hat{M}\}$  are indistinguishable. Using (3.4),

$$P\{D_t = t\} = P\{D_t = t, 0 \in \hat{M}^t\} = P\{D_t = t\}P_1\{0 \in \hat{\omega}_0\}.$$

Thus, the statement follows from (3.12).

(3.18) **PROPOSITION.** For any functions  $f \in bG^{\circ}$  and  $g \in bG^{\circ}_{t+}$  such that g = 0 on  $\{d_t = \infty\}$  and each i = 0, 1

$$P_{i} \{f_{\circ}\tau_{d_{t}} g\} = P_{i} \{g; d_{t} > t\} P_{0} \{f\} + P_{i} \{g; d_{t} = t\} P_{1} \{f\}.$$
(3.19)

Prec: For i = 1. Put T = t + s. By (3.4) and (3.12)

$$P_{1} \{ \mathbf{f} \circ \tau_{d_{t}} g \} = P \{ \mathbf{f} \circ M^{T} g \circ M^{S} \mid D_{s} = s \} / P \{ D_{s} = s \}.$$

Taking first conditional expectation with respect to  $G_{s+t}^{\prime}$ ,

$$P_{1} \{f \circ \tau_{d_{t}} g\} = P\{g \circ M^{S} 1_{$$

which is equivalent to  $(3.1^{\circ})$ .

Let -

$$\begin{split} \tilde{n}_{t} &\stackrel{\Delta}{=} \inf\{s > t \colon s \in \hat{\omega}^{\circ}\}, \qquad n_{t} \stackrel{\Delta}{=} \tilde{n}_{t} \circ M, \\ \tilde{\gamma}_{t} &\stackrel{\Delta}{=} \inf\{s > \tilde{n}_{t} \colon s \in \hat{\omega}^{\circ}\}, \qquad \gamma_{t} \stackrel{\Delta}{=} \tilde{\gamma} \circ M, \\ \tilde{\gamma}_{t} &\stackrel{\Delta}{=} \inf\{s > \tilde{\gamma}_{t}, s \in \hat{\omega}^{\circ}\}, \qquad \gamma_{t} \stackrel{\Delta}{=} \tilde{\gamma}_{v} \circ M, \\ \tilde{\gamma}_{t} &\stackrel{\Delta}{=} \inf\{s > \tilde{\gamma}_{t}, s \in \hat{\omega}^{\circ}\}, \qquad \gamma_{t} \stackrel{\Delta}{=} \tilde{\gamma}_{t} \circ M, \\ \tilde{\eta} &\stackrel{\Delta}{=} \tilde{\eta}_{0}, \quad \tilde{\gamma} \stackrel{\Delta}{=} \tilde{\gamma}_{0}, \quad \tilde{\gamma} \stackrel{\Delta}{=} \tilde{\gamma}_{0}, \qquad n \stackrel{\Delta}{=} \gamma_{0}, \quad \gamma \stackrel{\Delta}{=} \gamma_{0}. \end{split}$$
(3.20)

(5.21) **PROPOSITION.** For  $\tilde{y}$  and  $\tilde{r}$  defined by (3.20)

$$P_1(\tilde{\tau}_i = 0) = 1.$$
 (3.22)

and there exists a constant  $0 < x < \infty$  such that for each a

$$P_1\{\tilde{\gamma} > a\} = e^{-\lambda a} . \qquad (3.23)$$

Proof. (3.22) follows from (3.17). Let a, b > 0. Applying Proposition (3.18),

$$P_{1}\{\tilde{\gamma}>a+b\}=P_{1}\{\tilde{\gamma}>a,\tilde{\gamma}\circ\tau_{a}>b\}=P_{1}\{\tilde{\gamma}>a,d_{a}=a\}P_{1}\{\tilde{\gamma}>b\}+P_{1}\{\tilde{\gamma}>a,d_{a}>a\}P_{0}\{\tilde{\gamma}>b\} \quad (3.24)$$

If  $\tilde{\gamma} > a$  then  $a \in \hat{\omega}^{\circ}$  and  $d_a = a$ . Thus  $P_1\{\tilde{\gamma} > a, d_a > a\} = 0$  and (3.24) equals to  $P_1\{\tilde{\gamma} > a\} P_1\{\tilde{\gamma} > b\}$  whereas (3.23) follows.

Suppose (3.23) equals 1. Then  $P_1\{\mathbb{R}_+ \subset \omega^0\} = 1$ . The latter would imply that M is a mixture of a real line  $\mathbb{R}$  and a regenerative set with the law of regeneration  $P_0$ . This contradicts the assumption that M is a set of the third type. Likewise, if (3.23) equals 0, then this would imply that  $P\{d_a = a\} = 0$ . The latter is with a contradiction to (3.12).

Let  $\tilde{n}_+$ ,  $\tilde{\gamma}_+$ , ... etc. be given by (3.20). Define

$$\tilde{n}(0,t) \stackrel{\Delta}{=} \tilde{\gamma}(0,t) \stackrel{\Delta}{=} \tilde{v}(0,t) \stackrel{\Delta}{=} t , \qquad (3.25)$$

$$\tilde{n}(k,t) \stackrel{\Delta}{=} \tilde{n}_{\tilde{v}}(k-1,t) , \qquad (3.25)$$

$$\tilde{\gamma}(k,t) \stackrel{\Delta}{=} \tilde{\gamma}_{\tilde{v}}(k-1,t) , \qquad k=1, 2, ...$$

$$n(k,t) \stackrel{\Delta}{=} \tilde{n}(k,t) \circ M$$

$$\gamma(k,t) \stackrel{\Delta}{=} \tilde{\gamma}(k,t) \circ M$$

$$v(k,t) \stackrel{\Delta}{=} \tilde{v}(k,t) \circ M.$$

When t is fixed we will write for brevity  $\tilde{n}(k)$ ,  $\tilde{\gamma}(k)$ ,  $\gamma(k)$ , etc. instead of  $\tilde{n}(k,t)$ ,  $\tilde{\gamma}(k,t)$ ,  $\gamma(k,t)$ , etc.

The points n(k) and  $\gamma(k)$  mark the beginnings and the ends of the intervals which  $\hat{M} \cap [t, \infty[$  is composed of.

(3.26) **PROPOSITION.** The sequence  $\{(\gamma(k) - \eta(k), \nu(k) - \gamma(k), \eta(k+1) - \gamma(k))\}$  is a sequence of iid three-dimensionals vectors on  $(\Omega, F, P)$ . The sequences  $\{\gamma(k) - \eta(k)\}$  and  $\nu(k) - \gamma(k)\}$  are independent and for any a > 0

$$P{\gamma(k) - \eta(k) > a} = e^{-\lambda a},$$
 (3.27)

where  $\lambda$  is the same as in Proposition (3.21).

<u>The sequence</u> { $(\tilde{\gamma}(k) - \tilde{n}(k), \tilde{\nu}(k) - \tilde{\gamma}(k), \tilde{n}(k+1) - \tilde{\nu}(k))$ } <u>is a sequence of iid</u> <u>three-dimensional vectors on</u>  $(\Omega^{\circ}, G^{\circ}, P_{i})$ , i = 0, 1. <u>The sequences</u> { $\tilde{\gamma}(k) - \tilde{n}(k)$ } <u>and</u> { $\tilde{\nu}(k) - \tilde{\gamma}(k)$ } <u>are independent and for any</u> a > 0 <u>and any</u> i = 0, 1

$$P_{i}\{\tilde{\gamma}(k) - \tilde{\eta}(k) > a\} = e^{-\lambda a}.$$

 $p_{\text{toc}_0}$ . It follows from [2] Ch. VI that for each k the random variables n(k),  $\gamma(k)$  and  $\nu(k)$  are stopping times and if k < j then

$$v(j) \leq n(k) < \gamma(k) \leq v(k)$$
.

Let h be any bounded function of three variables. Since  $n(k) \in \hat{M}$ , using Proposition (3.5),

$$P\{h(\gamma(k)-\eta(k),\nu(k)-\gamma(k),\eta(k+1)-\nu(k)) \mid G_{\eta(k)+}\} = P_{1}\{h(\tilde{\gamma},\tilde{\nu}-\tilde{\gamma},\tilde{\eta}(2,0)-\tilde{\nu})\}.$$

The above shows independence of  $(\gamma(k) - \eta(k), \nu(k) - \gamma(k), \eta(k+1) - \nu(k))$  from the sequence  $\{\gamma(j) - \eta(j), \nu(j) - \gamma(j), \eta(j+1) - \nu(j)\}, j = 1, 2, ..., k - 1.$ 

Let g be a bounded function of one variable. Put  $f(\omega^0) = g(\tilde{\nu} - \tilde{\gamma})$ . Then using Proposition (3.5)

$$P\{g(v(k) - y(k)) | 1_{>b} (y(k) - n(k))\}$$

$$= P\{1_{>b}(y(k) - n(k))f_{\circ}\tau_{n(k)+b} \circ M\}$$

$$= P\{1_{>b}(y(k) - n(k))\}P_{1}\{f\}.$$
(5.28)

The last equality in (3.28) is due to

$$\{\gamma(k) - \eta(k) > b\} \in \{\eta(k) + b \in \mathbb{A}\}.$$

and (5.4). Likewise, setting  $h(\omega^0) = 1_{>b}(\tilde{\gamma})$ 

$$P{\gamma(k) - \eta(k) > b} = P{h \circ \tau_{\eta(k)} \circ M} = P_1{h} = P_1{\tilde{\gamma} > b}$$

and (3.27) follows from (3.23).

The proof of the second part of the proposition is done in a similar manner.

# 4. Deletion Operation and its Properties.

In this section we define an operator which removes parts of the set M in such a way that M becomes a regenerative set. Put

$$K(\omega^{\circ}) \stackrel{\Delta}{=} \omega^{\circ} \setminus \operatorname{closure}(\hat{\omega}^{\circ}) = \lim_{t \to -\infty} \omega^{\circ} \setminus \bigcup_{k=1}^{\infty} [\tilde{n}(k,t), \tilde{\gamma}(k,t)]$$

$$K(\omega^{\circ}) \stackrel{\Delta}{=} \overline{K(\omega^{\circ})}. \qquad (4.1)$$

The operator K removes closure of the interior of  $\omega^o$ , and the remaining set has no interior. Thus

$$\widehat{K(\omega^{\circ})} = \emptyset$$
.

(4.2) **PROPOSITION.** For any  $\omega^0$  and any t

$$\mathbf{d}_{\dagger} \circ \mathbf{K}(\omega^{\mathrm{o}}) \ \overline{\epsilon} \ \hat{\omega}^{\mathrm{o}} \ . \tag{4.3}$$

*Proci*. Suppose (4.3) is wrong, then for some  $k \ge 1$ 

$$\mathbf{d}_{+} \circ \mathbf{K}(\boldsymbol{\omega}^{0}) \in [\tilde{\mathbf{n}}(\mathbf{k}, \mathbf{t}), \, \tilde{\mathbf{\gamma}}(\mathbf{k}, \mathbf{t})] \tag{4.4}$$

Since  $d_t \in K(\omega^2) + K(\omega^2)$ , the only way that (4.4) can be true is

$$\mathbf{d}_{+} \circ \mathbf{K}(\boldsymbol{\omega}^{\mathsf{O}}) = \tilde{\eta}(\mathbf{k}, \mathbf{t}) \quad . \tag{4.5}$$

If  $\tilde{\pi}(k,t) = t$  then (4.5) fails because in this case  $d_t \in K(\omega^2) = \tilde{\tau}(k,t) \oplus \tilde{\pi}(k,t)$ . If  $t \in \tilde{\pi}(k,t)$  then (4.5) implies  $[]t, \tilde{\pi}(k,t)[ \in K(\omega^2)$ . Thus  $\tilde{\pi}(k,t) \in K(\omega^2)$  which contradicts (4.5).

(4.6) THEOREM. The set  $k \circ M$  is a stationary regenerative set.

Preof. From a trivial relation

$$\widehat{\omega^{0}+s} = \widehat{\omega}^{0}+s$$

it follows that

$$K \circ M + s = K(M + s).$$
 (4.7)

Likewise

$$\tau_{t} \circ K \circ M = K \circ \tau_{t} \circ M \quad . \tag{4.8}$$

Relation (4.7) shows that stationarity of M implies stationarity of  $K \circ M$ .

Put  $D_t = d_t \circ K \circ M$ . Then  $D_t'$  is a stopping time. By virtue of Proposition (3.5), Proposition (4.2) and (4.8), for any function  $f \in bG^\circ$ 

$$P\{\mathbf{f} \circ \tau_{D_{\mathbf{t}}} \circ K \circ M | \mathcal{G}_{D_{\mathbf{t}}}\} = P_{\mathbf{0}}\{\mathbf{f} \circ K \circ \tau_{D_{\mathbf{t}}} \circ M | \mathcal{G}_{D_{\mathbf{t}}}\} = P_{\mathbf{0}}\{\mathbf{f} \circ K\}$$
(4.9)

This proves that K ° M is regenerative with the law of regeneration

$$P = P_0 \circ K^{-1}$$
 (4.10)

(4.11) **REMARK.** The proof of Theorem (4.6) shows that  $K \circ M$  is regenerative with respect to the filtration  $G_t \stackrel{\Delta}{=} G_{D_t^{+}}$  which is larger than the natural filtration generated by  $K \circ M$ . It can be shown in a similar manner that

$${}^{P}_{0} \{ \mathbf{f} \circ \tau_{\mathbf{d}_{t}} \circ \mathbf{K} | G_{\mathbf{d}_{t}^{+}}^{0} \} = P \{ \mathbf{f} \circ \tau_{\mathbf{d}_{t}} | G_{\mathbf{d}_{t}^{+}}^{0} \} = P \{ \mathbf{f} \} .$$
 (4.12)

We will call  $K \circ M$  the regenerative part of the set M. By [6] the set  $K \circ M$  is either perfect or discrete.

According to [7] and [11] their exists a process  $z_0$  with independent increments such that  $K(\omega^0) = \overline{z_{R_1}}$  for  $P_0$  a.a.  $\omega^0$  and such that the local time

$$\theta_{s} = z_{s}^{-1} \stackrel{\Delta}{=} \inf\{t: z_{t} \ge s\}$$
(4.13)

is a continuous process adapted to the  $\sigma$ -field  $G_{t+}^{o}$  and for any  $u \in z_{\mathbf{R}}$ 

$$\theta_{u+s} = \theta_u + \theta_s \circ \tau_u . \qquad (4.14)$$

(4.15) **PROPOSITION.** If M is a stationary Markov set with a perfect regenerative part then

 $P_0 \{ 0 \in \hat{\omega}^0 \} = 0$ .

Proof. Put

$$T_n'(\omega^\circ) = \inf\{ae_n(s): s \ge 0, u \in \omega^\circ \text{ for all } s < u \le ae_n(s)\}$$

where  $x_n(s)$  is given by (3.8). Let

$$T_{n}(\omega^{O}) = \begin{cases} 0 \text{ if } 0 \in \hat{\omega}^{O} \\ T_{n}'(\omega^{O}) \text{ otherwise} \end{cases}$$

Then  $T_n$  is a sequence of stopping times such that

$$T_n = D_{T_n} = 0 \text{ on } \{0 \in \hat{\omega}^0\}.$$
 (4.16)

From [6] and [8] it follows that for any perfect regenerative set with the law of regeneration P

$$P\{0 \text{ is an isolated point in } \omega^0\} = 0$$
. (4.17)

On the other hand  $K(\omega^{\circ})$  and  $\omega^{\circ}$  coincide in a neighborhood of 0 on  $\{0 \in \hat{\omega}^{\circ}\}$ . From (4.10) and (4.17) follows

$$P_0\{0 \bar{\epsilon} \hat{\omega}^0, 0 \text{ is is loated point in } \omega^0\} = 0$$
. (4.18)

Combining (4.18) with (4.16) we get

$$d_{T_n} \neq 0$$
 a.s.  $P_0$ . (4.19)

Take  $f = g(r_{s_1}, r_{s_2}, \dots, r_{s_k})$ , where g is a positive bounded continuous function of k variables. By virtue of (4.19)

$$P_{0}{f} = \lim_{n} P_{0}{f \circ \tau_{d_{T_{n}}}} = \lim_{n} \{1_{T_{n}}(\hat{\omega}^{\circ}) P_{1}{f} + 1_{T_{n}}\bar{\epsilon}\hat{\omega}^{\circ} P_{0}{f}$$
(4.20)

Suppose  $P_0\{0 \in \hat{\omega}^\circ\} = \epsilon > 0$ . Then the right hand side of (4.20) converges to

 $\epsilon P_1{\mathbf{f}} + (1-\epsilon)P_0{\mathbf{f}}$ 

which implies  $P_0 = P_1$ . The latter implies M is a regenerative set, and this is in contradiction with our assumption that M is the set of the third type.

Let  $b(\omega^{\circ})$  be the set of accumulation from the left points of  $\omega^{\circ}$ , i.e.  $x \in b(\omega^{\circ})$  iff there exists a sequence  $\{x_n\}$  such that  $x_n < x$ ,  $x_n \in \omega^{\circ}$  and  $x_n \uparrow x$ .

(4.21) **PROPOSITION.** If M has a perfect regenerative part then for each k and t

$$P\{n(k,t) \in b(M)\} = 1$$
 (4.22)

*Proof.* Suppose (4.22) fails. Then with a positive probability there exists an interval contiguous to M whose right end coincide with  $\eta(k,t)$ . Fubini's theorem implies an existance of u for which

$$P\{D_{ij} = n(k,t), D_{ij} > u\} > 0$$
. (4.23)

Applying (3.4) to  $f = 1_0(\hat{\omega}^0)$  and using (4.23), we get

 $P_0\{0\in\hat{\omega}^\circ\}>0$ 

which is in contradiction with proposition (4.21).

Put

$$z_0 \stackrel{\Delta}{=} 0,$$
  
$$z_k \stackrel{\Delta}{=} \theta_{\tilde{\eta}(k)} \stackrel{\Xi}{=} \theta_{\tilde{\gamma}(k)} \stackrel{\Xi}{=} \theta_{\tilde{\nu}(k)},$$

where  $\theta_s$  is given by (4.13) and  $\tilde{n}(k)$ ,  $\tilde{\gamma}(k)$  and  $\tilde{\nu}(k)$  stand for  $\tilde{n}(k,0)$ ,  $\tilde{\gamma}(k,0)$ , and  $\tilde{\nu}(k,0)$ , given by (3.5).

(4.24) **PROPOSITION.** If M has a perfect regenerative part then  $\zeta_k - \zeta_{k-1}$ , k = 1, 2, ... are exponential iid on  $\{\Omega^o, G^o, P_0\}$ .

Proof. Let  $\tilde{n} = \tilde{n}(1)$ . Consider

$$P_0\{\zeta_1 > a+b\} = P_0\{\theta_{\tilde{\eta}} > a+b\} = P_0\{\theta_{\tilde{\eta}} > a, \theta_{\tilde{\eta}} - a>b\}.$$

Let

$$\sigma = \inf\{s: \theta_s \ge a\}. \tag{4.26}$$

Then  $\sigma$  is a stopping time with respect to  $G_{t+}^{\circ}$  and  $\theta_{\sigma} = a$ . Thus the right-hand side of (4.25) can be written as

$$P_{0}\{\theta_{\tilde{n}} > a, \theta_{\tilde{n}} - \theta_{\sigma} > b\} = P_{0}\{\theta_{\tilde{n}} > a, \theta_{\tilde{n}-\sigma} \circ \tau_{\sigma} > b\}$$
(4.27)  
$$= P_{0}\{\theta_{\tilde{n}} > a, \theta_{\tilde{n}\circ\tau_{\sigma}} \circ \tau_{\sigma} > b\}$$
  
$$= P_{0}\{P_{0}\{\theta_{\tilde{n}\circ\tau_{\sigma}} \circ \tau_{\sigma} > b | G_{\sigma+}^{\circ}\}; \theta_{\tilde{n}} > a\}$$
  
$$= P_{0}\{\theta_{\tilde{n}} > a\} P_{0}\{\theta_{\tilde{n}} > b\} .$$

The first equality in (4.27) is due to (4.14). The second equality holds because  $\sigma > \tilde{\eta}$  and for any s  $\tilde{\eta} \circ \tau_s = \tilde{\eta} - s$  on the set  $\{s < \tilde{\eta}\}$ . The last equality in (4.27) is a consequence of Proposition (3.10) and the equality

 $d_{\sigma} = \sigma$ 

which is true for any perfect regenerative set and any  $\sigma$  given by (4.26). Equally (4.27) shows that  $\zeta_1$  has exponential distribution.

Since for any k

the quantities  $\theta_{\tilde{n}(k)}$  and  $\theta_{\tilde{v}(k)}$  coincide. Thus, in a way similar to the one in which (4.27) was obtained,

$$P_{0} \{ \theta_{\tilde{\eta}(k+1)} - \theta_{\tilde{\eta}(k)} > a \mid G_{\tilde{\nu}(k)+}^{\circ} \}$$

$$= P_{0} \{ \theta_{\tilde{\eta}(k+1)} - \theta_{\tilde{\nu}(k)} > a \mid G_{\tilde{\nu}(k)+}^{\circ} \}$$

$$= P_{0} \{ \theta_{\tilde{\eta}(k+1)} - \tilde{\nu}(k) \circ \tau_{\tilde{\nu}(k)} > a \mid G_{\tilde{\nu}(k)+}^{\circ} \}$$

$$= P_{0} \{ \theta_{\tilde{\eta}\circ\tau_{\tilde{\nu}(k)}} \circ \tau_{\tilde{\nu}(k)} > a \mid G_{\tilde{\nu}(k)}^{\circ} \}$$

$$= P_{0} \{ \theta_{\tilde{\eta}\circ\tau_{\tilde{\nu}(k)}} \circ \tau_{\tilde{\nu}(k)} > a \mid G_{\tilde{\nu}(k)}^{\circ} \}$$

The above equality shows that  $\zeta_{k+1} - \zeta_k$  is independent of  $\zeta_n - \zeta_{n-1}$ , n = 1, 2, ..., kand have the same distribution as  $\zeta_1$ .

(4.28) **REMARK.** The proof of Proposition (4.24) also shows that  $\zeta_k - \zeta_{k-1}$  is independent of the sequence of random vectors  $(\tilde{v}(n) - \tilde{v}(n), \tilde{v}(n) - \tilde{n}(n)), n = 1, 2, ...$ 

### 5. Structure of Stationary Markov Set.

In this section we will show that each stationary set M is  $(\Pi, \alpha, \lambda, \mu)$ generated. This will be done separately for the case in which M has a perfect regenerative part and in the case in which M has a discreet regenerative part.

Suppose M is a set with a perfect regenerative part and  $P_0 \cdot K^{-1}$  is its law of regeneration. Consider the process  $V_t$  on  $(\Omega^0, G^0, P_0)$ 

$$V_{t} = \sum_{k:\zeta_{k} < t} (\tilde{v}(k) - \tilde{\eta}(k)),$$

where  $\zeta_k = \theta_{\tilde{\eta}(k)}$  with  $\theta$  given by (4.13). The process  $V_t$  is of a pure jump type. In view of Proposition (4.24)  $\zeta_k - \zeta_{k-1}$  are exponential iid. By virtue of the proposition (3.26) and Remark (4.28) the random variables ( $\tilde{\nu}(k) - \tilde{\eta}(k)$ ) are iid independent of the point process  $\zeta_k$ . Therefore, the process  $V_t$  is a process with independent increments.

Proposition (4.21) and Proposition (3.10) together with (4.18) show that  $\tilde{n}(k)$  and  $\tilde{v}(k)$  are points of accumulation of  $K \circ \omega^0$  a.s.  $P_{0}$ . This implies

$$z_{\zeta_{k}} = \tilde{v}(k), \quad z_{\zeta_{k}} = \tilde{n}(k), \quad (5.1)$$

where  $z_t$  is the process whose image is equal to  $K \circ \omega^0 \cap [0, \infty[$ . From (5.1) and the definition of  $V_t$  follows

$$V_{\zeta_k} - V_{\zeta_k} = z_{\zeta_k} - z_{\zeta_k}$$
 (5.2)

Put

$$W_t \stackrel{\Delta}{=} z_t - V_t$$
.

Since both  $z_t$  and  $V_t$  are processes with independent increments, so is  $W_t$ . The set  $K \circ \omega^0$  has Lebesgue measure zero, therefore the process  $z_t$  has translation constant equal to zero and is of a pure jump type. In view of (5.2),  $W_t$  is an increasing process of a pure jump type such that

$$W_{\zeta_k} = W_{\zeta_k}$$
 (5.3)

Relation (5.3) implies that  $V_{\circ}$  and  $W_{\circ}$  have no common points of discontinuity. Accordingly  $V_{\circ}$  and  $W_{\circ}$  are independent (see [10]).

(5.4) **THEOREM.** A stationary Markov set M with a perfect regenerative part is  $(\Pi, \alpha, \lambda, \mu)$ -generated.

Proof. For each t we need to find  $\phi_t$  such that  $M \cap [\phi_t, \infty]$  is a  $(\Pi, \alpha, \lambda, \mu)$ -set. In view of stationarity it is sufficient to consider only t = 0. Put  $\phi = \nu$ , where  $\nu$  is given by (3.20). Let

$$X_{k} \stackrel{\Delta}{=} \tilde{\gamma}(k) \circ \tau_{v} \circ M - \tilde{\eta}(k) \circ \tau_{v} \circ M \equiv \gamma(k+1,0) - \eta(k+1,0),$$

$$Y_{k} \stackrel{\Delta}{=} \tilde{v}(k) \circ \tau_{v} \circ M - \tilde{\gamma}(k) \circ \tau_{v} \circ M \equiv v(k+1,0) - \gamma(k+1,0),$$

$$x_{t} \stackrel{\Delta}{=} V_{t} \circ \tau_{v} + \phi,$$

$$y_{t} \stackrel{\Delta}{=} W_{t} \circ \tau_{v} + \phi.$$
(5.5)

Then for  $\Sigma_{t}$  given by (2.6) we have

 $Z_{t} = Z_{t} \circ \tau_{v} \circ M + \phi$  $\sigma_{k} = \zeta_{k} \circ \tau_{v} \circ M$ 

Let  $\Pi$  be the Levi's measure of the subordinator W. Let  $\alpha$  be the parameter

of the exponential distribution of  $\zeta_k - \zeta_{k-1}$ ,  $\lambda$  be the parameter of the exponential distribution of  $\gamma(k) - \eta(k)$  (see (3.27)) and

$$\mu(\Gamma) \stackrel{\Delta}{=} P\{\nu(k) - \gamma(k) \in \Gamma\} = P_0\{\tilde{\nu}(k) - \tilde{\gamma}(k) \in \Gamma\}.$$

ىدىمەكمەشمەشمەرمەتمەتمەتمەتمەتمەتمەتمەتمەتمەت باردى

We would like to show that the set  $M \cap [\phi, \infty[$  is a  $(\Pi, \lambda, \alpha, \mu)$ -set as defined by (2.6) - (2.8). Since  $\nu = D_{\nu}$  and  $\gamma \in \hat{M}$ , we can apply Proposition (3.5) and get

$$P\{f \circ M^{Y} | G_{\gamma^{+}}\} = P\{f \circ \tau_{\nu} \circ M | G_{\gamma^{+}}\} = P_{0}\{f\} .$$
 (5.6)

In particular, (5.6) shows that the law of  $(V_{\bullet} \circ \tau_{V} \circ M, W_{\bullet} \circ \tau_{V} \circ M)$  is the same as the law of  $(V_{\bullet}, W_{\bullet})$  on  $(\Omega^{\circ}, G^{\circ})_{\bullet}$ . It also shows independence of v and  $(V_{\bullet} \circ \tau_{V} \circ M, W_{\bullet} \circ \tau_{V} \circ M)_{\bullet}$ .

For  $M_1 \stackrel{\Delta}{=} \tau_v \circ M$  and for  $z_t = V_t + W_t$ 

$$K \circ M_1 = \overline{z_{\mathbb{R}_+} \circ M_1}$$
.

Thus

$$\overline{Z}_{\mathbb{R}_{+}^{\circ M}} \stackrel{\circ M}{\to} = K \circ M \cap [\phi, \infty[$$

The construction of the process  $V_t$  (and  $x_t$  by (5.5)) shows that L given by (2.7) coincides with the closure of  $\hat{M} \cap [\phi, \infty[$ . Since  $M = K \circ M \cup \hat{M}$ , we got the representation (2.8) with  $x_{\star}$  and  $y_{\star}$  given by (5.5). Proposition (4.24) shows that  $c_k = \zeta_k \circ \tau_y \circ M$  forms a Poisson point process. Proposition (3.26) and Remark (4.28) show the required independence of  $\{X_k\}$ ,  $\{Y_k\}$  and  $y_{\star}$  as well as independence of  $x_{\star}$  and  $y_{\star}$  given by (5.5). This concludes the proof that  $M \cap [\phi, \infty[$  is a (JI,  $\alpha, \lambda, \mu$ )-set.

A stationary Markov set with a discreet regenerative part cannot be treated in the same manner because Propositions (4.15) and (4.21) are no longer true in this case. As a result (5.1) and (5.2) as well as (5.3) might fail. The failure of (5.1) - (5.3) might result in dependence of the processes V. and W. . However, the case of a set with a discreet regenerative part can be treated "from scratch". The analysis of this case is rather simple, so we will only outline the main points without going into details.

Put

$$p = P_0 \{0 \in \hat{\omega}^0\}.$$
 (5.7)

It is easy to show that if M has a discreet regenerative part then 0 . $Let <math>\lambda$  be given by (3.23) and

$$\mu(\Gamma) = P_0\{\tilde{v} - \tilde{\gamma} \in \Gamma\}$$
(5.8)

$$\Pi(\Gamma) = P_0\{d_0 \in \Gamma | 0 \in \hat{\omega}^0\}$$
(5.9)

Proposition (3.5) shows that the right endpoint of each interval contiguous to M belongs to  $\hat{M}$  with probability p independently of the length of this interval. Thus  $\tau_v \circ M$  can be described by means of a Markov renewal process U(t) (see [1] Chapter 10) with three states. The holding time in the first state is exponential with parameter  $\lambda$ , the holding time in the second and third states have distribution  $\mu$  and  $\Pi$  respectively. The transition matrix of the imbedded discreet Markov chain is

0	1	0	
p	0	1-p	
( p	0	1-p	]

The set of times when U(t) undergoes transitions from one state to another or U(t) is in the first state corresponds to  $\tau_{11} \circ M$ .

It is easy to verify that the set  $M \cap [\nu, \infty] \equiv \nu + \tau_{\nu} \circ M$  is a  $(\Pi, \alpha, \lambda, \mu)$ -set where  $\alpha$  is such that

$$p = \frac{\alpha}{\alpha + 1}$$

(Note that if  $y_t$  is a (0, II)-process and  $x_t$  is the process defined by (2.6), then  $\alpha/(\alpha+1) = P\{\sigma_1 < \inf\{t: y_t \neq y_{t_t}\}\}$ ).

24

(5.9) **PROPOSITION.** If M is a stationary  $(\Pi, \alpha, \lambda, \mu)$ -generated set then  $\Pi$  and  $\mu$  satisfy (2.13), (2.14).

*Proof.* (For M with a perfect regenerative part, for M with a discreet regenerative part the proof is similar). Let  $x_t$  and  $y_t$  be the subordinators which generate  $(\Pi, \alpha, \lambda, \mu)$ -set (see (2.6)-(2.8)). Then

$$K \circ (\overline{Z_{\mathbb{R}_{+}}} \cup L) = \overline{Z_{\mathbb{R}_{+}}}.$$

The latter shows that the process Z = x + y generates stationary regenerative set  $K \circ M$ . If  $\Pi$ ' is the Levi's measure of Z then from [8] and [11]

$$\int_{0}^{\infty} x \Pi'(dx) < \infty \quad . \tag{5.10}$$

On the other hand it is known (see [10]) that

$$P\{Z_{t} - Z_{0}\} = t \int_{0}^{\infty} x \Pi'(dx)$$
 (5.11)

The left hand side of (5.11) can be rewritten as

$$P\{y_{t} - y_{0}\} + P\{x_{t} - x_{0}\} = t \int_{0}^{\infty} x \Pi(dx) + t \alpha^{-1} [\lambda^{-1} + \int_{0}^{\infty} x \mu(dx)] \quad (5.12)$$

Relations (5.10), (5.11) (5.12) imply (2.13) and (2.14).

(5.13) **PROPOSITION.** If M is a  $(\Pi, \alpha, \lambda, \mu)$ -generated set, then the quadruple  $(\Pi, \alpha, \lambda, \mu)$  is determined by M uniquely up to equivalency.

Proof. The compliment of M consists of a union of open intervals  $(L_t, R_t)$ . Since  $M \cap [\phi_0, \infty]$  is a  $(\Pi, \alpha, \lambda, \mu)$ -set we can write (recalling representation (2.6) - (2.8)).

$$P\{\sum_{\substack{\nu(1) \le t < \eta(2)}} f(R_{t}^{-L} L_{t}^{-}) R_{t}^{\neq L} t\}$$

$$= \lim_{k \to \infty} P\{\sum_{\substack{\nu(k) \le t < \eta(k+1)}} f(R_{t}^{-L} L_{t}^{-}) R_{t}^{\neq L} t\}$$
(5.14)

$$= Q\{\sum_{\sigma_{k} < t < \sigma_{k+1}} f(Z_{t} - Z_{t-}) I_{Z_{t} \neq Z_{t-}}\}$$

$$= Q\{\sum_{\sigma_{k} < t < \sigma_{k+1}} f(y_{t} - y_{t-}) I_{y_{t} \neq y_{t-}}\}$$

$$= \Pi\{f\} Q\{\sigma_{k+1} - \sigma_{k}\} = \alpha^{-1} \Pi\{f\}.$$
(5.14)

Here Q is the probability measure associated with the process x, y and Z in (2.6) - (2.8). Formula (5.14) shows that  $(\Pi, \alpha)$  is determined by M uniquely up to proportionality.

On the other hand, direct computations show that for any  $(\Pi, \alpha, \lambda, \mu)$ -set

$$P\{\gamma - \eta\} = [\lambda(1 - \alpha \mu\{0\} / (\alpha + \Pi(\mathbf{R}_{+}))]^{-1}$$
(5.15)

and

$$P\{\nu-\gamma \in \Gamma\} = \begin{cases} \mu(\Gamma) & \text{if } \Pi(\mathbb{R}_{+}) = \infty \\ \mu^{\dagger}\{\Gamma\} + \mu\{0\} \Pi(\mathbb{R}_{+})^{-1} \Pi(\Gamma), & \text{if } \Pi(\mathbb{R}_{+}) < \infty \end{cases}$$
(5.16)

Equalities (5.15) and (5.16) complete the proof of the proposition.

## 6. Markov Properties of the Residual Life Process

Consider the "residual life" process

$$R_{+} = \inf\{s - t: s > t, s \in M\}$$

$$(6.1)$$

associated with the stationary Markov set M. Markov property of M implies that  $R_{+}$  is a Markov (but not necessarily a strong Markov) process.

Consider a  $(\Pi, \alpha, \lambda, \mu)$ -set given by (2.7), (2.8) and the processes y., x. and Z. which generate it. Let

$$\mathbf{c}_{\mathbf{a}} \stackrel{\Delta}{=} \inf\{\mathbf{s} \ge 0: \mathbf{Z}_{\mathbf{s}} \ge \mathbf{a}\}, \qquad (6.2)$$

$$F_{a} = Z_{c_{a}},$$

$$H_{a} = Z_{c_{a}}.$$
(6.3)

Let N be the union of  $\sigma_k$  (see (2.6)). Then  $R_t$  given by (6.1) can be represented as

$$R_{t} = \begin{cases} 0 \text{ if } c_{t} \in N \text{ and } t \in y, \\ F_{t} - t \text{ otherwise}. \end{cases}$$
(6.4)

Let Q be the law of the subordinators x. and y. of (2.6) - (2.8). The transition function of  $R_t$  associated with a stationary  $(\Pi, \alpha, \lambda, \mu)$ -generated set is the same as transition function of  $R_t$  associated with any  $(\Pi, \alpha, \lambda, \mu)$ -set. Hence we can assume any distribution of  $y_0$  in (2.6) - (2.8), in particular

 $Q\{y_0 = 0\}$ .

The transition function of the process  $R_t$  given by (6.4) is

$$p(t,x;\Gamma) = 1_{\Gamma}(x) \quad \text{if} \quad x < t,$$

$$p(t,x;\Gamma) = Q\{F_{t} - t \in \Gamma, c_{t} \in N\} + \sum_{k=1}^{\infty} Q\{F_{t} - t \in \Gamma, c_{t} = \sigma_{k}, Z_{\sigma_{k}} - {}^{+}X_{k} \le t\}, x > 0, 0 \in \Gamma,$$

$$p(t,x;\{0\}) = \sum_{k=1}^{\infty} Q\{c_{t} = \sigma_{k}, Z_{\sigma_{k}} - {}^{+}X_{k} > t\}, x > 0,$$

$$p(t,0;\Gamma) = e^{-\lambda t} 1_{0}\{\Gamma\} + \int_{0}^{t} \lambda e^{-\lambda y} \mu(\Gamma + t - y) dy + \int_{0}^{t} \mu_{1}(dy) p(t - y, y;\Gamma),$$

$$(6.5)$$

where  $u_1$  is a distribution of the jumps of the process x. (i.e.,  $u_1$  is a convolution of  $u_1$  and an exponential distribution with parameter  $\lambda$ ).

 $A=(\Pi,\lambda,\lambda,\mu)\text{-generated set is stationary iff}$ 

$$m_t(\Gamma) \stackrel{\Delta}{=} P\{R_t \in \Gamma\}$$

does not depend on t

Inversely, if we are able to construct a probability measure m which is invariant with respect to  $p(t,x;\Gamma)$  then the stationary Markov process  $R_t$  with the one dimensional distribution m and the transition function p will yield a stationary  $(\Pi, \alpha, \lambda, \mu)$ -generated set by the formula

$$M = \overline{D}_{IR}$$
,

where  $D_t = t + R_t$ .

(6.6) THEOREM. If  $\Pi$  and  $\lambda$  are subject to (2.13), (2.14) then there exists a unique stationary probability measure m for the Markov process  $R_t$  associated with a  $(\Pi, \alpha, \lambda, \mu)$ -set. The measure m is given by

$$\mathfrak{m}(\mathbf{f}) = \mathbb{C}[\lambda^{-1}\mathbf{f}(0) + \int_{0}^{\infty}\mathbf{f}(\mathbf{t})\mu(]\mathbf{t},\infty[)d\mathbf{t} + \alpha^{-1}\int_{0}^{\infty}\mathbf{f}(\mathbf{t})\Pi(]\mathbf{t},\infty[)d\mathbf{t}].$$

For the proof of this theorem we need the following proposition.

(6.7) **PROPOSITION.** Let  $(y_s, Q)$  be a  $(0, \Pi)$ -process and let

$$c_a = \inf\{s: y_s \ge a\}.$$

Let S be an exponential random variable with parameter  $\alpha$  independent of the process  $y_t$ . Then

Prec 6. The right-hand side of (6.8) can be rewritten as

$$Q\{\sum_{y_{s}\neq y_{s-}, s\leq S} \int_{y_{s-}}^{y_{s}} f(y_{s}-u) du\}$$
  
= Q\{\sum\_{y\_{s}\neq y\_{s-}, s\leq S} \int\_{y\_{s-}}^{y\_{s}} f(u) du\}  
(6.9)

$$= Q\{\sum_{y_{s}\neq y_{s}}^{1} 1_{s \leq S} g(y_{s}-y_{s})\},$$

where  $g(t) = \int_{0}^{t} f(u) du$ . The right hand side of (6.9) is equal to

$$Q\{\Pi(g)S\} = \Pi(g)Q\{S\} = \alpha^{-1}\Pi(g)$$

(see [10] Section 3). By Fibini's theorem

$$\Pi(g) = \int_{0}^{\infty} g(x) \Pi(dx) = \int_{0}^{\infty} \{\int_{0}^{x} f(t)dt\} \Pi(dx) = \int_{0}^{\infty} f(t) \int_{t}^{\infty} \Pi(dx) = \int_{0}^{\infty} f(t) \Pi(]t, \infty[)dt$$

whereas (6.8) follows.

Precis of the Theorem (6.6). Let x. and y. be the processes (with  $Q\{y_0=0\}=1$ ) which generate a  $(\Pi, \alpha, \lambda, \mu)$ -set by formulae (2.6) - (2.8). Then the process  $R_t$ associated with this set by formula (6.4) is a regenerative process (see [1] Ch. 9) with the moments of regeneration  $\rho_1, \rho_2, \ldots, \rho_k, \ldots$ 

$$\sigma_k \stackrel{\Delta}{=} Z_{\sigma_k}$$
.

Really, by virtue of the strong Markov property

$$x_{s}^{k} \stackrel{\Delta}{=} x_{\sigma_{k}+s} - x_{\sigma_{k}}$$

and

$$y_{s}^{k} = y_{\sigma_{k}+s} - y_{\sigma_{k}}$$

have the same distribution as the processes  $x_s$  and  $y_s$  respectively and are independent of  $\{x_s, y_s; s \le \sigma_k\}$ . Since

$$R_{\rho_{k}+t} = R_{t} \circ (x_{\bullet}^{k}, y_{\bullet}^{k})$$

the process  $\mathbb{R}_{\rho_k}$  is independent of  $\{\mathbb{R}_s, s \le \rho_k\}$  and has the same distribution as  $\mathbb{R}_t$ . The same argument shows that the sequence  $\{\rho_k\}$  forms a renewal process.

Since

$$\mathcal{C}_{k+1} - \mathcal{D}_k \equiv \mathcal{Z}_{\sigma_{k+1}} - \mathcal{Z}_{\sigma_k} = (\mathbf{y}_{\sigma_{k+1}} - \mathbf{y}_{\sigma_k}) + \mathbf{x}_k + \mathbf{y}_k$$

and since  $X_k$  has a continuous (exponential) distribution,  $\rho_{k+1} - \rho_k$  has a continuous distribution as well. Thus the renewal process  $\{\rho_k\}$  is aperiodic and

$$E\{\rho_{k+1} - \rho_{k}\} = E\{X_{k}\} + E\{Y_{k}\} + E\{y_{\sigma_{k+1}} - y_{\sigma_{k}}\}$$

$$= \lambda^{-1} + \int_{0}^{\infty} t \mu(dt) + \alpha^{-1} \int_{0}^{\infty} t \Pi(dt) .$$
(6.10)

The right-hand side of (6.10) is finite by virtue of (2.13) and (2.14).

According to Theorem (2.25) of Chapter 9 of [1] there exists a unique stationary measure m for the regenerative process R<sub>•</sub>, given by

$$m(\Gamma) = C Q\{ \int_{0}^{\rho_{1}} 1_{\Gamma}(R_{t}) dt \}, \qquad (6.11)$$

where  $C^{-1}$  is equal to (6,10) (The expression in (6.11) is equivalent to the one given in Ch. 9, Theorem (2.25) of [1]). Since the process  $x_t$  is equal to 0 on the interval  $]0,\sigma_1[$  the process  $\mathbb{Z}_s$  coincides with  $y_s$  on  $]0,\sigma_1[$  and

$$R_{t} = y_{\sigma_{t}} - t \quad \text{for} \quad t \leq y_{\sigma_{1}}$$

$$R_{t} = 0 \quad \text{for} \quad y_{\sigma_{1}} \leq t < y_{\sigma_{1}} + X_{1}$$

$$R_{t} = Z_{\sigma_{1}} - t \equiv y_{\sigma_{1}} + X_{1} + Y_{1} - t \quad \text{for} \quad y_{\sigma_{1}} + X_{1} < t < Z_{\sigma_{1}}$$

(see (2.6), (2.8) and (6.4)). Thus

$$Q\left\{ \int_{0}^{\rho_{1}} 1_{\Gamma}(R_{t}) dt \right\} = Q\left\{ \int_{0}^{y_{\sigma_{1}}} 1_{\Gamma}(y_{c_{t}} - t) dt \right\} + Q\left\{ \int_{0}^{X_{1}} 1_{\Gamma}(0) dt \right\} + Q\left\{ \int_{0}^{y_{1}} 1_{\Gamma}(Y_{1} - t) dt \right\}$$
(6.12)

The first term in the right-hand side of (6.12) equals to

$$\alpha^{-1} \int_{0}^{\infty} \mathbf{1}_{\Gamma}(t) \Pi(]t, \infty]) dt \qquad (6.13)$$

by virtue of Proposition (6.7). The second term in (6.12) equals

 $1_{\Gamma}(0) \in \{X_1\} = 1_{\Gamma}(0)\lambda^{-1}$  (6.14)

$$Q\{\int_{0}^{Y_{1}} 1_{\Gamma}(Y_{1}-t) dt\} = Q\{\int_{0}^{Y_{1}} 1_{\Gamma}(t) dt\} = Q\{\int_{0}^{\infty} 1_{\Gamma}(t) 1_{t < Y_{1}} dt\} = \int_{0}^{\infty} 1_{\Gamma}(t) \mu(]t, \infty[) dt.$$

From (6.11) - (6.15) follows Theorem (6.6).

(6.16) COROLLARY. For each  $(\Pi, \alpha, \lambda, \mu)$  subject to (2.13), (2.14) there exists a stationary  $(\Pi, \alpha, \lambda, \mu)$ -generated set.

This completes the proof of the Theorem (2.12).

(6.17) **REMARK.** The proof of Theorem (6.6) shows that any  $(\Pi, \alpha, \lambda, \mu)$ -generated set M with

$$P\{\inf M = -\infty\} = 1$$

is stationary.

### 7. Reversability Properties of Stationary Markov Sets.

In this section we will prove Theorem (2.15). We consider a stationary  $(\Pi, \gamma, \gamma, \delta_0)$ -generated set with a perfect regenerative part. The proof of Theorem (2.15) for M with a discrete regenerative part is similar. The closure of  $\hat{M}$  consists of a union of closed intervals of iid exponential length and by virtue of Proposition (4.15) and (4.21) the endpoints of these intervals are points of accumulation of M- $\hat{M}$ . Therefore the endpoints of these intervals belong to K  $\circ$  M.

According to Theorem (4.6) the set  $K \circ M$  is stationary regenerative with Lebesgue measure zero. By Theorem 1 of [11] there exists a  $(0, \Pi^{+})$ -subordinator whose range coincides with  $K \circ M \cap [0, \infty[$ . If  $\mu = \delta_0$ , then (2.6) - (2.8) show that  $\Pi^{+}$  is the Levi's measure of the process Z and

$$\Pi' = \Pi + \alpha G_{\lambda}, \qquad (7.1)$$

where  $G_\lambda$  is an exponential distribution with parameter  $\lambda.$  In particular, M has

a perfect regenerative part iff

To show that -M has the same distribution as M we have to consider the two dimensional process

$$(L_t, R_t) = (\ell_t, r_t) \circ M$$
.

It follows from the Markov property of M that the process  $(L_t, R_t)$  is a stationary Markov process. If M is a  $(\Pi, \alpha, \lambda, \delta_0)$ -generated set then the transition function of  $(L_t, R_t)$  is

$$p(t,(u,v);\Gamma) = 1_{\Gamma}(u,v), \quad \text{if } v > t, \ \Gamma \in ]-\infty,0] \times [0,\infty[,$$

$$p(t,(u,v);\Gamma) = Q\{(H_{t-v},F_{t-v}) \in \Gamma, \ c_{t-v} \in N\}, \ v < t, \ \Gamma \in ]-\infty,0[\times]0,\infty[$$

$$p(t,(u,v); \ (0,0)) = Q\{c_{t-v} \in N\} \equiv \sum_{k} Q\{c_{t-v} = \sigma_{k}\}, \ v < t,$$

$$p(t,(0,0);\Gamma) = \int_{0}^{t} \lambda e^{-\lambda s} \ p(t,(0,s),\Gamma) \ ds, \ \Gamma \in ]-\infty,0] \times [0,\infty[$$

$$(7.2)$$

Here Q is the law of the processes  $x_s$ ,  $y_s$  and  $\tilde{z}_s$ , of (2.6) - (2.8),  $c_t$  is given by (6.2) and  $(H_t, F_t)$  are given by (6.3).

Note that when  $u = \delta_0$ , the length of each jump of the process Z. caused by a discontinuity of x. is exponentially distributed and the range of each jump belongs to the  $(\Pi, \lambda, \alpha, \delta_0)$ -generated set. This results in simplifications in (7.2) as compared to (6.5).

Let  $\Pi(x; \Gamma) = \Pi(\Gamma - x)$  where  $\Gamma - x = \{y - x; y \in \Gamma\}$ .

The process  $(L_t, R_t)$  is stationary due to stationarity of M. Repeating the proof of Theorem (6,6) for  $(L_t, R_t)$ , we can get that the one-dimensional distribution n of this process is given by

$$n(\Gamma \times \Delta) = c[\lambda^{-1} 1_{(0,0)}(\Gamma \times \Delta) + \alpha^{-1} \int_{-\infty}^{0} 1_{s}(\Gamma) \Pi(x;\Delta) ds], \ \Gamma \subset \mathbb{R}_{+}, \ \Delta \subset \mathbb{R}_{+}.$$

Let  $(u,v)^T = (v,u)$  and if  $\Delta \subset \mathbb{R} \times \mathbb{R}$  then  $\Delta^T$  should be understood similarly. Let

$$y_{t}^{*} \stackrel{\Delta}{=} -y_{t}, \qquad x_{t}^{*} \stackrel{\Delta}{=} -x_{t},$$

$$Z_{t}^{*} \stackrel{\Delta}{=} -Z_{t},$$

$$\Pi^{*}(\Gamma) \stackrel{\Delta}{=} \Pi(-\Gamma),$$

$$\Pi^{*}(x;\Gamma) \stackrel{\Delta}{=} \Pi^{*}(\Gamma-x) = \Pi(-x;-\Gamma).$$
(7.4)

Consider the set -M. The process

$$(L_t^{\star}, R_t^{\star}) \stackrel{\Delta}{=} (\ell_t, r_t) \circ (-M) = (-R_t, -L_t)$$

is a Markov process with the one-dimensional distribution (obtained by change of variables in (7.3))

$$(7.5)$$
$$n^{*}(\Gamma \times \Delta) = C[\lambda^{-1} 1_{(0,0)} ((-\Delta) \times (-\Gamma)) + \alpha^{-1} \int_{0}^{\infty} 1_{s}(\Delta) \Pi^{*}(s;\Gamma) ds], \Gamma \subset \mathbb{R}_{+}, \Delta \subset \mathbb{R}_{+}$$

and the backward transition function

$$p^*(s,(u,v);\Gamma) = p(s,(-v,-u);\Gamma^T), \Gamma \subset ]-\infty,0] \times [0,\infty[, u \le 0 \le v].$$
 (7.6)

Let

$$\Lambda(\Gamma) \stackrel{\Delta}{=} Q\{\int_{0}^{\infty} 1_{\Gamma}(Z_{s}) ds\},$$

$$\Lambda^{*}(\Gamma) \stackrel{\Delta}{=} Q\{\int_{0}^{\infty} 1_{\Gamma}(Z_{s}^{*}) ds\} = \Lambda(-\Gamma),$$

$$\Lambda_{b}(\Gamma) = \Lambda(\Gamma - b),$$

$$\Lambda_{b}^{*}(\Gamma) = \Lambda^{*}(\Gamma - b),$$
(7.7)

(7.8) **PROPOSITION.** For any :

For any function f of two variables

$$\mathbb{Q}\left\{\sum_{\substack{Z \\ t^{\neq Z} t}} f(Z_{t^{-}}, Z_{t}) \mathbf{1}_{t \in N}\right\} = \int_{0}^{\infty} \Lambda(dx) \int_{0}^{\infty} f(x, y) \Pi(x; dy)$$

$$Q\{_{Z_{t-}^{*}\neq Z_{t}^{*}}f(Z_{t-},Z_{t}) |_{t \in N}\} = \int_{-\infty}^{0} \int_{-\infty}^{0} f(x,y) \Pi^{*}(x;dy)$$

The proof of this proposition is well known.

(7.9) **PROPOSITION.** For any two functions g and h on IR

$$\int_{-\infty}^{\infty} g(x) \Pi(x,f) dx = \int_{-\infty}^{\infty} f(x) \Pi^{*}(x;g) dx \qquad (7.10)$$

$$\int_{-\infty}^{\infty} \Lambda_{\mathbf{x}}(\mathbf{f}) g(\mathbf{x}) d\mathbf{x} = \int_{-\infty}^{\infty} f(\mathbf{x}) \Lambda_{\mathbf{x}}^{*}(g) d\mathbf{x}$$
(7.11)

For the proof of this proposition see [11] Lemma 6.4.

(7.11) **PROPOSITION.** Measures n and n<sup>\*</sup> given by (7.3) and (7.5) respectively, coincide.

*Proof.* The first term in brackets in the right-hand side of (7.3) is equal to the first term in the right-hand side of (7.5). The integral term in the right-hand side of (7.3) equals to the corresponding term in (7.5) by virtue of (7.10).

(7.12) **PROPOSITION.** For any two sets  $\Gamma$ ,  $\Delta \subset \mathbb{R} \times \mathbb{R}$ .

$$\int_{\Gamma} n(du, dv) p(s, (u, v); \Delta) = \int_{\Delta} n^*(du, dv) p^*(s, (u, v); \Gamma)$$
(7.15)

Proof. Consider  $\Delta$  and  $\Gamma$  of the form

$$\Delta = \Delta' \times \Delta'', \qquad \Delta' < 0, \qquad \Delta'' > 0$$

$$\Gamma = \Gamma_1 \times \Gamma_2 \qquad \Gamma_1 < 0, \qquad \Gamma_2 > 0$$
(7.14)

Put  $\Delta_1 = \Delta^* + s$  and  $\Delta_2 = \Delta^{**} + s$  and assume

$$\Gamma_2 < \Delta_1 \tag{7.15}$$

(The inequality between two subsets of the real line means the corresponding inequality between any two points from the first and the second set respectively.)

For  $v < \Delta_1$  we can write

$$p(s, (u, v), \Delta) = Q\{H_{s-v} \in \Delta' + s - v, F_{s-v} \in \Delta'' + s - v, c_{s-v} \in N\}$$

$$= Q\{\sum_{Z_{t-} \neq Z_{t}} 1_{\Delta_{1} - v}(Z_{t-}) 1_{\Delta_{2} - v}(Z_{t}) 1_{t \in N}\}$$

$$= \int_{0}^{\infty} \Lambda(dy) 1_{\Delta_{1} - v}(y) \Pi(y; \Delta_{2} - v)$$

$$= \int_{0}^{\infty} \Lambda_{v}(dy) 1_{\Delta_{1}}(y) \Pi(y; \Delta_{2}).$$
(7.16)

We used Proposition (7.8) in the third equality in (7.16) and the identity  $\Pi(y - v; \Delta_2 - v) = \Pi(y, \Delta_2)$  in the fourth equality in (7.16). Thus the left hand side of (7.13) can be written as

$$\int_{-\infty}^{0} \mathbf{1}_{\Gamma_{1}}(s) ds \int_{s}^{\infty} \pi(s; dv) \mathbf{1}_{\Gamma_{2}}(v) \int_{v}^{\infty} \Lambda_{v}(dy) \mathbf{1}_{\Delta_{1}}(y) \pi(y; \Delta_{2}).$$
(7.17)

Applying successively (7.10) then (7.11) and then again (7.10) to (7.17) we get the following sequence of equalities:

$$\int_{\Gamma} n(du, dv) p(s, (u, v), \Delta) = \int_{0}^{\infty} \pi^{*}(v, \Gamma_{1}) \mathbf{1}_{\Gamma_{2}}(v) dv \int_{v}^{\infty} \Lambda_{v}(dy) \mathbf{1}_{\Delta_{1}}(y) \pi(y; \Delta_{2})$$

$$= \int_{-\infty}^{S} \pi(y; \Delta_{2}) \mathbf{1}_{\Delta_{1}}(y) dy \int_{-\infty}^{y} \lambda^{*}_{y}(dv) \mathbf{1}_{\Gamma_{2}}(v) \pi^{*}(v; \Gamma_{1})$$

$$= \int_{S}^{\infty} \mathbf{1}_{\Delta_{2}}(x) \pi^{*}(x; dy) \mathbf{1}_{\Delta_{1}}(y) \int_{-\infty}^{y} \lambda^{*}_{y}(dv) \mathbf{1}_{\Gamma_{2}}(v) \pi^{*}(v; \Gamma_{1}) .$$
(7.18)

From (7.5) and the analog of (7.16) for  $p^*(\cdot(\cdot, \cdot), -)$  we get that (7.18) equals to the right-hand side of (7.13).

The proof of (7.13) for arbitrary  $\Gamma$  and  $\Delta$  is done in a similar way.

(7.19) COROLLARY. The probability law of the set -M is the same as that of M. In particular M is left Markov. Proof. From proposition (7.11) we see that the processes

 $(\ell_t, r_t) \circ M$  and  $(\ell_t, r_t) \circ (-M)$  have the same one dimensional distribuitons. Proposition (7.12) shows that these two processes have the same backward transition function. Therefore, these two processes have the same law. The rest follows from representation

$$M = \overline{(r \circ M)}_{IR}$$

In the remainder of this section we show why  $(\Pi, \alpha, \lambda, \mu)$ -generated set is not left Markov if  $(\Pi, \alpha, \lambda, \mu)$  is not equivalent to  $(\Pi_1, \alpha_1, \lambda_1, \delta_0)$ .

If M has a perfect regenerative part (i.e.,  $\Pi(\mathbb{R}_{+}) = \infty$ ) then from (2.6) - (2.8) we see that the distribution of  $\nu(k) - \gamma(k)$  is equal to  $\mu$ . If  $\mu \neq \delta_{0}$  then with positive probability  $\nu(k) - \gamma(k) > 0$ . By Fubini's theorem there exists t such that

$$P\{L_{+} < t, L_{+} = \gamma(k)\} > 0$$
.

However, the latter contradicts to Proposition (4.21) (or, to be precise, to the analog of the Proposition (4.21) for left Markov sets.)

If M has a discreet regenerative part (i.e.  $\Pi(\mathbb{R}_+) < \infty$ ) and  $(\Pi, \alpha, \lambda, \mu)$  is not equivalent to  $(\Pi_1, \alpha_1, \lambda_1, \delta_0)$  then

$$\mu \neq c_1 \delta_0 + c_2 \Pi \quad (7.20)$$

From (2.6) - (2.8) we see that the distribution of the length of the jumps of the process y is  $\Pi(\mathbb{R}_{+})^{-1}\Pi$ . The distribution of  $\nu(k) - \gamma(k)$  is

$$\mu^{*} + (1 - \mu\{0\}) \Pi(\mathbb{R}_{+}^{-1}) \Pi$$
 (7.21)

where  $\mu'$  is a restriction of  $\mu$  on  $]0,\infty[$ . If (7.20) is true then (7.21) is not equal to  $\Pi(\mathbb{R}_{+})^{-1} \Pi$ . Elementary calculations show that in this case the conditional distribution of  $R_t - L_t$  given the event

### REFERENCES

- 1. Çinlar, E., Introduction to Stochastic Processes. Prentice-Hall, 1975.
- 2. Dellacherie, C., Capacites et processes stochastiques. Springer, 1972.
- Fitzsimmons, P.J., Fristedt, B., Maisonneuve, B.; Intersections and limits of regenerative sets. Z. Wahrscheinlichkeitstheorie verw. Gebiete 70, 157-173 (1985).
- 4. Hoffmann-Jørgensen, J.; Markov sets. Math. Scand. 24 (1969).
- 5. Krylov, N.V., Yushkevich, A.A.; Markov random sets. Trans. Mosc. Math. Soc. 13, 127-153 (1965).
- 6. Maisonneuve, B.; Ensembles régénératifs, temps locaux et subordinateurs. In Sèminaire de Probabilites V. Lecture Notes in Mathematics 191, Springer 1971.
- 7. Maisonneuve, B.; Systems regeneratifs. Asterisque 15, Société Mathematique de France 1974.
- 8. Maisonneuve, B.; Ensembles régénératifs de la droit. Z. Wahrscheinlichkeitstheorie verw. Gebiete 63, 501-510 (1983).
- 9. Meyer, P.A.; Ensemles régénératifs, d'après Hoffmann-Jørgensen. In Seminaire de Probabilités IV. Lecture Notes in Mathematics 124, Springer 1970.
- Skorohod, A.V.; Stochastic processes with independent increments (in Russian). Nauka, Moscow 1964.
- 11. Taksar, M.I.; Regenerative sets on real line. In Seminaire de Probabilités XIV. Lecture Notes in Mathematics 784, Springer, 1980.

