

AD-A170 211


SEMICONDUCTORS INVESTIGATED BY TIME RESOLVED  
SPECTROSCOPY USING FEMTOSECOND (U) CITY COLL NEW YORK  
ULTRAFAST SPECTROSCOPY AND LASER LAB R R ALFANO


1/1

UNCLASSIFIED

08 MAR 86 447215 AFOSR-TR-86-8500 F/G 20/12

NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

2

AD-A170 211

REPORT DOCUMENTATION PAGE

DTIC

ELECTE

S

JUL 23 1986


D

1a. SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS None	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) 447215		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 86-0500	
6a. NAME OF PERFORMING ORGANIZATION The City College	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Air Force Office of Scientific Research	
6c. ADDRESS (City, State and ZIP Code) New York, N.Y. 10031		7b. ADDRESS (City, State and ZIP Code) AFOSR/PKD Building 410 Bolling AFB, D.C. 20332-6448	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Air Force Office of Scientific Research	8b. OFFICE SYMBOL (If applicable) PKD NE	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR 85-0013	
8c. ADDRESS (City, State and ZIP Code) Bolling AFB, D.C. 20332-6448		10. SOURCE OF FUNDING NOS.	
		PROGRAM ELEMENT NO. 611025	PROJECT NO. 2305
		TASK NO. C1	WORK UNIT NO.
11. TITLE (Include Security Classification) Semiconductors Investigated by Time Resolved Spectroscopy... (U)			
12. PERSONAL AUTHOR(S) Robert R. Alfano			
13a. TYPE OF REPORT Final	13b. TIME COVERED FROM 12/1/84 TO 11/30/85	14. DATE OF REPORT (Yr., Mo., Day) 1986 March 8	15. PAGE COUNT 15
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB. GR.	
		Microstructures, Alloys, Bandgap Discontinuity, Carrier-Phonon Rates in 2D3D, Deformation Potential, Spin Dephasing Time, GaAs-GaAlAs, CdMnSe	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) AFOSR-85-0013			
Four major accomplishments have been achieved to help further the development of faster photonic and electronic devices.			
1. We have shown theoretically and experimentally how one can determine accurately one of the most important parameters in microstructures - the bandgap discontinuity in valence and conduction bands at the heterojunction from the photoluminescence measurements for ultrathin wells in the range of 15 to 80 Å for GaAs/AlGaAs and GaInAs/AlInAs structures.			
2. A model was developed using the electron degeneracy to describe the much slower carrier - optical phonon loss relaxation rate measurements in 2D as compared to 3D.			
(continued)			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION	
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. W.H.		22b. TELEPHONE NUMBER (Include Area Code) 202-767-4931	22c. OFFICE SYMBOL AFOSR/NE

3. The valence and conduction band deformation potentials were separately determined for the first time in semi-magnetic semiconductor alloys of  $\text{Cd}_{1-x}\text{Mn}_x\text{Se}$  from the shift in photoluminescence spectra versus  $x$ . The valence band deformation potential of wurtzite crystals is much larger compared to the one in zinc blende. This technique measures separately the values of valence and conduction band deformation potential instead of the difference between them.

4. The direct picosecond spin dephasing time and degree of spin alignment of photo-excited electrons in semi-magnetic semiconductor alloys of  $\text{Cd}_{1-x}\text{Mn}_x\text{Se}$  was measured. The fast dephasing times arise from the spin exchange between free carriers and the localized  $\text{Mn}^+$  ions.

AFOSR-TR- 86-0500



PROGRESS REPORT  
AFOSR - #85 0013

SEMICONDUCTORS INVESTIGATED BY TIME RESOLVED  
SPECTROSCOPY USING FEMTOSECOND AND  
PICOSECOND LASER TECHNOLOGY

Approved for public release;  
distribution unlimited.

ULTRAFAST SPECTROSCOPY AND LASER  
LABORATORY  
OF  
THE CITY COLLEGE OF NEW YORK  
NEW YORK, NY 10031

DEPARTMENT OF PHYSICS

PROFESSOR R. R. ALFANO

86 7 23 05 6

**PROGRESS REPORT  
AFOSR - #85 0013**

**SEMICONDUCTORS INVESTIGATED BY TIME RESOLVED  
SPECTROSCOPY USING FEMTOSECOND AND  
PICOSECOND LASER TECHNOLOGY**

**R. R. Alfano  
City College of New York  
Institute for Ultrafast Spectroscopy and Lasers  
New York, NY 10031**

**Program Manager: Dr. Gerald Witt**

**Report Date: March 8, 1986**

**RF: 447215**

## Introduction

This report summarizes the progress achieved during the 1984-1985 period in the research effort supported by AFOSR under contract #85 0013.

Four major accomplishments were made during this period in:

1. Microstructure semiconductor research.
2. Electron-degeneracy effects on electron optical phonon energy loss rate for two- and three-dimensional electron systems.
3. Measurement of deformation potential using photoluminescence in magnetic semiconductor alloys  $Cd_{1-x}Mn_xSe$ .
4. Spin polarization and dephasing times of carriers in alloy semiconductors  $Cd_{1-x}Mn_xSe$ .

The following describes the four areas in greater detail.

### 1. Microstructure Research

Band-engineering has produced spectacular photonic devices such as quantum lasers, photodetectors as well as high speed logic elements.<sup>1</sup> The intrinsic asymmetry associated with conduction-band-discontinuity ( $\Delta E_C$ ) and valence-band-discontinuity ( $\Delta E_V$ ) at the interface plays an important role in this "hot" field. For this reason, determining  $\Delta E_C$  and  $\Delta E_V$  accurately is of considerable interest and importance.

We have calculated the  $\Delta E$  vs  $L_z$  for various quantum well structures using their  $\Delta E_V$  as a parameter and various sets of masses for heavy-hole (hh) and light-hole (lh). It was found that there are always two distinct regions for the well widths  $L_z$  to find  $\Delta E_V$  accurately from those calculated curves. One region is called sensitive range ( $15\text{\AA} < L_z < 80\text{\AA}$ ) in which  $\Delta E$ 's are very sensitive to the chosen value of  $\Delta E_V$  in spite of various sets of masses for hh and lh. In this region, it is possible to determine  $\Delta E_V$  very accurately by fitting  $\Delta E$ 's to experimental data. The other region ( $L_z > 80\text{\AA}$ ) is called insensitive range for  $\Delta E$ 's to

the value of  $\Delta E_v$ . By this feature, we can state that it is almost impossible to determine  $\Delta E_v$  accurately using the optical transitions in thick quantum wells.

The calculated result<sup>2</sup> of  $\Delta E$  vs  $L_z$  for GaAs/(Al,Ga)As structure in fig. 1 was compared with the limited data available in the sensitive range yielding a good agreement with dingle's 15-85 rule:  $\Delta E_v = 0.15 \Delta E_g$ , where  $\Delta E_g$  is the difference of the bandgaps at the interface.

By measuring the  $\Delta E$ 's for two ultrathin 14.5-Å and 19.3-Å quantum wells from photoluminescence spectra<sup>3</sup> (inset of Fig. 1), we determined band-discontinuities for (Ga,In)As/(Al,In)As structure without any ambiguity which agrees with Dingle's rule.

Our experimental and theoretical work support Dingle's rule and question the recent work of Miller at Bell Telephone Laboratories who gives the relationship of  $\Delta E_v = 0.4 \Delta E_g$ .

The theoretical  $\Delta E$  vs  $L_z$  curves can be used to determine the band-discontinuities accurately for other microstructures. The key is to use ultrathin wells ( $L_z < 80 \text{Å}$ ).



Determination of the band-discontinuity of the heterojunctions




Fig. 1

$\Delta E$  - The calculated energy separation between heavy-hole and light-hole subbands. The width of the quantum well. ( $L_z$ )

2. Electron Degeneracy Effects on Electron-Energy-Loss Rate for two-and three-dimensional Electron Systems

Information about the hot carrier energy relaxation in semiconductors is important for the design and fabrication of high speed and high field ultrasmall devices. It is crucial to understand the underlying mechanisms which lead to a much smaller electron-energy-loss rate for two-dimensional (2D) electron system


much smaller electron-energy-loss rate for two-dimensional (2D) electron system than for three-dimensional (3D) electron system.

Previous studies<sup>5-9</sup> focused on how electron-phonon interaction is modified in 2D electron system in order to explain why electron-energy-loss rate for 2D case is much smaller than 3D case. These studies have generated conflicting interpretations regarding the influence of reduced dimensionality and plasma density on the electron-energy-loss rates. The conventional calculation of the rates from Fermi's Golden Rule as numerically calculated by Basu et al.<sup>8</sup> with and without the effects of degeneracy does not clear up this controversial issue.

We<sup>10</sup> primarily focused on the final available energy width for electrons to be scattered back for each electron system without involving intermediate states and the calculation of electron-phonon interaction matrix elements. The calculated ratio  $\beta$  of electron-energy-loss rates for 3D and 2D electron systems are shown in Fig. 2 based on difference of electron-degeneracy for 3D and 2D cases for different quasi-Fermi-energies (in meV).

The striking conclusions<sup>10</sup> extracted from Fig. 2 enable us to uniquely explain various experimental results.<sup>4-7</sup> The circles indicate our result<sup>4</sup> that  $\beta$  is about 4 obtained from detailed power balance for a set of single quantum well. The solid triangle and solid circle are the experimental data from Ryan et al.<sup>6</sup> as well as Xu and Tang<sup>7</sup>, respectively. Comparing our model to the most recent experimental data by Shah et al.<sup>5</sup>, not only the much lower electron-energy-rate for 2D than 3D can be explained, but also a quantitative agreement of the trend of  $\beta$  vs  $T_c$  can be achieved. This is shown in the inset of Fig. 2.

Fig. 2 Electron-degeneracy effects on electron-energy-loss rate for two- and three-dimensional electron systems



$\beta$  - The calculated ratio of electron-energy-loss rates for 3D and 2D electron systems.  $T_c$  - The electron temperature.

### 3. Measurement of Deformation Potential using Photoluminescence Cd<sub>1-x</sub>Mn<sub>x</sub>Se Alloys

For the first time, we have pointed out a unique method of measuring optical deformation potential in semiconductors using laser induced fluorescence. In CdMnSe (semi-magnetic semiconductor), the substitution of Mn at cation site (Cd), introduces a local pressurized environment. The pressure is equivalent to exter-

nally applied hydrostatic pressure on pure CdSe. This pressure manifested itself as optical deformation potential, will cause the conduction band and valence band to shift in opposite directions with respect to vacuum level. The increased or decreased bandgap could be monitored using the above bandgap laser photon excitation photoluminescence. This is a first time separate determination of conduction and valence band deformation potential. In the past only the difference was measured. This also points out a large valence band deformation potential in wurtzite crystals compared to the one in zinc blende.

In figure 3, the increase in the bandgap is shown as a function of Mn concentration. Up to Mn concentration of 30%, the Mn localized states remain above the conduction band and the photoluminescence peak shifts are corresponding to band-band transition. The slope for  $x < 0.3$  corresponds to joint (conduction and valence band ( $1.2 \text{ eV}/x$ )) optical deformation potential, for  $x > 0.3$  the photoluminescence peak is due to Mn ( $4G$ )-valence band transition. In this case, the slope ( $0.4 \text{ eV}/x$ ) corresponds only to the valence band as  $4G$  Mn is almost stationary with respect to the vacuum level. Using the x-ray data of lattice change with Mn concentration we deduced the conduction and valence band optical determination potential as shown in the figure.

Fig. 3 Bandgap shifts

ACCOMPLISHMENT

MEASUREMENT


OF DEFORMATION POTENTIAL

USING PHOTO LUMINESCENCE

CONDUCTION AND VALENCE

BAND SHIFTS IN  $CdMnSe$

Accomplishment-Measurment of deformation  
using photoluminescence in  $Cd_{1-x}Mn_xSe$



4. Spin Polarization and Dephasing Time of Carriers in Alloy Semiconductors  
Cd<sub>1-x</sub>Mn<sub>x</sub>Se

For over 10 years, it was known that the dipole allowed circularly polarized above bandgap photon can induce a net spin alignment of photogenerated electrons. The observed ratio of spin up:spin down is reflected in the recombination photoluminescence. Most past work used steady state techniques to estimate spin alignment. One can now determine the percentage of spin alignment and spin dephasing time. The experiments we carried out were with single pulse excitation using Nd:glass laser having pulse width of 6 psec. The time resolved detection was achieved using high resolution streak camera. This is a major accomplishment.





Here we review the first direct measurements of spin relaxation times in wurtzite CdSe and semi-magnetic semiconductor CdMnSe. Our past work was on spin relaxation and alignment in GaAs which was previously reported to you.

The figures 4a, b, and c show time resolved photoluminescence from CdSe and magnetic semiconductor Cd<sub>0.95</sub>Mn<sub>0.05</sub>Se and Cd<sub>0.9</sub>Mn<sub>0.1</sub>Se. The photoluminescence has the degree of polarization on the order of 30% at t=0, which also corresponds to spin polarization factor of ~ 30%. Spin polarization factor is defined as  $p(t) = \frac{n_{\uparrow} - n_{\downarrow}}{n_{\uparrow} + n_{\downarrow}}$  and decays in time with a single exponential. Each curve shows circular polarized luminescence for 6+ and 6- (right and left handed, respectively). The measured values of p(0) and spin relaxation time T<sub>S</sub> in CdSe are ~ 45 and 30 psec, respectively. The measured values of p(0) and spin relaxation time T<sub>S</sub> in CdMnSe with 5% Mn are 20% and 16 psec, respectively. The measured value of p(0) and spin relaxation time in CdMnSe with 10% Mn are 28% and 20 psec, respectively. The spin relaxation in pure CdSe is assigned to a mechanism due to non-centrosymmetric nature of crystal lattice. In CdMnSe the spin relaxation occurs due to two

equally important mechanisms - one due to non-centrosymmetry of lattice and the other due to spin exchange interaction between free electrons and localized  $Mn^{2+}$  ions.

The spin relaxation rates are plotted in the figure 4(d) and compared to a theoretical fit based on spin exchange interaction. More work is needed with higher concentration of Mn in CdMnSe in order to check the validity of the theoretical model.

Figure 4 Spin Dephasing time and polarization





References

1. Daniel S. Chemla, Phys. Today 57 (May 1985).
2. K. Shum, P. P. Ho, R. R. Alfano (submitted to Phys. Rev. for publication).
3. K. Shum, P. P. Ho, R. R. Alfano, D. F. Welch, G. W. Wicks, and L. F. Eastman, Phys. Rev. B32, 3806 (1985).
4. K. Shum, P. P. Ho, R. R. Alfano, D. F. Welch, G. W. Wicks, and L. F. Eastman (submitted to Phys. Rev.B for publication).
5. J. Shah, A. Pinczuk, A. C. Gossard, and W. Wiemann, Phys. Rev. Lett. 54, 2045 (1985).
6. J. F. Ryan, R. A. Taylor, A. J. Turberfield, Angela Maciel, J. M. Worlock, A. C. Gossard, and W. Wiemann, Phys. Rev. Lett. 53, 1841 (1984).
7. Z. Y. Xu and C. L. Tang, Appl. Phys. Lett. 44, 692 (1984).
8. P. K. Basu and Sudakshina Kundu, Appl. Phys. Lett. 47, 264 (1985).
9. F. A. Riddoch and B. K. Ridley, J. Phys. C 16, 6717 (1983) and J. P. Leburton, J. Appl. Phys. 56, 2850 (1984).
10. K. Shum and R. R. Alfano, Phys. Rev. (submitted).

## Publications

- Semiconductors Probed by Ultrafast Laser Spectroscopy, Vol 1 and 2, R. R. Alfano, Academic Press (1984).
- Photoluminescence of Spin-Polarized Electrons in Semiconductors, E. J. Johnson, R. J. Seymour, R. R. Alfano, in "Semiconductors Probed By Ultrafast Laser Spectroscopy", Vol. II, ed. R. R. Alfano, Academic Press, NY (1984).
- Techniques in time-Resolved Luminescence Spectroscopy, D. L. Rosen, A. G. Doukas, A. Katz, Y. Budansky, R. R. Alfano, in "Semiconductors Probed By Ultrafast Laser Spectroscopy", Vol. II, ed. R. R. Alfano, Academic Press, NY (1984).
- Optical Investigations of Chromium Chalcogenide Spinel in Magnetic Semiconductors, W. Miniscalco, A. Lempicki, S. S. Yao, R. R. Alfano, N. G. Stoffel, G. Margaritondo, in "Semiconductors Probed By Ultrafast Laser Spectroscopy", ed. R. R. Alfano, Academic Press, NY (1984).
- Colliding Pulse Mode Locking for an Antiresonant Cavity of a Nd:Glass Laser, J. M. Buchert, D. K. Basa, C. Tzu, and R. R. Alfano, J. Appl. Phys. 55, 683 (1984).
- Optical Properties and Photoexcitation of a Novel Liquid Form of Soluble Polyacetylene, R. Tubino, R. Dorsinville, W. Lam, R. R. Alfano, J. L. Birman, A. Bolognesi, S. Destri, M. Catellani and W. Porzio, Phys. Rev. B30, 6601 (1984).
- Photon Echo in Direct Gap Semiconductors, J. T. Manassah, R. R. Alfano, M. Conner and P. Ho, Physics Lett. 106A, 65-71 (1984).
- Ultrafast Supercontinuum Laser Source, J. T. Manassah, P. P. Ho, A. Katz and R. R. Alfano, Photonics, p53-59, Nov. (1984).
- Introduction to the Special Issue on Lasers in Biology and Medicine, R. R. Alfano and A. G. Doukas, IEEE J. of Quant. Elec. QE-20, 1342 (1984).
- Ultrafast Supercontinuum Laser Source, J. T. Manassah, P. P. Ho, A. Katz and R. R. Alfano, Photonics Spectra, p. 53-59 Nov. (1984).
- Photoluminescence Determination of the Pressure and Temperature of the Shock Wave Induced by a Picosecond Laser Pulse in the Layered Semiconductor GaSe, K. P. Leung, S. S. Yao, A. G. Doukas, R. R. Alfano, P. Harris, Phys. Rev. B31, 942 (1985).
- Nonlinear and Time Resolved Studies of Native Defects in CdSe, D. L. Rosen, Q. X. Li, R. R. Alfano, SPIES (1/85), 524, 95 (1985).
- Time Resolved Kinetics of e-h Plasmas in GaAsP under Intense Picosecond Laser Pulse Excitation, H. Zarrabi, W. B. Wang, R. R. Alfano, App. Phys. Lett. (1985).
- Time Resolved Photoluminescence Spectra of  $Ga_xIn_{1-x}P$  under Picosecond Laser Excitation Measured by a Streak Camera, H. Zarrabi, R. R. Alfano, Phys. Rev. B32, 3947 (1985).
- Picosecond Pulses Produced by Mode Locking an Nd:Glass Laser with Kodak Dye #26, N. H. Schiller, M. Foresti, R. R. Alfano, J. Opt. Society Physics B2, 729 (1985).
- Native Defects in Semi-insulating CdSe Studied by Photoluminescence and Absorption, D. L. Rosen, Q. X. Li, R. R. Alfano, Phys. Rev. B4, 2396 (1985).
- Picosecond Excite-and-Probe Absorption Measurement of the  $T_2$  State Nonradiative Lifetime in Ruby, S. K. Gayen, W. B. Wang, V. Petricevic, R. Dorsinville, R. R. Alfano, Appl. Phys. Lett. 47, 455 (1985).
- Infrared Photoinduced Absorption Studies in Soluble Trans-Polyacetylene, R. Dorsinville, R. Tubino, S. Krimchansky, R. R. Alfano, J. L. Birman, A. Bolognesi, S. Destri, M. Catellani, W. Porzio, Phys. Rev. B32, 3377 (1985).
- Determination of Temporal Correlation of Ultrafast Laser Pulses Using Phase Conjugation, J. Buchert, R. Dorsinville, P. Delfyett, S. Krimchansky, R. R. Alfano, Optics Communications 52, 433 (1985).

- Photoluminescence Determination of Well Depth of the  $\text{Ga}_{0.47}\text{In}_{0.53}\text{As}/\text{Ga}_{0.48}\text{In}_{0.52}\text{As}$  Single Quantum Well, K. Shum, P. P. Ho, R. R. Alfano, D. F. Welch, L. F. Eastman, Phys. Rev. B32, 3806 (1985).
- Time Response of Ultrafast Streak Camera System Using Femtosecond Laser Pulses, P. Ho, A. Katz, R. R. Alfano, N. Schiller, Opt. Comm. 54, 57 (1985).
- The Use of Short Lived Fluorescent Dyes to Correct for Artifacts in the Measurement of Fluorescence Lifetime, P. Ho, D. Mauzerall, R. R. Alfano, Photochem & Photobiol. 42, 183 (1985).
- Femtosecond Laser Pulse Compression Using Volume Phase Transmission Holograms, P. Ho, R. R. Alfano, T. Yang, A. Katz, R. Ferrante, App. Optics 24, 2021 (1985).
- Calibrating a Streak Camera, R. Dorsinville, R. R. Alfano, N. Schiller, The Photonics Design & Applications Handbook 31, H-18 (1985).
- Spectral Distribution of an Ultrafast Supercontinuum Laser Source, J. T. Manassah, R. R. Alfano and M. Mustafa, Physics Lett. 107A, 305 (1985).
- Generation of Infrared Supercontinuum Covering from 3  $\mu\text{m}$  to 14  $\mu\text{m}$  Wavelength in Dielectrics and Semiconductors, P. B. Corkum, P. P. Ho, R. R. Alfano, Optics Letters 54, (1985).
- Photoluminescence of Magnetic Semiconductor  $\text{CdCr}_2\text{Se}_4$ , P. P. Ho, R. R. Alfano et al., IEEE Qe-22, 205 (1986).

Ph.D. Thesis at the Institute for Ultrafast Spectroscopy  
and Lasers

The City College of New York

<u>TOPIC/DATE/FUNDING AGENCY</u>	<u>PRESENT POSITION</u>
1. "Picosecond and Steady State Spectroscopy of the Wurtzite Semimagnetic Semiconductor $Cd_{1-x}Mn_xSe$ " by Mahesh Junnarkar, 1986, AFOSR, NSF	IUSL
2. "Time Resolved Spectroscopy of Ternary Semiconductors $GaAs_{1-x}P_x$ and $Ga_xIn_{1-x}P_x$ under Picosecond Laser Pulse Excitation" by Hassan J. Zarrabi, 1985, AFOSR	General Optronics
3. "Picosecond and Steady State Spectroscopy of Defects in Semi-Insulating CdSe" by David L. Rosen, 1985, AFOSR	NRL
4. "A Study of Energy Transfer in the Photosynthetic Blue-Green Algae Nostoc Sp. probed by Picosecond Spectroscopy" by Aaron Dagen, 1985, NSF	Perkin Elmer
5. "Energy Transfer between Dye Molecules Investigated by Steady State and Time Resolved Spectroscopy" by Poyang Lu, 1982, AFOSR, NSF	IBM
6. "The Emission and Absorption Characteristics of Chalcogenide Semiconductors using Steady State and Time Resolved Picosecond Spectroscopy" by Shingshwang Yao, 1982, AFOSR	IBM
7. "Electron Spin and Energy Relaxation in Highly Photoexcited Gallium Arsenide" by Robert J. Seymour, 1981, AFOSR, NSF	GTE
8. "Carrier Transport in Amorphous Silicon utilizing Picosecond Photoconductivity" by Anthony M. Johnson, 1981, BTL	Bell Labs
9. "Energy Transfer in the Primary Stages of the Photosynthetic Process investigated by Picosecond Time Resolved Fluorescence Spectroscopy" by Francesco Pellegrino, 1981, NSF	Sperry
10. "Reorientational Relaxation Kinetics of Polyatomic Molecules in Different States of Condensed Media investigated by Picosecond Laser Pulse Induced Kerr Effect" by Ping-Pei Ho, 1978, NSF	CCNY Prof. EE

Ph.D. Thesis in Progress

1. "Ultrafast Transient Diffraction Gratings of Photoexcited Carriers in GaAs Structures" by A. Katz, AFOSR
2. "Ultrafast Quantum Well Physics" by Kai Shum, H. Chao, P. Ryerson, AFOSR
3. "Picosecond Raman Induced Phase Conjugation in Semiconductors and Polymers" by P. Delfyett, A. Walser, AFOSR
4. "Laser Induced Shock Wave Interaction Physics in Solids and Liquids" by S. Lee, B. Willman, ONR
5. "Nonradiative Relaxation Dynamics in Tunable Solid State Lasers" by V. Petricevic, B. Wang, ARO, NASA
6. "IR Phase Conjugation in Semiconductors" by N. Chen, AFOSR
7. "Ultrafast Interference Optical Computation" by Yao Li, AFOSR
8. "Spin Relaxation of Carriers in Semiconductors" by Winston Lam, X. Liu, NSF, AFOSR
9. "Supercontinuum Generation" by V. Caplan, X. Wang, NSF, AFOSR

END

10-86

DTIC