AD-A170 386	EXPLOITIN	NG AUXILIAR	Y INFOR	MATIO		T EXAM	INEES L TEST	IN THE	171	
UNCLASSIFIED	ETS-RR-8	RINCETON N 5-18-ONR NO	001 4-85	6-K-068	33	80	F/G 1	2/1	NL	
										í
		<i>.</i>				_,				



RR-86-18-ONR

170

28

7

86









EXPLOITING AUXILIARY INFORMATION ABOUT EXAMINEES IN THE ESTIMATION OF ITEM PARAMETERS

Robert J. Mislevy

This research was sponsored in part by the Personnel and Training Research Programs Psychological Sciences Division Office of Naval Research, under Contract No. N00014-85-K-0683

Contract Authority Identification Number NR No. 150-539

Robert J. Mislevy, Principal Investigator



, Educational Testing Service Princeton, New Jersey

May 1986

Reproduction in whole or in part is permitted for any purpose of the United States Government.

Approved for public release; distribution unlimited.

DITIC FILE COPY

		AD-A170306				
	REPORT DOCU	MENTATION PAGE				
REPORT SECURITY CLASSIFICATION		16 RESTRICTIVE MARKINGS				
SECURITY CLASSIFICATION AUTHORITY	<u></u>	3 DISTRIBUTION / AVAILABILITY OF REPORT				
DECLASSIFICATION / DOWNGRADING SCHE	DUUE	Approved for public release;				
		distribution unlimited.				
PERFORMING ORGANIZATION REPORT NUN RR-86-18-ONR	1BER(S)	5 MONITORING ORGANIZATION REPORT NUMBER(S)				
NAME OF PERFORMING ORGANIZATION	6b OFFICE SYMBOL	7a NAME OF MONITORING ORGANIZATION				
Educational Testing Service	(If applicable)	Personnel and Training Research Programs				
ADDRESS (City, State, and ZIP Code)		Office of Naval Research 7b ADDRESS (City, State, and ZIP Code)				
Princeton, NJ 08541		Code 442PT				
,		800 North Quincy Street Arlington, VA 22217-5000				
NAME OF FUNDING SPONSORING	8b OFFICE SYMBOL					
ORGANIZATION Office of Naval Research	(If applicable)	N00014-85-K-0683				
ADDRESS (City, State, and ZIP Code)		10 SOURCE OF FUNDING NUMBERS				
800 North Quincy Street		PROGRAM PROJECT TASK WORK UNIT ELEMENT NO NO NO ACCESSION N				
Arlington, VA 22217-5000		61153N RR04204 RR04204-01 4421539				
PERSONAL AUTHOR(S) Mislevy, Robert J.		14 DATE OF REPORT (Year, Month, Day) 15 PAGE COUNT				
echnical Report FROM_	TO	May 1986 42				
echnical Report FROM_						
echnical Report FROM_ SUPPLEMENTARY NOTATION	TO					
echnical Report FROM_ SUPPLEMENTARY NOTATION	18 SUBJECT TERMS EM-algorithm	(Continue on reverse if necessary and identify by block number) marginal maximum likelihood				
COSATI CODES	TOTOTOTOTO	(Continue on reverse if necessary and identify by block number) marginal maximum likelihood				
COSATI CODES FIELD GROUP SUB-GROUF	TO 18 SUBJECT TERMS EM-algorithm empirical Bay	May 1986 (Continue on reverse if necessary and identify by block number) marginal maximum likelihood res				
COSATI CODES FIELD GROUP SUB-GROUF ABSTRACT (Continue on reverse if necesse The precision of item	TO 18 SUBJECT TERMS EM-algorithm empirical Bay any and identify by block parameter estimat	May 1986 (Continue on reverse if necessary and identify by block number) marginal maximum likelihood res number) es can be increased by taking advantage of				
echnical Report FROM_ 5 SUPPLEMENTARY NOTATION 5 SUPPLEMENTARY NOTATION 7 COSATI CODES FIELD GROUP SUB-GROUF ABSTRACT (Continue on reverse if necessary The precision of item precision	TO 18 SUBJECT TERMS EM-algorithm empirical Bay any and identify by block parameter estimat tent proficiency	May 1986 42 (Continue on reverse if necessary and identify by block number) marginal maximum likelihood res number) es can be increased by taking advantage of variable and auxiliary examinee variables suc				
COSATI CODES FIELD GROUP SUB-GROUF ABSTRACT (Continue on reverse if necessa The precision of item dependencies between the lat as age, courses taken, and	TO 18 SUBJECT TERMS EM-algorithm empirical Bay any and identify by block parameter estimat tent proficiency years of schoolin	May 1986 (Continue on reverse if necessary and identify by block number) marginal maximum likelihood es number) es can be increased by taking advantage of variable and auxiliary examinee variables suc g. Gains roughly equivalent to two to six				
COSATI CODES FIELD GROUP SUB-GROUP ABSTRACT (Continue on reverse if necessa The precision of item p dependencies between the lat as age, courses taken, and y additional item responses ca applications. Empirical Bay	18 SUBJECT TERMS EM-algorithm empirical Bay any and identify by block parameter estimat tent proficiency years of schoolin an be expected in yes computational	May 1986 (Continue on reverse if necessary and identify by block number) marginal maximum likelihood res number) es can be increased by taking advantage of variable and auxiliary examinee variables suc g. Gains roughly equivalent to two to six typical educational and psychological procedures are presented, and illustrated				
echnical Report FROM	18 SUBJECT TERMS EM-algorithm empirical Bay any and identify by block parameter estimat tent proficiency years of schoolin an be expected in yes computational	May 1986 (Continue on reverse if necessary and identify by block number) marginal maximum likelihood res number) es can be increased by taking advantage of variable and auxiliary examinee variables suc g. Gains roughly equivalent to two to six typical educational and psychological procedures are presented, and illustrated				
COSATI CODES FIELD GROUP SUB-GROUP ABSTRACT (Continue on reverse if necessa The precision of item p dependencies between the lat as age, courses taken, and y additional item responses ca applications. Empirical Bay	18 SUBJECT TERMS EM-algorithm empirical Bay any and identify by block parameter estimat tent proficiency years of schoolin an be expected in yes computational	May 1986 (Continue on reverse if necessary and identify by block number) marginal maximum likelihood res number) es can be increased by taking advantage of variable and auxiliary examinee variables suc g. Gains roughly equivalent to two to six typical educational and psychological procedures are presented, and illustrated				

CCCC CO

and the second states of the

20 DISTRIBUTION / AVAILABILITY OF		21 ABSTRACT SECURITY CLASSIFICATION		
UNCLASSIFIED/UNLIMITED	SAME AS RPT DTIC USERS	Linclassified		
22a NAME OF RESPONSIBLE INDIVID	DUAL	22b TELEPHONE (Include Area Code) 22c OFFICE SY	MBOL	
Robert J. Mislevy		(609) 734-1271		
DD FORM 1473, 84 MAR	83 APR edition may be used un	SECURITY CLASSIFICATION (OF THIS PAGE	
	All other editions are ob	Unclassified	lassified	

1

EXPLOITING AUXILIARY INFORMATION ABOUT EXAMINEES IN THE

ESTIMATION OF ITEM PARAMETERS

Robert J. Mislevy

This research was sponsored in part by the Personnel and Training Research Programs Psychological Sciences Division Office of Naval Research, under Contract No. N00014-85-K-0683

Contract Authority Identification Number NR No. 150-539

Robert J. Mislevy, Principal Investigator

Educational Testing Service Princeton, New Jersey

May 1986

Reproduction in whole or in part is permitted for any purpose of the United States Goverment.

Approved for public release; distribution unlimited.

U	·
A	vailability Codes
Dilt	Avail and / or Special
A-1	

2

Abstract

The precision of item parameter estimates can be increased by taking advantage of dependencies between the latent proficiency variable and auxiliary examinee variables such as age, courses taken, and years of schooling. Gains roughly equivalent to two to six additional item responses can be expected in typical educational and psychological applications. Empirical Bayes computational procedures are presented, and illustrated with data from the <u>Profile of American Youth</u> survey.

Key words: EM-algorithm, empirical Bayes, marginal maximum likelihood

3

Exploiting Auxiliary Information about Examinees in the

Estimation of Item Parameters

A pervasive problem in item response theory (IRT) is the difficulty of simultaneously estimating large numbers of parameters from limited data. Even large samples of examinees may not eliminate the problem when each examinee responds to only a few items, as in educational assessment and adaptive testing. Certain improvements are obtained by using hierarchial models along the lines of Lindley and Smith (1972); treating examinee parameters as a sample from a common population enhances the stability and precision of item parameter as well as examinee parameter estimates. This approach has been applied to IRT by a number of researchers recently, including Bock and Aitkin (1981), Leonard and Novick (1985), Rigdon and Tsutakawa (1982), and Swaminathan and Gifford (1982).

For the most part, the aforementional writers consider all examinees to be members of a single, undifferentiated, population. This framework instantiates such beliefs as, "if the parameters of most examinees seem to lie between -3 and +3, then the parameter of an examinee who answered both of two hard math items correctly is probably somewhere between +1.5 to +3.5--even though his/her maximum likelihood estimate is $+\infty$." Additional stability and precision may yet be achieved if auxiliary information is

available about examinees, such as educational background or status on demographic variables. A statement like "the parameter of an examinee who answered both of two hard math items correctly <u>and</u> studied calculus in college is probably between +2.7 and +3.7," might result.

This paper addresses the utilization of auxiliary information about examinees in estimating item parameters. The following section reviews item parameter estimation when examinee parameters are known, then when examinee parameters are unknown and nothing is assumed about them. Attention then turns to the additional assumptions of first, an undifferentiated population, and second, a population differentiated with respect to auxiliary variables. Following this are sections that discuss anticipated gains in precision, outline computational procedures, and illustrate the approach with responses to four items from the Arithmetic Knowledge subtest of the Armed Services Vocational Aptitude Battery.

The Role of Auxiliary Information

The relevance of auxiliary examinee variables to item parameter estimation is not immediately obvious, since they play no role in the basic model for item responses. Letting $x_i = (x_{i1}, \dots, x_{in})$ represent the responses of examinee i to n test items and y_i represent values of auxiliary variables such as educational and demographic status, the standard IRT assumption of local

5

independence states that

$$p(\mathbf{x}_{i}|\boldsymbol{\theta}_{i}, \mathbf{y}_{i}, \boldsymbol{\beta}) = \prod_{j}^{n} p(\mathbf{x}_{ij}|\boldsymbol{\theta}_{i}, \boldsymbol{\beta}_{j}) , \qquad (1)$$

where θ_i is the examinee parameter, $\beta = (\beta_1, \dots, \beta_n)$ are possibly vector-valued item parameters, and the form of $p(x_{ij} | \theta_i, \beta_j)$ is specified a priori through the item response model. It follows that y_i would indeed be irrelevant to item parameter estimation if θ_i were known. The likelihood to be maximized with respect to β , given the data matrix $X = (x_1, \dots, x_N)$ of responses from N examinees with proficiencies $\theta = (\theta_1, \dots, \theta_N)$ and auxiliary variables $Y = (y_1, \dots, y_N)$, would be simply

$$L = \prod_{i=1}^{N} p(\mathbf{x}_{i} | \boldsymbol{\theta}_{i}, \boldsymbol{\beta}) \qquad (2)$$

The maximum likelihood estimate (MLE) β would satisfy the likelihood equations

$$D = \sum_{i} \partial \ell_{i}(\theta_{i}) / \partial \beta , \qquad (3)$$

where $\ell_i(\theta) = \log p(x_i | \theta, \beta)$, and the covariance matrix of

6

estimation error variances for β could be approximated by the inverse of the observed information matrix I:

$$I_{\theta} = \sum_{i} \left(\frac{\partial \ell_{i}(\theta_{i})}{\partial \beta} \right) \left(\frac{\partial \ell_{i}(\theta_{i})}{\partial \beta'} \right) \Big|_{\beta = \hat{\beta}}$$
(4)

But Equation 1 gives response probabilities conditioned on θ , and θ is not known in practice. The problem that must actually be solved is to maximize the marginal likelihood

$$L_{M} = \prod_{i} f p(\mathbf{x}_{i} | \boldsymbol{\theta}, \boldsymbol{\beta}) dF_{i}(\boldsymbol{\theta}) , \qquad (5)$$

where $F_i(\theta)$ is the distribution of the unknown proficiency of examinee i. This is an "incomplete data" problem, in the terminology of Dempster, Laird, and Rubin (1977), corresponding to the "complete data" problem of maximizing Equation 2 when θ is known. Assuming the required integrals exist, the likelihood equations become

$$0 = \sum_{i} p_{i}^{-1}(\mathbf{x}_{i}) \int \left[\frac{\partial \ell_{i}(\theta)}{\partial \beta} \right] dF_{i}(\theta)$$

where

7

$$p_{i}(x_{i}) = \int \ell_{i}(\theta) dF_{i}(\theta)$$

Louis (1982) shows that if Zack's (1971, Chapter 5) regularity conditions are met and if F_i is known for all i, the diagonal elements of the incomplete-data observed information matrix, namely

$$I_{x} = \sum_{i} p_{i}^{-1}(x_{i}) \int \left\{ \left(\frac{\partial \ell_{i}(\theta)}{\partial \beta} \right) \left(\frac{\partial \ell_{i}(\theta)}{\partial \beta'} \right) \right|_{\beta = \hat{\beta}} \right\} dF_{i}(\theta) , (6)$$

cannot exceed the diagonal elements I_{θ} . In other words, the precision with with elements of $\hat{\beta}$ would be estimated if θ were known provides an upper limit to the precision to be expected when θ is not known but must be inferred.

A similar phenomenon arises in the context of sample survey analysis when a clustered sampling design is employed to estimate a mean. If n units are sampled from each of N randomly-selected clusters, then the squared standard error of the mean, ignoring finite population corrections, is given as

8

$$SEM^{2} = \frac{\sigma^{2}}{nN} [1 + (n - 1)\rho]$$

where σ^2 is the population variance and ρ is the intraclass correlation coefficient indicating within-cluster homogeneity. If the number of clusters (N) is held constant, increasing the sample size (n) within clusters cannot decrease SEM² below $\rho\sigma^2/N$, the the value of SEM² obtained when the means of the sampled clusters are known without error.

The estimation of β in the context of IRT must also deal with uncertainty from two sources. First is the usual limitation of having data from only a finite sample of examinees. All other conditions remaining unchanged, increasing N leads to greater precision for $\hat{\beta}$. Second is the limitation that θ remains unknown even for sampled examinees. For a fixed sample of examinees, reducing uncertainty about θ leads to greater precision for $\hat{\beta}$. This can be achieved through (i) item responses, (ii) assumptions about the F_i's and (iii) auxiliary variables related to θ .

de Leeuw and Verhelst (1984) point out that finding maxima in terms of β and of each individual θ_i in the manner suggested by Birnbaum (1968) is equivalent to maximizing Equation 5 when each F_i concentrates its mass at the single (unknown) point θ_i . This joint maximum likelihood (JML) solution utilizes only information in responses x_i from examinee i to reduce uncertainty about θ_i .

9

Alternatively, one may consider the θ 's to be identically distributed, so that $F_i = F$ for all i. An auxiliary variable y is thereby implied for all examinees, an indicator signifying that each is a member of the population whose distribution is specified by F. Appearing in the literature are treatments that assume a completely specified form for F (e.g., Bock & Lieberman, 1970), others that assume parametric forms with unknown parameters α to be estimated along with β (e.g., Zwarts & Veldhuesen, 1985), and still others that provide nonparametric approximations (e.g., Tjur, 1982). Under the first of these three approaches, the assumed population distribution combines with x_i to produce $p(\theta_i | X)$, which in this case equals $p(\theta_i | x_i)$. Under the latter two approaches, responses from examinees other than examinee i also play a role in estimating F so that $p(\theta_i | x_i) \neq p(\theta_i | X)$.

A third alternative, falling between unique, unconstrained F_i 's and identical F_i 's, is to posit distributions that depend on auxiliary variables: that is, $F_i(\theta) = F_{y_i}(\theta)$. Examinees with identical y values are considered a random sample from a population indexed by that particular value of y, and these conditional distributions are allowed to vary with y. A following section gives details for two special cases, namely a linear model and a (quasi-) nonparametric mixture approximation.

10

How Much Can Be Gained?

Several factors contribute to the magnitude of the precision gains that can be achieved through population assumptions and auxiliary variables. One factor is the sensitivity of different model parameters to missing information. Mislevy's (1984) analysis of Bock and Lieberman's (1970) LSAT data showed that estimates of the population variance were more substantially improved by increases in test length than were estimates of the population mean. This might lead one to expect increased information about θ to have more effect on item slopes than on item thresholds in the context of item parameter estimation.

A second factor is the nature of the joint distribution of auxiliary variables with θ . An auxiliary variable adept at identifying low proficiency examinees, for example, adds information for those examinees most useful for estimating lower asymptote item parameters.

A third factor is the dependence of the estimated information upon estimated parameter values. Although a slope parameter may be consistently estimated under both the undifferentiated and undifferentiated population models, a higher estimate under the latter may <u>appear</u> less precise. This is because <u>estimated</u> standard errors for slopes are directly proportional to the values of the slope estimates, even though <u>true</u> standard errors depend on true

11

slope values and not their estimates. A slope estimated with the aid of auxiliary variables and obtaining a higher estimate can thus have a lower <u>true</u> standard error but a higher <u>estimated</u> standard error.

Since the same factors determine information gain from both increased test length and auxiliary variables, however, it is reasonable to consider the contribution of auxiliary variables in units of additional item responses. In the special case of dichotomous items, the amount of information conveyed by item responses alone is

$$i(\theta) = \sum_{j} \frac{P'(\theta)^{2}}{P_{j}(\theta)[1 - P_{j}(\theta)]}$$

where $P_j(\theta) = p(x_j = 1 | \theta)$ and $P_j(\theta) = dP_j(\theta)/d\theta$. For examinees with finite maximum likelihood estimates, Bayes theorem applied with a diffuse prior leads to the approximation $p(\theta | x_i) \stackrel{*}{\sim} N(\theta, \sigma_x^2)$ with $\sigma_x^2 = i^{-1}$. This follows by first rescaling the likelihood so that it integrates to one, then using its mode and curvature at the mode in a normal approximation.

Consider as an example the two-parameter logistic model, under which $P_j(\theta) \equiv p(x = 1 | \theta, a_j, b_j) = 1/\{1 + exp[-1.7a_j(\theta - b_j)]\}$.

12

The contribution of item j to information about θ is $2.89a_j^2P_j(\theta)[1 - P_j(\theta)]$, and the total information from n identical items for which $b_j \equiv \theta$ and $a_j \equiv a$ is simply 0.7225 na². Table 1 gives values of i and σ_x^2 in this simple case for selected test lengths and values of a. Note that where 1.7a = 1.0 (i.e., a = .588, corresponding to an item trait correlation of .7071 in a standard normal population), four additional items provide a unit gain in precision. The results provide an indication of the amount of information about θ that is employed in JML estimation of item parameters. It is apparent that as test length increases, information (i.e., precision) increases at a constant rate and the posterior variance decreases at a decreasing rate.

Insert Table 1 about here

The magnitude of gain in information about θ obtained by assuming an undifferentiated population (i.e., $F_i \equiv F$) can be gauged by extending the approximation employed for Table 1. If the normalized likelihood function induced by x_i is again approximated as $N(\hat{\theta}, \sigma_x^2)$ and if it is further assumed that examinee i has been selected at random from a population in which $\theta \sim N(\mu, \sigma^2)$, then

 $p(\theta | \mathbf{x}_1) \stackrel{*}{\sim} N(\tilde{\theta}, \tilde{\Sigma})$

where

$$\tilde{\theta} = \frac{\hat{\theta}\sigma_{\mathbf{x}}^{-2} + \mu\sigma^{-2}}{\sigma_{\mathbf{x}}^{-2} + \sigma^{-2}}$$

and

$$\tilde{\Sigma} = (\sigma^{-2} + \sigma_x^{-2})^{-1}$$

Table 2 shows values of the reciprocal of Σ (i.e., "precision") from various test lengths with identical items with 1.7a = 1 and a standard normal prior for θ . Note that for each test length, a unit gain in precision is achieved over the 1.7a = 1 column of Table 1. These tabled values fall within the ranges encountered in applied work, and suggest that the assumed distribution contributes about as much information about θ as four additional items. The corresponding value for 1.7a = .5 is sixteen items, and that for 1.7a = 1.5 is about one item. Since the <u>absolute</u> contribution is constant with respect to increasing test length, the <u>relative</u> contribution declines.

To gauge the additional impact of differentiating the population through auxiliary variables, we may consider numerical

14

values resulting from a regression model with homoscedastic residuals. Suppose y values account for $(100 \times z)$ -percent of the variance in a population with total variance 1.0, so that $F_y(\theta) \sim N(\mu_y, \sigma_e^2)$ with $\sigma_e^2 = 1 - r$. If the normalized likelihood induced by item responses is approximately $N(\hat{\theta}, \sigma_y^2)$, then

$$p(\theta | \mathbf{x}_{i}, \mathbf{y}_{i}) \stackrel{\circ}{\sim} \mathbb{N} \left[\frac{\hat{\theta} \sigma_{\mathbf{x}}^{-2} + \mu_{\mathbf{y}} \sigma_{\mathbf{e}}^{-2}}{\sigma_{\mathbf{x}}^{-2} + \sigma_{\mathbf{e}}^{-2}}, (\sigma_{\mathbf{e}}^{-2} + \sigma_{\mathbf{x}}^{-2})^{-1} \right]$$

Using the same simplified item response model and 'a' value as Table 2, Table 3 compares values of the inverse of the posterior variance for θ as determined by (i) item responses alone, (ii) with knowledge of membership in an undifferentiated population with unit variance, and (iii) with the additional knowledge of auxiliary variables that account for successively greater proportions of total variance. Values between 10- and 40-percent, a range typical of educational and psychological work, increase information (posterior precision) about θ by amounts roughly equivalent to one to three additional item responses. For items with 1.7a = .5, gains in item units would be doubled; for items with 1.7a = 1.5, gains in item units would be halved.

Insert Tables 2-3 about here

15

The Ignorability of p(y)

This section demonstrates that under reasonable assumptions, the population distribution of y can be ignored for the purposes of estimating item parameters β and population parameter α . Suppose that the distribution of y in a population of examinees is governed by the density function $p(y|\gamma)$, which depends on possibly unknown parameters γ but not upon item parameters β nor on the parameters α of the conditional distributions $f(\theta|y,\alpha)$. The probability of observing the data matrix (X,Y) from a random sample of N examinees is given by

$$P(X,Y|\beta,\alpha,\gamma)$$

 $= \prod_{i} f p(\mathbf{x}_{i} | \theta, \mathbf{y}_{i}, \beta, \alpha, \gamma) p(\theta | \mathbf{y}_{i}, \beta, \alpha, \gamma) p(\mathbf{y}_{i} | \beta, \alpha, \gamma) d\theta$ $= \prod_{i} f p(\mathbf{x}_{i} | \theta, \beta) p(\theta | \mathbf{y}_{i}, \alpha) p(\mathbf{y}_{i} | \gamma) d\theta$ $= \{ \prod_{i} f p(\mathbf{x}_{i} | \theta, \beta) p(\theta | \mathbf{y}_{i}, \alpha) d\theta \} \times \{ \prod_{i} p(\mathbf{y}_{i} | \gamma) \}$ $= P(\mathbf{X} | \mathbf{Y}, \beta, \alpha) P(\mathbf{Y} | \gamma) .$ (6)

16

Likelihood inferences about α and β are therefore independent of inferences about γ , and the conditional MLE's of α and β given Y are identical to MLE's obtained jointly with γ .

Models and Methods

This section presents two IRT models that differentiate examinees by means of auxiliary variables, and suggests computing approximations based on Bock and Aitkin's (1981) marginal maximum likelihood (empirical Bayes) procedures.

Mixtures of Finite Distributions

Mislevy (1984) decribes a nonparametric approximation of a continuous density function of a latent variable in terms of a distribution with mass at a finite number of prespecified points. The proficiency of each examinee, or θ_i , then, is assumed to take one of only Q known values. The "latent trait" problem is thereby replaced by an analogous "latent class" problem that is easier to solve. A single population was addressed in that presentation, and item parameters were assumed known. We now consider extensions to the simultaneous estimation of item parameters, and to multiple subpopulations indexed by an auxiliary variable y. This approach provides considerably flexibility in the distributions $F_i(\theta) = F_{y_i}(\theta)$. It lends itself well to discrete auxiliary variables with relatively few values.

17

It proves convenient to write such an auxiliary variable as a vector of 0/1 indicators. Define $y_i = (y_{i1}, \dots, y_{iK})$ by letting $y_{ik} = 1$ if examinees i is associated with the k'th of K exhaustive and mutually exclusive subpopulations, and zero otherwise. The probability of observing response pattern x_i from an examinee selected at random from a specified subpopulation is given by

$$p(x_i|y_i,\beta) = \prod_{k} \{ f p(x_i|\theta,\beta) dF_k(\theta) \}^{y_{ik}}, \qquad (7)$$

where F_k is the distribution in subpopulation k. This probability can be approximated by a finite distribution as

$$p(\mathbf{x}_{i}|\mathbf{y}_{i},\boldsymbol{\beta}) \approx \Pi \{ \Sigma p(\mathbf{x}_{i}|\boldsymbol{\Theta}_{q},\boldsymbol{\beta}) W_{qk} \}^{\mathbf{y}_{ik}}$$
(8)

where $\Theta_1, \ldots, \Theta_Q$ is a grid of points and W_{qk} is the weight or density at point q in subpopulation k. The weights W play the role of α in earlier notation. For the remainder of this subsection, we limit our attention to distributions of the form of the right-hand side of Equation 8. As demonstrated above, we may carry out the estimation of β and W conditional on Y.

18

Let (X,Y) be the data matrix observed from a sample of N examinees selected either randomly from the population as a whole or as random subsamples stratified on y. The probability of X given Y is proportional to

$$L_{M} = \prod_{i \ k} \prod_{q \ i \ k} \left\{ \sum_{q \ i \ k} p(\mathbf{x}_{i} | \Theta_{q}, \beta) W_{qk} \right\}^{\mathbf{y}_{ik}}$$

and its logarithm is

 $\ell_{M} = \log L_{M}$

$$= \sum_{i \in K} \sum_{j \in K} \frac{\log \sum_{i \in Q} p(\mathbf{x}_i | \Theta_{q,\beta})}{q} W_{qk}$$

Relative maxima with respect to β and W can be obtained by means of the EM algorithm, under the special case of missing indicators for a multinomial distribution (Dempster et al., 1977, Section 4.3). The expectation step of cycle t + 1 computes expected values of the following quantities:

- 19
- 1. The expected number of examinees with proficiency Θ_q from a sample of size N_k from subpopulation k, conditional on X, Y, $\hat{\beta}^t$, and \hat{W}^t :

$$\hat{\mathbf{N}}_{qk}^{t+1} = \sum_{i} y_{ik} \hat{\mathbf{p}}_{k}^{t} (\Theta_{q} | \mathbf{x}_{i})$$

where

$$\hat{p}_{k}^{t}(\Theta_{q}|\mathbf{x}_{i}) = p(\mathbf{x}_{i}|\Theta_{q},\beta = \hat{\beta}^{t})\hat{W}_{qk}^{t} / \sum_{r} p(\mathbf{x}_{i}|\Theta_{r},\beta = \hat{\beta}^{t})\hat{W}_{rk}^{t}$$

- an application of Bayes theorem, gives the posterior probability that the proficiency of examinee i is Θ_q , given provisional parameter estimates $\hat{\beta}^t$ and \hat{W}^t .
- 2. The expected number of correct responses to item j from examinees in subpopulation k with proficiency Θ_q , given a random sample of size N_k (again given $\hat{\beta}^t$ and \hat{W}^t):

$$\hat{R}_{jqk}^{t+1} = \sum_{i} y_{ik} x_{ij} \hat{p}_{k}^{t}(\Theta_{q} | x_{i})$$

20

The maximization step computes what would be MLE's of β and \hat{R} were observed quantities rather than conditional expectations. For W, we have simply

$$\hat{W}_{qk}^{t+1} = \hat{N}_{qk}^{t+1} / N_{k}$$

For $\boldsymbol{\beta}$, we solve conditional expectations of likelihood equations:

$$0 = \Sigma \qquad \frac{R_{jq+}^{t+1}R_{q+}^{t+1}P_{j}(\Theta_{q})}{P_{j}(\Theta_{q})[1 - P_{j}(\Theta_{q})]} \xrightarrow{\partial P_{j}(\Theta_{q})}{\partial \beta}, \qquad (9)$$

where $\hat{R}_{jq+}^{t+1} = \sum_{k} \hat{R}_{jqk}^{t+1}$ and \hat{N}_{q+}^{t+1} is similarly defined. Under the 2parameter logistic model, for example, Equation 9 simplifies as follows:

$$a_{j}: \qquad 0 = \sum_{q} \left[\hat{R}_{jq}^{t} - \hat{N}_{q+}^{t+1} P_{j}(\Theta_{q}) \right] (\Theta_{q} - b_{j})$$

$$b_{j}: \qquad 0 = \sum \left[\hat{R}_{jq+}^{t+1} - \hat{N}_{q+}^{t+1} P_{j}(\Theta_{q}) \right] a_{j}$$

In principle, the linear indeterminacy in the 1-, 2-, and 3- parameter logistic and normal IRT models presents no impediment to

21

the EM algorithm, which readily converges to one of the infinitely many solutions on a ridge. Numerical stability and the quality of the finite characterization of F are enhanced, however, by controlling the scaling of the solution at this point. One convenient way of doing so is to standardize the weighted average distribution. We have referred to the points 0_q as specified a priori; given the linear indeterminacy, we may conceive of only their relative spacing as prespecified. After each EM cycle, then, we may rescale the points as follows:

 $O_q = (O_q - \overline{O})/s$

where

$$\vec{D} = N^{-1} \sum_{k} N_{k} \sum_{q} O_{qk}^{kt}$$

and

$$s = N^{-1} \sum_{k} N_{k} \sum_{q} (O_{q} - \overline{O})^{2} \hat{w}_{qk}^{t}$$

Item parameters are adjusted accordingly. Under the 2- and 3-

parameter models, \hat{b}_j is replaced by $(\hat{b}_j - \overline{\Theta})/s$ and a_j is replaced by sa_j . Under 1-parameter models, rescaling takes place only with respect to $\overline{\Theta}$.

Iteration from several starting values helps to verify whether a given solution is indeed a global maximum. The observed information matrix for the item parameter estimates can then be approximated via Equation 6. Employing Louis's (1982) simplifications for "missing multinomial indicators" problems, we obtain

$$I_{X,Y}(\hat{\beta}) \approx \sum_{i k} \sum_{k} y_{ik} \sum_{q} \left(\frac{\partial \ell_{i}(\Theta_{q})}{\partial \beta} \right) \left(\frac{\partial \ell_{i}(\Theta_{q})}{\partial \beta} \right) \hat{p}_{k}(\Theta_{q}|x_{i}) ,$$
(10)

where $\hat{p}_k(\Theta_q | x_i)$ is evaluated at $\hat{\beta}$ and \hat{W} . A Linear Model

The unrestricted mixture solution described above becomes unwieldly as the number of potential values of the auxiliary variable increases. The more structured alternative of a linear model for $p(\theta | y)$ is suitable when y is vector-valued or is continuo rather than discrete. Assuming homoscedastic and normal residuals, we would have

$$\theta \sim N(y'\alpha,\sigma^2)$$

where auxiliary variables are coded so that the K columns of $Y = (y_1 : \cdots : y_K)'$, which are basis vectors for the K elements of α , are linearly independent. They may include values on measured variables such as previous test scores and dummy regression variables that encode selected contrasts among categorical auxiliary variables.

Maximum likelihood solutions for a and σ^2 in the special case of structured means for the cells of a multi-way design have been given by Mislevy (1985) under the assumption that item parameters are known, and by Zwarts and Veldhuesen (1985) under the assumption that $p(\mathbf{x}|\theta)$ is the Rasch model with unknown item parameters to be estimated jointly. These solutions are readily extended to the case of a general IRT model with unknown item parameters. This section describes an approximation over a grid of prespecified points so that computation is similar to the nonparametric solution described above. Attention is focused for convenience upon the 1-, 2-, and 3-parameter logistic and normal IRT models.

The linear indeterminacies of these models are again conveniently resolved by restrictions on the population parameters. First, we may without loss of generality fix σ^2 at unity to set the

24

unit-size of the scale. For 1-parameter models, a slope parameter common over items is then estimated. Second, we may set the origin by centering the elements of each column of Y at zero. All effects are thus cast as deviations around a grand mean of zero. This restriction, in conjunction with the independence of the basis vectors, completes the resolution of the scale.

The marginal likelihood for a sample of size N is written as

$$L = \prod \int p(\mathbf{x}_{1} | \boldsymbol{\theta}, \boldsymbol{\beta}) \phi(\boldsymbol{\theta} - \mathbf{y}_{1}^{\prime} \boldsymbol{\alpha}) d\boldsymbol{\theta}$$

where ϕ represents the standard normal density function. Approximation over a finite grid of points is accomplished by

 $L^{\star} = \prod_{i \neq q} \sum_{\alpha \neq i} p(x_i | 0_q, \beta) W_{qi}(\alpha)$

where

$$W_{qi}(\alpha) = \exp[-(\Theta_{q} - \frac{y_{i\alpha}}{z_{i\alpha}})^{2}/2] / \sum_{r} \exp[-(\Theta_{r} - \frac{y_{i\alpha}}{z_{i\alpha}})^{2}/2]$$

The weights W play the same role as those in the preceeding approximation. The difference is that they are no longer estimated

25

without restriction, but modeled as functions of the effect parameters α .

MML estimation can again proceed in EM cycles that solve the likelihood equations. Let $\hat{\beta}^{t}$ and $\hat{\alpha}^{t}$ be provisional estimates from cycle t. The E-step computes expected counts of examinees and correct responses at each point:

$$\hat{N}_{q}^{t+1} = \sum_{i} P(O_{q} | \mathbf{x}_{i}, \hat{\beta}^{t}, \hat{\alpha}^{t})$$

and

$$\hat{R}_{jq}^{t+1} = \sum_{i} x_{ij} P(\Theta_{q} | x_{i}, \hat{\beta}^{t}, \hat{\alpha}^{t})$$

where

$$P(\Theta_{q}|x_{i},\hat{\beta}^{t},\hat{\alpha}^{t}) = p(x_{i}|\Theta_{q},\hat{\beta}^{t})W_{qi}(\hat{\alpha}^{t})/\sum_{r} p(x_{i}|\Theta_{r},\hat{\beta}^{t})W_{ri}(\hat{\alpha}^{t}) .$$

It also computes the conditional expected value of each examinee's proficiency:

$$\hat{\theta}_{i}^{t+1} = \sum_{q} \Theta_{q} P(\Theta_{q} | \mathbf{x}_{i}, \hat{\beta}^{t}, \hat{\alpha}^{t})$$

26

The M-step pseudo-likelihood equations for item parameters can be written as in Equation 9. The equations for α simplify to

 $\hat{\alpha}^{t+1} = (Y'Y)^{-1}Y'\hat{\theta}^{t+1}$

where $\hat{\theta}_{1}^{t+1} = (\hat{\theta}_{1}^{t+1}, \dots, \hat{\theta}_{N}^{t+1})$. The posterior information matrix for $\hat{\beta}$ can again be approximated via Equation 10.

A Numerical Example

This section illustrates the procedures described above. The data are responses to four items from the Arithmetic Reasoning test of the Armed Services Vocational Aptitude Battery (ASVAB), Form 8A, as observed in a sample of 776 participants in the <u>Profile of American</u> <u>Youth</u> survey (U.S. Department of Defense, 1982). Table 4 gives counts of the sixteen possible response patterns occurring in each cell of a 2-by-2 design based on two background variables collected along with item responses. Because these variables are based on demographic information rather than the educationally-relevant information we would prefer, we will refer to the factors as simply Factor A and Factor B, nesting levels 1 and 2 within each.

Insert Table 4 about here

27

Four analyses were carried out on these data. In each, the 2-parameter logistic ogive was employed as the IRT model for conditional probabilities of correct response. The analyses differed in terms of the auxiliary information about examinees they employed. The first run used MML estimation of item parameters and densities over a grid of ten points, assuming examinees were drawn at random from a single undifferentiated population. The second and third runs differentiated the population via Factor A and Factor B respectively, and the fourth run employed both factors jointly.

Resulting item parameter estimates and standard errors, along with subpopulation means and standard deviations, are shown in Tables 5 through 8. The scale has been set in all solutions to standardize the total population. For each item parameter type, columns in Table 6 through 8 display the ratio of the squared standard error of the item parameter estimate under the undifferentiated model to the corresponding value in the differentiated model. The result can be interpreted as efficiency relative to the undifferentiated model, and the excess of a value above unity reflects the proportional increase in estimation precision. Geometric averages are also shown for the relative efficiency columns. The excess of such a value over unity, times

28

four, gives the increases of precision in the units of numbers of additional items of the same kind.

Insert Tables 5-8 about here

It is apparent that including auxiliary information had little effect on the <u>values</u> of the item parameter estimates. The differences between the estimates from the undifferentiated and the fully differentiated solutions occur only in the second decimal place. More significant differences exist in the accompanying (estimated) standard errors, however. The precision of threshold estimates was improved only modestly; an increase roughly equivalent to one additional item response per examinee was observed in the fully differentiated run. The precision of slope estimates was improved dramatically; an increase roughly equivalent to eight items was observed. It would appear that Factor A accounted for more increase in precision for slopes, while Factor B accounted for more increase in precision for thresholds.

Discussion

This paper has outlined procedures for incorporating auxiliary information about examinees into the IRT framework. Enhancing the precision of item parameter estimates was the primary focus. This section evaluates the value of improvements so attained, and discusses two additional aspects of the model.

29

The increase in information about item parameters in typical educational and psychological settings can be expected to lie in the range of two to six items. The numerical example suggests that the increase will vary by item parameter type, probably less for well-estimated parameters and greater for poorly-estimated parameters.

The expected increase is modest, to be sure, but in many applications it is free in the sense that it is already available for use. Because its incremental value decreases for longer tests, auxiliary information would be most useful in settings where relatively few response are solicited from each examinee. This would include two applications of great current interest, namely educational assessment and adaptive testing. In assessment, data that are sparse at the level of individuals--say, five items in a given scale--yield more efficient estimates of population parameters for a given total number of item responses. In adaptive testing, new items are calibrated using joint response patterns with previously-calibrated items while the number of old items is held to minimally acceptable levels--as few as, say, fifteen.

A side issue in the present paper but a fundamentally important result is that when examinees are indeed a random sample from a well-defined population, the estimated population

30

distributions and effect parameters are consistent within the limits of precision afforded by the numerical approximations (see Mislevy, 1984, 1985, on population estimation when item parameters are known). This stands in contrast to the asymptotically biased results obtained by using the distribution of $\hat{\theta}$ to approximate the distribution of θ . In fact, the discrepancy between the two distributions is largest in exactly those cases in which the present procedures offer most the benefit for item parameter estimation, namely short tests.

Finally, it is implicit in preceding discussions that auxiliary information about examinees can lead to improved estimates of individual proficiencies. Whether estimates that are improved in the sense of minimum mean squared error are unequivocally "better" for all applications is not clear, however. We have avoided advocating the use of auxiliary information when tests are used as contests--i.e., when important placement or selection decisions are made for individual examinees--because it would seem that in these situations the tester ought to gather enough data directly dependent upon proficiency (i.e., item reponses) to make satisfactorily precise decisions on that strength alone. In adaptive testing, for example, we would recommend the use of auxiliary information to improve item parameter estimation, but not to estimate scores that will be used to compare individual examinees.

References

Birnbaum, A. (1968). Some latent trait models and their use in inferring an examinee's ability. In F M. Lord and M. R. Novick, <u>Statistical theories of mental test scores</u> (pp. 397– 479). Reading, MA: Addison-Wesley.

Bock, R. D., & Aitkin, M. (1981). Marginal maximum likelihood estimation of item parameters: An application of an EM algorithm. Psychometrika, 46, 443-459.

Bock, R. D., & Lieberman, M. (1970). Fitting a response model for <u>n</u> dichotomously scores items. <u>Psychometrika</u>, <u>35</u>, 179-197.

de Leeuw, J., & Verhelst, N. (1984). Maximum likelihood estimation
in generalized Rasch models (Department of Data Theory
Research Report RR-84-11). Leiden, The Netherlands:
University of Leiden.

- Dempster, A. P., Laird, N. M., & Rubin, D. B. (1977). Maximum likelihood from complete data via the EM algorithm (with discussion. <u>Journal of the Royal Statistical Society</u>, <u>Series</u> <u>B</u>, <u>39</u>, 1-38.
- Leonard, T., & Novick, M. R. (1985). <u>Bayesian inference and</u> <u>diagnostics for the three-parameter logistic model</u> (ONR Technical Report 85-5). Iowa City, IA: CADA Research Group, University of Iowa.

32

Lindley, D. V., & Smith A. F. M. (1972). Bayes estimates for the linear model. <u>Journal of the Royal Statistical Society</u>, Series B, 34, 1-41.

Louis, T. (1982). Finding the observed information matrix when using the EM algorithm. <u>Journal of the Royal Statistical</u> Society, Series B, 44, 226-233.

Mislevy, R. J. (1984). Estimating latent distributions.

Psychometrika, 49, 359-381.

Mislevy, R. J. (1985). Estimate of latent group effects.

Journal of the American Statistical Association, 80, 993-997. Rigdon, S. E., & Tsutakawa, R. K. (1982). Parameter estimation in

latent trait models. <u>Psychometrika</u>, <u>48</u>, 567-574.

- Swaminathan, H., & Gifford, J. A. (1982). Bayesian estimation in the Rasch model. Journal of Educational Statistics, 9, 23-80.
- Tjur, T. (1982). A connection between Rasch's item analysis model and a multiplicative Poisson model. <u>Scandinavian Journal of</u> Statistics, 9, 23-30.
- U.S. Department of Defense (1982). <u>Profile of American youth</u> Washington, DC: Office of the Assistant Secretary of Defense (Manpower, Reserve Affairs, and Logistics).

Zacks, S. (1971). <u>The theory of statistical inference</u>. New York: Wiley.
33

Zwarts, M., & Veldhuesen, N. (1985). <u>Maximum marginal likelihood</u> <u>estimation in latent trait models</u>. Paper presented at the European meeting of the Psychometric Society, Cambridge, England.

34

Acknowledgment

This work was supported by contract no. N00014-85-K-0683, project designation NR 150-539, from Personnel and Training Research Programs, Psychological Sciences Division, Office of Naval Research. The author is grateful to Kathleen Sheehan and Martha Stocking for their comments and suggestions.

35

Table l

Posterior Precision for θ from Item Responses Only

	1.7a = .500		1.7a = 1.000		1.7a = 1.500	
<u>n</u>	i	σ ² _x	i	$\sigma_{\mathbf{x}}^{2}$	i	σ ² _x
2	.125	8.000	• 500	2.000	1.125	.889
4	.250	4.000	1.000	1.000	2.250	.444
8	•500	2.000	2.000	• 500	4.500	.222
16	1.000	1.000	4.000	•250	9.000	.111
32	2.000	• 500	8.000	.125	18.000	•056
64	4.000	•250	16.000	•063	36.000	•028
128	8.000	.125	32.000	.031	72.000	•014

n = number of identical items with a as noted and $b = \theta$.

i = information = posterior precision.

36

Table 2

Posterior Precision for $\boldsymbol{\theta}$ from Item Responses and Population Membership

			
<u>_n</u>	$i(\equiv (\Sigma^{-1}))$	Relative 2 Efficiency (σ_x^2/Σ)	Effective Gain
2	1.500	3.000	200.0%
4	2.000	2.000	100.0%
8	3.000	1.500	50.0%
16	5.000	1.250	25.0%
32	9.000	1.125	12.5%
64	17.000	1.063	6.3%
128	33.000	1.031	3.1%

1.7a = 1.000

n = number of identical items with a as noted and b = θ .

 $i = information \equiv posterior precision.$

37

Table 3

Precision Increases for $\boldsymbol{\theta}$ Resulting from the Use of

Auxiliary Information

Source	Increment in Posterior Precison	Precision Gain in Item Units	Gain over Undifferentiated Population
One-item response	•250	1.000	
Population membership	1.000	4.000	
Auxiliary information			
$R^2 = .10$	1.111	4.444	11.1%
$R^2 = .20$	1.250	5.000	25.0%
$R^2 = .30$	1.429	5.716	42.9%
$R^2 = .40$	1.667	6.668	66.7%
$R^2 = .50$	2.000	8.000	100.0%
$R^2 = .60$	2.500	10.000	150.0%
$R^2 = .70$	3.333	13.332	233.3%
$R^2 = .80$	5.000	20.000	400.0%
$R^2 = .90$	10.000	40.000	900.0%

Table 4

Item Response A1B2 A2B1 A2B2 1 2 3 4 A1B1 0 0 0 0 0 1 0 1 1 1 1 Total

Counts of Observed Response Patterns

39

Table 5

Item Parameter Estimates: Undifferentiated Population

Item	ĵ.	SE(b)	â	SE(a)		
1	422	•058	1.022	•171		
2	226	.072	•666	.094		
3	.152	.076	•705	.096		
4	.397	•080	•839	•114		
Population Mean: 0.000						
Populat	ion Standard	Deviation:	1.000			

Ě

40

Table 6

Item Parameter Estimates: Population Differentiated

Item	р b	SE(b)	Relative Efficiency	z a	SE(a)	Relative Efficiency	
1	436	•062	.875	.869	•069	6.142	
2	217	•077	.874	•622	•054	3.030	
3	•189	•072	1.114	•676	•056	2.939	
4	•465	•069	1.344	•775	•061	3.493	
Geometric average relative efficiency: 1.035						3.718	
Subpopulation means: .296,511							
Subpopulation standard deviations: .960, .850							

with Respect to Factor A Only

41

Table 7

Karaharan Salaharan S

Item Parameter Estimates: Population Differentiated

Item	 b	SE(b)	Relative Efficiency	â	SE(a)	Relative Efficiency
1	408	•057	1.035	•941	.073	5.487
2	211	•077	.874	•621	•056	2.818
3	.185	.071	1.146	•686	.058	2.740
4	•431	•064	1.563	•842	.067	2.895
	ric averag tive effic		1.128		<u> </u>	3.328
Subpop	ulation me	ans:		.136,147		
Subpop	Subpopulation standard deviations: 1.021, .955					

with Respect to Factor B Only

42

Table 8

Item Parameter Estimates: Population Differentiated

Item	Ъ	SE(b)	Relative Efficiency	â	SE(â)	Relative Efficiency
1	421	.052	1.244	1.006	•080	4.569
2	213	.071	1.028	•672	.059	2.538
3	.139	•065	1.367	•775	.063	2.311
4	•402	•066	1.469	•834	•066	2.983
Geometric average relative efficiency: 1.266						2.994
Subpopulation means:				.485, .073, -	.513,	502
Subpopulation standard deviations:				1.164, .855,	.642, .6	40

with Respect to Factors A and B

Educational Testing Service/Mislevy

Personnel Analysis Division, AF/MPXA 50360, The Pentagon Washington, DC 20330

Air Force Human Resources Lab AFHRL/MPD Brooks AFB, TX 78235

Dr. Earl A. Alluisi HQ, AFHRL (AFSC) Brooks AFB, TX 78235

Dr. Erling B. Andersen Department of Statistics Studiestraede 6 1455 Copenhagen DENMARK

Dr. Phipps Arabie University of Illinois Department of Psychology 603 E. Daniel St. Champaign, IL 61820

Technical Director, ARI 5001 Eisenhower Avenue Alexandria, VA 22333

Dr. Eva L. Baker UCLA Center for the Study of Evaluation 145 Moore Hall University of California Los Angeles, CA 90024

Dr. Isaac Bejar Educational Testing Service Princeton, NJ 08450

Dr. Menucha Birenbaum School of Education Tel Aviv University Tel Aviv, Ramat Aviv 69978 ISRAEL

Dr. Arthur S. Blaiwes Code N711 Naval Training Equipment Center Orlando, FL 32813 Dr. Bruce Bloxom Administrative Sciences Code 54B1 Navy Postgraduate School Monterey, CA 93943

Dr. R. Darrell Bock University of Chicago Department of Education Chicago, IL 60637

Cdt. Arnold Bohrer Sectie Psychologisch Onderzoek Rekruterings-En Selectiecentrum Kwartier Koningen Astrid Bruijnstraat 1120 Brussels, BELGIUM

Dr. Robert Breaux Code N-095R NAVTRAEQUIPCEN Orlando, FL 32813

Dr. Robert Brennan American College Testing Programs P. O. Box 168 Iowa City, IA 52243

Dr. Patricia A. Butler NIE Mail Stop 1806 1200 19th St., NW Washington, DC 20208

Mr. James W. Carey Commandant (G-PTE) U.S. Coast Guard 2100 Second Street, S.W. Washington, DC 20593

Dr. James Carlson American College Testing Program P.O. Box 168 Iowa City, IA 52243

Dr. John B. Carroll 409 Elliott Rd. Chapel Hill, NC 27514

Dr. Robert Carroll NAVOP 01B7 Washington, DC 20370

Educational Testing Service/Mislevy

Dr. Norman Cliff Department of Psychology Univ. of So. California University Park Los Angeles, CA 90007

Director, Manpower Support and Readiness Program Center for Navel Analysis 2000 North Beauregard Street Alexandria, VA 22311

Dr. Stanley Collyer Office of Naval Technology Code 222 800 N. Quincy Street Arlington, VA 22217-5000

Dr. Hans Crombag University of Leyden Education Research Center Boerhaavelaan 2 2354 EN Leyden The NETHERLANDS

CTB/McGraw-Hill Library 2500 Garden Road Monterey, CA 93940

Dr. Dattprasad Divgi Center for Naval Analysis 4401 Ford Avenue P.C. Box 16268 Alexandria, VA 22302-0268

Dr. Hei-Ki Dong Ball Foundation 800 Roosevelt Road Building C, Suite 206 Glen Ellyn, IL 60137

Defense Technical Information Center Cameron Station, Bldg 5 Alexandria, VA 22314 Attn: TC (12 Copies) Dr. Stephen Dunbar Lindquist Center for Measurement University of Iowa Iowa City, IA 52242

Dr. James A. Earles Air Force Human Resources Lab Brooks AFB, TX 78235

Dr. Kent Eaton Army Research Institute 5001 Eisenhower Avenue Alexandria, VA 22333

Dr. John M. Eddins University of Illinois 252 Engineering Research Laboratory 103 South Mathews Street Urbana, IL 61801

Dr. Susan Embretson University of Kansas Psychology Department Lawrence, KS 66045

ERIC Facility-Acquisitions 4833 Rugby Avenue Bethesda, MD 20014

Dr. Benjamin A. Fairbank Performance Metrics, Inc. 5825 Callaghan Suite 225 San Antonio, TX 78228

Dr. Leonard Feldt Lindquist Center for Measurement University of Iowa Iowa City, IA 52242

Dr. Richard L. Ferguson American College Testing Program P.O. Box 168 Iowa City, IA 52240

Dr. Gerhard Fischer Liebiggasse 5/3 A 1010 Vienna AUSTRIA

Educational Testing Service/Mislevy

Prof. Donald Fitzgerald University of New England Department of Psychology Armidale, New South Wales 2351 AUSTRALIA

Mr. Paul Foley Navy Personnel R&D Center San Diego, CA 92152

Dr. Carl H. Frederiksen McGill University 3700 McTavish Street Montreal, Quebec H3A 1Y2 CANADA

Dr. Robert D. Gibbons University of Illinois-Chicago P.O. Box 6998 Chicago, IL 69680

Dr. Janice Gifford University of Massachusetts School of Education Amherst, MA 01003

Dr. Robert Glaser Learning Research & Development Center University of Pittsburgh 3929 O'Hara Street Pittsburgh, PA 15260

Dr. Bert Green Johns Hopkins University Department of Psychology Charles & 34th Street Baltimore, MD 21218

Dr. Ronald K. Hambleton Prof. of Education & Psychology University of Massachusetts at Amherst Hills House Amherst, MA 01003

Ms. Rebecca Hetter Navy Personnel R&D Center Code 62 San Diego, CA 92152 Dr. Paul W. Holland Educational Testing Service Rosedale Road Princeton, NJ 08541

Prof. Lutz F. Hornke Universitat Dusseldorf Erziehungswissenschaftliches Universitatsstr. 1 Dusseldorf 1 WEST GERMANY

Dr. Paul Horst 677 G Street, #184 Chula Vista, CA 90010

Mr. Dick Hoshaw NAVOP-135 Arlington Annex Room 2834 Washington, DC 20350

Dr. Lloyd Humphreys University of Illinois Department of Psychology 603 East Daniel Street Champaign, IL 61820

Dr. Steven Hunka Department of Education University of Alberta Edmonton, Alberta CANADA

Dr. Huynh Huynh College of Education Univ. of South Carolina Columbia, SC 29208

Dr. Robert Jannarone Department of Psychology University of South Carolina Columbia, SC 29208

Dr. Douglas H. Jones Advanced Statistical Technologies Corporation 10 Trafalgar Court Lawrenceville, NJ 08148

Educational Testing Service/Mislevy

Dr. G. Gage Kingsbury Portland Public Schools Research and Evaluation Department 501 North Dixon Street P. O. Box 3107 Portland, OR 97209-3107

Dr. William Koch University of Texas-Austin Measurement and Evaluation Center Austin, TX 78703

Dr. Leonard Kroeker Navy Personnel R&D Center San Diego, CA 92152

Dr. Michael Levine Educational Psychology 210 Education Bldg. University of Illinois Champaign, IL 61801

Dr. Charles Lewis Faculteit Sociale Wetenschappen Rijksuniversiteit Groningen Oude Boteringestraat 23 9712GC Groningen The NETHERLANDS

Dr. Robert Linn College of Education University of Illinois Urbana, IL 61801

Dr. Robert Lockman Center for Naval Analysis 4401 Ford Avenue P.O. Box 16268 Alexandria, VA 22302-0268

Dr. Frederic M. Lord Educational Testing Service Princeton, NJ 08541

Dr. James Lumsden Department of Psychology University of Western Australia Nedlands W.A. 6009 AUSTRALIA Dr. William L. Maloy Chief of Naval Education and Training Naval Air Station Pensacola, FL 32508 Dr. Gary Marco Stop 31-E Educational Testing Service Princeton, NJ 08451 Dr. Clessen Martin Army Research Institute 5001 Eisenhower Blvd. Alexandria, VA 22333 Dr. James McBride Psychological Corporation c/o Harcourt, Brace, Javanovich Inc. 1250 West 6th Street San Diego, CA 92101 Dr. Clarence McCormick HQ, MEPCOM MEPCT-P 2500 Green Bay Road North Chicago, IL 60064 Mr. Robert McKinley University of Toledo Department of Educational Psychology Toledo, OH 43606 Dr. Barbara Means Human Resources Research Organization 1100 South Washington Alexandria, VA 22314 Dr. Robert Mislevy Educational Testing Service Princeton, NJ 08541 Headquarters, Marine Corps Code MPI-20 Washington, DC 20380 Dr. W. Alan Nicewander

University of Oklahoma Department of Psychology Oklahoma City, OK 73069

Educational Testing Service/Mislevy

Dr. William E. Nordbrock FMC-ADCO Box 25 APO, NY 09710

Dr. Melvin R. Novick 356 Lindquist Center for Measurement University of Iowa Iowa City, IA 52242

Director, Manpower and Personnel Laboratory, NPRDC (Code 06) San Diego, CA 92152

Library, NPRDC Code P201L San Diego, CA 92152

Commanding Officer, Naval Research Laboratory Code 2627 Washington, DC 20390

Dr. James Olson WICAT, Inc. 1875 South State Street Orem, UT 84057

Office of Naval Research, Code 1142PT 800 N. Quincy Street Arlington, VA 22217-5000 (6 Copies)

Special Assistant for Marine Corps Matters, ONR Code 00MC 800 N. Quincy St. Arlington, VA 22217-5000

Dr. Judith Orasanu Army Research Institute 5001 Eisenhower Avenue Alexandria, VA 22333

Wayne M. Patience American Council on Education GED Testing Service, Suite 20 One Dupont Circle, NW Washington, DC 20036 Dr. James Paulson Department of Psychology Portland State University P.O. Box 751 Portland, OR 97207 Dr. Roger Pennell Air Force Human Resources Laboratory Lowry AFB, CO 80230 Dr. Mark D. Reckase ACT P. O. Box 168 Iowa City, IA 52243 Dr. Malcolm Ree AFHRL/MP Brooks AFB, TX 78235 Dr. Carl Ross CNET-PDCD Building 90 Great Lakes NTC, IL 60088 Dr. J. Ryan Department of Education University of South Carolina Columbia, SC 29208 Dr. Fumiko Samejima Department of Psychology University of Tennessee Knoxville, TN 37916 Mr. Drew Sands NPRDC Code 62 San Diego, CA 92152 Dr. Robert Sasmor HQDA DAMA-ARL Pentagon, Room 3E516 Washington, DC 20310-0631 USA Dr. Mary Schratz Navy Personnel R&D Center San Diego, CA 92152 Dr. W. Steve Sellman OASD(MRA&L) 2B269 The Pentagon

Washington, DC 20301

Educational Testing Service/Mislevy

Dr. Kazuo Shigemasu 7-9-24 Kugenuma-Kaigan Fujusawa 251 JAPAN

Dr. William Sims Center for Naval Analysis 4401 Ford Avenue P.O. Box 16268 Alexandria, VA 22302-0268

Dr. H. Wallace Sinaiko Manpower Research and Advisory Services Smithsonian Institution 801 North Pitt Street Alexandria, VA 22314

Dr. Richard Sorensen Navy Personnel R&D Center San Diego, CA 92152

Dr. Paul Speckman University of Missouri Department of Statistics Columbia, MO 65201

Dr. Martha Stocking Educational Testing Service Princeton, NJ 08541

Dr. Peter Stoloff Center for Naval Analysis 200 North Beauregard Street Alexandria, VA 22311

Dr. William Stout University of Illinois Department of Mathematics Urbana, IL 61801

Maj. Bill Strickland AF/MPXOA 4E168 Pentagon Washington, DC 20330

Dr. Hariharan Swaminathan Laboratory of Psychometric and Evaluation Research School of Education University of Massachusetts Amherst, MA 01003 Mr. Brad Sympson Navy Personnel R&D Center San Diego, CA 92152

Dr. Kikumi Tatsuoka CERL 252 Engineering Research Laboratory Urbana, IL 61801

Dr. Maurice Tatsuoka 220 Education Bldg 1310 S. Sixth St. Champaign, IL 61820

Dr. David Thissen Department of Psychology University of Kansas Lawrence, KS 66044

Mr. Gary Thomasson University of Illinois Educational Psychology Champaign, IL 61820

Dr. Robert Tsutakawa The Fred Hutchinson Cancer Research Center Division of Public Health Sci. 1124 Columbia Street Seattle, WA 98104

Dr. Ledyard Tucker University of Illinois Department of Psychology 603 E. Daniel Street Champaign, IL 61820

Dr. Vern W. Urry Personnel R&D Center Office of Personnel Management 1900 E. Street, NW Washington, DC 20415

Dr. David Vale Assessment Systems Corp. 2233 University Avenue Suite 310 St. Paul, MN 55114

Dr. Frank Vicino Navy Personnel R&D Center San Diego, CA 92152

Educational Testing Service/Mislevy

Dr. Howard Wainer Division of Psychological Studies Educational Testing Service Princeton, NJ 08541

Dr. Ming-Mei Wang Lindquist Center for Measurement University of Iowa Iowa City, IA 52242

 Dr. Thomas A. Warm Coast Guard Institute
P. O. Substation 18 Oklahoma City, OK 73169

Dr. Brian Waters Program Manager Manpower Analysis Program HumRRO 1100 S. Washington St. Alexandria, VA 22314

Dr. David J. Weiss N660 Elliott Hall University of Minnesota 75 E. River Road Minneapolis, MN 55455

Dr. Ronald A. Weitzman NPS, Code 54Wz Monterey, CA 92152

Major John Welsh AFHRL/MOAN Brooks AFB, TX 78223

Dr. Rand R. Wilcox University of Southern California Department of Psychology Los Angeles, CA 90007

German Military Representative ATTN: Wolfgang Wildegrube Streitkraefteamt D-5300 Bonn 2 4000 Brandywine Street, NW Washington, DC 20016 Dr. Bruce Williams Department of Educational Psychology University of Illinois Urbana, IL 61801

Dr. Hilda Wing Army Research Institute 5001 Eisenhower Ave. Alexandria, VA 22333

Dr. Martin F. Wiskoff Navy Personnel R & D Center San Diego, CA 92152

Mr. John H. Wolfe Navy Personnel R&D Center San Diego, CA 92152

Dr. George Wong Biostatistics Laboratory Memorial Sloan-Kettering Cancer Center 1275 York Avenue New York, NY 10021

Dr. Wendy Yen CTB/McGraw Hill Del Monte Research Park Monterey, CA 93940

