

AD-A170 720

MULTIREFERENCE CI GRADIENTS AND MCSCF SECOND
DERIVATIVES(U) ARMY BALLISTIC RESEARCH LAB ABERDEEN
PROVING GROUND MD B H LENGFIELD ET AL. MAY 86
BRL-TR-2730 F/A 20/


141

UNCLASSIFIED

F/G 20/10

NL

[illegible]



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A170 720

AD

12




TECHNICAL REPORT BRL-TR-2730

MULTIREFERENCE CI GRADIENTS AND MCSCF SECOND DERIVATIVES

Byron H. Lengsfeld, III
Paul J. Saxe
George F. Adams
Michael J. Page

May 1986

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.



US ARMY BALLISTIC RESEARCH LABORATORY
ABERDEEN PROVING GROUND, MARYLAND

DTIC FILE COPY

86 7 21 057

Destroy this report when it is no longer needed.
Do not return it to the originator.

Additional copies of this report may be obtained
from the National Technical Information Service,
U. S. Department of Commerce, Springfield, Virginia
22161.

The findings in this report are not to be construed as an official
Department of the Army position, unless so designated by other
authorized documents.

The use of trade names or manufacturers' names in this report
does not constitute indorsement of any commercial product.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report BRL-TR-2730	2. GOVT ACCESSION NO. AD-A17072	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) MULTIREFERENCE CI GRADIENTS AND MCSCF SECOND DERIVATIVES		5. TYPE OF REPORT & PERIOD COVERED Final
7. AUTHOR(s) Byron H. Lengsfeld, III, Paul J. Saxe,* George F. Adams, Michael J. Page**		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS US Army Ballistic Research Laboratory ATTN: SLCBR-IB Aberdeen Proving Ground, MD 21005-5066		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Ballistic Research Laboratory ATTN: SLCBR-DD-T Aberdeen Proving Ground, MD 21005-5066		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 1L161102AH43
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE May 1986
		13. NUMBER OF PAGES 28
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE NA
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release; Distribution Unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES *Los Alamos National Laboratory, Los Alamos, NM 87545 **Naval Research Laboratory, Washington, D.C. 20375		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) CI Gradients MCSCF Second Derivatives		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) meg A set of simple and efficient formulas for the calculation of multireference CI gradients and MCSCF second derivatives is presented. The CI gradient formalism is extended to include a general class of references in the CI. This extension is necessary for the calculation of gradients for a commonly employed class of CI wavefunctions for which the reference configurations are selected from a generalized CAS MCSCF wavefunction. In addition, we report the first general multireference CI gradient calculations. The		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

20. Abstract (Cont'd):

Y-10

calculations are for the reaction $\text{Be} + \text{H}_2^+ \rightarrow \text{BeH}_2$ constrained to C_{2v} symmetry. Structures of the reactant and transition state and the activation energy calculated at the selected reference CI level compare favorably to the full second order CI results. MCSCF second derivatives are found to be useful for the optimization of the CI structures.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

TABLE OF CONTENTS

	<u>Page</u>
I. INTRODUCTION.....	5
II. COMPUTATIONAL DETAILS.....	6
A. First Derivatives.....	6
B. Coupled Perturbed MCSCF Equations.....	9
C. MCSCF Second Derivatives.....	10
D. Selected Reference CI Gradients.....	12
III. COMPARISON WITH OTHER WORK.....	13
IV. SAMPLE CALCULATIONS.....	14
V. CONCLUSION.....	16
REFERENCES.....	17
DISTRIBUTION LIST.....	21

[illegible]

I. INTRODUCTION

Recent advances in the efficient evaluation of integral derivatives¹⁻³ have resulted in a renewed interest in the analytical evaluation of the first and second derivatives of the energy with respect to nuclear coordinates for Hartree-Fock and post-Hartree-Fock wavefunctions. Much of this work has been pioneered by Pople and co-workers^{4,5} who developed efficient codes both for the evaluation of the second derivatives of the integrals and for solving the coupled-perturbed Hartree-Fock equations. The first and second derivatives are extremely useful tools for optimizing structures, characterizing stationary points, and calculating vibrational frequencies, as well as for more extensive studies of the Born-Oppenheimer potential energy surface away from the stationary points. The last several years have also seen tremendous advances made in CI⁶⁻¹⁰ and MCSCF¹¹⁻²⁷ methodologies, which coupled with the interest in derivatives, have led to powerful techniques for a wide range of wavefunctions. In this vein, we report our work on general, multireference CI gradients and MCSCF second derivatives.

Before presenting our work, it is appropriate to quickly review the previous activity in the area of CI gradients and MCSCF second derivatives. In 1980, both Krishnan, et al.⁵ and Brooks, et al.²⁸ presented both equations and algorithms for the analytical computation of single-reference UHF or closed-shell RHF CI gradients. Osamura, et al. extended the capabilities to include open-shell RHF reference functions,²⁹ and more recently published the equations for general multireference CI gradients.³⁰ Yamaguchi, et al.³¹ reported second derivatives for two-configuration SCF (TCSCF) wavefunctions and Camp, et al.³² developed the methodology and implemented second derivatives for CAS MCSCF wavefunctions. Independently, Pulay,³³ as well as Jorgensen and Simons,³⁴ have provided equations for MCSCF second derivatives. The work on both multireference CI gradients and MCSCF second derivatives is closely related since a major step in both types of calculation is the solution of the coupled-perturbed MCSCF (CPMCSCF) equations.

In this paper we report the first general multireference CI gradient calculations. We have extended the theory developed by Osamura, et al.³⁰ to include a general class of references in the CI. This extension is necessary for the calculation of gradients for a commonly employed class of CI wavefunctions for which the reference configurations are selected from a generalized CAS MCSCF wavefunction. By generalized CAS we mean that there is at least one partially occupied orbital subspace where the energy is invariant to rotations of the orbitals in space. The simplest example of this type of wavefunction is a full CI in a selected subspace of orbitals. The length of the CI expansion in such a calculation grows rapidly with the size of the active space. To avoid unreasonably large MCSCF expansions, it is desirable to partition the space of chemically active orbitals and perform CAS calculations in some or all of the subspaces, while a GVB type wavefunction might be used in the remaining subspaces. A MCSCF calculation where all single and double excitations are generated from one subspace into another subspace would also be considered a generalized CAS wavefunction. In order to calculate the gradient of a selected reference CI wavefunction in which the orbitals are obtained from a generalized CAS wavefunction, orbital derivatives not appearing in the CPMCSCF equations are needed and we develop the machinery necessary to obtain these quantities.

We also present the equations employed in our MCSCF second derivative calculations. Our equations are an extension of the open-shell SCF second derivative formulas of Osamura, et al.⁴² These equations are different from those presented by Pulay³³ and Jorgensen and Simons³⁴ principally in the manner in which the derivative overlap contributions are handled. We expand upon these differences later in the text. In addition, the open-shell second derivative formulas of Osamura, et al.⁴² are reformulated so that we need not contract Coulomb or exchange operators after the CPMSCF equations are solved.

The CPMSCF equations are also presented in a more compact and convenient manner than Osamura's. The gradient-like terms appearing in the CPMSCF are defined in terms of modified integrals involving derivative overlap terms, as well as integrals involving derivative atomic orbitals. This formulation is particularly convenient when MCSCF second derivatives are desired.

Finally, we present the results of sample calculations on the reaction, $\text{Be} + \text{H}_2 \rightarrow \text{BeH}_2$. Here, second derivatives obtained at the MCSCF level are used in the optimization of the geometry of the products and for locating the transition state at the multireference CI level. We compare the results obtained at the second-order CI level with the results obtained in a selected reference CI (SRCI) calculation.

II. COMPUTATIONAL DETAILS

A. First Derivatives

The energy of a general CI wavefunction

$$\psi = \sum_i C_i X_i \quad (1)$$

can be expressed as

$$E = \sum_{ij} \left(\sum_{p,q} C_p K_{pq}^{ij} C_q \right) h_{ij} + \sum_{ijkl} \left(\sum_{p,q} C_p K_{pq}^{ijkl} C_q \right) g_{ijkl} \quad (2a)$$

$$= \sum_{ij} D_{ij} h_{ij} + \sum_{ijkl} D_{ijkl} g_{ijkl} \quad (2b)$$

$$= \sum_{p,q} C_p H_{pq} C_q, \quad (2c)$$

where C is the CI vector, X is a configuration state function (CSF), h_{ij} and g_{ijkl} are one and two electron MO integrals, K_{pq}^{ij} and K_{pq}^{ijkl} are spin coupling constants, and H_{pq} is an element of the Hamiltonian matrix in our CSF basis. D_{ij} and D_{ijkl} are one-particle and two-particle density matrix elements, respectively. The first derivative of the CI energy, with respect to nuclear coordinates, is²⁸

$$E^a = \sum_{pq} C_p H_{pq}^a C_q + \sum_i^{\text{occ}} \sum_r^{\text{all}} L_{ri} U_{ri}^a \quad (3a)$$

$$= \sum_{ij} D_{ij} h_{ij}^a + \sum_{ijkl} D_{ijkl} g_{ijkl}^a + \sum_i^{\text{occ}} \sum_r^{\text{all}} L_{ri} U_{ri}^a \quad (3b)$$

In this equation h_{ij}^a and g_{ijkl}^a are derivatives of the one- and two-electron atomic orbital integrals transformed to the molecular orbital basis. The derivatives of the molecular orbital expansion coefficients corresponding to orbital i , ϕ_i^a , are expressed in terms of undifferentiated coefficient vectors ϕ_p as

$$\phi_i^a = \sum_p U_{pi}^a \phi_p \quad (4)$$

L_{ri} is a Lagrangian multiplier defined as

$$L_{ri} = 2 \left(\sum_j D_{ij} h_{rj} + 2 \sum_{jkl} D_{ijkl} g_{rjkl} \right) \quad (5)$$

For a MCSCF wavefunction

$$L_{ij} = L_{ji} \text{ for } ij \text{ occupied} \quad (6a)$$

and

$$L_{ri} = 0 \text{ for } i = \text{occupied and } r = \text{virtual}. \quad (6b)$$

These relations are simply a reflection of the variational conditions on the orbitals. Not all of the elements of U_{ij}^a are independent. By differentiating the orthonormality condition

$$U^\dagger S U = I \quad (7)$$

one obtains

$$U_{ij}^a + S_{ij}^a + U_{ji}^a = 0, \quad (8)$$

where

$$S_{ij}^a = \sum_{xy} \phi_{xi} S_{xy}^a \phi_{yj} \quad (9)$$

S_{xy}^a is the derivative of the x,y atomic orbital overlap integral with respect to nuclear coordinate a . U_{ij}^a can then be expressed as a sum of an antisymmetric matrix and an upper triangular matrix

$$U_{ij}^a = \Delta_{ij}^a + T_{ij}^a, \quad (10)$$

where

$$\Delta_{ij}^a = -(\Delta_{ij}^a)^+ \quad (11)$$

and T_{ij}^a is defined by

$$T_{ij}^a = -S_{ij}^a \text{ for } i < j \quad (12a)$$

$$= -\frac{1}{2} S_{ii} \text{ for } i=j \quad (12b)$$

$$= 0 \text{ for } i > j. \quad (12c)$$

Thus, the derivative of the MCSCF energy is

$$E^a = \sum_{ij} D_{ij} h_{ij}^a + \sum_{ijkl} D_{ijkl} g_{ijkl}^a + \sum_{i \text{ occ}} \sum_{j \text{ all}} L_{ij} T_{ij}^a \quad (13)$$

since

$$\sum_{i \text{ occ}} \sum_{j \text{ all}} L_{ri} \Delta_{ri}^a = 0. \quad (14)$$

Before proceeding with a discussion of the CPMCSCF equations, it is useful to further examine the contributions to the first derivative. We note that it is possible to rewrite the first derivative for a CI wavefunction as

$$E^a = \sum_{pq} C_p H_{pq}^a C_q + \sum_{pq} C_p H_{pq}^{U^a} C_q. \quad (15)$$

This equation is obtained by reordering the sums in Eq. (3a) as follows:

$$\begin{aligned} \sum_{i \text{ occ}} \sum_{j \text{ all}} L_{ri} U_{ri}^a &= \sum_{i \text{ occ}} \sum_{j \text{ all}} D_{ij} \left(\sum_{r \text{ occ}} 2 U_{ri}^a h_{rj} \right) \\ &+ \sum_{i \text{ occ}} \sum_{j \text{ all}} D_{ijkl} \left(\sum_{r \text{ occ}} 4 U_{ri}^a g_{rjkl} \right) \end{aligned} \quad (16a)$$

$$= \sum_{i \text{ occ}} D_{ij} \sum_{r \text{ occ}} (U_{ri}^a h_{rj} + U_{rj}^a h_{ri}) \quad (16b)$$

$$\begin{aligned} &+ \sum_{i \text{ occ}} \sum_{j \text{ all}} D_{ijkl} \sum_{r \text{ occ}} (U_{ri}^a g_{rjkl} + U_{rj}^a g_{irkl} \\ &+ U_{rk}^a g_{ijrl} + U_{rl}^a g_{ijkr}) \end{aligned}$$

$$= \sum_j D_{ij} h_{ij}^u + \sum_{ijkl} D_{ijkl} g_{ijkl}^u \quad (16c)$$

$$= \sum_{p,q} C_p H_{pq}^u C_q, \quad (16d)$$

where

$$h_{ij}^u = \sum_r (U_{ri}^a h_{rj} + U_{rj}^a h_{ri}) \quad (16e)$$

and

$$g_{ijkl}^u = \sum_r (U_{ri}^a g_{rjkl} + U_{rj}^a g_{irkl} + U_{rk}^a g_{ijrl} + U_{rl}^a g_{ijkr})$$

Similarly, the first derivative of the MCSCF energy is

$$E^a = \sum_p \sum_q C_p H_{pq}^a C_q + \sum_p \sum_q C_p H_{pq}^{Ta} C_q. \quad (17)$$

Partial-derivative integrals similar to h_{ij}^u and g_{ijkl}^u arise naturally in the quadratic SCF procedure of Bacskay³⁵ and in the atomic orbital based CPHF equations of Osamura, et al.³⁶ They have also been exploited by Olsen, et al.²³ in a cubic MCSCF procedure and by Lengsfeld²² in a quadratic MCSCF approach designed to handle large CI expansions. These integrals have also been used by Dupuis,³⁷ Pulay,³³ and Jorgensen and Simons³⁴ to simplify their derivative expressions. The partial-derivative Hamiltonian constructed from these quantities is particularly useful as it occurs in both the CPMCSCF equations and in the expressions for the MCSCF second derivatives.

B. Coupled Perturbed MCSCF Equations

The derivative of the molecular orbitals U_{ij}^a , which are needed to compute the CI gradient, are obtained by solving the CPMCSCF equations. These equations are generated by requiring that the wavefunction satisfy the MCSCF variational conditions to first order with a change in nuclear geometry. Thus, these equations result from requiring that the derivatives of the orbital and CI stationary conditions, with respect to a nuclear coordinate, vanish.

$$\frac{dG_{ij}}{da} = \frac{d(L_{ij} - L_{ji})}{da} = 0 \quad (18a)$$

$$= \frac{\partial G_{ij}}{\partial a} + \sum_{mn} \frac{\partial \Delta_{mn}}{\partial a} \frac{\partial G_{ij}}{\partial \Delta_{mn}} + \sum_p \frac{\partial C_p}{\partial a} \frac{\partial G_{ij}}{\partial C_p} + \sum_{\substack{\text{occ} \\ s}} \sum_{\substack{\text{all} \\ r}} \frac{\partial T_{rs}}{\partial a} \frac{\partial G_{ij}}{\partial T_{rs}} \quad (18b)$$

and

$$\frac{dG_p^{CI}}{da} = \frac{d}{da} \sum_q (H_{pq} - E\delta_{pq}) C_q = 0 \quad (19a)$$

$$= \frac{\partial G_p^{CI}}{\partial a} + \sum_{mn} \frac{\partial \Delta_{mn}}{\partial a} \frac{\partial G_p^{CI}}{\partial \Delta_{mn}} + \sum_q \frac{\partial C_q}{\partial a} \frac{\partial G_p^{CI}}{\partial C_q} + \text{ofc all } \sum_{rs} \frac{\partial T_{rs}}{\partial a} \frac{\partial G_p^{CI}}{\partial T_{rs}} \quad (19b)$$

The prime sign indicates that the sum only runs over the unique orbital rotations which change the energy. Gathering the terms which are known on the right-hand side of the equation, we obtain the CPMCSCF equations

$$\sum_{mn} \left(\frac{\partial G_{ij}^{CI}}{\partial \Delta_{mn}} \right) \Delta_{mn}^a + \sum_q \left(\frac{\partial G_{ij}^{CI}}{\partial C_q} \right) C_q^a = - (G_{ij}^a + G_{ij}^{Ta}) \quad (20)$$

$$\sum_{mn} \left(\frac{\partial G_p^{CI}}{\partial \Delta_{mn}} \right) \Delta_{mn}^a + \sum_q \left(\frac{\partial G_p^{CI}}{\partial C_q} \right) C_q^a = - [(G_p^{CI})^a + (G_p^{CI})^{Ta}] \quad (21)$$

where

$$\Delta_{mn}^a = \frac{\partial \Delta_{mn}}{\partial a} \text{ and } C_q^a = \frac{\partial C_q}{\partial a} \quad ,$$

with the condition that

$$\sum_p \frac{\partial C_p}{\partial a} C_p = 0 \quad .$$

In these expressions the superscript "a" in G_{ij}^a and $(G_p^{CI})^a$ is used to denote that derivative AO integrals are employed in the construction of these quantities. The left-hand side of these equations contains explicitly the Hessian matrix appearing in second order MCSCF theory. These equations are in fact similar in structure to the second order MCSCF equations, and can thus be most efficiently solved by expressing the CI variations in the CSF basis, as noted by Lengsfeld and Liu.^{21,22} These equations are also similar to those obtained by Osamura, et al.³⁰ However, we have reordered the sums appearing on the right-hand side of this equation in order that the quantities needed to compute the MCSCF second derivatives are readily available.

C. MCSCF Second Derivatives

The derivative with respect to a nuclear displacement can be expanded as follows:

$$\frac{d}{db} = \left(\frac{\partial}{\partial b} + \text{all } \sum_i \frac{\partial U_{ri}}{\partial b} \frac{\partial}{\partial U_{ri}} \right) + \sum_p \frac{\partial C_p}{\partial b} \frac{\partial}{\partial C_p} \quad (22a)$$

$$= \left(\frac{\partial}{\partial b} + \sum_i^{\text{all}} \sum_i^{\text{all}} U_{ri}^b \frac{\partial}{\partial U_{ri}} \right) + \sum_p C_p^b \frac{\partial}{\partial C_p} \quad (22b)$$

The second derivative of our MCSCF wavefunction is obtained by applying this operator to our expression for the first derivative. It simplifies matters if we make use of both expressions (Eqs. (13) and (15)) for the first derivative. Our second derivative expression is obtained by operating on Eq. (13) with the first two terms in Eq. (22) and then operating on Eq. (15) with the last term in Eq. (22).

We obtain

$$\begin{aligned} E^{ab} = & 2 \sum_{pq} C_p^b H_{pq}^a C_q + 2 \sum_{pq} C_p^b H_{pq}^{Ta} C_q + \sum_{ij} D_{ij}^{hab} + \sum_{ijkl} D_{ijkl}^{ab} \\ & + \sum_i^{\text{occ}} \sum_i^{\text{all}} L_{ri}^a U_{ri}^b + \sum_i^{\text{occ}} \sum_i^{\text{all}} L_{ri}^b T_{ri}^a + \sum_i^{\text{occ}} \sum_i^{\text{all}} L_{ri}^{Ub} T_{ri}^a \\ & + \sum_{ij}^{\text{occ}} L_{ij}^{Tab} + \sum_{ij}^{\text{occ}} L_{ij} \left(\sum_m^{\text{all}} U_{mi}^b T_{mj}^a + T_{im}^a U_{mj}^b \right) \end{aligned} \quad (23)$$

Note that only the first two terms are unique to MCSCF second derivatives. The remaining terms are equivalent to the open-shell formulas of Osamura, et al.⁴² It is also important to note that almost all of the quantities involving MO or CI derivatives are generated by setting up or solving the CPMCSCF equations. In particular, the vector

$$B_p^a = \sum_q (H_{pq}^a + H_{pq}^{Ta}) C_q$$

can be stored when $G_{CI}^a + G_{CI}^{Ta}$ is constructed. Moreover, the Lagrangian multipliers L_{ri}^a and L_{ri}^{Ta} are used to obtain the gradient terms G_{ij}^a and G_{ij}^{Ta} in the CPMCSCF equations. The trace of the density matrix with the second derivative integrals is calculated by transforming the density matrix elements to the AO basis. The most time consuming step in this transformation only requires mn^4 multiplications where m is the number of active orbitals and n is number of basis functions. The remaining contributions can be obtained from the product of one-particle density matrices.

The only term which requires further consideration is $\sum_{ir} L_{ri}^{Ub} T_{ri}^a$. This term formally requires one to contract Coulomb and exchange operators (or more efficiently to contract Osamura's Y_{nimj} matrix) with the solution to the CPMCSCF equations, U^b , to form L_{ri}^{Ub} . However, we can eliminate this step by noting that

$$\begin{aligned}
& \sum_r^{\text{all}} \sum_i^{\text{occ}} L_{ri}^b T_{ri}^a = \sum_r^{\text{all}} \sum_i^{\text{occ}} L_{ri} T_{ri}^a U_{ri}^b \\
& + \sum_j^{\text{occ}} L_{ji} \sum_p^{\text{all}} (U_{rp}^b T_{pi}^a - T_{rp}^a U_{pi}^b) .
\end{aligned}
\tag{24}$$

The right-hand side of this equation is particularly convenient as all of these terms are generated when we set up the CPMCSCF equations. Using this relation and Eq. (10), we now obtain the final second derivative expression.

$$\begin{aligned}
E^{ab} = & 2 \sum_{pq} C_p^b (H_{pq}^a + H_{pq}^{Ta}) C_q + \sum_{ij} D_{ij} h_{ij}^{ab} + \sum_{ijkl} D_{ijkl} g_{ijkl}^{ab} \\
& + \sum_r^{\text{all}} \sum_i^{\text{occ}} L_{ri}^b T_{ri}^a + \sum_r^{\text{all}} \sum_i^{\text{occ}} (L_{ri}^a + L_{ri}^{Ta}) U_{ri}^b \\
& + \sum_j^{\text{occ}} L_{ji} [T_{ji}^{ab} + \sum_p^{\text{all}} (T_{pi}^a T_{pj}^b + T_{pi}^a T_{jp}^b)] .
\end{aligned}
\tag{25}$$

Equation (23) can be decomposed in another fashion if the orthonormality conditions are expressed in such a way that T is symmetric. Camp, King, McIver, and Mullally³² have expressed the orthonormality conditions in this way. However, there is an advantage in employing an upper triangular T matrix as only the first p (p is the number of occupied orbitals) columns are needed to obtain the integrals h_{ij}^{Ta} and g_{ijkl}^{Ta} . Thus, the transformations needed to obtain h_{ij}^{Ta} and g_{ijkl}^{Ta} can be performed very efficiently. This fact is also exploited in Osamura's equations.

D. Selected Reference CI Gradients

The gradient of the energy for a CI wavefunction requires knowledge of the first order variations in the molecular orbitals. The CPMCSCF scheme does not uniquely define a transformation of the orbitals, but only specifies a transformation of the variational parameters. This is enough information to define an orbital transformation excluding an arbitrary orthogonal mixing within the invariant subspaces. If the CI wavefunction has the same invariant subspaces as the reference wavefunction, then the gradient is well defined. This is the case, e.g., with a SDCI using closed-shell Hartree-Fock orbitals. Both wavefunctions are invariant to mixings among the doubly occupied core orbitals and information concerning such mixings is not required for the CI gradient.

For a CI wavefunction constructed as all excitations of a given order from selected references of a generalized CAS wavefunction, this is not the case. The CI is not invariant to mixings of orbitals within the partially

occupied subspaces. The most common way to address this problem in calculating the CI energy itself is to require that the orbitals be natural orbitals of the one-particle density matrix. For the calculation of the CI gradient, we then remove the ambiguity by requiring the natural orbital conditions to be satisfied to first order with a change in nuclear geometry. The derivative of the natural orbital condition must vanish in analogy with Eqs. (18) and (19) for the variational conditions.

$$\frac{d}{da} (DW - W\lambda) = 0 \quad . \quad (26)$$

Here D is a subblock of the one particle density matrix and λ is a diagonal matrix of occupation numbers. At $a=0$, $W=I$, and thus $D(0)=\lambda(0)$.

Evaluating Eq. (26) leads to expressions for the derivatives of parameters which mix orbitals within the invariant subspace.

$$\Delta_{ij}^a = w_{ij}^a = \frac{-D_{ij}^a}{\lambda_i - \lambda_j} \quad . \quad (27)$$

In this equation, $D_{ij}^a = 2 \sum_{pq} C_p^{aKij} D_{pq}$ and C_p^a is obtained from the solution to the CPMCSCF equations. Δ_{ij}^a is then obtained with the overlap derivative portion T_{ij}^a as in Eq. (10) to obtain U_{ij}^a .

III. COMPARISON WITH OTHER WORK

As noted in the Introduction, our second derivative expressions differ from those of Pulay³³ and Jorgensen and Simons³⁴ principally in the manner in which the derivative overlap terms are included in the CPMCSCF equations, and subsequently in the final expression for the second derivatives. Jorgensen and Simons include an overlap term in the definition of their AO basis. They are then able to derive a very compact set of formulas. However, the formulas for the derivatives of their AO integrals are involved and the overall efficiency of their method depends very strongly on how their derivative AOs are computed. We should also note that Jorgensen and Simons' method was developed to describe second derivatives of CI and coupled-cluster wavefunctions, as well as MCSCF wavefunctions. The comparative efficiency of their method must also be judged on how well it treats highly correlated wavefunctions, and this analysis is beyond the scope of the present study.

Pulay has derived the CPMCSCF equations in such a manner that the orbital variations are expressed in the AO basis (as opposed to the MO basis used by Osamura) and the CI variations are in the CSF basis. His CPMCSCF equations do not require a contraction of Coulomb and exchange operators with overlap terms, but his CPMCSCF equations are larger than Osamura's because he is working in the AO basis. We feel that the overall efficiency of both methods should be about the same. However, we also note that Pulay's equations neglect projection operator terms appearing in Osamura's equations (see also Lengsfeld and Liu²¹). These terms are needed if the Hessian appearing in the CPMCSCF equations is to be nonsingular.

The recent communication by Camp, et al.,³² provides few details of their second derivative equations. However, they express the variation of the molecular orbitals with nuclear displacement in terms of a product of an exponential and a Hermitian operator. The orthonormality conditions at first order are treated by this Hermitian operator as opposed to the upper triangular matrix appearing in Eq. (12). Further details are needed to determine if the simplicity of their final equations offset the computational expense of working with a Hermitian matrix as opposed to an upper triangular matrix in constructing terms like L^{T^a} appearing in our second derivative expression.

IV. SAMPLE CALCULATIONS

The reaction $\text{Be} + \text{H}_2 \rightarrow \text{BeH}_2$ was studied in C_{2v} symmetry. The geometry of the products and transition state was stabilized at the MCSCF and multi-reference CI level. The MCSCF wavefunction employed in this study was a four-electron in four-orbital CAS. This wavefunction correctly describes the cleavage of the BeH bonds, but does not contain all of the configurations needed to describe s^2 and p^2 near degeneracy in Be. Second order CI calculations, based on the 4 in 4 CAS, are compared to the results of selected reference singles and doubles CI (SRCI) calculations. In the SRCI calculations, the references were selected on the basis of their cumulative weight in the natural orbital representation of the MCSCF wavefunction. The weight, W , was defined as follows:

$$W = \sum_i C_i^2 \quad (28)$$

The basis set used for beryllium was Dunning's 5s contraction³⁸ of Huzinaga's 10s primitive set³⁹ and Bartlett's p function⁴⁰ composed of three primitives which we augmented with an uncontracted p function of exponent 0.057 181. The hydrogen basis was Dunning's 2s contraction⁴¹ of Huzinaga's 4s primitive set³⁹ with a scale factor of 1.2. The results of these calculations are presented in the following four tables. Table 1 lists the references employed in the SRCI calculations. The stable geometries are given in Table 2, and the MCSCF vibrational frequencies in Table 3. Our calculated MCSCF and CI energies are given in Table 4.

The reaction is symmetry forbidden along the C_{2v} reaction path and this results in a transition state with two dominant CSFs.

The results of the SRCI calculations accurately reproduce the second order CI results while only requiring a fraction of the computational effort. The MCSCF calculations provide a good description of the reaction (in this basis) but the activation energy is a bit high as expected.

We found that the multireference CI gradient calculations converged very rapidly to stable points when MCSCF second derivatives and starting geometries were employed. In the most unfavorable case, four iterations were required for convergence (largest component of the gradient less than 1.0×10^{-4} a.u./bohr).

Table 1. Dominant Configurations in the MCSCF Wavefunction

<u>Equilibrium Geometry Configuration</u>		<u>Coefficient</u>
1	$1a_1^2 2a_1^2 1b_2^2$	0.989
2	$1a_1^2 2a_1 1b_2 3a_1 2b_2$	0.092
Weight = 0.98766		
<u>Transition State Geometry Configuration</u>		<u>Coefficient</u>
1	$1a_1^2 2a_1^2 1b_2^2$	0.740
2	$1a_1^2 2a_1^2 3a_1^2$	-0.623
3	$1a_1^2 2a_1 1b_2 3a_1 2b_2^*$	-0.197 0.117

*The two coefficients associated with this configuration correspond to the two spin couplings.

Table 2. Geometries of BeH₂ Stationary Points

Equilibrium	<u>MCSCF</u>	<u>SRCI</u>	<u>SDCI</u>
R _{BeH}	2.61 876 bohr	2.60 512 bohr	2.60 637 bohr
θ	180.00°	180.00°	180.00°
C _{2v}	3.0875 bohr	3.0687 bohr	3.0693 bohr
θ	47.38°	48.68°	48.76°

Table 3. MCSCF Vibrational Frequencies

<u>Equilibrium</u>	<u>Transition State</u>
790.1 cm ⁻¹	4044.9i cm ⁻¹
790.1 cm ⁻¹	
1827.5 cm ⁻¹	993.4 cm ⁻¹
2069.4 cm ⁻¹	4514.7 cm ⁻¹

Table 4. MCSCF and CI Energies for BeH₂

	<u>MCSCF</u>	<u>SRCI</u>	<u>SDCI</u>
Equilibrium	-15.773 651 a.u.	-15.798 597 a.u.	-15.798 965 a.u.
Transition State	-15.597 013 a.u.	-15.628 925 a.u.	-15.629 315 a.u.
ΔE	0.176 638 a.u. (110.8 kcal)	0.169 672 a.u. (106.5 kcal)	0.169 650 a.u. (106.45 kcal)

V. CONCLUSION

We have presented a set of simple and efficient formulas for the calculation of multireference CI gradients and MCSCF second derivatives. The additional machinery needed to compute selected-reference CI gradients was developed. Sample calculations were presented in which SRCI structures and activation energies compared very favorably with the full second order CI results. In the reaction investigated in this work, MCSCF second derivatives were found to be very useful in the stabilization of the CI structures.

REFERENCES

1. M. Dupuis and H.F. King, "Molecular Symmetry and Closed-Shell SCF Calculations," Int. J. Quantum Chem., Vol. 11, p. 613, 1977.
2. P. Saxe, Y. Yamaguchi, and H.F. Schaefer, III, "Analytical Second Derivatives in Restricted Hartree-Fock Theory. A Method for High-Spin Open-Shell Molecular Wavefunctions," J. Chem. Phys., Vol. 77, p. 5647, 1982.
3. H.B. Schlegel, "An Efficient Algorithm for Calculating Ab Initio Energy Gradients Using s,p Cartesian Gaussians," J. Chem. Phys., Vol. 77, p. 3675, 1982.
4. J.A. Pople, R. Krishnan, H.B. Schlegel, and J.S. Binkley, "Derivative Studies in Hartree-Fock and Moller-Plesset Theories," Int. J. Quantum Chem. Symp., Vol. 13, p. 225, 1979.
5. R. Krishnan, H.B. Schlegel, and J.A. Pople, "Derivative Studies in Configuration-Interaction Theory," J. Chem. Phys., Vol. 72, p. 4654, 1980.
6. H. Lischka, R. Shepard, F.B. Brown, and I. Shavitt, "New Implementation of the Graphical Unitary Group Approach for Multireference Direct Configuration Interaction Calculations," Int. J. Quantum Chem. Symp., Vol. 15, p. 91, 1981.
7. P. Saxe, D.J. Fox, H.F. Schaefer, III, and N.C. Handy, "The Shape-Driven Graphical Unitary Group Approach to the Electron Correlation Problem. Application to the Ethylene Molecule," J. Chem. Phys., Vol. 77, p. 5584, 1982.
8. B. Liu and M. Yoshimine, "The Alchemy Configuration Interaction Method. I. The Symbolic Matrix Method for Determining Elements of Matrix Operators," J. Chem. Phys., Vol. 74, p. 612, 1982.
9. P.E.M. Siegbahn, "Generalizations of the Direct CI Method Based on the Graphical Unitary Group Approach. I. Single Replacements from a Complete CI Root Function of Any Spin, First Order Wavefunctions," J. Chem. Phys., Vol. 70, p. 5391, 1979.
10. P.E.M. Siegbahn, "Generalizations of the Direct CI Method Based on the Graphical Unitary Group Approach. II. Single and Double Replacements from Any Set of Reference Configurations," J. Chem. Phys., Vol. 72, p. 1647, 1980.
11. L.G. Yaffe and W.A. Goddard, III, "Orbital Optimization in Electronic Wavefunctions; Equations for Quadratic and Cubic Convergence of General Multiconfiguration Wavefunctions," Phys. Rev. A, Vol. 13, p. 1682, 1976.
12. E. Daalgard and P. Jorgensen, "Optimization of Orbitals for Multiconfiguration Reference States," J. Chem. Phys., Vol. 69, p. 3833, 1978.

13. D. Yeager and P. Jorgensen, "Convergency Studies of Second and Approximate Second Order Multiconfigurational Hartree-Fock Procedures," J. Chem. Phys., Vol. 71, p. 755, 1979.
14. B.H. Lengsfeld, III, "General Second Order MCSCF Theory: A Density Matrix Directed Algorithm," J. Chem. Phys., Vol. 73, p. 382, 1980.
15. H.J. Werner and W. Meyer, "A Quadratically Convergent Multiconfiguration Self-Consistent Field Method with Simultaneous Optimization of Orbitals and CI Coefficients," J. Chem. Phys., Vol. 73, p. 2342, 1980.
16. B.O. Roos, P.R. Taylor, and P.E.M. Siegbahn, "A Complete Active Space SCF (CASSCF) Using a Density Matrix Formulated Super-CI Approach," Chem. Phys., Vol. 48, p. 157, 1980.
17. P.E.M. Siegbahn, J. Almlof, A. Heiberg, and B.O. Roos, "The Complete Active Space SCF (CASSCF) Method in a Newton-Raphson Formulation with Application to the HNO Molecule," J. Chem. Phys., Vol. 74, p. 2384, 1981.
18. B.O. Roos, "The Complete Active Space SCF Method in a Fock-Matrix-Based Super-CI Formulation," Int. J. Quantum Chem. Symp., Vol. 14, p. 175, 1980.
19. H.J. Werner and W. Meyer, "A Quadratically Convergent MCSCF Method for the Simultaneous Optimization of Several States," J. Chem. Phys., Vol. 74, p. 5794, 1981.
20. R.N. Diffenderfer and D.R. Yarkony, "Application of the State-Averaged MCSCF Procedure: Application to Radiative Transitions of MgO," J. Phys. Chem., Vol. 86, p. 5098, 1980.
21. B.H. Lengsfeld, III and B. Liu, "A Second Order MCSCF Method for Large CI Expansions," J. Chem. Phys., Vol. 75, p. 478, 1981.
22. B.H. Lengsfeld, III, "General Second-Order MCSCF Theory for Large CI Expansions," J. Chem. Phys., Vol. 77, p. 4073, 1982.
23. J. Olsen, P. Jorgensen, and D.L. Yeager, "Cubic Contributions in Multiconfigurational Self-Consistent Field (MCSCF) Calculations," J. Chem. Phys., Vol. 77, p. 356, 1982.
24. R.N. Camp and H.F. King, "Stable Methods for Achieving MCSCF Convergence," J. Chem. Phys., Vol. 77, p. 3056, 1982.
25. G. Das, "A Quadratically Convergent MCSCF Scheme Using Fock Operators," J. Chem. Phys., Vol. 74, p. 5775, 1980.
26. P.E.M. Siegbahn, A. Heiberg, B.O. Roos, and B. Levy, "A Comparison of the Super-CI and the Newton-Raphson Scheme in the Complete Active Space SCF Method," Phys. Scr., Vol. 21, p. 323, 1980.

27. R. Shepard and J. Simons, "Multiconfiguration Wavefunction Optimization Using the Unitary Group Method," Int. J. Quantum Chem. Symp., Vol. 14, p. 211, 1980.
28. B.R. Brooks, W.D. Laidig, P. Saxe, J.D. Goddard, Y. Yamaguchi, and H.F. Schaefer, III, "Analytic Gradients from Correlated Wavefunctions Via the Two-Particle Density Matrix and the Unitary Group Approach," J. Chem. Phys., Vol. 72, p. 4652, 1980.
29. Y. Osamura, Y. Yamaguchi, and H.F. Schaefer, III, "Analytic Configuration Interaction (CI) Gradient Techniques for Potential Energy Hypersurfaces. A Method for Open-Shell Molecular Wavefunctions," J. Chem. Phys., Vol. 75, p. 2919, 1981.
30. Y. Osamura, Y. Yamaguchi, and H.F. Schaefer, III, "Generalization of Analytic Configuration Interaction (CI) Gradient Methods for Potential Energy Hypersurfaces, Including a Solution to the Coupled-Perturbed Hartree-Fock Equations for Multiconfiguration SCF Molecular Wavefunctions," J. Chem. Phys., J. Chem. Phys., Vol. 77, p. 383, 1982.
31. Y. Yamaguchi, Y. Osamura, G. Fitzgerald, and H.F. Schaefer, III, "Analytical Force Constants for Post-Hartree-Fock Wavefunctions: The Simplest Case," J. Chem. Phys., Vol. 78, p. 1607, 1983.
32. R.N. Camp, H.F. King, J.W. McIver, Jr., and D. Mullally, "Analytical Force Constants for MCSCF Wavefunctions," J. Chem. Phys., Vol. 79, p. 1088, 1983.
33. P. Pulay, "Second and Third Derivatives of Variational Energy Expressions: Application to Multiconfiguration Self-Consistent Field Wavefunctions," J. Chem. Phys., Vol. 78, p. 5043, 1983.
34. P. Jorgensen and J. Simons, "Ab Initio Analytical Molecular Gradients and Hessians," J. Chem. Phys., Vol. 79, p. 334, 1983.
35. G.B. Bacskay, "A Quadratically Convergent Hartree-Fock (QC-SCF) Method. Application to Open Shell Orbital Optimization and Coupled Perturbed Hartree-Fock Calculations," Chem. Phys., Vol. 65, p. 383, 1982.
36. Y.. Osamura, Y. Yamaguchi, P. Saxe, D.J. Fox, M.A. Vincent, and H.F. Schaefer, III, "Analytic Second Derivative Techniques for Self-Consistent-Field Wavefunctions. A New Approach to the Solution of the CPHF Equations," Theochem., Vol. 103, p. 183, 1983.
37. M. Dupuis, "Energy Derivatives for Configuration Interaction Wavefunctions," J. Chem. Phys., Vol. 74, p. 5758, 1981.
38. T.H. Dunning, Jr., "Gaussian Basis Sets for Use in Molecular Calculations. III. Contraction of (10s 6p) Atomic Basis Sets for First-Row Atoms," J. Chem. Phys., Vol. 55, p. 716, 1971. In our calculations, the large exponent in the 2s contraction on beryllium was 3.6880 instead of Dunning's 3.6680.

39. S. Huzinaga, "Gaussian Type Functions for Polyatomic Systems. I.," J. Chem. Phys., Vol. 42, p. 1293, 1965.
40. G.D. Purvis, III, R. Shepard, F.D. Brown, and R.J. Bartlett, " C_{2v} Insertion Pathway for BeH_2 : A Test Problem for the Coupled-Clusters Single and Double Excitation Model," Int. J. Quantum Chem., Vol. 23, p. 835, 1983.
41. T.H. Dunning, Jr., "Gaussian Basis Functions for Use in Molecular Calculations. I. Contraction of (9s 5p) Atomic Basis Sets for the First-Row Atoms," J. Chem. Phys., Vol. 53, p. 2823, 1970.
42. Y. Osamura, Y. Yamaguchi, P. Saxe, M.A. Vincent, J.F. Gaw, and H.F. Schaefer, III, "Unified Theoretical Treatment of Analytic First and Second Energy Derivatives in Open-Shell Hartree-Fock Theory," Chem. Phys., Vol. 72, p. 131, 1982.

DISTRIBUTION LIST

<u>No. Of Copies</u>	<u>Organization</u>	<u>No. Of Copies</u>	<u>Organization</u>
12	Administrator Defense Technical Info Center ATTN: DTIC-DDA Cameron Station Alexandria, VA 22304-6145	1	Commander US Army Aviation Research and Development Command ATTN: AMSAV-E 4300 Goodfellow Blvd. St. Louis, MO 63120
1	HQ DA DAMA-ART-M Washington, DC 20310	1	Director US Army Air Mobility Research and Development Laboratory Ames Research Center Moffett Field, CA 94035
1	Commander US Army Materiel Command ATTN: AMCDRA-ST 5001 Eisenhower Avenue Alexandria, VA 22333-0001	4	Commander US Army Research Office ATTN: R. Ghirardelli D. Mann R. Singleton R. Shaw P.O. Box 12211 Research Triangle Park, NC 27709-2211
10	Central Intelligence Agency Office of Central Reference Dissemination Branch Room GE-47 HQS Washington, DC 20505		
1	Commander Armament R&D Center US Army AMCCOM ATTN: SMCAR-TSS Dover, NJ 07801	1	Commander US Army Communications - Electronics Command ATTN: AMSEL-ED Fort Monmouth, NJ 07703
1	Commander Armament R&D Center US Army AMCCOM ATTN: SMCAR-TDC Dover, NJ 07801	1	Commander ERADCOM Technical Library ATTN: DELSD-L, Reports Section Fort Monmouth, NJ 07703-5301
1	Director Benet Weapons Laboratory Armament R&D Center US Army AMCCOM ATTN: SMCAR-LCB-TL Watervliet, NY 12189	2	Commander Armament R,D&E Center US Army AMCCOM ATTN: SMCAR-LCA-G, D.S. Downs J.A. Lannon Dover, NJ 07801
1	Commander US Army Armament, Munitions and Chemical Command ATTN: SMCAR-ESP-L Rock Island, IL 61299	1	Commander Armament R,D&E Center US Army AMCCOM ATTN: SMCAR-LC-G, L. Harris Dover, NJ 07801

DISTRIBUTION LIST

<u>No. Of Copies</u>	<u>Organization</u>	<u>No. Of Copies</u>	<u>Organization</u>
1	Commander Armament R,D&E Center US Army AMCCOM ATTN: SMCAR-SCA-T, L. Stiefel Dover, NJ 07801	1	Commander US Army Development and Employment Agency ATTN: MODE-TED-SAB Fort Lewis, WA 98433
1	Commander US Army Missile Command Research, Development and Engineering Center ATTN: AMSMI-RD Redstone Arsenal, AL 35898	1	Office of Naval Research Department of the Navy ATTN: R.S. Miller, Code 432 800 N. Quincy Street Arlington, VA 22217
1	Commander US Army Missile and Space Intelligence Center ATTN: AMSMI-YDL Redstone Arsenal, AL 35898-5000	1	Commander Naval Air Systems Command ATTN: J. Ramnarace, AIR-54111C Washington, DC 20360
2	Commander US Army Missile Command ATTN: AMSMI-RK, D.J. Ifshin W. Wharton Redstone Arsenal, AL 35898	2	Commander Naval Ordnance Station ATTN: C. Irish P.L. Stang, Code 515 Indian Head, MD 20640
1	Commander US Army Missile Command ATTN: AMSMI-RKA, A.R. Maykut Redstone Arsenal, AL 35898-5249	1	Commander Naval Surface Weapons Center ATTN: J.L. East, Jr., G-23 Dahlgren, VA 22448-5000
1	Commander US Army Tank Automotive Command ATTN: AMSTA-TSL Warren, MI 48397-5000	2	Commander Naval Surface Weapons Center ATTN: R. Bernecker, R-13 G.B. Wilmot, R-16 Silver Spring, MD 20902-5000
1	Director US Army TRADOC Systems Analysis Activity ATTN: ATAA-SL White Sands Missile Range, NM 88002	1	Commander Naval Weapons Center ATTN: R.L. Derr, Code 389 China Lake, CA 93555
1	Commandant US Army Infantry School ATTN: ATSH-CD-CSO-OR Fort Benning, GA 31905	2	Commander Naval Weapons Center ATTN: Code 3891, T. Boggs K.J. Graham China Lake, CA 93555

DISTRIBUTION LIST

<u>No. Of Copies</u>	<u>Organization</u>	<u>No. Of Copies</u>	<u>Organization</u>
5	Commander Naval Research Laboratory ATTN: L. Harvey J. McDonald E. Oran J. Shnur R.J. Doyle, Code 6110 Washington, DC 20375	1	NASA Langley Research Center Langley Station ATTN: G.B. Northam/MS 168 Hampton, VA 23365
1	Commanding Officer Naval Underwater Systems Center Weapons Dept. ATTN: R.S. Lazar/Code 36301 Newport, RI 02840	4	National Bureau of Standards ATTN: J. Hastie M. Jacox T. Kashiwagi H. Semerjian US Department of Commerce Washington, DC 20234
1	Superintendent Naval Postgraduate School Dept. of Aeronautics ATTN: D.W. Netzer Monterey, CA 93940	1	Aerojet Solid Propulsion Co. ATTN: P. Micheli Sacramento, CA 95813
4	AFRPL/DY, Stop 24 ATTN: R. Corley R. Geisler J. Levine D. Weaver Edwards AFB, CA 93523-5000	1	Applied Combustion Technology, Inc. ATTN: A.M. Varney P.O. Box 17885 Orlando, FL 32860
1	AFRPL/MKPB, Stop 24 ATTN: B. Goshgarian Edwards AFB, CA 93523-5000	2	Applied Mechanics Reviews The American Society of Mechanical Engineers ATTN: R.E. White A.B. Wenzel 345 E. 47th Street New York, NY 10017
2	AFOSR ATTN: L.H. Caveny J.M. Tishkoff Bolling Air Force Base Washington, DC 20332	1	Atlantic Research Corp. ATTN: M.K. King 5390 Cherokee Avenue Alexandria, VA 22314
1	AFWL/SUL Kirtland AFB, NM 87117	1	Atlantic Research Corp. ATTN: R.H.W. Waesche 7511 Wellington Road Gainesville, VA 22065
1	Air Force Armament Laboratory ATTN: AFATL/DLODL Eglin AFB, FL 32542-5000	1	AVCO Everett Rsch. Lab. Div. ATTN: D. Stickler 2385 Revere Beach Parkway Everett, MA 02149

DISTRIBUTION LIST

<u>No. Of</u> <u>Copies</u>	<u>Organization</u>	<u>No. Of</u> <u>Copies</u>	<u>Organization</u>
1	Battelle Memorial Institute Tactical Technology Center ATTN: J. Huggins 505 King Avenue Columbus, OH 43201	1	General Motors Rsch Labs Physics Department ATTN: R. Teets Warren, MI 48090
1	Cohen Professional Services ATTN: N.S. Cohen 141 Channing Street Redlands, CA 92373	2	Hercules, Inc. Allegany Ballistics Lab. ATTN: R.R. Miller E.A. Yount P.O. Box 210 Cumberland, MD 21501
2	Exxon Research & Eng. Co. Government Research Lab ATTN: A. Dean M. Chou P.O. Box 48 Linden, NJ 07036	1	Hercules, Inc. Bacchus Works ATTN: K.P. McCarty P.O. Box 98 Magna, UT 84044
1	Ford Aerospace and Communications Corp. DIVAD Division Div. Hq., Irvine ATTN: D. Williams Main Street & Ford Road Newport Beach, CA 92663	1	Honeywell, Inc. Government and Aerospace Products ATTN: D.E. Broden/ MS MN50-2000 600 2nd Street NE Hopkins, MN 55343
1	General Applied Science Laboratories, Inc. ATTN: J.I. Erdos 425 Merrick Avenue Westbury, NY 11590	1	IBM Corporation ATTN: A.C. Tam Research Division 5600 Cottle Road San Jose, CA 95193
1	General Electric Armament & Electrical Systems ATTN: M.J. Bulman Lakeside Avenue Burlington, VT 05401	1	IIT Research Institute ATTN: R.F. Remaly 10 West 35th Street Chicago, IL 60616
1	General Electric Company ATTN: A. Wait 2352 Jade Lane Schenectady, NY 12309	2	Director Lawrence Livermore National Laboratory ATTN: C. Westbrook M. Costantino P.O. Box 808 Livermore, CA 94550
1	General Electric Ordnance Systems ATTN: J. Mandzy 100 Plastics Avenue Pittsfield, MA 01203		

DISTRIBUTION LIST

<u>No. Of Copies</u>	<u>Organization</u>	<u>No. Of Copies</u>	<u>Organization</u>
1	Lockheed Missiles & Space Co. ATTN: George Lo 3251 Hanover Street Dept. 52-35/B204/2 Palo Alto, CA 94304	1	Science Applications, Inc. ATTN: R.B. Edelman 23146 Cumorah Crest Woodland Hills, CA 91364
1	Los Alamos National Lab ATTN: B. Nichols T7, MS-B284 P.O. Box 1663 Los Alamos, NM 87545	1	Science Applications, Inc. ATTN: H.S. Pergament 1100 State Road, Bldg. N Princeton, NJ 08540
1	Olin Corporation Smokeless Powder Operations ATTN: V. McDonald P.O. Box 222 St. Marks, FL 32355	3	SRI International ATTN: G. Smith D. Crosley D. Golden 333 Ravenswood Avenue Menlo Park, CA 94025
1	Paul Gough Associates, Inc. ATTN: P.S. Gough 1048 South Street Portsmouth, NH 03801	1	Stevens Institute of Tech. Davidson Laboratory ATTN: R. McAlevy, III Hoboken, NJ 07030
2	Princeton Combustion Research Laboratories, Inc. ATTN: M. Summerfield N.A. Messina 475 US Highway One Monmouth Junction, NJ 08852	1	Teledyne McCormack-Selph ATTN: C. Leveritt 3601 Union Road Hollister, CA 95023
1	Hughes Aircraft Company ATTN: T.E. Ward 8433 Fallbrook Avenue Canoga Park, CA 91303	1	Textron, Inc. Bell Aerospace Co. Division ATTN: T.M. Ferger P.O. Box 1 Buffalo, NY 14240
1	Rockwell International Corp. Rocketdyne Division ATTN: J.E. Flanagan/HB02 6633 Canoga Avenue Canoga Park, CA 91304	1	Thiokol Corporation Elkton Division ATTN: W.N. Brundige P.O. Box 241 Elkton, MD 21921
4	Sandia National Laboratories Combustion Sciences Dept. ATTN: R. Cattolica S. Johnston P. Mattern D. Stephenson Livermore, CA 94550	1	Thiokol Corporation Huntsville Division ATTN: R. Glick Huntsville, AL 35807
		3	Thiokol Corporation Wasatch Division ATTN: S.J. Bennett P.O. Box 524 Brigham City, UT 84302

DISTRIBUTION LIST

<u>No. Of Copies</u>	<u>Organization</u>	<u>No. Of Copies</u>	<u>Organization</u>
1	United Technologies ATTN: A.C. Eckbreth East Hartford, CT 06108	1	University of California, Berkeley Mechanical Engineering Dept. ATTN: J. Daily Berkeley, CA 94720
3	United Technologies Corp. Chemical Systems Division ATTN: R.S. Brown T.D. Myers (2 copies) P.O. Box 50015 San Jose, CA 95150-0015	1	University of California Los Alamos Scientific Lab. ATTN: T.D. Butler P.O. Box 1663, Mail Stop B216 Los Alamos, NM 87545
2	United Technologies Corp. ATTN: R.S. Brown R.O. McLaren P.O. Box 358 Sunnyvale, CA 94086	2	University of California, Santa Barbara Quantum Institute ATTN: K. Schofield M. Steinberg Santa Barbara, CA 93106
1	Universal Propulsion Company ATTN: H.J. McSpadden Black Canyon Stage 1 Box 1140 Phoenix, AZ 85029	1	University of Southern California Dept. of Chemistry ATTN: S. Benson Los Angeles, CA 90007
1	Veritay Technology, Inc. ATTN: E.B. Fisher 4845 Millersport Highway P.O. Box 305 East Amherst, NY 14051-0305	1	Case Western Reserve Univ. Div. of Aerospace Sciences ATTN: J. Tien Cleveland, OH 44135
1	Brigham Young University Dept. of Chemical Engineering ATTN: M.W. Beckstead Provo, UT 84601	1	Cornell University Department of Chemistry ATTN: E. Grant Baker Laboratory Ithaca, NY 14853
1	California Institute of Tech. Jet Propulsion Laboratory ATTN: MS 125/159 4800 Oak Grove Drive Pasadena, CA 91103	1	Univ. of Dayton Rsch Inst. ATTN: D. Campbell AFRPL/PAP Stop 24 Edwards AFB, CA 93523
1	California Institute of Technology ATTN: F.E.C. Culick/ MC 301-46 204 Karman Lab. Pasadena, CA 91125	1	University of Florida Dept. of Chemistry ATTN: J. Winefordner Gainesville, FL 32611

DISTRIBUTION LIST

<u>No. Of Copies</u>	<u>Organization</u>	<u>No. Of Copies</u>	<u>Organization</u>
3	Georgia Institute of Technology School of Aerospace Engineering ATTN: E. Price W.C. Strahle B.T. Zinn Atlanta, GA 30332	2	Princeton University Forrestal Campus Library ATTN: K. Brezinsky I. Glassman P.O. Box 710 Princeton, NJ 08540
1	University of Illinois Dept. of Mech. Eng. ATTN: H. Krier 144MEB, 1206 W. Green St. Urbana, IL 61801	1	Princeton University MAE Dept. ATTN: F.A. Williams Princeton, NJ 08544
1	Johns Hopkins University/APL Chemical Propulsion Information Agency ATTN: T.W. Christian Johns Hopkins Road Laurel, MD 20707	1	Purdue University School of Aeronautics and Astronautics ATTN: J.R. Osborn Grissom Hall West Lafayette, IN 47906
1	University of Michigan Gas Dynamics Lab Aerospace Engineering Bldg. ATTN: G.M. Faeth Ann Harbor, MI 48109-2140	2	Purdue University School of Mechanical Engineering ATTN: N.M. Laurendeau S.N.B. Murthy TSPC Chaffee Hall West Lafayette, IN 47906
1	University of Minnesota Dept. of Mechanical Engineering ATTN: E. Fletcher Minneapolis, MN 55455	1	Rensselaer Polytechnic Inst. Dept. of Chemical Engineering ATTN: A. Fontijn Troy, NY 12181
3	Pennsylvania State University Applied Research Laboratory ATTN: K.K. Kuo H. Palmer M. Micci University Park, PA 16802	1	Stanford University Dept. of Mechanical Engineering ATTN: R. Hanson Stanford, CA 94305
1	Polytechnic Institute of NY Graduate Center ATTN: S. Lederman Route 110 Farmingdale, NY 11735	1	University of Texas Dept. of Chemistry ATTN: W. Gardiner Austin, TX 78712
		1	University of Utah Dept. of Chemical Engineering ATTN: G. Flandro Salt Lake City, UT 84112

DISTRIBUTION LIST

<u>No. Of Copies</u>	<u>Organization</u>
1	Virginia Polytechnic Institute and State University ATTN: J.A. Schetz Blacksburg, VA 24061

Aberdeen Proving Ground

Dir, USAMSAA
ATTN: AMXSY-D
AMXSY-MP, H. Cohen
Cdr, USATECOM
ATTN: AMSTE-TO-F
Cdr, CRDC, AMCCOM
ATTN: SMCCR-RSP-A
SMCCR-MU
SMCCR-SPS-IL

USER EVALUATION SHEET/CHANGE OF ADDRESS

This Laboratory undertakes a continuing effort to improve the quality of the reports it publishes. Your comments/answers to the items/questions below will aid us in our efforts.

1. BRL Report Number _____ Date of Report _____

2. Date Report Received _____

3. Does this report satisfy a need? (Comment on purpose, related project, or other area of interest for which the report will be used.) _____

4. How specifically, is the report being used? (Information source, design data, procedure, source of ideas, etc.) _____

5. Has the information in this report led to any quantitative savings as far as man-hours or dollars saved, operating costs avoided or efficiencies achieved, etc? If so, please elaborate. _____

6. General Comments. What do you think should be changed to improve future reports? (Indicate changes to organization, technical content, format, etc.) _____

	Name

	Organization
CURRENT	_____
ADDRESS	Address

	City, State, Zip

7. If indicating a Change of Address or Address Correction, please provide the New or Correct Address in Block 6 above and the Old or Incorrect address below.

	Name

	Organization
OLD	_____
ADDRESS	Address

	City, State, Zip

(Remove this sheet along the perforation, fold as indicated, staple or tape closed, and mail.)

----- FOLD HERE -----

Director
U.S. Army Ballistic Research Laboratory
ATTN: SLCBR-DD-T
Aberdeen Proving Ground, MD 21005-5066




NO POSTAGE
NECESSARY
IF MAILED
IN THE
UNITED STATES

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300

BUSINESS REPLY MAIL
FIRST CLASS PERMIT NO 12062 WASHINGTON, DC
POSTAGE WILL BE PAID BY DEPARTMENT OF THE ARMY

Director
U.S. Army Ballistic Research Laboratory
ATTN: SLCBR-DD-T
Aberdeen Proving Ground, MD 21005-9989



----- FOLD HERE -----

END

DTIC

9-86