

	.0	4.5 50	2.8	2.5
		50 03	<u>3.2</u> 3.6	2.2
			4.0	2.0
				1.8
1.2	5 ∥	<u> </u>	4	1.6

MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS-1963-A

:

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

TECHNICAL REPORT No. 12

Laser-Linewidth Effects on the Photon-Phonon Conversion Rate at a Gas-Solid Interface

by

Henk F. Arnoldus and Thomas F. George

Prepared for Publication

in

Journal of the Optical Society of America B

Departments of Chemistry and Physics State University of New York at Buffalo Buffalo, New York 14260



043

14

8

86

0

July 1986

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

UTIC FILE COPY

	REPORT DOCUME	NTATION PAGE			
1. REPORT SECURITY CLASSIFICATION Unclassified		16. RESTRICTIVE MARKINGS			
28. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY	OF REPORT		
26. DECLASSIFICATION/DOWNGRADING SCHED	ULE	Approved for public	release; distribution unlimited		
4. PERFORMING ORGANIZATION REPORT NUM	BERIS	5. MONITORING ORGANIZATION	REPORT NUMBER(S)		
UBUFFAL0/DC/86/TR-	12				
6. NAME OF PERFORMING ORGANIZATION	66. OFFICE SYMBOL	78. NAME OF MONITORING ORG	NIZATION		
State University of New York	(if applicable)				
6c. ADDRESS (City, State and XIP Code)	1	7b. ADORESS (City, State and ZIP ('odei		
Fronczak Hall, Amherst Campus		Chemistry Program			
Buffalo, New York 14260		800 N. Quincy Street			
		Arlington, Virgin	a 22217		
8. NAME OF FUNDING/SPONSORING ORGANIZATION	85. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT	IDENTIFICATION NUMBER		
Office of Naval Research		Contract N00014-86-K-0043			
8c. ADDRESS (City, State and ZIP Code)		10. SOURCE OF FUNDING NOS.			
Chemistry Program		PROGRAM PROJECT ELEMENT NO. NO.	TASK WORK UNIT		
Arlington, Virginia 22217					
11. TITLE	Photon_Phonon (Conversion Pate at a G	s-Solid Interface		
12. PERSONAL AUTHORIS) Henk F A	rnoldus and Thom	las F. George			
134 TYPE OF REPORT		14 DATE OF REPORT /Yr We I	IS PAGE COUNT		
FROM	TO	July 1986	17		
16. SUPPLEMENTARY NOTATION					
Prepared for publicati	on in Journal of	the Optical Society	of America B		
17. COSATI CODES	18. SUBJECT TERMS (C	ontinue on reverse if necessary and ide THEEECTC MAC	ntify by block number) TED ENIINTINN		
FIELD GROOP SUB. GR.	PHOTON-PHONON	CONVERSION LON	G-TIME LIMIT		
	GAS-SOLID INTE	RFACE COH	RENCES		
19. ABSTRACT (Continue on reverse if necessary and	l identify by block number	1			
An atom, bounded to the s	urface of a crys	tal, absorbs photons	rom a strong incident		
laser field and subsequently e	mits phonons int	cothe solid. In this	paper we study the		
influence of the laser linewid	th on this photo	on-phonon conversion p	ocess. The appearance		
of coherences obscures the int	erpretation of 1	the equation of motion	in terms of transition		
rates and prohibits the deriva	tion of a master	is assumed to be reso	ant with a set of two		
levels only results from the	work done by the	assumed to be reso external field on th	transition dipole.		
In the steady state, the energy absorption from the radiation field equals the energy					
emission in the phonon field. It is shown that in this long-time limit the coherences					
can be expressed in terms of the populations, and hence the system is again described by					
a master equation. Specific aspects of the finite laser bandwidth are pointed out.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRAC	ст	21. ABSTRACT SECURITY CLASS	FICATION		
UNCLASSIFIED/UNLIMITED 🖾 SAME AS RPT. 🖾 DTIC USERS 🗍		Unclassified			
228. NAME OF RESPONSIBLE INDIVIDUAL		225. TELEPHONE NUMBER	220. OFFICE SYMBOL		
Dr. David L. Nelson		(202) 696-4410			

Journal of the Optical Society of America B

To appear in the feature issue entitled "Laser-Induced Molecular Physics at Surfaces"

LASER-LINEWIDTH EFFECTS ON THE PHOTON-PHONON

CONVERSION RATE AT A GAS-SOLID INTERFACE

Henk F. Arnoldus and Thomas F. George

Departments of Physics & Astronomy and Chemistry 239 Fronczak Hall State University of New York at Buffalo Buffalo, New York 14260

ABSTRACT

An atom, bounded to the surface of a crystal, absorbs photons from a strong incident laser field and subsequently emits phonons into the solid. In this paper we study the influence of the laser linewidth on this photon-phonon conversion process. The appearance of coherences obscures the interpretation of the equation of motion in terms of transition rates and prohibits the derivation of a master equation in the transient region. The absorbed energy from the driving field, which is assumed to be resonant with a set of two levels only, results from the work done by the external field on the transition dipole. In the steady state, the energy absorption from the radiation field equals the energy emission in the phonon field. It is shown that in this long-time limit the coherences can be expressed in terms of the populations and hence the system is again described by a masterlike equation. Specific aspects of the finite laser bandwidth are pointed out.

PACS: 71.36, 78.90

Dist. ib	Dist.ibution /				
A	Availability Codes				
Dist	Avail and/or Special				
A-1					

I. INTRODUCTION

Irradiation of dye-coated surfaces or adsorbed atoms on a crystal with strong infrared laser light amounts to photon-phonon conversion reactions. The internal vibrational modes of the molecule or the atom-surface bond absorbs photons from the radiation field (stimulated excitation), and the subsequent spontaneous decay (coupling to the phonon field of the crystal) is accompanied by a phonon emission into the solid. Additional processes like thermal or radiative desorption and spontaneous emission of radiation are neglibible in comparison with the thermal transitions in the adbond. In this fashion, the resonant coupling of vibrational modes of adsorbed species with the incident electromagnetic field provides an effective method for heating a crystal, which itself might be transparent for the impinging light. In this paper, we study the effect of the laser linewidth on the adbond-mediated energy transport from a laser field into the phonon field.

The interaction of an adsorbed atom in a vibrational bond with a harmonic crystal is commonly described by a Pauli-type master equation.¹⁻⁶ The distribution of the population over the vibrational levels is determined by the phonon absorption and emission processes, which occur at a rate $n_k(t)a_{kl}$ for a transition from the initial state $|k\rangle$ to the final state $|l\rangle$. Here the rate constants a_{kl} depend on the temperature of the crystal and the shape of the potential well in which the atom is bounded (apart from an overall k, l-independent constant). The a_{kl} 's gain in general contributions from single-phonon and multiphonon processes. The population $n_k(t)$ of level $|k\rangle$ follows from the master equation and the initial state at t=0, whereas thermal relaxation to a unique steady state is inherent to the structure of the master equation.

Incorporation of the presence of a weak driving laser-field is easily done in a perturbative way.⁷⁻¹² Stimulated absorption and emission rates are included in the master equation as additional transitions between optically-coupled levels. The energy absorption rate from the radiation field is then simply the difference between the number of absorptions and emissions per unit time, multiplied by the photon energy $\hbar\omega_{\rm r}$. This approach does not apply anymore for an intense laser field, since the appearance of multiphoton processes prohibits such a simple interpretation of radiative transitions.¹³⁻¹⁵ In two previous papers,^{16,17} we showed that transparent mechanisms can be recovered, provided that we diagonalize the laser-atom interaction first. This was accomplished by a transformation to the famous dressed-states representation.¹⁸ With respect to these joint atomic and laser states, including the interaction, the phonon relaxation processes were understood as transitions between dressed states. Then every transition $|k \rightarrow | l \rangle$, which occurs at a rate $n_k(t)a_{k,l}$ (the subscripts k, l now refer to dressed states), corresponds to a phonon absorption from or emission into the crystal, where the phonon energy equals the level separation of the states $|k\rangle$ and $|l\rangle$. Summation over all transitions then yields the net energy flux into the crystal.

The derivation of these energy-transport equations hinges on the existence of a master equation for the level populations, both in the lowintensity limit and in the dressed-state picture. This enables us to identify the occurring transitions, the rate constants and the energies of the involved phonons and photons. A simple balance then gives the expression for the energy transport from the laser into the atomic bond, and finally into the crystal. It is not so obvious that this procedure can be employed in general. This is due to the fact that the equations for the

populations might couple to the time evolution of the coherences, which obscures any direct interpretation. We shall show that this problem already emerges if a finite laser linewidth is taken into account. A more sophisticated approach to the energy-transport problem, which does not rely on a specific structure of the equation of motion, will be used to tackle this apparent complication.

II. DRIVEN ADBOND

An atom is bounded to the surface of a harmonic-lattice crystal by attractive electromagnetic forces. If we denote the non-degenerate vibrational eigenstates by $|k\rangle$, we can write the Hamiltonian of the adbond as

$$H_{a} = \sum_{k} \hbar \omega_{k} P_{k}$$
(2.1)

in terms of the energy eigenvalues M_{k} and the projectors $P_{k} = |k\rangle \langle k|$ onto the eigenstates. This system is irradiated by a laser field, with electric component

$$E(t) = E_0 Re \varepsilon_L e$$
(2.2)

at the position of the atom. Here \mathbf{E}_{o} is the real amplitude, $\underline{\mathbf{\varepsilon}}_{L}$ is the normalized polarization vector ($\underline{\mathbf{\varepsilon}}_{L} * \cdot \underline{\mathbf{\varepsilon}}_{L} = 1$) and $\phi(t)$ is a stochastically fluctuating phase, which broadens the laser line around its central frequency $\mathbf{\omega}_{L}$. The coupling to the atomic bond is established by a dipole interaction $\mathbf{H}_{ar}(t) = -\underline{\mu} \cdot \underline{\mathbf{E}}(t)$, with $\underline{\mu}$ the transition dipole-moment operator between the eigenstates $|\mathbf{k}\rangle$. Phonon absorption/emission is considered as a relaxation process, and hence it is most conveniently accounted for by a Liouville operator Γ , which acts on the density operator ρ of the adbond according to 19

$$\Gamma \rho = \frac{1}{2} \int_{k\ell} a_{k\ell} (P_k \rho + \rho P_k - 2P_\ell \langle k | \rho | k \rangle). \qquad (2.3)$$

Every term k, ℓ pertains to a transition $|k\rangle \rightarrow |\ell\rangle$, and $a_{k\ell}$ is the inverse lifetime of level $|k\rangle$, due to its spontaneous decay $(\omega_k > \omega_\ell)$ or excitation $(\omega_k < \omega_\ell)$ to level $|\ell\rangle$. Then the laser-driven and thermally-damped atomic bond is represented by the density operator $\rho(t)$, whose time evolution is governed by the full equation of motion

$$i\hbar \frac{d\rho}{dt} = [H_{a},\rho] + [H_{ar}(t),\rho] - i\hbar \Gamma \rho, \rho^{\dagger} = \rho, Tr \rho = 1.$$
 (2.4)

The laser field $\underline{E}(t)$ is a stochastic process, due to its randomly fluctuating phase, and this turns Eq. (2.4) into a stochastic differential equation for $\rho(t)$.

We will assume that only two levels $|e\rangle$ (excited) and $|g\rangle$ (ground) of the bond are significantly coupled by the incident radiation. The detuning between the optical frequency $\omega_{\rm L}$ and the level separation $\omega_{\rm O} = \omega_{\rm e} - \omega_{\rm g} > 0$ of the driven transition will be indicated by

and the coupling strength is expressed in the Rabi frequency

$$\Omega = \mathbf{E}_{\mathbf{A}} |\langle \mathbf{e} | \boldsymbol{\mu} \cdot \boldsymbol{\epsilon}_{\mathbf{L}} | \mathbf{g} \rangle | / \mathbf{h}.$$
(2.6)

Then the interaction Hamiltonian with the radiation field attains the form

$$-i(\omega_{L}t + \phi(t))$$

$$H_{ar}(t) = -\frac{1}{2}\hbar\Omega |e\rangle < g|e + Hermitian conjugate. (2.7)$$

III. TRANSFORMATION AND AVERAGE

The interaction Hamiltonian $H_{ar}(t)$ oscillates with frequency ω_{L} and is stochastic through the phase $\phi(t)$. With a stochastic transformation to the

rotating frame, we can eliminate the $exp(\pm i\omega_L t)$ -dependence and facilitate the appearance of $\phi(t)$.^{20,21} We define the density operator $\sigma(t)$ as

which has the equation of motion

$$i\frac{d\sigma}{dt} = \hbar^{-1}[H_d,\sigma] + \dot{\phi}(t)[P_g,\sigma] - i\Gamma\sigma, \qquad (3.2)$$

where the time-independent dressed-atom Hamiltonian is explicitly

$$H_{d} = \sum_{i \neq e,g} \hbar \omega_{i} P_{i} + \frac{1}{2} \hbar (\omega_{e} + \omega_{g} + \omega_{L}) (P_{e} + P_{g}) - \frac{1}{2} \hbar \Delta (P_{e} - P_{g})$$

$$i \neq e,g$$

$$- \frac{1}{2} \hbar \Omega (|e > \langle g| + |g > \langle e|). \qquad (3.3)$$

Now the time derivative $\dot{\phi}(t)$ of the phase enters the equation, and in a multiplicative way (e.g., as $\dot{\phi}(t)\sigma(t)$). Notice that no information is lost in the transformation, since Eq. (3.1) can be inverted.

The solution of the equation of motion (3.2) for $\sigma(t)$ refers to a single realization of the process $\phi(t)$, but only the average over the stochastics of the phase have a significance. In general, this average would depend on the details of the stochastics of $\phi(t)$, but it can be shown²² that for Eq. (3.2) the average is quite insensitive to the precise specification of the process, as long as the phase fluctuations represent the broadening of a single-mode laser line. For any Markovian diffusion process $\phi(t)$ and for the uniform random-jump process, the average of Eq. (3.2) acquires the simple form

$$i \frac{d\Pi}{dt} = \chi^{-1}[H_d,\Pi] - i \Pi, \qquad (3.4)$$

where we have introduced the abbreviation

$$\Pi(t) = \{\sigma(t)\}, \tag{3.5}$$

with the slashed angle brackets denoting a stochastic average (in order to distinguish from a quantum-mechanical average <...>). The operator W, which accounts for the laser linewidth, equals

$$W\Pi = \lambda[P_{\rho}, [P_{\rho}, \Pi]], \qquad (3.6)$$

where λ is the half-width at half-maximum of the Lorentzian laser line. Alternatively, we can write W as

$$W\Pi = \lambda (P_{g}\Pi + \Pi P_{g} - 2P_{g} \langle g | \Pi | g \rangle), \qquad (3.7)$$

which is reminiscent of the structure of the phonon-relaxation operator Γ from Eq. (2.3). Just as a_{kl} can be considered as a contribution to the width of level $|k\rangle$, we can interpret 2λ as the additional width of the ground state, due to phase fluctuations. This can also be inferred from Eq. (2.2), if we rewrite it as

$$E(t) = E_0 \operatorname{Re}_{L} e^{0} \qquad (3.8)$$

The laser frequency $\omega_{\rm L}$ can be regarded to be shifted instantaneously by the amount $\dot{\phi}(t)$, which diminishes or increases the detuning $\Delta = \omega_{\rm L} - \omega_{\rm O}$ by $\dot{\phi}(t)$. Effectively, this gives rise to a width 2λ .

IV. MATRIX ELEMENTS

Equation (3.4) is an operator equation for the density matrix $\Pi(t)$ of the driven adbond. This determines the populations, abbreviated as

$$\Pi_{L}(t) = \langle k | \Pi(t) | k \rangle, \tag{4.1}$$

and the coherences $\langle k | \Pi(t) | l \rangle$, $k \neq l$, with respect to the eigenstates $| k \rangle$ of the unperturbed bond. After taking the diagonal part of Eq. (3.4), we find for the time evolution of the populations

$$\frac{d}{dt} \Pi_{\varrho} = -A_{\varrho} \Pi_{\varrho} + \sum_{k} \Pi_{k} a_{k\varrho}, \quad \varrho \neq e,g, \qquad (4.2)$$

$$\frac{d}{dt} \Pi_{e} = \Omega \operatorname{Im} \langle e | \Pi | g \rangle - A_{e} \Pi_{e} + \sum_{k} \Pi_{k} a_{ke}, \qquad (4.3)$$

$$\frac{d}{dt} \Pi_{g} = -\Omega \operatorname{Im} \langle e | \Pi | g \rangle - A_{g} \Pi_{g} + \sum_{k} \Pi_{k} a_{kg}, \qquad (4.4)$$

where we have introduced the total-decay constant of level $|k\rangle$ by

$$\mathbf{A}_{\mathbf{k}} = \sum_{\boldsymbol{\ell}} \mathbf{a}_{\mathbf{k}\boldsymbol{\ell}}, \qquad (4.5)$$

and we have used the Hermiticity of II(t). The set (4.2) - (4.4) resembles a master equation, but it contains the inhomogeneous term $\Omega Im \langle e | I | g \rangle$, and hence it is not a closed set. The equation for the coherence between the excited state and the ground state is readily found to be

$$i\frac{d}{dt} < e|\Pi|g\rangle = \frac{1}{2}\Omega(\Pi_e - \Pi_g)$$

- $(\Delta + \frac{1}{2}i(A_e + A_g + 2\lambda)) < e|\Pi|g\rangle.$ (4.6)

Eqs. (4.2) - (4.4) and (4.6), accompanied by the normalization

$$\sum_{k} \Pi_{k}(t) = 1,$$
 (4.7)

which is $Tr\Pi(t) = 1$, determine the populations $\Pi_k(t)$ and the coherence $\langle e | \Pi(t) | g \rangle$. This generalizes the master equation for the case of an arbitrarily strong incident field with a finite bandwidth. These are simple linear first-order differential equations, and can be solved immediately for any configuration of levels. Notice that the laser linewidth λ only enters in the last term on the right-hand side of Eq. (4.6), and that the coupling of Eq. (4.6) with the set (4.2) - (4.4) is brought about by the inversion $\Pi_e - \Pi_g$, rather than by the separate populations.

V. ABSORPTION RATE

Since the coherence $\langle e | \Pi | g \rangle$ enters the equations for the populations Π_k , it is not so obvious which transitions occur, and at which rates. Therefore, we have to start from more elementary principles. The absorbed energy by the atomic bond from the external field equals the work that is done on the dipole by the field. Hence, we can write for the absorption rate (energy per unit time)²³

$$\frac{dZ}{dt} = \underbrace{E}(t) \cdot \frac{d}{dt} \langle \mu(t) \rangle, \qquad (5.1)$$

where $\mu(t)$ is the dipole moment of the system in the Heisenberg picture. This expression accounts for absorptions and stimulated emissions of photons from and into the field, and the balance between these gain and loss terms is the net energy-absorption rate dZ/dt from the radiation field into the adbond. In the steady state, there can be no accumulation of energy in the bond, so then dZ/dt equals the balance between the emission and absorption rates of phonon energy into the crystal. Effectively, this is a conversion of radiation energy into the thermal energy, or, a photon-phonon conversion reaction, mediated by the optically-active atomic bond.

Transformation of Eq. (5.1) to the Schrödinger picture gives

$$\frac{dZ}{dt} = Tr(\underline{E}(t) \cdot \underline{\mu}) \frac{d\rho}{dt}, \qquad (5.2)$$

and with the equation of motion (2.4) for $\rho(t)$, the stochastic transformation (3.1) to the σ -representation and with $\sigma^{\dagger} = \sigma$, we obtain

$$\frac{dZ}{dt} = \hbar \Omega \frac{d}{dt} \operatorname{Re} \langle \mathbf{e} | \sigma(t) | \mathbf{g} \rangle + \hbar \Omega (\omega_{L} + \dot{\phi}(t)) \operatorname{Im} \langle \mathbf{e} | \sigma(t) | \mathbf{g} \rangle.$$
(5.3)

This relation reveals that the absorption rate is basically determined by the coherence $\langle \mathbf{e} | \sigma | \mathbf{g} \rangle$, rather than by the populations of the levels. From a slightly different point of view, we can interpret this as the fact that the

occurrence of coherent transitions between states is reflected in the presence of off-diagonal matrix elements. This is in contrast to the incoherent thermal decay, which is goverened by a master equation for the populations only.

The appearance of $\dot{\phi}(t)$ in the second term on the right-hand side of Eq. (5.3) is not very convenient. With the aid of the equation of motion (3.2) for $\sigma(t)$, we can eliminate the $\dot{\phi}(t)$ -term and the time derivative in the first term simultaneously. We find

$$\frac{dZ}{dt} = -\frac{1}{2}\hbar\Omega(A_e + A_g) \operatorname{Re}\langle e | \sigma | g \rangle + \hbar\Omega\omega_o \operatorname{Im}\langle e | \sigma | g \rangle.$$
(5.4)

Comparison with Eq. (5.3) shows that the time differentiation of the real part of the coherence is replaced by $-\frac{1}{2}(A_e + A_g)$, and the instantaneous laser frequency $\omega_L + \dot{\phi}(t)$, which multiplies the imaginary part, is altered into the atomic resonance frequency ω_0 . This displays that the contribution to dZ/dt from the real part originates from the cooperation of thermal decay and coherent excitation, whereas the imaginary part is pure optical, since it persists even in the absence of the crystal. The stochastic average of dZ/dt is now easily found. We simply replace $\sigma(t)$ by $\Pi(t)$ in Eq. (5.4).

VI. STEADY STATE

Due to the thermal relaxation, the density matrix $\Pi(t)$ will reach a steady state after a time of the order of a_{kl}^{-1} , elapsed from the instant of preparation (for instance, the switch-on of the laser). The long-time solution will be indicated by

$$\bar{\Pi} = \lim_{t \to \infty} \Pi(t).$$
(6.1)

This $\overline{\Pi}$ then obeys the set of equations (4.2) - (4.4) and (4.6) with the left-hand sides set equal to zero. Taking the imaginary part of Eq. (4.6),

we obtain

$$\operatorname{Re} \langle \mathbf{e} | \overline{\mathbf{\Pi}} | \mathbf{g} \rangle = \frac{-\Delta}{\frac{1}{2} (\mathbf{A}_{\mathbf{e}} + \mathbf{A}_{\mathbf{g}}) + \lambda} \operatorname{Im} \langle \mathbf{e} | \overline{\mathbf{\Pi}} | \lambda \rangle.$$
(6.2)

This relation between the real and imaginary parts of the coherence holds only in the steady state. Subsequently, we consider the real part of Eq. (4.6). Application of Eq. (6.2) then yields

$$\operatorname{Im} \langle \mathbf{e} | \overline{\mathbf{n}} | \mathbf{g} \rangle = \frac{1}{2} \Omega \frac{\frac{1}{2} (\mathbf{A}_{\mathbf{e}} + \mathbf{A}_{\mathbf{g}}) + \lambda}{\left[\frac{1}{2} (\mathbf{A}_{\mathbf{e}} + \mathbf{A}_{\mathbf{g}}) + \lambda\right]^{2} + \Delta^{2}} (\overline{\mathbf{n}}_{\mathbf{g}} - \overline{\mathbf{n}}_{\mathbf{e}}).$$
(6.3)

The results (6.2) and (6.3) show that the coherence can be expressed entirely in terms of the population inversion $\overline{\Pi}_{e} - \overline{\Pi}_{g}$ in this limit.

With (6.2) and (6.3) we can rewrite Eqs. (4.2) - (4.4) in the steady state as

$$\sum_{\mathbf{k}} \overline{\Pi}_{\mathbf{k}} \mathbf{a}_{\mathbf{k}} \mathbf{f} = \mathbf{A}_{\mathbf{k}} \overline{\Pi}_{\mathbf{k}}, \quad \mathbf{k} \neq \mathbf{e}, \mathbf{g}, \quad (6.4)$$

$$\sum_{\mathbf{k}} \overline{\Pi}_{\mathbf{k}} (\mathbf{a}_{\mathbf{k}\mathbf{e}} + \mathbf{a}_{\mathbf{k}\mathbf{g}}) = \mathbf{A}_{\mathbf{e}} \overline{\Pi}_{\mathbf{e}} + \mathbf{A}_{\mathbf{g}} \overline{\Pi}_{\mathbf{g}}, \qquad (6.5)$$

$$\sum_{k} \overline{\Pi}_{k} (a_{ke} - a_{kg}) = A_{e} \overline{\Pi}_{e} - A_{g} \overline{\Pi}_{g}$$

$$- \Omega^{2} \frac{\frac{1}{2}(A_{e} + A_{g}) + \lambda}{\left[\frac{1}{2}(A_{e} + A_{g}) + \lambda\right]^{2} + \Delta^{2}} (\bar{\Pi}_{g} - \bar{\Pi}_{e}), \qquad (6.6)$$

where the coherence $\langle e | \bar{\Pi} | g \rangle$ has disappeared. This set of equations only contains populations, and it can be regarded as a master-like equation for a coherently-driven system. Notice that the optical parameter Ω only enters via Ω^2 , which is proportional to the laser intensity. The coherence (6.2), (6.3), however, also depends on the field amplitude Ω .

With the use of Eq. (6.2), the stochastic average of the absorption rate, Eq. (5.4), can be written as

$$\{\frac{dZ}{dt}\} = \frac{1}{4} \{\omega_{L} - \Delta \frac{2\lambda}{A_{e} + A_{g} + 2\lambda}\} \Omega Im \langle e|\bar{I}|g \rangle.$$
(6.7)

If the laser linewidth λ is small in comparison with the relaxation constant $A_e + A_g$, the second term in curly brackets vanishes, and the absorption rate reduces to the $M_{u_L}\Omega Im \langle e | \overline{\Pi} | g \rangle$. Since M_{u_L} is the energy of a laser photon, this implies that $\Omega Im \langle e | \overline{\Pi} | g \rangle$ equals the number of photons per unit time, which is converted into thermal energy. For $\lambda \gg A_e + A_g$, the term in curly brackets becomes $\omega_L - \Delta = \omega_0$. This is consistent with the fact that for a large bandwidth the energy of a photon is no longer well-defined. Now the radiation excites the system from $|g\rangle$ to $|e\rangle$, and subsequent thermal decay corresponds to an effective gain of phonon energy M_{ω_0} . This process occurs again at a rate $\Omega Im \langle e | \overline{\Pi} | g \rangle$.

Substitution of Eq. (6.3) into Eq. (6.7) finally yields

$$\left\{\frac{dZ}{dt}\right\} = \Re\left\{\omega_{L} - \Delta \frac{2\lambda}{A_{e} + A_{g} + 2\lambda}\right\}$$

$$x \frac{1}{2}\Omega^{2} \frac{\frac{1}{2}(A_{e} + A_{g}) + \lambda}{\left[\frac{1}{2}(A_{e} + A_{g}) + \lambda\right]^{2} + \Delta^{2}} (\bar{\pi}_{g} - \bar{\pi}_{e}). \qquad (6.8)$$

The master-like equation (6.4) - (6.6), together with the normalization (4.7), constitutes a simple set of linear algebraic equations, which is easily solved for a particular case. This determines $\bar{\Pi}_{g} - \bar{\Pi}_{e}$, and thereby the energy-absorption rate (6.8).

For $\lambda << A_e + A_g$, the laser linewidth has no significance at all, since it disappears from Eqs. (6.4) - (6.6) and Eq. (6.8). If we then take $|\Delta| >>$ $A_e + A_g$, we recover our previous results.^{16,17} In the situation $\lambda >> A_e + A_g$, the energy-absorption rate reduces to

$$\left\{\frac{dZ}{dt}\right\} = \hbar\omega_{0} \cdot \frac{1}{2}\Omega^{2} \frac{\lambda}{\lambda^{2} + \Delta^{2}} \left(\bar{\pi}_{g} - \bar{\pi}_{e}\right).$$
(6.9)

If the system is driven close to resonance (Δ^{-0}), the prefactor is proportional to ω_0/λ , and hence the absorption is diminished by an increasing linewidth. This can be understood from the representation (3.8) of the laser field. The probability for a photon absorption is largest if $\omega_L = \omega_0$, but due to $\dot{\phi}(t) \neq 0$, the optical frequency is temporarily shifted out of resonance. This reduces the absorption. Conversely, for $|\Delta|$ large, the phase fluctuations can shift ω_L into resonance, and thereby enhance the absorption. From Eq. (6.9) we see that this is indeed the case, since $\langle dZ/dt \rangle$ becomes proportional to $\omega_0 \lambda/\Delta^2$.

VI. CONCLUSIONS

We have considered the irradiation of an atom, bounded to a crystal, by intense non-monochromatic laser light. The single-mode laser line is broadened by stochastic phase fluctuations, which turns the equation of motion for the density operator $\sigma(t)$ of the atomic bond into a stochastic differential equation. The average over the stochastics of $\phi(t)$ was performed in a rotating-frame representation, yielding the equation of motion (3.4) for $\Pi(t) = \{\sigma(t)\}$. It appears that the laser linewidth gives rise to an additional width 2λ of the ground state. This is a consequence of the specific form of our transformation (3.1) to the rotating frame. We remark that other kinds of transformations²⁴, which also eliminate the $\exp(\pm i\omega_{L}t)$ time dependence in the interaction Hamiltonian, would result in effective widths of both $|e\rangle$ and $|g\rangle$.

The combination of laser-linewidth damping and thermal relaxation with coherent excitation by an arbitrarily strong radiation field prohibits the derivation of a master equation for the populations of the vibrational levels. It becomes inevitable to take the coherences between the opticallycoupled states into consideration. This gives the set of equations (4.2) -

(4.4) and (4.6), which can be regarded as a generalized master equation. Only in the long-time limit we can express the coherence $\langle e | \overline{\Pi} | | g \rangle$ in the population-difference $\overline{\Pi}_{g} - \overline{\Pi}_{e}$, which implies the emergence of a genuine master equation.

The energy absorption from the laser field by the adbond is evaluated by calculating the work done on the dipole moment of the bond. In general, the absorption rate is determined by the coherence $\langle e | \Pi(t) | g \rangle$ (Eq. (5.4)), and in the steady state it can be expressed in the inversion $\overline{\Pi}_e - \overline{\Pi}_g$, which follows from the solution of the master equation. The transport of energy from the bond to the phonon field is brought about by thermal relaxation. This can be inferred directly from Eqs. (6.6) and (6.8). For $a_{kl} = 0$ for all k, t, we find $\overline{\Pi}_e = \overline{\Pi}_g$ in the steady state, and hence $\{dZ/dt\} = 0$. We have shown how the overall factor changes from $M\omega_L$ to $M\omega_O$ with increasing linewidth, and we see that a finite λ enhances (diminishes) the energy flux if the detuning is large (small).

ACKNOWLEDGEMENTS

We wish to thank Professor H. Metiu for reading this manuscript and making a number of helpful suggestions. This research was supported by the Office of Naval Research, the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009, and the National Science Foundation under Grant CHE-8519053. The United States Government is authorized to reproduce and distribute reprints notwithstanding any copyright notation hereon.

REFERENCES

1.	S. Efrima, K.F. Freed, C. Jedrzejek and H. Metiu, Chem. Phys. Lett. 74, 43 (1980).
2.	Z.W. Gortel, H.J. Kreuzer and R. Teshima, Phys. Rev. B 22, 5655 (1980).
3.	Z.W. Gortel, H.J. Kreuzer, R. Teshima and L.A. Turski, Phys. Rev. B 24, 4456 (1981).
4.	C. Jedrzejek, K.F. Freed, S. Efrima and H. Metiu, Chem. Phys. Lett. <u>79</u> , 227 (1981) and Surf. Sci. <u>109</u> , 191 (1981).
5.	E. Goldys, Z.W. Gortel and H.J. Kreuzer, Surf. Sci. <u>116</u> , 33 (1982).
6.	S. Efrima, C. Jedrzejek, K.F. Freed, E. Hood and H. Metiu, J. Chem. Phys. <u>79</u> , 2436 (1983).
7.	J. Lin and T.F. George, Surf. Sci. <u>100</u> , 381 (1980).
8.	J. Lin and T.F. George, J. Phys. Chem. <u>84</u> , 2957 (1980).
9.	G. Korzeniewski, E. Hood and H. Metiu, J. Vac. Sci. Technol. <u>20</u> , 594 (1982).
10.	J. Lin, X.Y. Huang and T.F. George, Z. Phys. B <u>48</u> , 355 (1982).
11.	X.Y. Huang, T.F. George and J.M. Yuan, J. Opt. Soc. Am. B <u>2</u> , 985 (1985).
12.	Z.W. Gortel, P. Piercy, R. Teshima and H.J. Kreuzer, Surf. Sci. <u>165</u> , L12 (1986).
13.	M.S. Slutsky and T.F. George, Chem. Phys. Lett. 57, 474 (1978).
14.	2.W. Gortel, H.J. Kreuzer, P. Piercy and R. Teshima, Phys. Rev. B <u>27</u> , 5066 (1983).
15.	A.C. Beri and T.F. George, J. Vac. Sci. Technol. B 3, 1529 (1985).
16.	H.F. Arnoldus, S. van Smaalen and T.F. George, Phys. Rev. B (1986), in press.
17.	S. van Smaalen, H.F. Arnoldus and T.F. George, Phys. Rev. B (1986), submitted.
18.	C. Cohen-Tannoudji, in <u>Frontiers in Laser Spectroscopy</u> , Proc. 27th Les Houches Summer School, edited by R. Balian, S. Haroche and S. Liberman (North-Holland, Amsterdam, 1977), p.3 ff.
19.	W.H. Louisell, <u>Quantum Statistical Properties of Radiation</u> (Wiley, New York, 1973), Ch. 6.
20.	G.S. Agarwal, Phys. Rev. A <u>18</u> , 1490 (1978).

- 21. H.F. Arnoldus and G. Nienhuis, J. Phys. B: At. Mol. Phys. <u>16</u>, 2325 (1983).
- 22. H.F. Arnoldus and G. Nienhuis, J. Phys. B: At. Mol. Phys. <u>19</u>, 873 (1986).
- 23. L. Allen and J.H. Eberly, Optical Resonance and Two-Level Atoms (Wiley, New York, 1975), p. 142.
- 24. G. Nienhuis, Acta Phys. Pol. A <u>61</u>, 235 (1982).

DL/413/83/01 GEN/413-2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

10053563

いたちちちちの

	No. Copies		No. Copi
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 5042 Crane, Indiana 47522	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 2770	1 9
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 1911	1 .2
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

0L/413/83/01 056/413-2

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai Department of Chemistry University of California Berkeley, California 94720

Dr. J. Murday Naval Research Laboratory Surface Chemistry Division (6170) 455 Overlook Avenue, S.W. Washington, D.C. 20375

Dr. J. B. Hudson Materials Division Rensselaer Polytechnic Institute Troy, New York 12181

Dr. Theodore E. Madey Surface Chemistry Section Department of Commerce National Bureau of Standards Washington, D.C. 20234

Dr. J. E. Demuth IBM Corporation Thomas J. Watson Research Center P.O. Box 218 Yorktown Heights, New York 10598

Dr. M. G. Lagally Department of Metallurgical and Mining Engineering University of Wisconsin Madison, Wisconsin 53706

Dr. R. P. Van Duyne Chemistry Department Northwestern University Evanston, Illinois 60637

Dr. J. M. White Department of Chemistry University of Texas Austin, Texas 78712

Dr. D. E. Harrison Department of Physics Naval Postgraduate School Monterey, California 93940 Dr. W. Kohn Department of Physics University of California, San Diego La Jolla, California 92037

Dr. R. L. Park Director, Center of Materials Research University of Maryland College Park, Maryland 20742

Dr. W. T. Peria Electrical Engineering Department University of Minnesota Minneapolis, Minnesota 55455

Dr. Keith H. Johnson Department of Metallurgy and Materials Science Massachusetts Institute of Technology Cambridge, Massachusetts 02139

Dr. S. Sibener Department of Chemistry James Franck Institute 5640 Ellis Avenue Chicago, Illinois 60637

Dr. Arnold Green Quantum Surface Dynamics Branch Code 3817 Naval Weapons Center China Lake, California 93555

Dr. A. Wold Department of Chemistry Brown University Providence, Rhode Island 02912

Dr. S. L. Bernasek Department of Chemistry Princeton University Princeton, New Jersey 08544

Dr. P. Lund Department of Chemistry Howard University Washington, D.C. 20059

0L/413/83/01 056/413-2

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter Code 6132 Naval Research Laboratory Washington, D.C. 20375

Dr. Richard Colton Code 6112 Naval Research Laboratory Washington, D.C. 20375

Dr. Dan Pierce National Bureau of Standards Optical Physics Division Washington, D.C. 20234

Dr. R. Stanley Williams Department of Chemistry University of California Los Angeles, California 90024

Dr. R. P. Messmer Materials Characterization Lab. General Electric Company Schenectady, New York 22217

Dr. Robert Gomer Department of Chemistry James Franck Institute 5640 Ellis Avenue Chicago, Illinois 60637

Dr. Ronald Lee R301 Naval Surface Weapons Center White Oak Silver Spring, Maryland 20910

Dr. Paul Schoen Code 5570 Naval Research Laboratory Washington, D.C. 20375

Dr. John T. Yates Department of Chemistry University of Pittsburgh Pittsburgh, Pennsylvania 15260 Dr. Richard Greene Code 5230 Naval Research Laboratory Washington, D.C. 20375

Dr. L. Kesmodel Department of Physics Indiana University Bloomington, Indiana 47403

Dr. K. C. Janda California Institute of Technology Division of Chemistry and Chemical Engineering Pasadena, California 91125

Dr. E. A. Irene Department of Chemistry University of North Carolina Chapel Hill, Northc Carolina 27514

Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974

Dr. Martin Fleischmann Department of Chemistry Southampton University Southampton **S**09 5NH Hampshire, England

Dr. John W. Wilkins Cornell University Laboratory of Atomic and Solid State Physics Ithaca, New York 14853

Dr. Richard Smardzewski Code 6130 Naval Research Laboratory Washington, D.C. 20375

Dr. H. Tachikawa Chemistry Department Jackson State University Jackson, Mississippi 39217

DL/413/83/01 056/413-2

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis Department of Physics University of California Irvine, California 92664

22,22

Dr. D. Ramaker Chemistry Department George Washington University Washington, D.C. 20052

Dr. J. C. Hemminger Chemistry Department University of California Irvine, California 92717

Dr. T. F. George Chemistry Department University of Rochester Rochester, New York 14627

Dr. G. Rubloff IBM Thomas J. Watson Research Center P.O. Box 218 Yorktown Heights, New York 10598

Dr. Horia Metiu Chemistry Department University of California Santa Barbara, California 93106

Dr. W. Goddard Division of Chemistry California Institute of Technology Pasadena, California 91125

Dr. J. T. Keiser Department of Chemistry University of Richmond Richmond, Virginia 23173 Dr. R. W. Plummer Department of Physics University of Pennsylvania Philadelphia, Pennsylvania 19104

Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 41106

Dr. N. Winograd Department of Chemistry Pennsylvania State University University Park, Pennsylvania 16802

Dr. Roald Hoffmann Department of Chemistry Cornell University Ithaca, New York 14853

Dr. A. Steckl Department of Electrical and Systems Engineering Rensselaer Polytechnic Institute Troy, NewYork 12181

Dr. G. H. Morrison Department of Chemistry Cornell University Ithaca, New York 14853

Dr. P. Hansma Physics Department University of California Santa Barbara, California 93106

Dr. J. Baldeschwieler California Institute of Technology Division of Chemistry Pasadena, California 91125

0L/413/83/01 056/413-2

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen Hughes Research Laboratory 3011 Malibu Canyon Road Malibu, California 90265

Dr. J. H. Weaver Department of Chemical Engineering and Materials Science University of Minnesota Minneapolis, Minnesota 55455

Dr. W. Goddard Division of Chemistry California Institute of Technology Pasadena, California 91125

Dr. A. Reisman Microelectronics Center of North Carolina Research Triangle Park, North Carolina 27709

Dr. M. Grunze Laboratory for Surface Science and Technology University of Maine Orono, Maine 04469

Dr. J. Butler Naval Research Laboratory Code 6115 Washington D.C. 20375

Dr. L. Interante Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181

Dr. Irvin Heard Chemistry and Physics Department Lincoln University Lincoln University, Pennsylvania 19352

Dr. K.J. Klaubunde Department of Chemistry Kansas State University Manhattan, Kansas 66506 Dr. W. Knauer Hughes Research Laboratory 3011 Malibu Canyon Road Malibu, California 90265

Dr. C. B. Harris Department of Chemistry University of California Berkeley, California 94720

Dr. F. Kutzler Department of Chemistry Box 5055 Tennessee Technological University Cookesville, Tennessee 38501

Dr. D. DiLella Chemistry Department George Washington University Washington D.C. 20052

Dr. R. Reeves Chemistry Department Renssaeler Polytechnic Institute Troy, New York 12181

