

AD-A171 528

AN UPPER LIMIT ON SHEPHERDING SATELLITES AT SATURN'S

1/1


RING G(U) IOWA UNIV IOWA CITY DEPT OF PHYSICS AND
ASTRONOMY J A VAN ALLEN JUL 86 U. OF IOWA-86-30


UNCLASSIFIED

N00014-86-K-0404

F/G 3/2

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A



AD-A171 520

An Upper Limit on Shepherding Satellites
at Saturn's Ring G

by

JAMES A. VAN ALLEN



FILE COPY

Department of Physics and Astronomy
THE UNIVERSITY OF IOWA

Iowa City, Iowa 52242

86 9 02 140

①①

U. of Iowa 86-30

An Upper Limit on Shepherding Satellites
at Saturn's Ring G

by

JAMES A. VAN ALLEN

Department of Physics and Astronomy
The University of Iowa
Iowa City, Iowa 52242

PHYSICS
LIBRARY
JUL 10 1986

July 1986

1986
digital version is available

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER U. of Iowa 86-30	2. GOVT ACCESSION NO. AD-A171520	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) An Upper Limit on Shepherding Satellites at Saturn's Ring G		5. TYPE OF REPORT & PERIOD COVERED Progress, July 1986
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) JAMES A. VAN ALLEN		8. CONTRACT OR GRANT NUMBER(s) N00014-86-K-0404
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Physics and Astronomy The University of Iowa Iowa City, Iowa 52242		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Electronics Program Office Arlington, VA 22217		12. REPORT DATE July 1986
		13. NUMBER OF PAGES 33
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Submitted for publication in <u>J. Geophys. Res.</u>		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Saturn Magnetosphere Saturn's Ring G		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) See page following.		

DD FORM 1473
1 JAN 73EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ABSTRACT

A simplified analytical theory of the absorption of energetic magnetospheric particles by an inert satellite is developed for the case in which the radius b of the satellite is much less than the equatorial gyroradius r_g of the particle which, in turn, is very much less than the radius r of the satellite's orbit. This theory and the observed particle-absorption signatures of Saturn's Ring G are used to establish an upper limit on shepherding satellites associated with the ring. The resulting upper limit, ignoring the absorption of the optically observed particulate matter, is $b = 3.3$ km (accurate to a factor of two) for a single satellite or $\sum_{i=1}^n b_i = 11.2 \text{ km}^2$ (accurate to a factor of four) for an assemblage of n satellites of various radii b_i . No shepherding satellites at Ring G have been detected optically.

case of a ring system

Accession For	
DTIC GRA&I	<input checked="" type="checkbox"/>
DMC TAB	<input type="checkbox"/>
Unpublished	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist _____	
A-1	



INTRODUCTION

During Pioneer 11's passage through the inner radiation belts of Saturn on 1 September 1979, a number of distinctive absorption features were observed in curves of the radial dependence of the intensity of energetic, trapped particles [Van Allen et al., 1980a, 1980b; Simpson et al., 1980a, 1980b; Fillius et al., 1980; Fillius and McIlwain, 1980; Trainor et al., 1980; and McDonald et al., 1980]. One such feature was a well defined inflection (not a dip) in curves of the intensity of protons ($E_p > 80$ MeV) vs. radial distance. In the vicinity of this feature the inbound and outbound curves (at a time lapse of nearly four hours and at quite different longitudes) were virtually identical, suggesting that the feature was a time-stationary one. Van Allen et al. [1980b] designated this absorption feature 1979 S3 and labeled it as a "suspected [satellite] but interpretation of [absorption] signature ambiguous". The relevant ambiguity stems from the failure of such a limited set of data to distinguish the absorption macro-signature of one or more small satellites from that of a continuous circular ring of particulate matter [Van Allen,

1982]. The ambiguity was resolved to some extent in November 1980 by Voyager 1 images [Smith et al., 1981], which revealed a thin continuous ring of radius about 170,000 km, normal optical opacity $\approx 3 \times 10^{-5}$, and radial width about 500 km (after estimated correction for imaging "smear"). The Voyager investigators designated it Ring G. A detailed interpretation of the energetic particle data associated with Ring G has been reported [Van Allen, 1983]. The mean radius of Ring G was found to be $170,200 \pm 900$ km, in agreement with the optical value. Other conclusions of this work, which ignored the possibility of ring-associated satellites, were that "the particulates in Ring G have an effective radius $R \geq 0.035$ cm, an areal mass density $\sigma > 1.4 \times 10^{-6}$ g cm $^{-2}$, and an areal number density $n < 8 \times 10^{-3}$ cm $^{-2}$."

In related work, Goldreich and Tremaine [1979, 1982] suggested that small, as yet discovered, satellites might be closely associated with each of the narrow rings of Uranus and, if so, would act as agents (shepherds) for dynamically maintaining the observed distribution of particulate matter. This theoretical suggestion received resounding confirmation with the Voyager 1 discovery [Smith et al., 1981] of two small satellites (1980 S26 and 1980 S27) that apparently act as shepherds for Saturn's Ring F [Gehrels et al., 1980]. More recently, Smith et al. [1986] have observed two small satellites associated

with Uranus' Ring ϵ . However, there are many, many narrow rings at Jupiter, Saturn, and Uranus that have no thus-far detected shepherds associated with them. Saturn's Ring G is one such example; but, in this case, no comprehensive search in longitude for small (radii ≤ 10 km), nearby satellites was made by either Voyager 1 or Voyager 2 [B. A. Smith, private communication]. Hence, there is no definitive optical evidence either for or against the presence of such satellites at Ring G.

On the basis of a crude comparative analysis of the energetic particle intensity at Ring G and in the inner slot region (caused by Janus and Epimetheus), Van Allen [1983] reported that "there are no satellites having radii of the order of a kilometer or larger in or near Ring G." This comparison ignored all of the optically observed particulate matter in Ring G and attributed its entire absorption signature to one or more small satellites, thus attempting to establish extreme upper limits.

The present paper has the same purpose but employs an absolute, rather than a comparative, analysis and probably provides superior results.

ABSORPTION OF A MAGNETICALLY-TRAPPED, GYRATING PARTICLE
BY A PLANETARY SATELLITE

The absorption of magnetospheric particles by planetary satellites has been discussed by a number of authors for various ranges of the relevant parameters [Singer, 1962; Mead and Hess, 1973; Mogro-Campero and Fillius, 1976; Thomsen, 1977; Van Allen et al., 1980a, 1980b; Thomsen et al., 1977; Van Allen, 1982, 1983; Rairden, 1980; Hood, 1981; Schardt and McDonald, 1983; Paonessa and Cheng, 1985; and Bell and Armstrong, 1986]. In its full generality, the absorption problem is a very complex geometrical one. Most of the above cited authors have treated it by approximate analytical methods. The two important exceptions are Rairden [1980] and Bell and Armstrong [1986], who used computerized sampling (Monte Carlo) methods.

The present treatment is specialized to the magnetosphere of Saturn and to the case in which the radius b of the (spherical, inert) satellite is very much less than the gyroradius r_g of the particle and in which r_g is, in turn, very much less than r , the radius of the satellite's orbit. It is further supposed that the magnetic moment of the planet is centered and aligned with its

rotational axis [Smith et al., 1980; Acuña and Ness, 1980; Acuña et al., 1980; and Connerney et al., 1984]; that the satellite is moving in a prograde circular Keplerian orbit of radius r in the planet's equatorial plane; and that the guiding center of the particle's helical motion drifts in longitude in a circle of approximately the same radius and oscillates in latitude between mirror latitudes $\pm \lambda_m$. With all of these simplifications, the absorption problem can be treated by a simple, analytical method having reasonable accuracy. Remarks on the effects of departures from the simplifying conditions will be made later. Other basic parameters are the kinetic energy E and species of the particle, its velocity vector \vec{V} , its equatorial pitch angle $\alpha_o = \arccos(\vec{V}_o \cdot \vec{B}_o / V_o B_o)$ (where \vec{B}_o is the local magnetic field vector and the zero subscripts denote equatorial values), the equatorial values of its gyroradius r_g and gyroperiod T_g , and its latitudinal bounce period T_B . The relationship connecting α_o and λ_m is $\sin^2 \alpha_o = \cos^6 \lambda_m (4 - 3 \cos^2 \lambda_m)^{-\frac{1}{2}}$. Extensive use is made of the formulae of Thomsen and Van Allen [1980]. For numerical examples of the Ring G problem, we assume that a representative particle is a proton $E_p = 200$ MeV and that a representative value of α_o is 70° . The other relevant parameters then have values as follows: $r = 170,200$ km; $T_g = 9.084 \times 10^{-2}$ s; $T_B = 3.085$ s; $r_g = 2,305$ km;

$\lambda_m = 9.6^\circ = 0.168 \text{ rad}$; and $V = 169,740 \text{ km s}^{-1}$. The angular velocity ω of the satellite relative to the longitude of the particle's guiding center is $2.120 \times 10^{-2} \text{ rad s}^{-1}$ and thus $\omega r = 3,608 \text{ km s}^{-1}$; and the interval of time T_E between encounters of the satellite with the gyrating particle is $2\pi/\omega = 296 \text{ s}$. The overall results are relatively insensitive to the value of E_p within the energy range 80-300 MeV. In the course of the analysis, numerical values of calculated quantities are given in square brackets as an aid to quantitative understanding.

During a particular encounter, there are two opportunities for the particle to hit the satellite. The probability of a hit during each opportunity is now calculated.

The origin of the adopted Cartesian coordinate system is the point at which the guiding center of a particular gyrating particle pierces the equatorial plane; the origin moves in longitude at the same average rate as does the guiding center. The X-axis is tangent to the satellite's orbit, the Y-axis is directed to the center of the planet, and the Z-axis is parallel to the rotational axis of the planet. In this frame of reference the particle moves along a helical path on the surface of a truncated, circular tube of magnetic flux, whose axis at the equator is along the Z-axis, whose length is $\approx 2r\lambda_m$ and whose equatorial radius is r_g . The satellite drifts parallel to the X-axis through this surface with velocity $v_x = \omega r$, $v_y = 0$, and $v_z = 0$ and with its center at $y = y_0$ and $z = 0$. A hit can occur only if $|y_0| \leq (r_g + b)$.

The X-Y plane section through the surface is shown in Figures 1(a) and 1(b). For the case shown in Figure 1(a), two separate opportunities for a hit occur within the two intervals of time during which the center of the satellite drifts from A to B (first to second contact) and from C to D (third to fourth contact). Let $l = \overline{AB} = \overline{CD}$. Then from the geometry of the figure,

$$l = \left\{ \sqrt{(r_g + b)^2 - y_o^2} - \sqrt{(r_g - b)^2 - y_o^2} \right\}, \quad (1)$$

for $0 \leq |y_o| \leq (r_g - b)$

For the case shown in Figure 1(b), the two separate segments \overline{AB} and \overline{CD} merge into a single segment \overline{AD} ; denoting $l = \overline{AD}/2$ for this case,

$$l = \left\{ \sqrt{(r_g + b)^2 - y_o^2} \right\}, \quad (2)$$

for $(r_g - b) \leq |y_o| \leq (r_g + b)$

A sample, composite plot of l/r_g vs y_o/r_g is shown in Figure 2.

It is noted that:

$$l_{\max} = 2\sqrt{br_g} \quad \text{at} \quad |y_o| = (r_g - b) \quad (3)$$

and that the mean value of l ,

$$\langle l \rangle = \frac{\int l(y_0) dy_0}{\int dy_0} = \frac{\pi b}{(1+b/r_g)} \approx \pi b \quad (4)$$

with integration over the range $0 \leq |y_0| \leq (r_g + b)$ [cf. Schardt and McDonald, 1983]. The center of the satellite spends time Δt in transversing l where

$$\Delta t = \frac{l}{\omega r} \quad (5)$$

and the radius of the circular patch of intersection between the cylindrical surface and the satellite during Δt varies from zero to b and back to zero with a mean value:

$$\langle b \rangle = \pi b/4, \quad (6)$$

except for y_0 nearly equal to r_g .

In the vicinity of the equator, the particle's motion may be visualized as composed of two components: a Z-component

$$V_z \approx V \cos \alpha_0 \quad [58,000 \text{ km s}^{-1} \text{ for } \alpha_0 = 70^\circ], \quad (7)$$

and a circumferential component:

$$v_{\phi} = \frac{2\pi r}{T_g} \cdot [159,400 \text{ km s}^{-1}] \quad (8)$$

The random probability that the particle lies within the azimuth required for a hit during Δt is

$$p_1 = \frac{\Delta t}{T_g} = \frac{\ell}{\omega r T_g} \cdot \quad (9)$$

If p_1 as calculated by (9) is less than unity for all values of y_0 , then

$$\langle p_1 \rangle = \frac{\langle \ell \rangle}{\omega r T_g} = \frac{\pi b}{\omega r T_g} \quad [9.6 \times 10^{-3} b, \quad (10) \\ \text{with } b \text{ in km}]$$

By (9) for $\ell_{\max} = 2 \sqrt{b r_g}$, p_1 exceeds unity if

$$b > 12 \text{ km.}$$

However, even if b exceeds this value by as much as a factor of 2, the mean value of $\langle p_1 \rangle$ as given by (10) is only 5 percent greater than its properly calculated value.

The independent (assuming no correlation between gyrophase and bounce phase) probability that the particle lies within $\pm \pi b/4$ of the equator at an arbitrary moment is

$$\langle p_2 \rangle = \left(\frac{\pi b/2}{V \cos \alpha_0} \right) \left(\frac{2}{T_B} \right)$$

$$\langle p_2 \rangle = \frac{\pi b}{V T_B \cos \alpha_0} \quad (11)$$

[For $\alpha_0 = 70^\circ$ and previously quoted values of V and T_B ,
 $\langle p_2 \rangle = 1.75 \times 10^{-5} b$, with b in km.]

A typical mean value of Δt (for $b = 5$ km) is 0.0044 s; during this time interval the equatorial projection of the particle's trace in space (when near the equator) travels through an azimuthal arc of 0.30 radian or 0.048 of a circle. A typical time lapse between centers of the two opportunities is 1.1 s or about 12 gyro-periods of the particle, and therefore the probability for a hit in either opportunity, insofar as it is small, is independent of that in the other (except for singular cases). When $y_0 \approx r_g$ the probabilities of a hit during the two opportunities are not statistically independent but for $b \ll r_g$ this qualification has a trivial effect on the overall analysis and I take the probability for a hit on a given encounter to be twice that for one of the two opportunities therein.

Thus the mean probability of a hit (averaged over y_0) during either of the two opportunities is:

$$\begin{aligned} \langle P \rangle &= \langle p_1 \rangle \cdot \langle p_2 \rangle \\ \langle P \rangle &= \frac{\pi^2 b^2}{\omega r V T_g T_B \cos \alpha_0} \end{aligned} \quad (12)$$

For a complete encounter (two opportunities, assumed independent)

$$\langle P \rangle = \frac{2\pi^2 b^2}{\omega r V T_g T_B \cos \alpha_0} \quad (13)$$

Putting my representative numerical values in (13)

$$\langle P \rangle = 3.36 \times 10^{-7} b^2 \quad (14)$$

with b in km. Equations (13) and (14) are appropriate for $b \leq 20$ km and are the principal result of the foregoing analysis

$\langle P \rangle$ in equations (13) and (14) may also be interpreted as the probability of a hit within T_E , the time interval between encounters. If there are n satellites of radii b_1, b_2, \dots, b_n moving in random phases in the same orbit, then the probabilities of a hit are additive and

$$\langle P \rangle = 3.36 \times 10^{-7} \sum_{i=1}^n b_i^2. \quad (15)$$

Equation (15) shows that $\langle P \rangle$ per T_E is proportional to the sum of the cross-sectional areas of the n satellites, provided that all are large enough to absorb the specified particle in a single hit and small enough ($b \leq 20$ km) to meet the criterion for applicability of (13).

The number of encounters per day is $86,400/T_E = 292$ in the case at hand. Hence the mean lifetime τ of a representative particle against absorption by a satellite of radius b is given by

$$\tau = \frac{1.02 \times 10^4}{b^2} \text{ days} = \frac{28}{b^2} \text{ years} \quad (16)$$

or for n satellites having radii b_1, b_2, \dots, b_n ,

$$\tau = \frac{28}{\sum_{i=1}^n b_i^2} \text{ years.} \quad (17)$$

In (14), (15), (16), and (17), b and b_i are in km.

COMMENTS ON VALIDITY AND ACCURACY
OF EQUATION (13)

The derivation of equation (13), though novel in detail, is akin to those of previous authors [in particular, the derivation of Schardt and McDonald, 1983] but has, I think, greater clarity. Nonetheless, it is not exact even for $b \leq 10$ km. A discussion of its shortcomings follows.

By equation (11), the value of $\langle p_2 \rangle$ becomes equal to 1.0 if

$$\frac{\pi b}{V T_B \cos \alpha_0} = 1. \quad (18)$$

For $b = 10$ km, $V = 169,740 \text{ km s}^{-1}$ and $T_B = 3.085 \text{ s}$, this condition corresponds to $\cos \alpha_0 \leq 6 \times 10^{-5}$ or α_0 differing from 90° by only 0.004° . Within this range of α_0 , $\langle p_2 \rangle$ must be set equal to 1.0 and $\langle P \rangle$ is very much greater than it is for values of α_0 that differ substantially from 90° . It is therefore clear that particles having α_0 near 90° are preferentially absorbed. The dependence of $\langle P(\alpha_0) \rangle$ on α_0 is illustrated by Table 1 (all for $E_p = 200 \text{ MeV}$).

For an angular distribution of unidirectional particle intensity of the form $\sin^n \alpha_0$, the weighted value of $\langle P \rangle$ over the range $0^\circ \leq \alpha_0 \leq 85^\circ$ (i.e., ignoring the short-lived range $85^\circ \leq \alpha_0 \leq 90^\circ$) is equal to $\langle P(\alpha_0) \rangle$ with $\alpha_0 = 74^\circ$ to 77° for $n = 1, 2$, and 3 ; and over the range $0^\circ \leq \alpha_0 \leq 80^\circ$, the weighted value of $\langle P \rangle$ is equal to $\langle P(\alpha_0) \rangle$ with $\alpha_0 = 64^\circ$ to 70° for $n = 1, 2$, and 3 . These calculations provide the justification for adopting $\alpha_0 = 70^\circ$ as a representative value for simplified considerations which do, of course, ignore refined features of the absorption process (i.e., the angular dependence of $\langle P(\alpha_0) \rangle$). It is noted that the absorption signatures of Ring G were measured with omnidirectional detectors.

If the magnetic axis of the planet is tilted with respect to its rotational axis, and/or if the orbit of the hypothetical satellite is inclined to the planet's equatorial plane, there are additional considerations. However, if the tilt and inclination angles are small ($\leq 1^\circ$), then the consequent effects are significant only for α_0 near 90° and are otherwise swallowed up in the distribution of particle intensity with α_0 . For Saturn, the tilt of the magnetic axis is $\leq 1^\circ$ [Connerney et al., 1984] and the inclination of the orbits of known inner satellites is also $\leq 1^\circ$ [Synnott et al., 1981, 1983]. Further, if the magnetic dipole moment of the planet is offset from the geometrical center of the planet and/or

if the hypothetical satellite is moving in an eccentric orbit, the effect is to reduce the probability of a hit per T_E for a specified particle. The ratio r_g/r (≈ 0.0135) gives a measure of the magnitude of the combined eccentricities at which this effect becomes significant. By the references just cited, the offset of the dipole is < 600 km (i.e., $< 0.26 r_g$) and the eccentricities of the orbits of the known inner satellites are ≤ 0.009 . Hence, the potential defects in the simplified derivation of equation (13) as listed in this paragraph are probably trivial. Observational support for this conclusion is provided by Figures 4 and 5 of Van Allen [1983], which show that the radial width of the absorption signature of Ring G is, in fact, not significantly greater than $2 r_g$. The guiding centers of particles do, of course, diffuse back and forth in r ; if the characteristic time constant for such diffusion is less than the lifetime for absorption, then the signature is broadened. The above observational fact apparently shows that this effect is also not important -- a result that is more persuasive than, but consistent to an order of magnitude with, an estimate of the time constant for diffusion, which is $r_g^2/4D \approx 7$ months.

In Van Allen [1982] it is shown that the energy distribution of protons in the vicinity of Ring G is such that about half of the particles have energy E_p greater than 200 MeV, and half have energy

E_p less than 200 MeV. This is the foundation for adopting 200 MeV as a representative value of E_p in the foregoing numerical examples.

Overall, I estimate that the numerical values in equations (14) and (16) are trustworthy to a factor of two.

ADDITIONAL COMMENTS

A noteworthy feature of equation (13) is that, for b sufficiently small (i.e., ≤ 10 km), $\langle P \rangle$ is proportional to the sum of the cross-sections of the assemblage of absorbing satellites as written in (15). The latter equation is applicable to particulate matter in the form of small spheres of radius b and volumetric mass density ρ provided that $4 \rho b/3$ is greater than the range of the relevant protons ($\approx 26 \text{ g cm}^{-2}$) in the particulate material. If a sphere is pulverized into n equal smaller spheres and if the latter are dispersed widely (no "shadowing"), then the projected area of the assemblage is increased by the factor $n^{1/3}$. If, however, the radii of the small objects are such that $4 \rho b/3$ is much less than the range of the protons, a given quantity of material becomes an even more effective absorber. The latter case is found to be applicable to the particulate matter in Ring G, if its absorption signature is attributed entirely to small particulates of uniform size [Van Allen, 1983]. The inferred sum of the volumes of all particulates in the ring is equivalent to the volume of a single sphere of radius only 0.1 km. The inferred total number of particulates is 4.3×10^{16} . By the pulverization theorem, the

radius of a single satellite having the same absorption for energetic protons would be 8 km if, contrary to my previous analysis, the radii b of the particulates were greater than $4 \rho b/3$.

For a single satellite with $b \geq 100$ km, $\langle p_1 \rangle$ must be set equal to unity for each of the two opportunities during an encounter. Then

$$\langle P \rangle = 2 \langle p_2 \rangle = \frac{2\pi b}{V T_B \cos \alpha_0} \quad (19)$$

or numerically

$$\langle P \rangle = 3.5 \times 10^{-5} b \quad (20)$$

for $\alpha_0 = 70^\circ$ and b in km. In this case, $\langle P \rangle$ is proportional to the first power of b and, corresponding to (16) and (17)

$$\tau = \frac{196}{b} \text{ years} \quad (21)$$

for a single satellite or

$$\tau = \frac{196}{\sum b_i} \text{ years} \quad (22)$$

for more than one.

The intersection of (20) and (14) occurs at

$$b = 100 \text{ km.} \quad (23)$$

There is a gradual transition from the quadratic regime for $b \leq 20$ km to the linear regime for $b \geq 100$ km. As b increases further, $\langle P \rangle$ approaches 1.0 asymptotically.

CONCLUSIONS

The analysis in Van Allen [1983] finds that the lifetime against absorption of 200 MeV protons whose guiding centers lie within $\pm r_g$ of the center of Ring G lies in the range 1.6 to 3.5 years. If one adopts an intermediate value of 2.5 years, then by (16)

$$b = 3.3 \text{ km} \quad (24)$$

or by (17)

$$\sum_{i=1}^n b_i^2 = 11.2 \text{ km}^2. \quad (25)$$

The reader is reminded that these estimates are in the nature of extreme upper limits for the radii of one or more shepherding satellites because they ignore the absorbing effects of the optically-observed distribution of particulate matter.

It is difficult to assign a rigorous uncertainty to the foregoing values but I estimate that equation (24) is uncertain by less than a factor of two and equation (25), by less than a factor of four.

ACKNOWLEDGEMENTS

This work has been supported by the Ames Research Center/
NASA under Contract NAS2-12327, by the U. S. Office of Naval
Research under Contract N00014-85-K-0404, and by the National
Aeronautics and Space Administration under Headquarters Grant
NGL 16-001-002, all of which are gratefully acknowledged.

REFERENCES

- Acuña, M. H., and N. F. Ness, The magnetic field of Saturn: Pioneer 11 observations, Science, 207, 444-446, 1980
- Acuña, M. H., N. F. Ness, and J.E.P. Connerney, The magnetic field of Saturn: Further studies of Pioneer 11 observations, J. Geophys. Res., 85, 5675-5678, 1980
- Bell II, E. V., and T. P. Armstrong, Monte Carlo simulation of charged particle impact on the satellites of Jupiter and Saturn, J. Geophys. Res., 91, 1397-1403, 1986
- Connerney, J.E.P., L. Davis, Jr., and D. L. Chenette, Magnetic field models. In Saturn, ed. by T. Gehrels and M. S. Matthews, pp. 354-377, University of Arizona Press, Tucson, AZ, 1984
- Fillius, W., W.-H. Ip, and C. E. McIlwain, Trapped radiation belts of Saturn: First look, Science, 207, 425-431, 1980
- Fillius, W., and C. McIlwain, Very energetic protons in Saturn's radiation belt, J. Geophys. Res., 85, 5803-5811, 1980
- Gehrels, T., et al., Imaging photopolarimeter on Pioneer Saturn, Science, 207, 434-439, 1980
- Goldreich, P., and S. Tremaine, Toward a theory for the Uranian rings, Nature, 277, 97-99, 1979

- Goldreich, P., and S. Tremaine, The dynamics of planetary rings,
Ann. Rev. Astron. Astrophys., 20, 249-283, 1982
- Hood, L. L., A comparison of characteristic times for
satellite absorption of energetic protons trapped in the
Jovian and Saturnian magnetic field, Geophys. Res. Lett., 8,
976-979, 1981
- McDonald, F. B., A. W. Schardt, and J. H. Trainor, If you've seen
one magnetosphere, you haven't seen them all: Energetic
particle observations in the Saturn magnetosphere,
J. Geophys. Res., 85, 5813-5830, 1980
- Mead, G. D., and W. N. Hess, Jupiter's radiation belts and the
sweeping effects of its satellites, J. Geophys. Res., 78,
2793-2811, 1973
- Mogro-Campero, A., and W. Fillius, The absorption of trapped
particles by the inner satellites of Jupiter and the radial
diffusion coefficient of particle transport, J. Geophys.
Res., 81, 1289-1295, 1976
- Paonessa, M. T., and A. F. Cheng, A theory of satellite sweeping,
J. Geophys. Res., 90, 3428-3434, 1985
- Rairden, R. L., Satellite sweeping of electrons and protons in
Saturn's inner magnetosphere, Res. Rep. 80-29, Univ. of
Iowa, Iowa City, July 1980

- Schardt, A. W., and F. B. McDonald, The flux and source of energetic protons in Saturn's inner magnetosphere, J. Geophys. Res., 88, 8923-8935, 1983
- Simpson, J. A., T. S. Bastian, D. L. Chenette, G. A. Lentz, R. B. McKibben, K. R. Pyle, and A. J. Tuzzolino, Saturnian trapped radiation and its absorption by satellites and rings: The first results from Pioneer 11, Science, 207, 411-415, 1980a
- Simpson, J. A., T. S. Bastian, D. L. Chenette, R. B. McKibben, and K. R. Pyle, The trapped radiations at Saturn and their absorption by satellites and rings, J. Geophys. Res., 85, 5731-5762, 1980b
- Singer, S. F., Radiation belts of Venus and Mars (with consideration of sweeping effect of Phobos). In Space Age Astronomy, ed. by A. J. Deutsch and W. G. Klemperer, pp. 444-461, Academic Press, New York, 1962
- Smith, E. J., L. Davis, Jr., D. E. Jones, P. J. Coleman, Jr., D. S. Colburn, P. Dyal, and C. P. Sonett, Saturn's magnetic field and magnetosphere, Science, 207, 407-410, 1980
- Smith, B. A., et al., Encounter with Saturn: Voyager 1 imaging science results, Science, 212, 163-191, 1981
- Smith, B. A., et al., Voyager 2 in the Uranian system: Imaging science results, Science, 233, 43-64, 1986

- Synnott, S. P., C. F. Peters, B. A. Smith, and L. A. Morabito,
Orbits of the small satellites of Saturn, Science, 212,
191-192, 1981
- Synnott, S. P., R. J. Terrile, R. A. Jacobson, and B. A. Smith,
Orbits of Saturn's F ring and its shepherding satellites,
Icarus, 53, 156-158, 1983
- Thomsen, M. F., On determining a radial diffusion coefficient
from the observed effects of Jupiter's satellites, Ph.D.
dissertation, University of Iowa, 1977
- Thomsen, M. F., C. K. Goertz, and J. A. Van Allen, On determining
magnetospheric diffusion coefficients from the observed
effects of Jupiter's satellite Io, J. Geophys. Res., 82,
5544-5550, 1977
- Thomsen, M. F., and J. A. Van Allen
Motion of trapped electrons and protons in Saturn's inner
magnetosphere, J. Geophys. Res., 85, 5831-5834, 1980
- Trainor, J. H., F. B. McDonald, and A. W. Schardt, Observations
of energetic ions and electrons in Saturn's magnetosphere,
Science, 207, 421-425, 1980
- Van Allen, J. A., Findings on rings and inner satellites of
Saturn by Pioneer 11, Icarus, 51, 509-527, 1982
- Van Allen, J. A., Absorption of energetic protons by Saturn's
ring G, J. Geophys. Res., 88, 6911-6918, 1983

Van Allen, J. A., B. A. Randall, and M. F. Thomsen, Sources and sinks of energetic electrons and protons in Saturn's magnetosphere, J. Geophys. Res., 85, 5679-5694, 1980a

Van Allen, J. A., M. F. Thomsen, B. A. Randall, R. L. Rairden, and C. L. Grosskreutz, Saturn's magnetosphere, rings and inner satellites, Science, 207, 415-421, 1980b

Table 1. Pitch-Angle Dependence of the Probability of Absorption

-- Relative to That at 70°.

α_0	$\frac{\langle P(\alpha_0) \rangle}{\langle P(70^\circ) \rangle}$	α_0	$\frac{\langle P(\alpha_0) \rangle}{\langle P(70^\circ) \rangle}$
89°	20.36	45°	0.42
85	4.07	40	0.37
80	2.03	35	0.34
75	1.34	30	0.31
70	1.00	25	0.28
65	0.79	20	0.26
60	0.65	15	0.24
55	0.55	10	0.22
50	0.48	5	0.21


FIGURE CAPTIONS

Figures 1(a) and 1(b).

The origin of the adopted Cartesian coordinate system is the point at which the guiding center of a particular gyrating particle pierces the equatorial plane; the origin moves in longitude at the same average rate as does the guiding center. The Y-axis is directed toward the planet's center and the Z-axis is parallel to its axis of rotation. In each of the two drawings, the large circle represents the equatorial projection of the segment of the particle's helical trajectory that lies near the equatorial plane. The small circles represent successive positions of the equatorial cross-section of a spherical satellite as it moves at constant speed v_x parallel to the instantaneous X-axis with its center at $y = y_0$, $z = 0$. Note that for the cases of interest in this paper $b/r_g \leq 0.001$. A much larger value of this parameter, namely 0.1, was chosen for these drawings in the interest of graphical clarity.


Figure 2

A composite plot of equations (1) and (2) of the text in dimensionless form. Note that this illustrative plot is for $b/r_g = 0.01$, a value of the order of ten times those of relevance to the cases of interest in this paper.



$$0 \leq |y_0| \leq (r_g - b)$$

(a)



$$b/r_g = 0.1$$

$$(r_g - b) \leq |y_0| \leq (r_g + b)$$

(b)

Figure 1





Figure 2



END

10-86

DT/C