

AD-A171 585

HIGH TEMPERATURE TEMPORAL STABILITY OF SELECTED
OXIDIZERS AS SOLIDS AND IN AQUEOUS SOLUTIONS(U) NAVAL
RESEARCH LAB WASHINGTON DC R E PELLENBARG ET AL

1/1


UNCLASSIFIED

88 AUG 86 NRL-MR-5824

F/G 15/2

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963 A

AD-A171 585

High Temperature Temporal Stability of Selected Oxidizers as Solids and in Aqueous Solutions

ROBERT E. PELLENBARG AND ELIZABETH SMIROLODO

*Combustion and Fuels Branch
Chemistry Division*

DTIC
ELECTE
SEP 4 1986
S B

DTIC FILE COPY

Approved for public release. distribution unlimited

86

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				
1a REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b RESTRICTIVE MARKINGS		
2a SECURITY CLASSIFICATION AUTHORITY		3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.		
2b DECLASSIFICATION/DOWNGRADING SCHEDULE				
4 PERFORMING ORGANIZATION REPORT NUMBER(S) NRL Memorandum Report 5824		5 MONITORING ORGANIZATION REPORT NUMBER(S)		
6a NAME OF PERFORMING ORGANIZATION Naval Research Laboratory	6b OFFICE SYMBOL (If applicable) Code 6182	7a NAME OF MONITORING ORGANIZATION		
6c ADDRESS (City, State, and ZIP Code) Washington, DC 20375-5000		7b ADDRESS (City, State, and ZIP Code)		
8a NAME OF FUNDING/SPONSORING ORGANIZATION Naval Sea Systems Command	8b OFFICE SYMBOL (If applicable)	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c ADDRESS (City, State, and ZIP Code) Washington, DC 29362		10 SOURCE OF FUNDING NUMBERS		
		PROGRAM ELEMENT NO. STP4213	PROJECT NO.	TASK NO. SF64-561
				WORK UNIT ACCESSION NO. DN080-124
11 TITLE (Include Security Classification) High Temperature Temporal Stability of Selected Oxidizers as Solids and in Aqueous Solutions				
12 PERSONAL AUTHOR(S) Pellenberg, Robert E. and Smiroldo, Elizabeth				
13a TYPE OF REPORT Interim	13b TIME COVERED FROM _____ TO _____	14 DATE OF REPORT (Year, Month, Day) 1986 August 8	15 PAGE COUNT 16	
16 SUPPLEMENTARY NOTATION				
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP		
19 ABSTRACT (Continue on reverse if necessary and identify by block number) <p>Various potential decontamination agents were examined as solids and in aqueous solutions for long term stability at high temperatures. The following oxidizers were assayed iodometrically: the hypochlorite salts of calcium and lithium, sodium dichloroisocyanurate (PACE) and the peroxygen compounds sodium perborate, sodium peroxydisulfate, sodium percarbonate, and magnesium monoperoxyphthalate (H-48). The inorganic peroxide solids and the solid sodium dichloroisocyanurate were stable at 80°C, while the organic peroxide solids and the hypochlorite salts deteriorated markedly within 72 hours. In freshwater solutions of 0.01 N or less, the inorganic hypochlorite and peroxide salts decomposed slowly at 60°C. Conversely, the sodium dichloroisocyanurate, magnesium monoperoxyphthalate, and sodium percarbonate solutions exhibited near complete decomposition in 24 hours.</p>				
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED-UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21 ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a NAME OF RESPONSIBLE INDIVIDUAL Robert E. Pellenberg		22b TELEPHONE (Include Area Code) 202-767-0063	22c OFFICE SYMBOL	

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted
All other editions are obsolete

SECURITY CLASSIFICATION OF THIS PAGE

CONTENTS

	<u>Page</u>
INTRODUCTION	1
EXPERIMENTAL	1
RESULTS AND DISCUSSION	3
Oxidizers as Solids	3
Oxidizers in Solutions	4
CONCLUSIONS	4
RECOMMENDATIONS	5
ACKNOWLEDGEMENTS	5
REFERENCES	6
TABLES 1 - 4	7 - 10
FIGURES 1 - 7	11 - 12

S DTIC
ELECTE **D**
SEP 4 1988
B



Accession	
NEWS GRA	<input checked="" type="checkbox"/>
DEIC TAB	<input type="checkbox"/>
Unannounc	<input type="checkbox"/>
Justified	<input type="checkbox"/>
By	
Distrib	
Availa	
Dist	
A-1	

HIGH TEMPERATURE TEMPORAL STABILITY OF SELECTED OXIDIZERS AS
SOLIDS AND IN AQUEOUS SOLUTIONS (U)

INTRODUCTION

Naval elements are potential targets for chemical warfare (CW) or biological warfare (BW), especially during armed hostilities. An effective CW/BW defense posture requires that impacted elements maintain operational status even in a toxic environment. This posture requires the ability to neutralize as necessary CW and/or BW agents on critical surfaces. An element in this decontamination (decon) process is fluids, based on sea or fresh water, which can neutralize agents/toxins.

Most agents/toxins will be oxidized/hydrolyzed by washdown with water containing appropriate additives. These additives include the traditional decontaminant of choice, calcium hypochlorite. Specifically, hypochlorite ion serves to hydrolyze, and oxidize G-agents, a class of CW compounds (Block and Davis, 1978). Furthermore, hypochlorous acid in aqueous solution at pH 7 appears effective in countering BW materials such as spores (Hoffman and Spiner, 1962; Fielding, et al., 1967, 1968; for $\text{HOCl} \rightleftharpoons \text{H}^+ + \text{OCl}^-$, $K = 2.95 \times 10^{-5}$ and $\text{pK} = 7.5$).

In spite of proven effectiveness of hypochlorite for decontamination applications, hypochlorite-based decon presents problems. These problems are in part associated with the chemical nature of hypochlorite. Specifically, in the case of calcium hypochlorite, the solid tends to cake and decompose in storage, and the salt disperses slowly into water, yielding a chalky suspension. Ideally, a decon reagent should be as chemically effective as calcium hypochlorite. Additionally, this ideal reagent should be stable when stored as a solid, be stable in aqueous solution (fresh or sea water) and dissolve rapidly to give a true solution. On-site generation of the ideal decon reagent would also be desirable, to reduce logistics of storing decon reagents.


The evaluation process for selecting suitable supplements/replacements for calcium hypochlorite will consider the factors listed above. This report examines the long-term stability at high temperature of selected hypochlorite and peroxide reagents as dry solids, and as aqueous solutions. Such data will help define storage stability of candidate decon reagents, and their stability in decon fluids prepared prior to actual need.

Stability is defined as retention of oxidative capacity with time. For this test, residual oxidative capacity was monitored iodometrically.

EXPERIMENTAL

Candidate oxidizers were examined as dry solids, and as aqueous solutions in distilled water. Multiple aliquots of the reagents (Table 1) were weighed into tared Pyrex petri dishes and placed in an oven set to maintain 80° C. At various times, one solid aliquot of each of the candidates was removed from the oven. The solids were placed in 100 mL of distilled water to yield approximately 0.01 N solutions, which were assayed iodometrically (see below). Aliquots of solid retained at room temperature served as controls in this sequence. Aqueous solutions of the candidate oxidizers were prepared in distilled water, giving 200 mL of approximately 0.01N solution. These solutions were placed in 250 mL round-bottomed flasks, and fitted to a Soxhlet extraction apparatus set to maintain the fluids at 60° C. The condensers of the Soxhlet apparatus were used to minimize solution concentration changes caused by evaporation. At various times, 2.0 mL aliquots of these hot, stored solutions were taken. The oxidizer content of these aliquots was measured iodometrically.

Iodometric titration was used to monitor the oxidizer content of all samples tested. These samples consisted of 2.0 mL aliquots of solution made from dry, heat stressed solid oxidizer, or 2.0 mL aliquots of heat stressed aqueous solution. The following equations describe the iodometric reactions:



Essentially, an oxidizer [hypochlorite ion in (1) above] oxidizes iodide to iodine, which then oxidizes thiosulfate ion. The amount of thiosulfate consumed (added as a standardized solution) corresponds to the amount of oxidizer present in the original aliquot. For the titrations, 2 mL aliquots of the various test solutions (from the stored solid, or stored solution) were added to 50 mL distilled water. The pH of each solution was then adjusted to 4.5 ± 0.5 with 2 drops of 2 N sulfuric acid in distilled water. To the acidified solution were added 2.0 mL of 10% wt./wt. potassium iodide in distilled water. This addition yielded gold color solutions if there was oxidizer in the test aliquot. The kinetics of the reaction between oxidizer and iodide are crucial in this titration. The chlorine based oxidizer reacted essentially instantaneously with added potassium iodide, so that the titration could proceed as described below. However, the peroxygen reagents reacted much more slowly with added iodide. Therefore these test solutions

were allowed to stand prior to titration with standard thiosulfate solution. These standing times are cited in Table 4, and were selected to allow the oxidizer-iodide interactions to proceed, but not necessarily to completion. In some cases (e.g. peroxydisulfate ion), reaction to completion would take so long that one would begin to lose gaseous iodine from the titration vessel upon standing; in these cases, a long, but not infinite reaction time was chosen. (Pellenbarg, 1985, addresses the reaction kinetics between iodide and peroxygen species in more detail). After reaction between iodide and oxidizer, the magnetically stirred solutions were titrated with 0.0375 N sodium thiosulfate solution (Fisher) dispensed from a 10 mL burette, to give a pale yellow color. Two to several drops of soluble starch solution (1.0 g. soluble starch [triturated to a paste with cold water] added slowly with stirring to 100 mL boiling distilled water containing 1 g boric acid [Fisher] as a preservative, cooled and stored in a brown glass bottle) were added to the titration mix to give a dark blue color. Additional thiosulfate solution was added until a clear, colorless endpoint resulted. The total volumes (titer) of thiosulfate solution used to neutralize oxidizer liberated iodine was recorded, and served as a monitor of residual oxidizer in a given sample as a function of time: the more oxidizer present in a given sample, the more thiosulfate solution consumed.

RESULTS AND DISCUSSIONS

Oxidizers as Solids

At 80° C, PACE, a chlorine based oxidizer, exhibited excellent long term stability as a solid, retaining almost 100% of initial oxidizer content after 24 days. The other chlorine containing materials were much less stable as heat-stressed solids. Lithium hypochlorite lost all its oxidizer capacity after only 3 days, and the oxidizer content of calcium hypochlorite fell by approximately two-thirds after 10 days. (Figures 1 - 3)

The peroxygen reagents exhibited mixed stability, compared to the chlorine bearing materials. Sodium perborate and peroxydisulfate were fairly stable with time at 80° C, losing some 50% of initial oxidizer content in 24 days. Conversely, H-48 and sodium percarbonate were markedly unstable as solids at 80° C, and became essentially inactive after only 3 days. (Figures 4 -7).

The solids analyzed above required dissolution prior to measurement of initial or residual oxidizer content. This dissolution process yielded several significant observations. PACE dissolved almost instantly, and gave a no-residue solution at all times in the test sequence. However, the hypochlorite salts, especially after aging at 80° C, dissolved slowly, and

left undissolved precipitates/sediments. These suspended/sedimented solids could prove troublesome when dispersing any decon solution containing a hypochlorite salt; such problems, as clogging of application equipment, should be investigated in more detail.

Like PACE, heat-stressed H-48 and sodium percarbonate solids dissolved rapidly and cleanly, but gave solutions with little residual oxidizer content. The sodium perborate and sodium peroxodisulfate solids tended to dissolve well, and gave solution with fair residual oxidizer content. These latter solutions though, reacted slowly with added iodide, indicating the possibility of a different reaction mechanism between oxidizer and substrate (iodide ion) for the chlorine-based, and some oxygen-based oxidizers. This possible kinetic difference should be examined in more detail to indicate potential new approaches to decon chemistry.

Oxidizers as Solutions

One CW/BW defense scenario could involve the preparation and storage of a decon fluid prior to actual need, so that decon materials would be on hand and ready for use as necessary. Any candidate which rapidly decomposes in solution may prove less than useful in such a field setting. With this possibility in mind, 60° C solutions of candidate oxidizers in distilled water were monitored with time.

At 60° C in solution, PACE decomposed markedly in a few hours, and lost most of its oxidizer capacity in less than one day. Upon prolonged storage of the aged PACE solutions, the initially clear solution yielded yellowish particles as a precipitate. These particulates could cause problems with solution dispensing. In solution, both hypochlorite salts were more stable with time than was the PACE, and indeed, were more stable for a longer time than were the dry, heat-stress solids (Figures 1 - 3). It would be useful to examine mechanisms which could account for the differing stabilities observed in this testing sequence.

The tested peroxygen salts exhibited variable stability in solution, but all decomposed within about 4 days at 60° C (Figures 5 - 8). Thus, sodium perborate retained approximately 60% of its initial oxidizer content after 4 days, while the other candidates were essentially gone in this time. In solution, sodium percarbonate gave a precipitate/sediment, sodium perborate yielded long crystalline needles, while the sodium peroxydisulfate did not change appreciably. The changes in the chemical/physical characteristics of the candidate solutions have implications as to the solutions' ultimate utility as a decon fluid.

CONCLUSIONS

The study described in this report examined the stability in storage at high temperature of seven candidate decon materials tested as dry powders and as aqueous solutions in distilled water. As solids, only PACE, of the chlorine-based oxidizers, exhibited oxidizer capacity for more than 24 hours. Both lithium and calcium hypochlorites decomposed rapidly at high temperatures when stored as solids. The inorganic peroxygen materials sodium perborate and sodium peroxydisulfate were stable as solids at high temperature, while solid H-48 and solid sodium percarbonate lost all oxidizer capacity in approximately 72 hours.

In solution, however, PACE was much less stable than the other chlorine-based oxidizers, and decomposed markedly in less than 24 hours. However, even the lithium and calcium hypochlorites decomposed appreciably in 36 - 48 hours. The peroxygen compounds in solution also exhibited variable, but short lifetimes, with sodium perborate losing only about half its oxidizing capacity after 4 days. The other peroxygen materials were less stable with time in solution. These observations indicate that, in general, the oxidizers studied exhibit instability at high temperatures whether in solution, or as the solid. This fact dictates appropriate logistics when storing solid materials to be added to decontamination fluids, and suggests caution with regard to the use of hot solutions of decon reagents stored for more than a few hours prior to use.

RECOMMENDATIONS

This study has identified several topics which impact directly on the preparation of decon fluids from solid, stored precursors. Specifically needed now are data which:

1. Compare the decon efficiency of the better candidates examined in the study against appropriate CW and BW simulants, and ultimately, against live agents and toxins.
2. Study effective packaging/storage procedures for those candidates which prove effective in decon tests with simulants/agents.
3. Examine the effects of various additives to the oxidizer solutions studied with regards to the residual oxidizer capacity. AFFF is a potential additive in this context, because of its useful surfactant properties and its availability aboard Navy ships.
4. Investigate the degradation of Navy materials (including composites, rubber, etc.) in contact with solutions of the candidates in fresh and salt water, with and without additives such as AFFF.

ACKNOWLEDGEMENTS

The authors wish to thank the Naval Sea Systems Command, especially NAVSEASYSKOM Code 05R16, for partial support of this project.

REFERENCES

1. Block, F. and Davis, G. T., "Survey of Decontamination Methods Related to Field Decontamination of Vehicles and Material (Abridged)," Report #ARCSL-TR-79002, 28 p., 1989.
2. Fielding, G. H., Neihof, R. A. and Echols, W. H., "Biological Warfare Decontamination with Hypochlorite. Part I- Exploratory Studies: Application to the STOP System," NRL Report 6506, 1967.
3. Fielding, G. H., Neihof, R. A., "Bio-Decontamination with Hypochlorite," in Report of NRL progress, p. 7-9, Oct. 1968.
4. Hoffman, R. K., and Spiner, D. R., "Decontamination Procedures for Personnel Handling Biological Weapons," TM12 (AD285543), U.S. Army Biological Laboratories, Ft. Detrick, MD, Sept. 1962.
5. Pellenbarg, R. E., "pH Dependent Interactions Between Aqueous Iodide Ion and Selected Oxidizers," NRL Memorandum Report 5693, December 1985.

TABLE 1: OXIDIZERS TESTED

Name:	Formula Weight	Salt, grams/liter (Normality)
Lithium Hypochlorite *(Lithco)	58.4	6.56 (0.03)
Calcium Hypochlorite (Fisher)	141.1	3.58 (0.025)
Sodium Dichloroisocy- anurate (PACE) (Olin)	255.0	6.41 (0.025)
Sodium Perborate (Alfa)	153.8	1.55 (0.01)
Sodium Peroxydisulfate (Alfa)	238.0	2.37 (0.01)
Magnesium Bis (2-Car- boxylato mono peroxy- benzoic acid) hexahy- drate (H-43) (Interox)	496.3	4.96 (0.01)

* Sources

Lithco: Lithium Corporation of America, Bessemer City, NC:
Lithco product is approximately 30% lithium hypochlorite.

Fisher: Fisher Scientific Company, Fairlawn, NJ

Olin: Olin Chemicals, Stanford, CT

Alfa: Alfa Products, Danvers, MA

Interox: Interox America, Houston, TX

TABLE 2: HYPOCHLORITE TEST RESULTS

LITHIUM HYPOCHLORITE

Solids*		Solutions**	
Time (days)	Titer (mL.)	Time (days)	Titer (mL.)
0	2.53	0	2.53
1	0.01	1/8	2.52
3	0.04	1	2.56
10	0.01	6	0.99
18	0.003	8	1.01

CALCIUM HYPOCHLORITE

0	2.93	0	2.93
3	2.21	1/8	2.69
10	0.66	1	2.52
14	0.96	2	2.34
18	1.14	6	1.04
24	0.55	8	1.24

FACE

0	4.30	0	4.77
3	4.22	1/8	3.89
10	3.60	1	0.23
14	3.80	2	0.04
18	4.70	6	0.00
24	4.18	8	0.00

* Solids stored as dry material at 80° C.

** Solutions: Candidates dissolved in distilled water, retained at 60° C.

TABLE 3: PEROXYGEN TEST RESULTS

SODIUM PERBORATE

Solids*		Solutions**	
Time (days)	Titer (mL.)	Time (days)	Titer (mL.)
0	0.44	0	0.33
1	0.36	1	0.24
3	0.31	4	0.18
10	0.38		
14	0.42		
18	0.24		
24	0.22		

SODIUM PEROXYDISULFATE

0	0.18	0	0.18
3	0.21	1	0.07
10	0.14	4	0.05
14	0.22		
18	0.17		
24	0.11		

H-48

0	1.34	0	1.34
3	0.04	1	0.18
10	0.06	4	0.00
14	0.00	8	0.00
18	0.00		
24	0.00		

SODIUM PERCARBONATE

0	0.83	0	0.83
3	0.05	1	0.20
10	0.01	4	0.10
14	0.04		
18	0.00		
24	0.02		


* Solids stored as dry materials at 80° C.

** Solutions: Candidates dissolved in distilled water, retained at 60° C.


TABLE 4: PHYSICAL STATE OF OXIDIZER

Oxidizer	Physical State		Standing Time before Titration (min)
	as solid	in solution	
Lithium hypochlorite	dissolves slowly	slight sediment formation	0
Calcium hypochlorite	yields a chalky suspension, dissolves slowly	yields chalky suspension/sediment	0
PACE	dissolves quickly	yellowish precipitate forms	0
Sodium perborate	dissolves well	forms long crystalline needles when heated	25
Sodium peroxydisulfate	dissolves well	remained clear	50
Sodium percarbonate	dissolves well	yields precipitation/sediment	5
H-48	dissolves well; changed from white to tan with time	hot stored solution changes from clear to brown when heated	0- as solid 10- in solution


LITHIUM HYPOCHLORITE
FIGURE 1



CALCIUM HYPOCHLORITE
FIGURE 2




PAGE
FIGURE 3




OXIDIZER STABILITY DATA
(for chlorine-based oxidizers)


SODIUM PERBORATE




SODIUM PEROXYDISULFATE



SODIUM PERCARBONATE



H-48
FIGURE 7



OXIDIZER STABILITY DATA
 (for oxygen-based oxidizers)

END

10-86

DT/C