

AD-A171 998

DYNAMICAL - CHEMICAL COUPLING IN THE MESOSPHERE AND
LOWER THERMOSPHERE(U) BOSTON UNIV MA DEPT OF ELECTRICAL
COMPUTER AND SYSTEMS ENGINE. J M FORBES 31 JAN 85
AFOSR-TR-86-0653 AFOSR-84-0182


1/1

UNCLASSIFIED

F/G 4/1

NL





1000

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

28

(2)

REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS BEFORE COMPLETING FORM

1. REPORT NUMBER AFOSR-TR-86-0653		2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Dynamical - Chemical coupling in the Mesosphere and Lower Thermosphere		5. TYPE OF REPORT & PERIOD COVERED Final Report	
7. AUTHOR(s) Prof. J. M. Forbes		8. CONTRACT OR GRANT NUMBER(s) AFOSR-84-0182	
9. PERFORMING ORGANIZATION NAME AND ADDRESS College of Engineering, Boston University Boston, MA 02215		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102^F 2310/A2	
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research Building 410 Bolling AFB, DC 20332		12. REPORT DATE 31 January 1985	
		13. NUMBER OF PAGES 3	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified	
		15a. DECLASSIFICATION, DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public releas; distribution unlimited			
17. DISTRIBUTION STATEMENT (of abstract entered in Block 20, if different from Report)	18. SUPPLEMENTARY NOTES		
	B		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Ionosphere Dynamics Arecibo Tides			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The dynamic ionosphere over Arecibo is simulated using a finite element technique. It is shown that the so-called 'collapse of the Arecibo F-Layer is caused by the upward-propagating semidiurnal tide excited in the upper stratosphere by ozone heating. The steep underside density gradients observed in conjunction with the collapse are shown to be due to the shear in the meridional wind field of the semidiurnal tide. These gradients are capable of triggering the gradient drift plasma instability.			

AD-A171 998

DTIC FILE COPY

DTIC ELECTED

SEP 15 1986

DD FORM 1473 JAN 73

Unclassified SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Copy 16
↓

and accounting for plasma irregularity formation and observations of VHF scintillations associated with the collapse phenomenon.

↑

Boston University
Department of Electrical, Computer, and Systems Engineering

Final Technical Report
1 June 1984 - 1 December 1984
Grant No: AFOSR - 84 - 0182

Entitled
Dynamical - Chemical Coupling In The Mesosphere
And Lower Thermosphere

Professor Jeffrey M. Forbes
Principal Investigator

Approved for public release.
distribution unlimited

ARMY OFFICE OF SCIENTIFIC RESEARCH (AFSC)
OFFICE OF TRANSMITTAL TO DTIC
This technical report has been reviewed and is
approved for public release IAW AFR 190-12.
Distribution is unlimited.
MATTHEW J. KERPER
Chief, Technical Information Division

1. INTRODUCTION

Due to the transfer of the Principal Investigator, Prof. Jeffrey M. Forbes, from Boston College to Boston University during January 1984, Grant AFOSR-81-0090 to Boston College was terminated and the remaining funds (\$19,326 total) were re-awarded to the P.I. at Boston University to complete the work. This report constitutes the final report under the re-award grant AFOSR-84-0182 to Boston University.

2. FINITE-ELEMENT SIMULATION OF THE "MIDNIGHT COLLAPSE" OF THE IONOSPHERE OVER ARECIBO.

A numerical simulation code using the finite element technique developed and utilized by Prof. M. Mendillo and the Astronomy Department Group at Boston University for modelling F-region chemical releases with characteristic times of less than an hour was modified for the simulation of longer period phenomena with special emphasis on the dynamical behavior of the F-region ionosphere. Modifications for the study of F-region dynamical behavior included making provisions for (a) photoionization; (b) winds; (c) electric fields; (d) airglow calculations; and (e) protonospheric replenishment at night. All of these efforts required much debugging and testing against "known results" at many levels of development.

The main result of the simulations was that the dynamic behavior of the Arecibo ionosphere results directly from upward propagating semidiurnal tidal components of the neutral wind excited below

thermospheric levels. The addition of terdiurnal and higher - order components in the wind field improves the agreement with experiment and it is hypothesized that these result from the non-linear coupling between the neutral wind and the diurnal variation in ion-drag. In addition, the steep underside density gradients occurring in conjunction with the collapse are shown to be due to the shear in the meridional wind field, which is expected to be associated with upward propagating components of the semidiurnal wind described above. These gradients are capable of triggering the gradient drift plasma instability, and accounting for plasma irregularity formation and observations of VHF scintillations associated with the collapse phenomenon.

Furthermore, we have shown the importance of an abatement, rather than reversal, of the neutral wind velocities in determining the level of h_{max} in our model. This is especially germane in the case of the pre-sunrise increase in h_{max} when an increase in the height of the peak occurs when vertical drifts due to neutral winds are large in magnitude and downward.

The above results are discussed in detail in the manuscript "The Dynamic Ionosphere over Arecibo: A Theoretical Investigation" by D.J. Cray and J.M. Forbes, J. Geophys. Res., to appear in late 1985 or early 1986.



<input checked="" type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	
Distributed to Authority Available for Dist Special	
A-1	

END

10-86

DTIC