

AD-R172 017

THE SHAPES OF THE CIRCUMSTELLAR 'SILICATE' FEATURES(U)

1/1

AIR FORCE GEOPHYSICS LAB HANSCOM AFB MA


I R LITTLE-MARENIN ET AL. 08 SEP 86 AFGL-TR-86-0183

UNCLASSIFIED

F/G 3/2

NL





1057

# AD-A172 017

## T DOCUMENTATION PAGE

1a. R Un		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for Public Release; Distribution Unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) AFGL-TR-86-0183		7a. NAME OF MONITORING ORGANIZATION	
6a. NAME OF PERFORMING ORGANIZATION Air Force Geophysics Laboratory	6b. OFFICE SYMBOL (if applicable) OPI	7b. ADDRESS (City, State, and ZIP Code)	
6c. ADDRESS (City, State, and ZIP Code) Hanscom AFB Massachusetts, 01731		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (if applicable)	10. SOURCE OF FUNDING NUMBERS	
8c. ADDRESS (City, State, and ZIP Code)		PROGRAM ELEMENT NO. 62101F	PROJECT NO. 7670
		TASK NO. 06	WORK UNIT ACCESSION NO. 13
11. TITLE (Include Security Classification) The Shapes of the Circumstellar "Silicate" Features			
12. PERSONAL AUTHOR(S) Irene R. Little-Marenin *, Stephen D. Price			
13a. TYPE OF REPORT Reprint	13b. TIME COVERED FROM 2 Jul 86 TO 8 Jul 86	14. DATE OF REPORT (Year, Month, Day) 1986 September 8	15. PAGE COUNT 2
16. SUPPLEMENTARY NOTATION *Wellesley College Reprinted from Summer School on Interstellar Processes July 2-8, 1986 at Grand Teton National Park			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	IR Emission , Circumstellar dust shells	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The spectral energy distributions of circumstellar dust around cool O-rich stars have been analyzed. Three general categories of spectral shapes have been identified.			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL Stephen D. Price		22b. TELEPHONE (Include Area Code) (617) 377-4552	22c. OFFICE SYMBOL OPT

DTIC FILE COPY



Availability Codes  
Avail and/or Special

S

SEP 17 1986

A

THE SHAPES OF THE CIRCUMSTELLAR "SILICATE" FEATURES

Irene R. Little-Marenin (AFGL and Wellesley C.)

Stephan D. Price (AFGL)

*micron features*  
→ Around oxygen-rich stars we find that the spectra of most long-period variables (LPV) show an excess infrared emission which is attributed to circumstellar silicate dust grains. These grains produce emission features at about 10 and 18  $\mu\text{m}$  due to bending and stretching modes of SiO respectively. It has been known (Forrest, Gillett and Stein 1975) that the spectral energy distribution of the 10  $\mu\text{m}$  emission shows variations from star to star. With the availability of many IRAS Low Resolution Spectra (LRS) in the 8-22  $\mu\text{m}$  region of M stars, we can now study the 10  $\mu\text{m}$  feature to determine its uniformity (or lack thereof). For this analysis we assume that the 8-22  $\mu\text{m}$  emission from these stars is produced by a) the stellar photosphere, b) a continuum emission from the dust grains and c) a strongly wavelength dependent dust grain emission term. By representing the first two terms with blackbody energy distributions and subtracting them from the observed spectrum, we are left with a remaining strongly wavelength dependent emission feature which we call the excess silicate or 10  $\mu\text{m}$  emission.




Fig. 1. Two M star LRS spectra (V Cam and W Hor) are plotted together with black body energy distributions fitted to either side of the emission feature. The 10  $\mu\text{m}$  excess (observed - local black body continuum) is plotted above the wavelength axis. The difference in the shape of the excess between the two stars is obvious.

The excess silicate emissions from about 130 LPVs can be divided into three groups characterized by similar spectral shapes. All three shapes are shown in Figure 2. The average silicate feature of semiregular (SRa,b,c) and irregular (Lb) variables is shown as a solid line. This feature extends from 8.4 to 14.5  $\mu\text{m}$  with a peak at  $10 \pm 0.05 \mu\text{m}$ . The FWHM is  $2.1 \pm 0.15 \mu\text{m}$  but the feature is asymmetric with the ratio of the widths of the short wavelength (rising) branch compared to the long wavelength (falling) branch of 0.6:1.5 at half intensity. The rising branch shows only minor variations in wavelength from star to star whereas the wavelength at the FWHM point of the falling branch varies by  $\pm 0.15 \mu\text{m}$ . The average silicate excess of Mira variables is shown as a dashed line in Figure 2. The average feature extends from 8 to 14.5  $\mu\text{m}$  with a peak at 9.75  $\mu\text{m}$  and a FWHM of 2.3  $\mu\text{m}$ . The feature is asymmetric but on the average has a less

AFGL 12-86-0183

steeply rising branch than the other LPVs giving a ratio of the rising width to the falling width of 0.75:1.5 at half intensity. However, the spectral shape of the silicate emission feature among the Miras shows much greater variation from star to star than that of the other LPVs. The shape of the emission from Miras ranges from one identical to the SRs and Lbs to a much broader one, extending from  $<8 \mu\text{m}$  (the limit of the LRS detectors) to  $\sim 14.5 \mu\text{m}$  with a corresponding shift in peak emission from  $10$  to  $9.6 \mu\text{m}$  and an increase in the FWHM from  $2.1$  to  $2.4 \mu\text{m}$ . The long wavelength edge of the feature appears to vary very little among these Miras. Hence the difference in spectral shape between the Miras and the other LPVs is primarily due to the fact that in Miras the rising branch varies in wavelengths accompanied by a shift in the peak emission to shorter wavelengths. This shift to shorter wavelength correlates with the strength of the silicate excess. In general the greater the strength of the feature the shorter the wavelength of the rising branch. Unlike the results of DeGioia-Eastwood et al (1981), we find that the strength correlates only very weakly with period. This corroborates the conclusion reached by Vardya, De Jong and Willems (1981). At a given period the excess can vary by a factor of 4. The  $18 \mu\text{m}$  emission feature is very similar in both types of profiles and extends from about  $15$  to  $>22 \mu\text{m}$ . Both these  $\sim 10 \mu\text{m}$  and  $18 \mu\text{m}$  features have been attributed to silicates.

The most interesting  $10 \mu\text{m}$  emission occurs in stars which tend to show weak excesses but includes a few stars which have excesses comparable in strength to the stars with the other types of features (see Fig. 1a and 1b). There are relatively few stars in this group, but they constitute almost half of the stars with weak emission irrespective of their variability type. The feature has three components (dotted line Figure 2) with peaks at  $10$ ,  $11$  and  $13.1 \mu\text{m}$ . The  $10 \mu\text{m}$  peak is strongest in M stars and agrees in wavelength with the silicate peak of the SRs but it is narrower. The intensities of the  $11$  and  $13.1 \mu\text{m}$  peaks vary greatly being at times quite weak. These stars show an emission excess at long wavelength which is significantly different from the  $18 \mu\text{m}$  emission. It appears to extend from about  $16$  to  $22 \mu\text{m}$  with a peak at about  $19.5 \mu\text{m}$ . If the 3 component feature is also due to silicates is not yet known. The peak at  $8 \mu\text{m}$  seen in Figure 2 appears to be an artifact of our method of analysis. It disappears if photospheric temperatures are used to fit the shortest wavelengths of the LRS. This research is supported in part by a University Resident Research Fellowship from the Air Force Office of Scientific Research to the Air Force Geophysics Laboratory.




Fig 2. The three types of  $10 \mu\text{m}$  excess.

#### REFERENCES

- DeGioia-Eastwood, K., Hackwell, J.A., Grasdalen, G.L., and Gehrz, R.D. 1981, *Ap.J.*, 245, L75.  
 Forrest, W.J., Gillett, F.C., and Stein, W.A. 1975, *Ap.J.*, 175, 423.  
 Vardya, M.S., De Jong, T., and Willems, F.J. 1986, *Ap.J.*, 304, L29.

END

10-86

DTIC