

AD-A172 038


A WELL-POSEDNESS PROPERTY OF A CLASS OF VARIATIONAL
PROBLEMS AND ITS APPL. (U) TEXAS UNIV AT AUSTIN DEPT OF
ELECTRICAL AND COMPUTER ENGINEER. J H MORRISON ET AL.
25 APR 86 AFOSR-TR-86-0794 AFOSR-81-0047 F/G 12/1

1/1

UNCLASSIFIED

NL





AD-A172 038

(2)

JRT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY NA			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE NA					
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 86 - 0794		
6a. NAME OF PERFORMING ORGANIZATION University of Texas at Austin		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION AFOSR/NM		
6c. ADDRESS (City, State and ZIP Code) Dept. of Electrical and Computer Engineering Austin, TX 78712			7b. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL (If applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-81-0047 and AFOSR-86-0026		
8c. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448			10. SOURCE OF FUNDING NOS.		
			PROGRAM ELEMENT NO. 6.1102F	PROJECT NO. 2304	TASK NO. 75
11. TITLE (Include Security Classification) A Well-Posedness Property of a Class of Variational Problems and Its Application to Nonlinear Estimation (UNCLASSIFIED)					
12. PERSONAL AUTHOR(S) J. M. Morrison and G. L. Wise					
13a. TYPE OF REPORT Reprint		13b. TIME COVERED FROM 10/1/80 to 10/31/86		14. DATE OF REPORT (Yr., Mo., Day) 1986 April 25	
15. PAGE COUNT 7					
16. SUPPLEMENTARY NOTATION Proceedings of the Twenty-Third Annual Allerton Conference on Communication, Control, and Computing, Monticello, Illinois, October 2-4, 1985, pp. 28-34.					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) nonlinear estimation, Orlicz spaces, convergence of estimators		
FIELD	GROUP	SUB. GR.			
19. ABSTRACT (Continue on reverse if necessary and identify by block number) In this paper we consider a class of nonlinear estimators that are the solutions to a certain variational problem. These estimators generalize the concept of conditional expectation; we investigate their continuity and convergence properties.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☐ DTIC USERS ☐					
21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED					
22a. NAME OF RESPONSIBLE INDIVIDUAL Major Brian W. Woodruff			22b. TELEPHONE NUMBER (Include Area Code) (202) 767-5027		22c. OFFICE SYMBOL AFOSR/NM

DTIC FILE COPY

DTIC
ELECTE
SEP 19 1986
S D

A WELL-POSEDNESS PROPERTY OF A CLASS OF VARIATIONAL PROBLEMS AND ITS APPLICATION TO NONLINEAR ESTIMATION

JOHN M. MORRISON
 Department of Mathematics
 University of Texas at Austin
 Austin, Texas 78712

Approved for public release;
 distribution unlimited.

and

GARY L. WISE
 Departments of Electrical and Computer Engineering and Mathematics
 University of Texas at Austin
 Austin, Texas 78712

ABSTRACT

In this paper we consider a class of nonlinear estimators that are the solutions to a certain variational problem. These estimators generalize the concept of conditional expectation; we investigate their continuity and convergence properties.

I. INTRODUCTION

In this paper we will develop some machinery that generalizes some well known approximation and continuity properties of minimum mean square error analysis. These will allow us to examine estimation theory in a much more general context and give us a much wider choice of criteria for analyzing and penalizing error. We will be required to forego some of the main convenient accoutrements of minimum mean square error analysis: the Hilbert space structure of L^2 , the projection theorem, and the fact that minimum norm projection onto a closed subspace is a nonexpansive linear operator. However, in return we receive a method that allows us to customize our notion of error to our particular model. Oftentimes, mean square error is not the appropriate fidelity criterion.

II. PRELIMINARIES

Throughout this paper let $(\Omega, \mathcal{F}, P)$ denote a probability space and let $L^0(\Omega, \mathcal{F}, P)$ denote the set of all random variables $X: \Omega \rightarrow \mathbb{R}$ modulo a.s. equivalence equipped with the topology of convergence in probability.

Let $\phi: [0, \infty) \rightarrow [0, \infty)$ be convex, increasing, and satisfy $\phi(0)=0$. We define the Orlicz space

$$L^\phi(\Omega, \mathcal{F}, P) = \{X \in L^0(\Omega, \mathcal{F}, P): \int_\Omega \phi(|X|) dP < \infty\}.$$

The function ϕ is said to satisfy the doubling condition if there exist $C, M > 0$ such that $x \geq M \Rightarrow \phi(2x) \leq C\phi(x)$. If ϕ satisfies the doubling condition, $L^\phi(\Omega, \mathcal{F}, P)$ is a vector space. Moreover, it is a Banach space with the Luxemburg norm

$$\|X\|_{L^\phi} = \inf \left\{ \lambda > 0: \int_\Omega \phi\left(\frac{|X|}{\lambda}\right) dP \leq \phi(1) \right\}.$$

Note that this is simply the Minkowski functional of a certain subset of L^ϕ . Also, if $\{X_n\}_{n=1}^\infty$ and X are in L^ϕ , $X_n \xrightarrow{L^\phi} X$ if and only if $\lim_{n \rightarrow \infty} \int_\Omega \phi(|X_n - X|) dP = 0$.

For the basic facts on Orlicz spaces see [3]. Details on Minkowski functions can be found in [1], pp. 294-295 or [6], pp. 23-26.

Let $\phi: [0, \infty) \rightarrow [0, \infty)$ be increasing and have a strictly increasing first derivative ϕ' with $\phi'(x) \rightarrow \infty$ as $x \rightarrow \infty$. Since $\phi: [0, \infty) \rightarrow [0, \infty)$ is a homeomor-

phism, $\psi = \phi^{-1}$ is also continuous, strictly increasing, and satisfies $\lim_{x \rightarrow \infty} \psi(x) = \infty$. We define the conjugate function Ψ to Φ to be

$$\Psi(x) = \int_0^x \psi(t) dt, \quad x \geq 0.$$

This is somewhat more restrictive than the definition employed in [3], but it eliminates certain annoying technicalities. It also allows us to dispense with the customary nonatomicity assumption made about the measure spaces used in this context. We call the pair Φ, Ψ an Orlicz pair.

The Banach space L^Φ is reflexive if and only if Ψ and Φ satisfy the doubling condition. A simple calculation shows that the strict convexity of Φ implies that of the Luxemburg norm on L^Φ and that of the convex functional $X \rightarrow \int_\Omega \Phi(|X|) dP$. In addition, if the Banach space L^Φ is reflexive, we have $L^{\Phi*} \cong L^\Psi$ [3]. Hereafter we posit all Orlicz spaces are reflexive.

Let B be a Banach space. We say B is (i) locally uniformly convex if whenever $\{x_n\}_{n=1}^\infty$ and $\{y_n\}_{n=1}^\infty$ are sequences in the unit ball of B with $\lim_{n \rightarrow \infty} \|x_n + y_n\| = 2$, $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$; (ii) uniformly convex if $\forall \epsilon > 0$ there exists $\delta > 0$ so that for any $x, y \in B$ so that $\|x\| = \|y\| = 1$, $\|x + y\| \geq 2(1 - \delta) \Rightarrow \|x - y\| < \epsilon$. Uniform convexity is a much stronger condition than local uniform convexity. Every uniformly convex space is reflexive [6], pp. 126-128. For more about these sorts of conditions, see [2].

If B is a Banach space, $Q \subseteq B$ is said to be proximal if $\forall x \in B$, the problem $\min_{z \in Q} \|x - z\|$

possesses a solution. If B is reflexive we may apply the Smul'lyan theorems to see that every closed convex subset is proximal. Moreover, if B is strictly convex the norm minimizer problem has at most one solution. In this case for $x \in B$ and $K \subseteq B$ that is closed and convex let $p_K(x)$ denote the unique solution to $\min_{z \in K} \|x - z\|$.

The map p_K is called the metric projection of B onto K . Shortly we will see that if B is locally uniformly convex and reflexive then for any closed convex $K \subseteq B$, p_K is norm continuous. In [5] we showed that p_K is weakly sequentially continuous if B is reflexive and strictly convex.

Let M be a separable metric space. A Borel measurable map $Q: M \rightarrow M$ is said to be a round off map if Q has finite range, say $\{p_1, \dots, p_n\} = \text{range } Q$ and $Q(p_k) = p_k$, $1 \leq k \leq n$. The set $\{Q^{-1}(p_1), \dots, Q^{-1}(p_n)\}$ is called the partition of M defined by Q . A sequence $\{Q_n\}_{n=1}^\infty$ of round off maps is called a round off scheme if (i) $\forall x \in M$, $\lim_{n \rightarrow \infty} \text{dia } Q_n^{-1}(Q_n(x)) = 0$; (ii) the partition of M defined by Q_{n+1} refines that defined by Q_n , $n \in \mathbb{N}$. Note $\sigma(Q_n) \subseteq \sigma(Q_{n+1})$, $n \in \mathbb{N}$. See [5] for details on round off schemes.

The primary result of [5] is contained in the Theorem 1: Let Φ, Ψ be an Orlicz pair, $(\Omega, \mathcal{F}, P)$ be a nonatomic probability space, M be a separable metric space and $\{X(t): t \in [0, T]\}$ be a stochastically continuous process on $(\Omega, \mathcal{F}, P)$ taking values in M . Then for any round off scheme $\{Q_n\}_{n=1}^\infty$ on M and any increasing sequence $\{P_m\}_{m=1}^\infty$ of partitions of $[0, T]$ whose meshes decrease to zero we have for any $Y \in L^\Phi(\Omega, \mathcal{F}, P)$



☒
☐
☐
Codes
or
11

A-1

$$\|E_{\Phi}(Y|Q_n(X(t)): teP_m) - E_{\Phi}(Y|X(t): te[0,T])\|_{L_{\Phi}} \rightarrow 0$$

as $m, n \rightarrow \infty$; or, equivalently,

$$0 = \lim_{m, n \rightarrow \infty} \int_{\Omega} \Phi(|E_{\Phi}(Y|Q_n(X(t)): teP_m) - E_{\Phi}(Y|X(t): te[0,T])|) dP$$

where $E_{\Phi}(\cdot|\mathcal{F}): L^{\Phi}(\Omega, \mathcal{F}, P) \rightarrow L^{\Phi}(\Omega, \mathcal{F}, P)$ denotes metric projection and $\mathcal{F}$ is any σ -subalgebra of $\mathcal{P}$.

III. A CONVEXITY CONDITION FOR CONVEX FUNCTIONALS ON REFLEXIVE BANACH SPACES

Let B be a reflexive Banach space and $\psi: B \rightarrow \mathbb{R}$ be a convex, norm continuous map satisfying

$$\lim_{\|x\| \rightarrow \infty} \psi(x) = +\infty.$$

Then using the Smul'lyan theorems we see that $\forall x \in B$

$$\min_{z \in K} \psi(x-z) \tag{1}$$

possesses a closed convex nonvoid subset L_x of K of solutions. Furthermore, if ψ is strictly convex each L_x is a singleton $\{p(x)\}$. We call the map $p: B \rightarrow K$ the solution map of (1). Let $\delta > 0$; define

$$K_{\delta, x} = \{z \in K: \psi(x-z) < \delta + \psi(x-p(x))\}.$$

We say ψ is locally uniformly convex if $\lim_{\delta \rightarrow 0} \text{dia } K_{\delta, x} = 0$ for all $x \in B$ and closed convex $K \subseteq B$.

Proposition 2: If $\psi: B \rightarrow \mathbb{R}$ is locally uniformly convex and uniformly continuous on bounded subsets of B , $K \subseteq B$ is nonvoid convex and closed and $p: B \rightarrow K$ is the solution map of (1) then p is norm continuous.

Proof: Let $x_n \rightarrow x$ in B and $\delta > 0$. Take $z \in K_{\delta, x}$. Then

$$\psi(x-z) < \delta + \psi(x-p(x)).$$

Since $x_n \rightarrow x$ there exists $N \in \mathbb{N}$ so that $n \geq N \Rightarrow$

$$\psi(x_n - z) < \delta + \psi(x_n - p(x)).$$

By minimality

$$\begin{aligned} \psi(x_n - p(x_n)) &\leq \psi(x_n - z) \\ &< \delta + \psi(x_n - p(x)). \end{aligned}$$

Taking limit suprema,

$$\limsup_{n \rightarrow \infty} \psi(x_n - p(x_n)) \leq \psi(x - p(x)). \tag{2}$$

Suppose by way of the contrapositive that $\{p(x_n)\}_{n=1}^{\infty}$ is unbounded.

Since $x_n \rightarrow x$, $\{x_n\}_{n=1}^{\infty}$ is bounded; this forces $\{x_n - p(x_n)\}_{n=1}^{\infty}$ to be unbounded.

Observing that $\lim_{\|x\| \rightarrow \infty} \psi(x) = +\infty$ it now follows that

$$\limsup_{n \rightarrow \infty} \psi(x_n - p(x_n)) = +\infty,$$

a flagrant violation of (2). This insures that $\{p(x_n)\}_{n=1}^{\infty}$ is bounded.

Suppose that for some $z_0 \in B$, $p(x_n) \rightarrow z_0$. Then $x_n - p(x_n) \rightarrow x - z_0$; ψ is convex and norm continuous and therefore weakly lowersemicontinuous. Thus

$$\begin{aligned}\psi(x-z_0) &\leq \liminf_{n \rightarrow \infty} \psi(x_n - p(x_n)) \\ &\leq \limsup_{n \rightarrow \infty} \psi(x_n - p(x_n)) \leq \psi(x - p(x)).\end{aligned}$$

Next observe that K is closed and convex and therefore weakly closed. Thus $p(x_n) \in K$, $n \in \mathbb{N}$, $p(x_n) \rightarrow z_0 \Rightarrow z_0 \in K$. By minimality, $z_0 = p(x)$.

Since B is reflexive and $\{p(x_n)\}_{n=1}^{\infty}$ is bounded each subsequence of $\{p(x_n)\}_{n=1}^{\infty}$ must have a further weakly convergent subsequence. But the foregoing discussion tells us that this further subsequence must converge to $p(x)$. Immediately it follows $p(x_n) \rightarrow p(x)$.

We may apply the weak lower semicontinuity of ψ to see that

$$\psi(x - p(x)) \leq \liminf_{n \rightarrow \infty} \psi(x_n - p(x_n)).$$

Combine this with (2) to get

$$\psi(x - p(x)) = \lim_{n \rightarrow \infty} \psi(x_n - p(x_n)).$$

Put $M = 1 + \sup_n \|x_n - p(x_n)\|$; note $M < \infty$. Take $\epsilon > 0$, choose $\eta > 0$ s.t. $\eta < 1$ and $\|x - y\| < \eta$, $\|x\|, \|y\| \leq M \Rightarrow |\psi(x) - \psi(y)| < \epsilon$. Because $x_n \rightarrow x$ there exists $N \in \mathbb{N}$ such that $n \geq N \Rightarrow \|x_n - x\| < \eta$. Then $n \geq N \Rightarrow$

$$\|x_n - p(x_n) - (x - p(x_n))\| = \|x_n - x\| < \eta$$

so $n \geq N \Rightarrow$

$$|\psi(x_n - p(x_n)) - \psi(x - p(x))| < \epsilon.$$

This forces $\lim_{n \rightarrow \infty} \{\psi(x_n - p(x_n)) - \psi(x - p(x_n))\} = 0$ and hence $\lim_{n \rightarrow \infty} \psi(x - p(x_n)) = \psi(x - p(x))$.

Noting $\lim_{n \rightarrow \infty} \psi(x - p(x_n)) = \psi(x - p(x))$ there exists $N \in \mathbb{N}$ so that $n \geq N \Rightarrow$

$$\psi(x - p(x_n)) < \delta + \psi(x - p(x)).$$

But this says that for $n \geq N$, $p(x_n) \in K_{\delta, x}$. Obviously $p(x) \in K_{\delta, x}$ so for $n \geq N$, $\|p(x_n) - p(x)\| \leq \text{dia } K_{\delta, x}$. Since $\lim_{\delta \rightarrow 0} \text{dia } K_{\delta, x} = 0$ we must have $p(x_n) \rightarrow p(x)$. QED

Proposition 3: Suppose $\{K_n\}_{n=1}^{\infty}$, K are nonvoid closed convex subsets of B

with $K_n \subseteq K_{n+1}$, $n \in \mathbb{N}$ and $K = \bigcup_{n=1}^{\infty} K_n$. Let $\psi: B \rightarrow \mathbb{R}$ be locally uniformly

convex and norm continuous. Finally denote by p_n the solution map of (1) corresponding to K_n , $n \in \mathbb{N}$ and p denote the solution map of K . Then for $x \in B$, $p_n(x) \rightarrow p(x)$.

Proof: Let $\delta > 0$, $x \in B$. By the continuity of ψ , $K_{\delta, x}$ is a relative neighborhood of $p(x)$ in K . Noting $K = \bigcup_{n=1}^{\infty} K_n$, we see there must be an $N \in \mathbb{N}$ so that $n \geq N \Rightarrow K_n \cap K_{\delta, x} \neq \emptyset$.

Fix $n \geq N$, $z \in K_{\delta, x} \cap K_n$. Then we know that (i) $\psi(x - z) < \delta + \psi(x - p(x))$;
(ii) $\psi(x - p_n(x)) \leq \psi(x - z)$. Assembling the facts, $n \geq N \Rightarrow$

$$\psi(x - p_n(x)) \leq \delta + \psi(x - p(x)).$$

Letting $n \rightarrow \infty$,

$$\limsup_{n \rightarrow \infty} \psi(x-p_n(x)) \leq \psi(x-p(x)). \quad (3)$$

We may imitate a familiar argument in Proposition 2 to see that $\{p_n(x)\}_{n=1}^{\infty}$ is bounded.

Since $K_n \subseteq K_{n+1} \subseteq K$, $n \in \mathbb{N}$ it is easy to see that

$$\psi(x-p_n(x)) \geq \psi(x-p_{n+1}(x)) \geq \psi(x-p(x)).$$

With the assistance of (3) we obtain

$$\psi(x-p(x)) = \lim_{n \rightarrow \infty} \psi(x-p_n(x)).$$

Pick $\delta > 0$. There exists $N \in \mathbb{N}$ such that $n \geq N \Rightarrow$

$$\psi(x-p_n(x)) < \psi(x-p(x)) + \delta.$$

We may now conclude $p_n(x) \in K_{\delta, x}$, $n \geq N$. Since $\text{dia } K_{\delta, x} \rightarrow 0$ as $\delta \rightarrow 0$ we conclude that $p_n(x) \rightarrow p(x)$ in norm. QED

IV. SOME FACTS ABOUT ORLICZ SPACES

Theorem 4: Let $(\Omega, \mathcal{P}, P)$ be a probability space and ϕ, ψ be an Orlicz pair.

Then $L^{\phi}(\Omega, \mathcal{P}, P)$ is uniformly convex if and only if

$\forall \epsilon$ such that $0 < \epsilon < \frac{1}{4}$ there exists $R_{\epsilon} > 0$ so that

$$\liminf_{n \rightarrow \infty} \frac{\psi(x)}{\psi((1-\epsilon)x)} > R_{\epsilon}.$$

Proof: This is the central result of [4].

Define $\xi: L^{\phi}(\Omega, \mathcal{P}, P) \rightarrow [0, \infty)$ by $\xi(X) = \int_{\Omega} \phi(|X|) dP$. Then ξ is a strictly convex norm continuous functional and $\lim_{\|X\|_{L^{\phi}} \rightarrow \infty} \xi(X) = +\infty$. Let $0 < \epsilon < 1$,

$X, Y \in L^{\phi}(\Omega, \mathcal{P}, P)$ and $X \neq Y$ a.s. $[P]$. Then $\|X-Y\|_{L^{\phi}} \neq 0$. Using the convexity of ϕ ,

$$\begin{aligned} \phi(|X-Y|) &= \phi\left(\epsilon \left(\frac{1}{\epsilon} |X-Y|\right) + (1-\epsilon)(0)\right) \\ &\leq \epsilon \phi\left(\frac{|X-Y|}{\epsilon}\right) + (1-\epsilon)\phi(0) \\ &= \epsilon \phi\left(\frac{|X-Y|}{\epsilon}\right). \end{aligned}$$

Integrating,

$$\int_{\Omega} \phi(|X-Y|) dP \leq \epsilon \int_{\Omega} \phi\left(\frac{|X-Y|}{\epsilon}\right) dP.$$

Now suppose $\|X-Y\|_{L^{\phi}} \leq 1$. Then

$$\begin{aligned} \int_{\Omega} \phi(|X-Y|) dP &\leq \|X-Y\|_{L^{\phi}} \int_{\Omega} \phi\left(\frac{|X-Y|}{\|X-Y\|_{L^{\phi}}}\right) dP \\ &= \|X-Y\|_{L^{\phi}} \phi(1), \end{aligned}$$

amply demonstrating the global uniform continuity of ξ .

We will not address the question of the local uniform convexity of ξ here. The hypothesis of Theorem 4 or possibly something similar might

guarantee the local uniform convexity of ξ .

V. A CONVERGENCE PRINCIPLE FOR A GENERALIZED CONDITIONAL EXPECTATION

Definition: Let $(\Omega, \mathcal{P}, P)$ be a probability space, $\mathcal{F}$ be a sub σ -algebra of $\mathcal{P}$ and ϕ, ψ be an Orlicz pair. For $X \in L^\phi(\Omega, \mathcal{P}, P)$, we define $\hat{E}_\phi(X|\mathcal{F})$ to be the unique solution to the variational problem

$$\min \left\{ \int_{\Omega} \phi(|X-Z|) dP : Z \in L^\phi(\Omega, \mathcal{F}, P) \right\}.$$

Recall from [5] that if $\phi(x) = \frac{x^2}{2}$ for $x \geq 0$, then $\hat{E}_\phi(\cdot|\mathcal{F})$ is ordinary conditional expectation given $\mathcal{F}$.

Remark: If the functional $X \mapsto \int_{\Omega} \phi(|X|) dP$ is locally uniformly convex then $\hat{E}_\phi(\cdot|\mathcal{F})$ is norm continuous on $L^\phi(\Omega, \mathcal{P}, P)$.

Theorem 5: Let $(\Omega, \mathcal{P}, P)$ be a probability space, $\{\mathcal{F}_n\}_{n=1}^{\infty}$ be an increasing family of sub σ -algebras of $\mathcal{P}$ and put $\mathcal{F} = \bigvee_{n=1}^{\infty} \mathcal{F}_n$. Suppose ϕ, ψ is an Orlicz pair so that $X \mapsto \int_{\Omega} \phi(|X|) dP$ is locally uniformly convex. Then for $X \in L^\phi(\Omega, \mathcal{P}, P)$,

$$\hat{E}_\phi(X|\mathcal{F}_n) \xrightarrow{L^\phi} \hat{E}_\phi(X|\mathcal{F}) \text{ as } n \rightarrow \infty,$$

or equivalently,

$$\lim_{n \rightarrow \infty} \int_{\Omega} \phi(|\hat{E}_\phi(X|\mathcal{F}_n) - \hat{E}_\phi(X|\mathcal{F})|) dP = 0.$$

Proof: This is an immediate consequence of Proposition 3. QED

We may now proceed as in [5] to get

Theorem 6: Let $(\Omega, \mathcal{P}, P)$ be a probability space, $\{X(t): t \in [0, T]\}$ be a process on $(\Omega, \mathcal{P}, P)$ taking values in a separable metric space M and ϕ, ψ be an Orlicz pair with $X \mapsto \int_{\Omega} \phi(|X|) dP$ locally uniformly convex. Then if $\{Q_n\}_{n=1}^{\infty}$ is a round off scheme for M and $\{P_m\}_{m=1}^{\infty}$ is an increasing sequence of partitions of $[0, T]$ whose meshes decrease to zero, we have for $Y \in L^\phi(\Omega, \mathcal{P}, P)$

$$\hat{E}_\phi(Y|Q_n(X(t)): t \in P_m) \xrightarrow{L^\phi} \hat{E}_\phi(Y|X(t): t \in [0, T])$$

as $m, n \rightarrow \infty$, or equivalently,

$$\lim_{m, n \rightarrow \infty} \int_{\Omega} \phi(|\hat{E}_\phi(Y|Q_n(X(t)): t \in P_m) - \hat{E}_\phi(Y|X(t): t \in [0, T])|) dP = 0.$$

VI. CONCLUSIONS AND A PARTING SHOT AT EARLIER WORK

The abstract principle developed in Section III subsumes the Banach space principle developed in [5]. Simply take the functional to be the norm.

We have enhanced the feasibility of studying the operators $\hat{E}_\phi$ numerically in two ways. First, knowing that $\hat{E}_\phi$ is continuous assures us that

it will tolerate small L^Φ -perturbations. Second, the result of Theorem 6 shows us we may approximate $\hat{E}_\phi(Y|X(t): t \in [0, T])$ by $\hat{E}_\phi(Y|Q_n(X(t)): t \in P_m)$. Since $\sigma(Q_n(X(t)): t \in P_m)$ is finite, calculation of $\hat{E}_\phi(Y|Q_n(X(t)): t \in P_m)$ consists of solving a finite dimensional nonlinear optimization problem. A large body of knowledge now exists about the numerical solution of such problems.

ACKNOWLEDGEMENTS

This research was supported by the Air Force Office of Scientific Research under Grants AFOSR-81-0047 and its successor. The authors would like to thank the Banach Space and Approximation Theory groups at the University of Texas at Austin for many helpful suggestions.

REFERENCES

1. Brown, A., and Pearcy, C., *Introduction to Operator Theory I*. New York: Springer-Verlag, 1977.
2. Day, M., *Normed Linear Spaces*. New York: Springer Verlag, 1973.
3. Krosnoselskii, M., and Rutickii, Y., *Convex Functions and Orlicz Spaces*. Groningen: P. Noordhoff, 1961.
4. Milnes, W., "Convexity of Orlicz Spaces," *Pacific Journal of Mathematics*, Vol. 7, No. 3, 1957, pp. 1451-1483.
5. Morrison, J., and Wise, G., "A Stability Property of Conditional Expectations," *Proc. Nineteenth Annual Conference on Information Sciences and Systems*, Baltimore, MD, March 27-29, 1985, pp. 226-229.
6. Yosida, K., *Functional Analysis*, 6th edition. New York: Springer-Verlag, 1980.
7. Zeidler, E., *Nonlinear Functional Analysis*, Vol. III. New York: Springer-Verlag, 1985.

END

10-86

DTIC