AD-A172 561 FINAL TECHNICAL REPORT ON CONTRACT F08671-83-0517 FOR 1 1/1 FEBRUARY 1983 - 31 JANUARY 1996(U) EVE RESEARCH INST OF RETINA FOUNDATION BOSTON MA L E AREND 30 MAR 86 UNCLASSIFIED AF0SR-TR-86-0845 F08671-83-0517 F/G 5/10 NL									
		•••							



ł

A MARTINE AND A CARACTERIA CARA

	BEPORTOCUM	ENTATION PAGE					
1. REPORT SECURITY OF ACCOUNTS	110	15. RESTRICTIVE M.	ARKINGS				
	ECTE	3. DISTRIBUTION/A	VAILABILITY C	OF REPORT			
AD-A172 561	0.1 1986	Approved fo	r public r	release;			
MU-MILL JUI		aistributio	n unlimite	=u			
L. UN NERDRINU	MBERD D	5. MONITORING OR	GANIZATION R	EPORT NUMBER	(S)		
		AFOSR	-18-8	6-084	5		
LAWRENCE E. ARENI	6b. OFFICE SYMBOL (If applicable)	78. NAME OF MONIT		NIZATION			
EYE RESEARCH INSTITUTE		AIR FORCE	RESEARC	r H/Ni			
6c. ADDRESS (City State and ZIP Code)		75. ADDRESS (City,	State and ZIP Co	ide)			
BOSTON, MA 02114		BUILDING 4	10 ID DC 000	30			
		BOLLING AF	в DC 203	52			
B. NAME OF FUNDING SPONSORING	86. OFFICE SYMBOL	9. PROCUREMENT	INSTRUMENT IS	DENTIFICATION	NUMBER		
AFOSR	NL	FQ8671-8	300517				
8c ADDRESS (City, State and ZIP Code)		10. SOURCE OF FUI	10. SOURCE OF FUNDING NOS.				
Building 410		PROGRAM ELEMENT NO	PROJECT NO.	TASK NO.	WORK UNI NO.		
Bolling AFB DC 20332							
11. TITLE (Include Security Classification)	61102F	2313	A5	N/A			
FINAL REPORT. TECHNICAL			L				
12. PERSONAL AUTHOR(S) AREND, 134. TYPE OF REPORT FINAL FROM 2	LAWRENCE E. covered /1/83 to 1/31/	14. DATE OF REPO 86 86 MAR	вт (Yr., Mo., Da 30	15. PAGE	COUNT		
12. PERSONAL AUTHOR(S) AREND, 134. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTATION	LAWRENCE E. covered /1/83 to 1/31/	14. DATE OF REPO 86 86 MAR	RT (Yr., Mo., Da 30	y) 15. PAGE	COUNT		
12. PERSONAL AUTHOR(S) AREND, 13. TYPE OF REPORT 13b. TIME FINAL FROM ½ 16. SUPPLEMENTARY NOTATION 17 COSATI CODES	LAWRENCE E. covered /1/83 to 1/31/	14. DATE OF REPO 86 86 MAR	RT (Yr., Mo., Da 30 necessary and ider	15. PAGE	E COUNT		
12. PERSONAL AUTHOR(S) AREND, 130. TYPE OF REPORT 130. TIME FINAL FROM 2 16. SUPPLEMENTARY NOTATION 17 COSATI CODES FIELD GROUP SUB. GR	LAWRENCE E. covered /1/83 to 1/31/ 18. Subject terms EYE MOVEME	14. DATE OF REPO 86 86 MAR Continue on reverse if n CNTS, SPATIAI	RT (Yr., Mo., Da 30 lecessary and ider	15. PAGE 15. PAGE 15. PAGE 15. PAGE	E COUNT		
12. PERSONAL AUTHOR(S) AREND, 134. TYPE OF REPORT 136. TIME FINAL FROM ½ 16. SUPPLEMENTARY NOTATION 17 COSATI CODES FIELD GROUP	LAWRENCE E. COVERED /1/83 to 1/31/ 18. SUBJECT TERMS EYE MOVEME VISUAL ILI	14. DATE OF REPO 86 86 MAR Continue on reverse if n NTS, SPATIAI JUSIONS, COL(RT (Yr., Mo., Da 30 eccessory and iden 2 PATTERN DR CONST/	15. PAGE 15. PAGE	COUNT		
12. PERSONAL AUTHOR(S) AREND, 13a. TYPE OF REPORT 13b. TIME FINAL FROM 2 16. SUPPLEMENTARY NOTATION 17 COSATI CODES FIELD GROUP SUB. GR 19. ABSTRACT (Continue on recerse if necessary)	LAWRENCE E. COVERED /1/83 to 1/31/ 18. SUBJECT TERMS EYE MOVEME VISUAL ILI and identify by block numi	14. DATE OF REPO 86 86 MAR "Continue on reverse if n NTS, SPATIAI JUSIONS, COLO	BT (Yr., Mo., Da 30 Recemery and Ider D PATTERN DR CONSTA	19) 15. PAGE 16. PAGE 16. PAGE 16. PAGE 15. PAGE 15. PAGE 15. PAGE 15. PAGE 15. PAGE 15. PAGE 15. PAGE 16. PAGE 1	E COUNT		
12. PERSONAL AUTHOR(S) AREND, 13a. TYPE OF REPORT 13b. TIME FINAL FROM 2 16. SUPPLEMENTARY NOTATION 17 COSATI CODES FIELD GROUP SUB. GR 19. ABSTRACT (Continue on release of necessary) LOW CONTrast, low Dattorns	LAWRENCE E. COVERED /1/83 to 1/31/ 18. SUBJECT TERMS EYE MOVEME VISUAL ILL and identify by block number v spatial frech ance staircas	14. DATE OF REPO 86 86 MAR Continue on reverse if n NTS, SPATIAI JUSIONS, COLO Deri quency lumina Ses. with no	BT (Yr. Mo., Da 30 Recessory and ider PATTERN DR CONSTA ance saw1 brightne	15. PAGE 1817y by block num VISION, ANCY, COLO tooth ess change	COUNT DR VISION es over t		
I2. PERSONAL AUTHOR(S) AREND, 13b. TIME FINAL FROM 2 16. SUPPLEMENTARY NOTATION 17 COSATI CODES FIELD GROUP SUB GR 19. ABSTRACT (Continue on receive if necessary) Low contrast, low patterns look like lumin shallower luminance slor	LAWRENCE E. COVERED /1/83 to 1/31/ 18. SUBJECT TERMS EYE MOVEME VISUAL ILL and identify by block number visual frechance staircas be. Brightnes	14. DATE OF REPO 86 86 MAR "Continue on reverse if n CNTS, SPATIAI USIONS, COLO Der, guency lumina ses, with no as measurement	BT (Yr. Mo., Da 30 PATTERN DR CONSTA ance sawt brightne nts at co	15. PAGE 11. 15. PAGE 11. 17 by block num 1 VISION, ANCY, COLO tooth ess change orrespond	COUNT CR VISION es over t ing point		
AREND, AREND, 130. TIME FINAL FROM 2 16. SUPPLEMENTARY NOTATION 17 COSATI CODES FIELD GRCUP SUB GR 19 ABSTRACT (Continue on receive if necessary) Low contrast, low patterns look like lumir shallower luminance slop in different cycles of t	LAWRENCE E. COVERED /1/83 to 1/31/ 18. SUBJECT TERMS EYE MOVEME VISUAL ILL and identify by block number v spatial frequence staircas be. Brightness these patterns	14. DATE OF REPO 86 86 MAR Continue on reverse if n NTS, SPATIAL JUSIONS, COLO Deri guency lumina ses, with no ss measuremet s showed sub-	RT (Yr. Mo., De 30 eccessory and iden DATTERN DR CONSTA ance sawt brightne nts at co stantial	VISION, ANCY, COLO tooth ess change orrespond: illusory	COUNT CR VISION es over t ing point brightne		
12. PERSONAL AUTHOR(S) AREND, 13. TYPE OF REPORT 13b. TIME FINAL FROM 2 16. SUPPLEMENTARY NOTATION 17 COSATI CODES FIELD GROUP SUB_GR 19. ABSTRACT (Continue on release if necessary) Low contrast, low patterns look like lumin shallower luminance slop in different cycles of t differences. Other meas strictly_Subthroshold	LAWRENCE E. COVERED /1/83 to 1/31/ 18 SUBJECT TERMS EYE MOVEME VISUAL ILI and identify by block number visual frequences be Brightness these patterns surements show impance gradi	14. DATE OF REPO 86 86 MAR Continue on reverse if n NTS, SPATIAL USIONS, COLO Der, Juency lumina Ses, with no as measurements showed sub- wed that the ients but of	RT (Yr. Mo., Da 30 eccessory and iden D PATTERN DR CONSTA ance sawt brightne nts at co stantial illusion ccur with	y) 15. PAGE 10/1/2 by block num VISION, ANCY, COLO tooth ess change orrespond: illusory n is not o h slightly	COUNT CR VISION es over t ing point brightne confined y supra-		
12. PERSONAL AUTHOR(S) AREND, 13a. TYPE OF REPORT 13b. TIME FINAL FROM 2 16. SUPPLEMENTARY NOTATION 17 COSATI CODES FIELD GROUP SUB_GR 19. ABSTRACT (Continue on referse if necessary) Low contrast, low patterns look like lumin shallower luminance slop in different cycles of t differences. Other meas strictly subthreshold lu threshold gradients as w	LAWRENCE E. COVERED /1/83 to 1/31/ 18. SUBJECT TERMS EYE MOVEME VISUAL ILL and identify by block number v spatial frequences these patterns Surements show uminance gradi vell.	14 DATE OF REPO 86 86 MAR Continue on reverse if n NTS, SPATIAL JUSIONS, COLO Der, quency lumina ses, with no s measurements s showed subsissed wed that the ients, but of	RT (Yr. Mo. De 30 eccenary and iden D PATTERN DR CONSTA ance sawt brightne nts at co stantial illusion ccur with	y) 15. PAGE NUTY by block num VISION, ANCY, COLO tooth ess change orrespond: illusory n is not o h slightly	COUNT COUNT DR VISION es over t ing point brightne confined y supra-		
12. PERSONAL AUTHOR(S) AREND, 13. TYPE OF REPORT 13b. TIME FINAL FROM 2 16. SUPPLEMENTARY NOTATION 17 COSATI CODES FIELD GROUP SUB_GR 19. ABSTRACT (Continue on referent if necessary) Low contrast, low patterns look like lumin shallower luminance slop in different cycles of t differences. Other meas strictly subthreshold lu threshold gradients as w In models which a	LAWRENCE E. COVERED /1/83 to 1/31/ 18. SUBJECT TERMS EYE MOVEME VISUAL ILL and identify by block number v spatial frequences these patterns surements show uminance gradi vell. attempt to exp	14 DATE OF REPO 85 86 MAR Continue on reverse if n NTS, SPATIAL USIONS, COLO Deri quency lumina Ses, with no ss measurements showed subside wed that the tents, but of on these	RT (Yr. Mo. Da 30 eccessory and iden DATTERN DR CONSTA ance sawt brightne nts at co stantial illusion ccur with illusion	y) 15. PAGE 10/y by block num VISION, ANCY, COLO tooth ess change orrespond: illusory n is not o h slightly s the visu	COUNT CR VISION OR VISION DR VISION brightne confined y supra- ual syste		
AREND, AREND, 13a TYPE OF REPORT 13b TIME FINAL FROM 2 16. SUPPLEMENTARY NOTATION 17 COSATI CODES FIELD GROUP SUB GR 19. ABSTRACT (Continue on receive if necessary) Low contrast, low patterns look like lumin shallower luminance slop in different cycles of t differences. Other meas strictly subthreshold lu In models which a integrates over the three	LAWRENCE E. COVERED /1/83 to 1/31/ 18. SUBJECT TERMS EYE MOVEME VISUAL ILL and identify by block numble v spatial frequences ance staircas be. Brightness these patterns surements show uminance gradi well. attempt to expense esholded grad: ers problems of	14. DATE OF REPO 86 86 MAR Continue on reverse if n NTS, SPATIAL USIONS, COLO Deri- guency lumina ses, with no ss measurements showed subside wed that the tents, but of obtain these ient of the due to curl	RT (Yr. Mo. De 30 eccenary and iden DATTERN DR CONSTA ance sawt brightne nts at co stantial illusion ccur with illusions stimulus introduce	y) 15. PAGE 1607 by block num VISION, ANCY, COLO tooth ess change orrespond: illusory n is not of h slightly s the visit distributed ed by the	COUNT Count DR VISION Prightne confined y supra- ual syste tion. nonlinea		
12. PERSONAL AUTHOR(S) AREND, 13a. TYPE OF REPORT 13b. TIME FINAL FROM 2 16. SUPPLEMENTARY NOTATION 17 COSATI CODES FIELD GRCUP SUB. GR 19. ABSTRACT (Continue on receive if necessary) Low contrast, low patterns look like lumin shallower luminance slop in different cycles of t differences. Other meas strictly subthreshold lu threshold gradients as w In models which a integrates over the three The integration encounted threshold operator.	LAWRENCE E. COVERED /1/83 to 1/31/ 18. SUBJECT TERMS EYE MOVEME VISUAL ILL and identify by block number v spatial frequences these patterns surements show uminance gradi vell. attempt to expension esholded grad: ers problems of ghtness measure	¹⁴ DATE OF REPO <u>85</u> 86 MAR ^{(Continue on reverse if n NTS, SPATIAL USIONS, COLO ^{Jer} Juency lumina Ses, with no ses measurements showed subs wed that the lents, but of blain these ient of the due to curl rements indi}	RT (Yr. Mo. Da 30 eccessory and iden PATTERN DR CONSTA ance sawt brightne nts at co stantial illusion ccur with illusion stimulus introduce cated the	y) 15. PAGE 160/y block num VISION, ANCY, COLO tooth ess change orrespond: illusory n is not of h slightly s the visit distribut ed by the at these	COUNT COUNT DR VISION PS OVER t ing point brightne confined y supra- ual syste tion. nonlinea problems		
AREND, AREND, 130. TYPE OF REPORT 130. TIME FINAL FROM 2 16. SUPPLEMENTARY NOTATION 17 COSATI CODES FIELD GROUP SUB GR IS ABSTRACT (Continue on release if necessary of Low contrast, low patterns look like lumin shallower luminance slop in differences. Other meas strictly subthreshold luthreshold gradients as w In models which a integrates over the three	LAWRENCE E. COVERED /1/83 to 1/31/ 18. SUBJECT TERMS EYE MOVEME VISUAL ILL and identify by block number visual frequences be Brightness these patterns surements show uninance gradi vell. attempt to expension sholded grad; ers problems of phtness measure rt, further su	¹⁴ DATE OF REPO <u>86</u> 86 MAR ^{(Continue on reverse if n (NTS, SPATIAL USIONS, COLO ^(Continue on reverse if n) (USIONS, COLO ^(Continue on reverse if n) ^{(Continue on reverse}}	RT (Yr. Mo. Da 30 eccessory and iden D PATTERN DR CONSTA ance sawt brightne nts at co stantial illusion ccur with illusions stimulus introduce cated the he model:	y) 15. PAGE 100/y block num VISION, ANCY, COLO tooth ess change orrespond: illusory n is not of h slightly s the visit distributed ed by the at these s.	DR VISION DR VISION DS OVER t ing point brightne confined y supra- ual syste tion. nonlinea problems		
AREND, AREND, 136. TYPE OF REPORT TIGE TYPE OF REPORT TYPE OF REPORT TYPE OF REPORT TIGE TYPE OF REPORT TIME TIME TYPE OF REPORT TYPE OF REPORT TIGE TYPE OF REPORT TYPE OF REPORT <td colspan="2" of="" repor<="" td="" type=""><td>LAWRENCE E. COVERED /1/83 to 1/31/ 18. SUBJECT TERMS EYE MOVEME VISUAL ILL and identify by block number r spatial frequence staircas be. Brightness these patterns surements show uminance gradi vell. attempt to expension ers problems of ghtness measur rt, further su</td><td>^{(Continue on reverse if n SS 86 MAR (Continue on reverse if n NTS, SPATIAL JUSIONS, COLO (Jerr) (uency lumina Ses, with no ses, with no ses, with no ses, with no ses, with no ses, with no ses, with no late to curl cements, but of blain these lue to curl cements indi- upport for t}</td><td>RT (Yr. Mo. De 30 eccenary and iden DATTERN DR CONSTA ance sawt brightne nts at co stantial illusions cur with illusions stimulus introduce cated the he model:</td><td>y) 15. PAGE 16. If y block num VISION, VISION, ANCY, COLO tooth ess change orrespond: illusory n is not of h slightly s the visit distributed by the at these y s.</td><td>COUNT CO</td></td>	<td>LAWRENCE E. COVERED /1/83 to 1/31/ 18. SUBJECT TERMS EYE MOVEME VISUAL ILL and identify by block number r spatial frequence staircas be. Brightness these patterns surements show uminance gradi vell. attempt to expension ers problems of ghtness measur rt, further su</td> <td>^{(Continue on reverse if n SS 86 MAR (Continue on reverse if n NTS, SPATIAL JUSIONS, COLO (Jerr) (uency lumina Ses, with no ses, with no ses, with no ses, with no ses, with no ses, with no ses, with no late to curl cements, but of blain these lue to curl cements indi- upport for t}</td> <td>RT (Yr. Mo. De 30 eccenary and iden DATTERN DR CONSTA ance sawt brightne nts at co stantial illusions cur with illusions stimulus introduce cated the he model:</td> <td>y) 15. PAGE 16. If y block num VISION, VISION, ANCY, COLO tooth ess change orrespond: illusory n is not of h slightly s the visit distributed by the at these y s.</td> <td>COUNT CO</td>		LAWRENCE E. COVERED /1/83 to 1/31/ 18. SUBJECT TERMS EYE MOVEME VISUAL ILL and identify by block number r spatial frequence staircas be. Brightness these patterns surements show uminance gradi vell. attempt to expension ers problems of ghtness measur rt, further su	^{(Continue on reverse if n SS 86 MAR (Continue on reverse if n NTS, SPATIAL JUSIONS, COLO (Jerr) (uency lumina Ses, with no ses, with no ses, with no ses, with no ses, with no ses, with no ses, with no late to curl cements, but of blain these lue to curl cements indi- upport for t}	RT (Yr. Mo. De 30 eccenary and iden DATTERN DR CONSTA ance sawt brightne nts at co stantial illusions cur with illusions stimulus introduce cated the he model:	y) 15. PAGE 16. If y block num VISION, VISION, ANCY, COLO tooth ess change orrespond: illusory n is not of h slightly s the visit distributed by the at these y s.	COUNT CO
AREND, AREND, I30. TYPE OF REPORT ISD. TIME FINAL FROM 2 ISD. TIME FIELD GROUP SUB. GR IN ABSTRACT (Continue on receive if necessary of LOW contrast, low patterns look like lumin shallower luminance slop in different cycles of t differences. Other meas strictly subthreshold lut threshold gradients as w In models which a integrates over the three	LAWRENCE E. COVERED /1/83 to 1/31/ 18. SUBJECT TERMS EYE MOVEME VISUAL ILL and identify by block number > spatial frequences ance staircas be. Brightness these patterns surements show uminance gradi vell. attempt to expenses sholded gradi ers problems of phtness measure of, further staircas PT. GOTIC USERS []	14 DATE OF REPO <u>85</u> 86 MAR Continue on reverse if n NTS, SPATIAL USIONS, COLO Ver, (uency lumina Ses, with no ss measurements showed subs ved that the lents, but of blain these lent of the lent of the lent of the lent of the lent of the lent of the lent of	RT (Yr. Mo. Da 30 eccessory and iden PATTERN DR CONSTA ance sawt brightne nts at co stantial illusion ccur with illusions stimulus introduce cated the he models CURITY CLASSI ied	y) 15. PAGE 1607 PAGE 1607 PAGE 1607 PAGE 1607 PAGE 1607 PAGE 1607 PAGE 15. PAGE 16. PAGE 15. PAGE 15. PAGE 15. PAGE 15. PAGE 15. PAGE 15. PAGE 15. PAGE 15. PAGE 15. PAGE 16. PAGE 15. PAGE 15. PAGE 16. PA	COUNT COUNT DR VISION PS OVER ti ing point brightne confined y supra- ual syste tion. nonlinea problems		
AREND, AREND, I30. TYPE OF REPORT ISD. TIME FIELD GROUP SUB.GR IP ABSTRACT (Continue on receise if necessary) LOW CONTRAST, low patterns look like lumin Shallower luminance slop In different cycles of t differences. Other meas strictly subthreshold lu threshold gradients as v In models which a integrates over the three The integration encountee threshold operator. Lig have a visual counterpar 20. DISTRIBUTION/AVAILABILITY OF ABSTI UNCLASSIFIED/UNLIMITED SAME AS RI 222. NAME OF RESPONSIBLE INDIVIDUAL	LAWRENCE E. COVERED /1/83 to 1/31/ IB. SUBJECT TERMS EYE MOVEME VISUAL ILL and identify by block number / spatial frequences be Brightness these patterns Surements show Iminance gradi vell. attempt to expense sholded gradi ers problems of phtness measure of the sure as a sub- should the sub- ers problems of the sure as a sub- the sure as a sub- control of the sub- should the sub- sub- should the sub- should the sub- s	14. DATE OF REPO 85 86 MAR (Continue on reverse if n NTS, SPATIAL USIONS, COLO Per) [uency lumina Ses, with no is measurements showed subsy wed that the lents, but of Dain these lient of the lient o	RT (Yr. Mo., De 30 eccenary and iden _ PATTERN DR CONSTA ance sawt brightne nts at co stantial illusion ccur with illusions stimulus introduce cated that he models CURITY CLASSI ied	y) 15. PAGE 16. If y block num VISION, ANCY, COLO tooth ess change orrespond: illusory n is not of h slightly s the visit distributed by the at these y s. IFICATION	DR VISION DR VISION DR VISION DR VISION Drightne confined y supra- ual syste tion. nonlinea problems		

.

SECURITY CLASSIFICATION OF THIS PAGE

19. (cont.)

> Several new illusion were found to result from this nonlinear threshold for spatial gradient, including patterns where gradients were perceived in uniform regions and nonuniform stimulus regions were perceived as uniform.

Color constancy was measured for multiple regions of the visual field under different illuminants. Little hue constancy was observed, but subjects could accurately predict the appearance of a colored surface under the various illuminants. The results were little affected by the number of colored surfaces in the scene.



Accesio	on For				
NTIS DTIC Unann Justific	CRA&I TAB ouriced sation				
By Distrib	ution/				
Availability Codes					
Dist	Avail and for Special				
A-1	، ۲۰۰ ۲				



SECURITY CLASSIFICATION OF THIS PAGE

AFOSR . TR. 86-0845

١

FINAL TECHNICAL REPORT

CONTRACT PQ8671-8300517

2/1/83 - 1/31/86

Eye Research Institute 20 Staniford Street Boston, MA 02114

Principal Investigator: Lawrence E. Arend

Affinite and the second of the second (NESC) Approved for an is distributed and is the second second

FQ8671-8300517

REPORT: 2/1/83 TO 1/31/86

I. OBJECTIVES

The overall objective of the project was further development of a model (Arend, 1973) describing the role of retinal image motion in the human visual system's processing of edge information in patterns. Several lines of investigation were proposed, including:

A. Eye Movements and Stabilized Retinal Images

Recent experiments using grating patterns held stationary on the retina have raised questions concerning the role of retinal image motion in spatial pattern vision. The fact that high contrast gratings remain visible when stabilized with the SRI Double Purkinje Image Eyetracker (Kelly, 1979) has been interpreted as meaning that the human visual system can process pattern information in the absence of temporal change of the retinal image, contradicting previous views. This issue is a critical part of the most basic understanding of human pattern vision. One goal of the project was stabilization work in which we 1) replicated Kelly's basic observations of stabilized gratings and 2) reinterpreted all grating stabilization data in light of new calculations of the sensitivity of the human visual system to small image motion.

B. Spatial Gradients and Integration over Edge Information

A large component of the project was investigation of how information from sharp and shallow spatial gradients of intensity and color gets assembled into a stable and reliable visual array of lightness and color. Goals for this portion of the project were 1) assembly and calibration of the display system consisting of minicomputer, image processor and color video monitor 2) software development for displaying patterns and controlling experiments and 3) experiments on gradient illusions.

C. Lightness and Color Constancy in Complex Patterns

The third major goal was investigation of lightness and color constancy in complex patterns. One of the properties of the model is encoding of information about relative lightnesses and colors. Relatively little work has been done on color perception with complicated spatial patterns. Further development of the model requires information about the roles of several parameters, including complexity (number of different colors under single illuminant, range of different colors) and spatial distribution of reflectances and illuminants.

II. STATUS OF RESEARCH EFFORT

A. Stabilized Retinal Images

Substantial progress was made in evaluating Purkinje Image Eyetracker stabilization of gratings. The principal conclusions are:

1) Most of the gross form of Kelly's grating stabilization data replicates in our lab using a similar eyetracker, different subjects and improved methods.

2) Calculations from stabilized grating contrast sensitivities show that the P.I. tracker (in fact, probably no existent eyetracker) can eliminate all meaningful retinal image motion and therefore:

a) Residual visibility of stabilized objects does not necessarily imply that the visual system contains mechanisms capable of responding to temporally unchanging inputs and
b) Only subjective pattern fading and disappearance can be relied upon as a criterion of stabilization adequacy, and then only with qualification.

In light of current evidence, therefore, it is likely that all pattern information is derived from retinal temporal changes. We therefore must have a model like that under investigation here, describing the extraction of spatial pattern information from temporal information.

A paper describing our stabilized grating experiments and motion-sensitivity calculations was published in the Journal of the Optical Society of America A.

B. Spatial Gradients and Integration of Edge Information

1. Display System. During the first year a display system consisting of a VAX 11/750 minicomputer, an Adage 3000 image processor and a Tektronix 690 RGB monitor was selected, purchased, and installed. The display system was calibrated in the second year to provide accurate specification of luminance and chromaticity over the full output range of the monitor.

The spectral emitances of the red, green, and blue phosphors of the monitor were obtained by a grating-monochromator multichannel analyzer and the CIE chromaticity coordinates of the phosphors were calculated (fig. 1).

Luminance vs. digital-input-data curves were obtained for the three guns singly and together at nine screen locations. Unlike published data on other monitors (Cowan, 1984) the curves for all three guns were well fit by a single power function (after multiplication by a constant equating the maximum outputs for the guns). Thus the outputs can all be linearized with one power function. The function in use linearizes the light output to better than one percent at all values (fig. 2). The output of the phosphors is stable; repeatability is excellent from week to week. Figure 1. Relative power spectra of phosphors of Tektronix 690 monitor. Right peak: red phosphor. Middle peak: green phosphor. Left peak: blue phosphor.

Fig. 2. a) Luminance of Tektronix 690 phosphors as function of digital data applied to D/A converters. Scaled to maximum luminance of white (all three guns at maximum)=94.0 Cd/m^2 . Curves, top to bottom: white (all guns on), green, red, blue.

b) Curves of fig. 2a each scaled to its own maximum luminance. Power function is the same for all three guns.

c) Luminance curves of phosphors after linearization, as function of image data level. Each curve scaled to its own maximum.



ŀ

CARACTER.



.

SACAL COURSE SELECTION STRUCTURE PUBLICATION

2. <u>Measurement of Gradient Illusions</u>. The first step in studying gradient illusions (Craik-O'Brien-Cornsweet and related illusions) was development of satisfactory techniques for measuring the appearance of these complex spatial patterns.

We explored several techniques and used them to obtain more quantitative detail than previously available A wide variety of shallow gradient patterns yield illusory brightnesses. We chose initially to study a circular ring pattern with a radial luminance distribution consisting of three cycles of a low contrast sawtooth (fig. 3). Even though the three cycles have identical luminance the figure appears as a nearly uniform brightness central disk surrounded by two nearly uniform rings of increasing (3a) or decreasing (3b) brightness relative to the central disk.

The subject could vary the luminance of a small square in the corner of the display to make its brightness match the centers of the inner and outerbands (fig. 3, 1 and 2).

In figs. 4 and 5 are brightness matches from two subjects for several sawtooth contrasts in each polarity. While there are important differences of detail the subjects generally agree that substantial brightness differences occurred at all contrasts. This was a somewhat surprising result. In the simplest form of the model the illusory brightness differences require that the shallow gradient be below threshold and not perceived. At our lowest contrast the bands were very uniform in brightness and the edges had low apparent contrast. At the highest physical contrast the rings had pronounced brightness gradients between high contrast edges. It was initially somewhat surprising therefore that large brightness differences occurred even in the presence of the visible gradient. In an attempt to quantify the appearances at the various contrasts the subjects were asked to judge the fraction ofthe ring which appeared uniform in brightness. While there are obviously criterion definition problems with this task, in practice the judgements were not difficult and had reasonable error bars. Means of five judgements are shown in figs. 6 and 7. Both subjects judged the lowest contrast patterns to have only narrow regions of nonuniformity adjacent to the contours. At the highest contrast both saw brightness gradients over most of the ring, but RG adopted a less strict criterion than LA.

Upon closer consideration there are two simple possible explanations for the brightness differences at the highest contrasts: 1) Even though less of each ring appears uniform, the uniform portion spans a greater luminance range due to the higher physical contrast. 2) The threshold for gradient need not be a step function wherein increases of luminance gradient change abruptly from no visibility to weight equal to that given steep gradients; a more reasonable function would be sigmoid with physical gradients passing gradually from no representation of shallow gradients. While the second option seems likely, the data in figs. 4-7 are also consistent with the first.

The graded threshold account is, however, supported by a further experiment. Patterns with the radial luminance profiles

Figure 3. Luminance and brightness as a function of radial distance. Inset: display configuration. Left: luminance along radius of inset (zero = center). Right: sketch of brightness distribution along radius of inset pattern. Subject adjusted luminance of rectangle in inset pattern to match brightness at points 1 and 2.

Figure 4. Brightness matches for patterns of fig. 3 as a function of pattern contrast $(=(L_{max}-L_{min})/(L_{max}+L_{min}))$. Circles, top: point 1, top configuration of fig. 3. Circles, bottom: point 2, top configuration. Squares, top: point 2, bottom configuration of fig. 3. Squares, bottom: point 1, bottom configuration. Subject LA.

Figure 5. Same as fig. 4, subject RG.

Figure 6. Estimates of percentage of width of band which appeared uniform in brightness as a function of pattern contrast. Patterns of fig. 3, top configuration. Circles: outer band. Squares: inner band. Subject LA.

Figure 7. Same as fig. 6, subject RG.









personal manager pressess pressess and the





Fig. 3









فللمتعادية

CARACTER PERSONAL







of fig. 8 have no subthreshold gradients if the small steps are detectable. Nevertheless, for patterns with many small steps per ring there is a clear brightness difference between the bands. In fig. 9 are brightness matching data from two subjects for 5-11 steps per ring. In all the stimuli the small steps were visible. No brightness differences occurred for few steps, but clear differences occurred at 9 and 11 steps. The brightness differences in fig. 9 clearly indicate that the sum of the small brightness steps within the rings did not equal the sum of the large brightness steps between rings. We therefore conclude that the threshold for spatial gradient is not abrupt but graded. More detailed experiments will allow determination of the form of the weighting curve transferring luminance gradients to brightness gradients.

3. Gradient and Curl A major issue proposed for second and third year study was inconsistent integrals in two dimensional applications of Land and McCann's (1971) and Arend's (1973) essentially one-dimensional models. The problem is illustrated in fig. 10. According to both models the brightness of B relative to that of A is obtained by the visual system by taking the gradient of log retinal illuminance, setting all gradients below some threshold to zero and integrating over the resulting Explicitly in Land and McCann's model and distribution. implicitly in Arend's the differentiation and integration are done along arbitrarily chosen curves between A and B. The models make the correct prediction along the straight line joining A and B in fig. 10. The shallow gradient is below threshold so the only derivatives going into the integral are the downward steps at the contours, indicating that B is darker than A. Along the path crossing the outer boundary, however, we get a different answer. There is no change between A and the outer boundary, a downward step of known magnitude to the background luminance, no change over the background segment, an upward step on crossing back into B's band equal and opposite to the initial crossing into the background, and no change between the outer boundary and The sum of these derivatives is zero indicating that A and B в. are equal in brightness.

These inconsistent integrals will occur with any model which accounts for gradient illusions by taking derivatives or differences, thresholding, and then summing or integrating. This includes second-derivative models like Horn (1974) and Marr's (1974). In all cases the presence of path-dependent line integrals means there is no true two-dimensional integral and some special process must be proposed for resolving the inconsistencies (unless one wishes to propose that the appearance of such patterns is unstable or multistable, changing over time, or perhaps beyond description at any given time).

Not all patterns produce inconsistent integrals in the models. The radial sawtooth distributions of fig. 3, for example, produce no inconsistent integrals. All paths between a point in the inner ring and a point in the outer ring cross the same subthreshold and suprathreshold gradients. It seemed possible that distributions with inconsistent integrals might produce smaller illusory brightness differences and larger Figure 8. Cube root of luminance as function of radial position.

Figure 9. a) Brightness matches of centers of three bands of pattern of fig. 8 as function of pattern contrast (same definition as fig. 4). Top curve: inner band. Middle curve: middle band. Bottom curve: outer band. Subject AR.

b) Same as fig. 9a, subject LA.

۰.

.



brightness gradients than consistent patterns. To test this hypothesis we matched brightnesses in radial-sawtooth disks (see 4) and radial-sawtooth half disks (fig. 11). The latter fig. are merely curved versions of the pattern in fig. 10 and have Data from two subjects the same inconsistent integral problems. As hypothesized the half-disk are shown in figs. 12 and 13. brightness difference (and hence the size of the illusion) is consistently smaller than for the full-disk for both subjects. The difference is less pronounced for RG than for LA, and background luminance had a larger effect on LA's half-disk data than RG's.

Analysis of inconsistent integral problems led us to two new paradoxical patterns, the circumferential sawtooth pattern and the drifting sawtooth illusion.

As noted in the original proposal, one way of resolving the problems in the integration is to perceive brightness gradients which reduce or remove the curl responsible for the inconsistent integrals, i.e., essentially, lower one's threshold for the spatial luminance gradients. While no quantitative data have yet been collected documenting the appearance of the full and half ring patterns, their appearances are consistent with this type of solution. Brightness gradients are much more pronounced in the half-ring patterns than in the corresponding full-ring patterns.

There is, however, at least one other method of resolving The offending curl occurs where the border with the the problem. background slices across the subthreshold luminance gradient as, for example, at the top and bottom of fig. 10. If the integration is carried out only over areas which do not cross this border there are no inconsistent integrals. Thus the could be dealt with by segmenting the visual field into problem separate areas of integration, each internally consistent. The visual problem then reduces to combining the output of the separate integrations to give a brightness distribution extending over the entire visual field. This segmentation could resemble the segmentation required for veridical perception of areas under very different illuminations.

Some evidence that the visual system is sensitive to the geometrical relations among subregions of the image comes from observations of "circumferential sawtooths", displays with a sawtooth luminance distribution around a circumference rather In this distribution a line than along a radius (fig. 14). integral extending from point A around the circumference back to A is not zero (indicating that A is brighter than A or that A is darker than A, depending on the direction of rotation). Visually this pattern is unstable (or possibly multistable) with the brightness distribution varying from moment to moment, depending upon such factors as eye movements and point of Two adjacent sectors frequently look different in fixation. brightness and relatively homogeneous within sectors (though never as homogeneous as in the bands in fig. 3 even when the luminance gradient is less). The other two sectors, however, are nonuniform and their brightness relative to the other three sectors remains vague, even under close attention. If one sector is now replaced by the dark background there is still curl at the Figure-10. Schematic of integrals computed in brightness models along various paths through two sawtooth luminance patterns. a) Left panel: Linear sawtooth pattern. Right panel: Sketch of integral of thresholded derivative along paths 1 and 2 between points A and B. b) Left panel: Radial sawtooth pattern. Right panel: Sketch of integral of thresholded spatial derivative of luminance along paths 3 and 4 between points C and D. Shallow luminance gradient is below threshold, abrupt steps are not.

Figure 11. Display for brightness matches to half of radial sawtooth pattern. Subject adjusts luminance of rectangle to match brightness of center of inner ring or outer ring.

Figure 12. Brightness matches of centers of rings of radial sawtooth luminance patterns as function of luminance of surround $(L_{max}=94.0 \text{ Cd/m}^2)$. a) Complete pattern. Squares: inner ring. Circles: outer ring. b) Half pattern. Squares: inner ring. Circles: outer ring. Subject LA.

Figure 13. Same as Fig. 12, subject RG.



Fig. 10

F.'g. 11

Figure 14. Luminance as a function of angular position in annular circumferential sawtooth pattern. Dashed line: path of problematic integral of thresholded spatial derivative of luminance in brightness models. Shallow gradient within sectors is below threshold, abrupt steps are not. Thus integral after 360° counterclockwise rotation is sum of four downward steps, indicating that brightness of A does not equal brightness of A.

Figure 15. Sketch of truncated circumferential sawtooth pattern.

Figure 16. CIE diagram indicating Red/Green isoluminous gratings of saturation CSF experiment. Closed curve: spectrum locus. Triangle: color space of Tektronix monitor, corners are phosphor loci. Upper point: experimentally determined unique green. Lower point: D_{6500} white of monitor. Gratings were formed by modulating sinusoidally as indicated in the figure along the line passing through unique green and D_{6500} .

Figure 17. Logarithm of inverse of excitation purity of green peak of just detectable isoluminous chromaticity grating as function of spatial frequency. Purity = (green peak - D_{6500})/(510 nm - D_{500})



No.

AND DESCRIPTION AND DESCRIPTION AND ADDRESS AND ADDRESS ADDRESS ADDRESS ADDRESS ADDRESS ADDRESS ADDRESS ADDRESS



inner and outer borders of each sector (fig. 15). Nevertheless, the three remaining sectors immediately appear uniform and different in brightness.

We have also discovered a remarkable illusion in which stripes successively drift onto the display, each stripe darker than the previous, but with no change in the overall brightness of the display. Two cycles of a low contrast luminance sawtooth initially fill the screen, appearing quite homogeneous with the left brighter than the right. The entire pattern is then slowly drifted to the left with new cycles of sawtooth appearing on the right edge to replace the portions disappearing at the left. Each cycle as it appears is darker than those to its left. Under passive viewing there is no obvious change in the brightness of the band as it moves across the screen, giving the overall impression that darker bars are being added to the screen. It is only with mental effort that one notices that the overall brightness of the screen is not changing. Under careful scrutiny the brightness of the individual cycle can be seen to increase during its traverse.

These experiments with inconsistent integral patterns provide further demonstrations that the visual system represents patterns not in terms of local luminances or luminance ratios, but in terms of relatively global integrals over spatial derivative information.

C. Color Constancy

Color constancy has been attributed to three distinct mechanisms: adaptation, edge-ratio normalization, and perceptual processes such as discounting the illuminant.

Clearly, color constancy may partly result from adaptation, whereby the sensitivities of the chromatic mechanisms of the visual system change over time in response to the changed illuminant. Such processes have been modelled by the Von Kries coefficient law.

Constancy may, however, also stem from simultaneous mechanisms, such as normalization within neural edge-ratio computations like those of the Arend model or Land's retinex theory which operate independently of temporal (fast or slow) adaptation. The empirical foundation for theories of simultaneous color constancy is, however, rather sparse.

Finally, in the Helmholzian tradition, color constancy may also be due to 'discounting the illuminant.' In this case the local hue, saturation and lightness are affected by a change of illuminant, but the apparent surface color remains constant due to accurate perception of the illuminant change.

We conducted experiments to examine the relative contributions of the latter two mechanisms to color constancy. Our subjects viewed computer simulations of colored paper arrays. A standard patch of colored paper was displayed under one illuminant and a test patch under a different illuminant. The subject was asked either to adjust the test patch to match its hue and saturation to those of the standard patch (hue match) or to adjust the test patch to look as if it were "cut from the same piece of paper" as the standard, i.e., to match its apparent surface color (paper match). The two tasks were performed with two different spatial paper arrays.

A number of authors have argued that color constancy depends upon the complexity of the scene. Calculation of the unknown illuminant and unknown surface spectral reflectances from the retinal image may strongly depend upon the number of different color patches viewed. To examine this hypothesis we used cente -surround arrays (two surfaces) and complex arrays (32 surfaces) (1'g. 16).

Observers matched patches (simulated Munsell papers) in two simultaneously- presented computer-controlled displays, a standard array presented under 6500 K illumination and a test array under 4000 or 10000 K. The adjusted patch was surrounded by a single color (annulus display) or by many colors (mondrian display). Observers made either hue and saturation matches, or 'paper' matches, in which the subject was asked to make the test patch look as if it were cut from the same piece of paper as the standard patch.

Results for the annulus condition are shown in CIE (x,y) coordinates in Figure 17. The stimuli are plotted in the same fashion in each panel, with the (x,y) coordinates of the 6500 K standard patches that were matched represented by circles. Coordinates of patches under 4000 K and 10000 K are indicated by squares and triangles, respectively. The mean chromaticities set by the observers are indicated by closed symbols with error bars of +/- one standard error. The open symbols indicate the chromaticity of the paper being matched under the test illuminants. In this representation of the data, perfect color constancy would occur if the subject set the test patch chromaticity to the chromaticity the standard patch would have under the test illuminant, i.e., if open and closed symbols coincided. Results for the hue match are shown on the left of Figure 17, and those for the paper match on the right. The paper matches show quite good color constancy for most of the test patches, although the shift is typically slightly erroneous in The hue match data show much less magnitude or direction. constancy by comparison; the data depart much less from the 6500 deg chromaticity being matched, though the small shifts obtained are typically in the general direction of constancy. Similar results are shown for the mondrian patterns in Figure 18. As with the annuli, the hue matches shifted substantially less than required by color constancy.

In summary, we found little simultaneous color constancy for hue and saturation for any of our spatial configurations, when we asked for direct hue and saturation matches. The 4000 K and 10000 K test illuminants were always easily discriminated from the 6500 K standard display and from each other. All of the test pattern's hues shifted systematically from those of their standard pattern counterparts, i.e., there was a substantial preservation of illuminant information.

Pattern complexity had little effect on the amount of simultaneous color constancy. The hue matches in the mondrian condition showed no more deviation from the 6500 K chromaticity point than in the annulus condition.

These experiments do not rule out the possibility that



10YR 5/4 R 5/2 P 5/10 G 5/2 108 5/4 BG 10 P B 5/6 5/2 Y RP 5/4 10 P R 5/10 GΥ 5/10-5/12 В 5/4 R 5/6 5/8 YR 5/4 10BG 5/4 GΥ G P 5/4 5/4 PB 5/6 101 10 G 5/6 5/4 5/4 В Ν YR Y 5/2 5/ 10 RP 5/10 5/6 5/2 Ρ BG 5/2 10R 5/6 5/4 RР 5/6 10GY 5/6

7. 16

STANDARD

TEST



ľ.























shallow chromatic illumination gradients may be unperceived or underweighted perceptually, producing partial or full simultaneous color constancy for hue matches in such scenes.

Under our viewing conditions the illumination difference was always clearly perceptible. We therefore reject any model which inflexibly discards, through some normalizing process, information about abrupt illumination changes within a single scene. The fact that humans can compute reflectance information while retaining some illumination information indicates that the visual system includes a process capable of parsing retinal illuminance changes into overlaid illumination and reflectance gradient fields. Further experiments will be required to delineate the responsible mechanisms.

REFERENCES

- Arend, L. E. (1973) Spatial and differential operators in human vision. **Psychol. Rev., 80,** 374-395.
- Horn, B. K. P. (1974) Determining lightness from an image. Computer Graphics and Image Processing, 3, 277-299.
- Kelly, D. H. (1979) Motion and Vision. II. Stabilized spatiotemporal threshold surface. J. Opt. Soc. Amer., 69, 1340-1349.
- Kelly, D. H. (1981) Disappearance of stabilized chromatic gratings. <u>Science</u>, 214, 1257-1258.
- Land, E. H. and McCann, J. J. (1971) Lightness and Retinex theory. J. Opt. Soc. Amer., 61, 1-11.
- Marr, D. (1974) The computation of lightness by the primate retina. Vision Research, 14, 1377-1388.

III. PAPERS

- Arend, L.E. "Filling-In" between edges. <u>Behav. Brain Sci.</u> 6, 657, 1983.
- Arend, L.E. Spatial gradient illusions and inconsistent integrals. <u>Invest. Ophthal. & Vis. Sci.</u>, April Supple., 280, 1985.
- Arend, L.E. and Timberlake, G.T. What is psychophysically perfect image stabilization? Do perfectly stabilized images always disappear? J. Opt. Soc. Amer. A, 3, 235-241, 1986.
- Arend, L.E. and Goldstein, R. Lightness constancy in complex patterns. <u>Invest. Ophthal. & Vis. Sci.</u>, April Supple., 280, 1986.

Arend, L. and Reeves, A. Simultaneous color constancy. Submitted to J. Opt. Soc. Amer. A.

- Arend, L.E. and Goldstein, R. Spatial gradientillusionsand inconsistent integrals. In Preparation. Probably submitted to <u>Perception and Psychophysics.</u>
- Arend, L.E. Cornsweet illusion without subthreshold gradients. In Preparation. Probably submitted to Vision Research.
- Arend, L.E. and Goldstein, R. Lightness constancy in complex patterns. In Preparation. Probably submitted to <u>J. Opt.</u> <u>Soc. Amer. A.</u>
- Arend, L. E. and Lange, R. V. Apparent contrast of suprathreshold spatiotemporal gratings. (Probably to be submitted to Vision Research).

IV. PROFESSIONAL PERSONNEL

Arend, Lawrence E., Principal Investigator

Goldstein, Robert, Research Assistant

Timberlake, George T., nonsalaried part-time collaborator

Reeves, Adam, nonsalaried part-time collaborator

V. PROFESSIONAL INTERACTIONS

- Smith-Kettlewell Institute for Visual Sciences, colloquium on eyetracker stabilization, November, 1983.
- Stanford Research International, informal presentation of eyetracker data to C. Burbeck and D. Kelly, November, 1983.
- Brown University, Psychology Department Colloquium, "Inconsistent integrals in brightness illusions", November, 1984.
- Visual Information Processing Research Review, AFOSR, "Stabilized Retinal Gratings", May, 1984.
- Arend, L.E. "Spatial gradient illusions and inconsistent integrals," Paper presented at Spring meeting of Assoc. for Research in Vision and Ophthal., April, 1985.
- Invited talk at AFOSR Human Resources Laboratory, Williams AFB, AZ, "Color in complex displays," Feb., 1986.

VI. INVENTIONS

There were no patentable inventions under this project.

LUU PRASSEN DEPUTY RELACED DEPENDENT SEEDEN

