

AD-A179 117

MOLECULAR MODELING IN DRUG DESIGN FOR THE DEVELOPMENT
OF ORGANOPHOSPHORUS ANTIDOTES/PROPHYLACTICS(U) ARIZONA
UNIV TUCSON DEPT OF BIOCHEMISTRY W CHIU 30 APR 85
DAND17-84-C-4110


1/1

UNCLASSIFIED

F/G 6/15

NL





MICROCOPY RESOLUTION TEST CHART
100% COPY

AD-A179 117

28

2

AD _____

Molecular Modeling in Drug Design for the
Development of Organophosphorus Antidotes/Prophylactics

DTIC FILE COPY

Annual Report

Wah Chiu, Ph.D.

April 30, 1985

Supported by

U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND
Fort Detrick, Frederick, Maryland 21701-5012

Contract No. DAMD17--84-C-4110

University of Arizona
Tucson, Arizona 85721

DOD DISTRIBUTION STATEMENT

Approved for public release; distribution unlimited

The findings in this report are not to be construed as
an official Department of the Army position unless so
designated by other authorized documents.

DTIC
ELECTE
APR 09 1987
S D
E

87-4-7-845

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		7a. NAME OF MONITORING ORGANIZATION	
6a. NAME OF PERFORMING ORGANIZATION University of Arizona	6b. OFFICE SYMBOL (if applicable)	7b. ADDRESS (City, State, and ZIP Code) Dept. of Biochemistry, Bio Sciences West Tucson, Arizona 85721	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION U.S. Army Medical Research & Development Command	8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DAMD17-84-C-4110	
8c. ADDRESS (City, State, and ZIP Code) Fort Detrick Frederick, Maryland 21701-5012	10. SOURCE OF FUNDING NUMBERS		
	PROGRAM ELEMENT NO. 61102A	PROJECT NO. 3M161102BS11	TASK NO. EB WORK UNIT ACCESSION NO. 024
11. TITLE (Include Security Classification) Molecular Modeling in Drug Design for the Development of Organophosphorus Antidotes/ Prophylactics			
12. PERSONAL AUTHOR(S) Wah Chiu, Ph.D.			
13a. TYPE OF REPORT Annual Report	13b. TIME COVERED FROM 84/5/1 TO 85/4/30	14. DATE OF REPORT (Year, Month, Day) 1985 April 30	15. PAGE COUNT 14
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
06	03	Acetylcholinesterase, Computer, Electron Microscopy	
07	03	Crystal, Structure	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Our long-term goal is to design drug against the acetylcholinesterase inhibitors by molecular modeling techniques and theoretically predicted action of drug. Prior to this effort, we need to have more knowledge of the three-dimensional structure of the acetylcholinesterase. In the current year, we have tested various biochemical procedures for purifying acetylcholinesterase from electrical eel in the form suitable for structural analysis by electron diffraction method. Preliminary result suggests the feasibility of crystallizing the acetylcholinesterase. New computer algorithm has been implemented for processing electron micrograph data for structural investigation. <i>Key words</i>			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION	
22a. NAME OF RESPONSIBLE INDIVIDUAL Mrs. Virginia Miller		22b. TELEPHONE (Include Area Code) 301/663-7325	22c. OFFICE SYMBOL SGRD-RMS

Summary

Our long-term goal is to design drug against the acetylcholinesterase inhibitors by molecular modeling techniques and theoretically predicted action of drug. Prior to this effort, we need to have more knowledge of the three-dimensional structure of the acetylcholinesterase. In the current year, we have tested various biochemical procedures for purifying acetylcholinesterase from electrical eel in the form suitable for structural analysis by electron diffraction method. Preliminary result suggests the feasibility of crystallizing the acetylcholinesterase. New computer algorithm has been implemented for processing electron micrograph data for structural investigation.

Application For	
NEWS RELEASE	<input checked="" type="checkbox"/>
DISPATCH	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Mail and/or Special
A-1	



Foreword

Citations of commercial organizations and trade names in this report do not constitute an official Department of the Army endorsement or approval of the products or services of these organizations.

Table of Contents

	Page
Cover Page	i
DD Form 1473	ii
Summary	iii
Foreword	iv
Table of Contents	1
Annual Report	
(I) Purification and preliminary structural studies of electric eel acetylcholinesterase	2
Introduction	2
Experimental Methods	3
Results	3
Conclusion	4
(II) Installation of Major Equipment	5
(III) Research Plan for Year 2	5
(IV) References	5
Figure 1. SDS-polyacrylamide gel electrophoresis of electric eel AchE	7
Figure 2. Native electrophoresis of electric eel AchE	7
Figure 3. Electron micrograph of purified AchE	8
Figure 4. Light micrograph of AchE crystals	9
Distribution List	10

Annual Report for May 1, 1984 - April 30, 1985

Contract No. DAMD 17-84-C-4110

Name of Contractor: Arizona Board of Regents
University of Arizona, Tucson, AZ 85721

Project Title: Molecular Modeling in Drug Design for the Development of
Organophosphorus Antidotes/Prophylactics

Term of Contract: May 1, 1984 - April 30, 1987

Principal Investigator: Dr. Wah Chiu
Telephone No. (602)-621-7524 or 621-1773

(I) Purification and preliminary structural studies of electric eel
acetylcholinesterase

Introduction

Acetylcholinesterase is an enzyme which plays an integral role in the function of cholinergic synapses. This enzyme catalyzes the hydrolysis of the neurotransmitter acetylcholine to choline and acetic acid.

Several molecular forms of acetylcholinesterase have been identified. Asymmetric forms have been identified in the electric organs of Torpedo and Electrophorus and in mammalian muscle and brain (Dudaí et al., 1973; Rieger et al., 1973; Lwebuga-Mukasa et al., 1976; Rieger et al., 1980). These molecules consist of clusters of globular catalytic subunits which are anchored to the basement membrane by a collagen-like tail. At least three different sizes differing in the number of catalytic subunits have been identified by density gradient centrifugation.

A second form of the enzyme has been identified in Torpedo electric organ (Bon and Massoulie, 1980; Viratelle and Bernhard, 1980; Lee et al., 1982) and mammalian erythrocyte membranes (Grossman and Lieflander, 1975; Ott et al., 1975). This form of the enzyme consists of a disulfide linked dimer of catalytic subunits which is anchored to the plasma membrane. Early evidence indicated that a small hydrophobic peptide anchored the esterase to the membrane, but recent evidence suggests that the enzyme is linked directly to phosphatidylinositol (Stieger et al., 1984; Stieger and Brodbeck, 1985; Futerman et al., 1985).

The structural information available on acetylcholinesterase is limited to biochemical characterizations and several electron micrographs (for review see Rosenberry, 1975; Massoulié and Bon, 1982). Density gradient centrifugation and electron microscopy established the asymmetric nature of the 9S, 14S, and 18S forms and the apparent tetrameric nature of the 11S form produced by proteolysis of the asymmetric forms (Dudai *et al.*, 1973; Rieger *et al.*, 1976). Circular dichroism studies and an amino acid analysis revealed the collagen-like nature of the tail fragment of the asymmetric forms (Mays and Rosenberry, 1981). Amino acid analysis of the globular 11S form has also been reported (Leuzinger and Baker, 1967; Bon *et al.*, 1976). Peptide mapping has demonstrated that the catalytic subunits of the dimeric membrane form and the globular 11S form are different (Lee *et al.*, 1982). Crystallization of the 11S form and preliminary X-ray diffraction results have been reported, but no structural information has been obtained (Leuzinger *et al.*, 1968; Chothia and Leuzinger, 1976).

Experimental Methods

Numerous acetylcholinesterase purification schemes have been published. Nearly all involve affinity purification using quaternary ammonium esterase inhibitors as the affinity ligand (Berman and Young, 1971; Massoulié and Bon, 1976; Webb and Clark, 1978; Ralston *et al.*, 1983). The methods developed by these investigators provided the groundwork from which we have developed our purification procedure. We have so far focused our effort in the purification and structural determination of the 11S form of the enzyme.

The purified enzyme is evaluated by SDS and nondenaturing polyacrylamide gels. Electron microscopy of the purified enzyme was done initially in a Hitachi electron microscope in the College of Agriculture and subsequently in our newly acquired Philips 420 electron microscope.

We have set up several crystallization conditions of different pH, ionic strength and protein concentration with our purified AChE (McPherson, 1976). Light microscopy was used to monitor the growth of the crystals.

We have been developing a set of computer programs for 3 dimensional reconstruction of noncrystalline objects and of objects with a helical symmetry. We expect to use these computer algorithms to determine the molecular structures of the catalytic subunits of acetylcholinesterase and of the tail part of this enzyme which appears to have a helical arrangement.

Results

Crystallization of proteins for structural determination requires milligram quantities of pure protein (McPherson, 1976). These quantities of acetylcholinesterase are available commercially and, therefore, several of these preps were evaluated. Sigma, Worthington, and Boehringer Mannheim esterase preparations of 11S AChE were analyzed on both SDS and nondenaturing polyacrylamide gels. Multiple bands were observed in all of these preps, even after further purification of the esterase by affinity chromatography in our laboratory. Figures 1a and 2b show the banding pattern observed from a Sigma prep. The SDS gel pattern suggests some proteolytic degradation during the preparation by Sigma.

In order to better control the treatment of the esterase, we developed a purification procedure using electric organ from Electrophorus electricus as the starting material. This procedure involves affinity purification and trypsinization of the asymmetric forms to produce a more soluble globular 11S form. The best preparation we have obtained shows two major bands in a polyacrylamide gel stained for esterase activity (Figure 2a). The SDS gel pattern of the same prep shows a single major band around 70,000 daltons (Figure 1b). The performance of the affinity columns has been found to be somewhat erratic. We have found that contaminating proteins sometimes co-purify with the esterase. Some of the heterogeneity observed in our preps by gel electrophoresis (as in Figure 1c) may be the result of nonuniform tryptic cleavage. Standardization of the tryptic digestion procedure and optimization of our affinity column performance will alleviate these problems. Progress is being made in this regard. Our purification protocol should be capable of providing the quantities of pure acetylcholinesterase of 11S form needed for the structural studies.

We examined some of our purified enzyme of 11S in the electron microscope. Figure 3 is a typical micrograph of this preparation and shows various sizes. These may represent various types of aggregates of the monomer. More work is needed to sort out whether the aggregation is inherent to this form of the enzyme or whether it represents an artifact resulting from specimen preparation techniques.

In our preliminary trials, we have obtained several crystal forms of 11S AchE as shown in the light microscope. Figure 4 shows the largest crystal to be 300 microns on one edge of the crystal. Unfortunately, this crystal form is too big for electron diffraction and too small for X-ray diffraction analysis. We are certain that this crystal is formed by the enzyme because we have run an activity gel of the protein prior to its use for crystallization. We plan to refine the crystallization procedure to make the crystal suitable for electron diffraction analysis.

We have completed the implementation of the computer program for reconstructing the 3 dimensional structure of an object with a helical symmetry. We tested this program initially with simulated data and subsequently with an electron micrograph of a T4 phage tail which is known to have helical symmetry (DeRosier and Klug, 1968). The reconstruction of T4 phage tail agrees well with the published results. This confirms the correctness of our computer algorithm. We also used this set of programs to test a presumably helical object of unknown structure. The reconstruction result looks quite promising. With this algorithm providing the groundwork, we will develop in the next year the computer software necessary for the 3-dimensional reconstruction of an object with no crystallographic symmetry.

Conclusion

In summary, we have accomplished our proposed task in year 01 to demonstrate our capability of purifying the eel acetylcholinesterase, to record a preliminary electron image of this enzyme and to show the feasibility of forming a crystal from this enzyme.

(II) Installation of Major Equipment

We have installed two major instruments which are essential for the proposed research. The first instrument is the Perkin Elmer 1010M microdensitometer which was installed in July 1984. The instrument was tested by our professional staff to be performing as specified by the manufacturer. We have used it to digitize our electron microscopic data prior to its processing by computer. The second instrument is the Philips 420 electron microscope which was installed in March 1985. This equipment is performing as specified at room temperature. We are waiting for the delivery of the specimen rotation holder and the cold specimen holder from the manufacturer around midsummer 1985.


(III) Research Plan for Year 02 (May 85 - April 86)

We plan to refine our biochemical preparation procedures to make it a reliable and reproducible one, to scale up for the production of a large quantity of this enzyme for structural study, and to refine the crystallization condition of this enzyme. We will try to optimize the tryptic treatment of the whole enzyme. We plan to improve the affinity column used for the purification step. We will try to use different columns as suggested by the COTR and some other investigators (Ralston *et al.*, 1983; Webb and Clark, 1978). The variations of the affinity resins to be tested include different sizes of spacer arms and different types of ligands. For example, procainamide and acridinium will be used as ligands in the column. The possible spacer arms for trial are diaminodipropylamine, hexanediamine and aminocaproic acid-diaminopropane. We plan to continue the development of computer software for reconstructing 3 dimensional structure of individual protein molecules. We anticipate collection of preliminary 3 dimensional data of this enzyme either in noncrystalline or in crystalline form with our newly acquired electron microscope.

(IV) References

- Berman, J. D. and Young, M. (1971) *Proc. Nat. Acad. Sci. (U.S.A.)* 68: 395.
- Bon, S. and Massoulie, J. (1980) *Proc. Nat. Acad. Sci. (U.S.A.)* 77: 4464.
- Bon, S., Huet, M., Lemonnier, M., Rieger, F. and Massoulie, J. (1976) *Eur. J. Biochem.* 68: 523.
- DeRosier, D. and Klug, A. (1968) *Nature* 217: 130.
- Dudai, Y., Herzberg, M. and Silman, I. (1973) *Proc. Nat. Acad. Sci. (U.S.A.)* 70: 2473.
- Futerman, A. H., Fiorini, R. M., Roth, E., Low, M. G. and Silman, I. (1985) *Biochem. J.* 226: 369.
- Grossmann, H. and Lieflander, M. (1975) *Hoppe-Seyler's Z. Physiol. Chem. Bd.* 356: 663.

- Lee, S. L., Camp, S. J. and Taylor, P. (1982) *J. Biol. Chem.* 257: 12302.
- Leuzinger, W. and Baker, A. L. (1967) *Proc. Nat. Acad. Sci. (U.S.A.)* 57: 446.
- Lwebuga-Mukasa, J. S., Lappi, S. and Taylor, P. (1976) *Biochem.* 15: 1425.
- Massoulie, J. and Bon, S. (1976) *Eur. J. Biochem.* 68: 531.
- Massoulie, J. and Bon, S. (1982) *Ann. Rev. Neurosci.* 5: 57.
- Mays, C. and Rosenberry, T. L. (1981) *Biochem.* 20: 2810.
- McPherson, A. J. (1976) *Methods of Biochem. Anal.* 23: 249.
- Ott, P., Jenny, B. and Brodbeck, U. (1975) *Eur. J. Biochem.* 57: 469.
- Ralston, J. S., Main, A. R., Kilpatrick, B. F. and Chasson, A. L. (1983) *Eur. J. Biochem.* 211: 243.
- Rieger, F., Bon, S., Massoulie, J. and Cartaud, J. (1973) *Eur. J. Biochem.* 34: 539.
- Rieger, F., Bon, S., Massoulie, J., Cartaud, J., Picard, B. and Benda, P. (1976) *Eur. J. Biochem.* 68: 513.
- Rieger, F., Chetelat, R., Nicolet, M., Kamal, L. and Pouillet, M. (1980) *FEBS Lett.* 121: 169.
- Rosenberry, T. L. (1975) *Adv. Enzym.* 43: 103.
- Stieger, S. and Brodbeck, U. (1985) *J. Neurochem.* 44: 48.
- Stieger, S., Brodbeck, U., Reber, B. and Brunner, J. (1984) *FEBS Lett.* 168: 231.
- Viratelle, O. M. and Bernhard, S. A. (1980) *Biochem.* 19: 4999.
- Webb, G. and Clark, D. G. (1978) *Arch. Biochem. Biophys.* 191: 278.



1a

1b

1c

2a

2b

Figure 1

Figure 2

Figure 1: SDS-Polyacrylamide Gel Electrophoresis of Electric Eel AChE. Polyacrylamide gel stained for protein with Coomassie Blue. Lane a: AChE (Sigma) after further purification; specific activity = 4000 U/mg. Lane b: AChE prepared from electric organ (this laboratory); specific activity = 4000 U/mg. Lane c: Second preparation of AChE from electric organ (this laboratory); specific activity = 2000 U/mg. 1 Unit = 1 mole acetylthiocholine/min.

Figure 2: Native Electrophoresis of Electric Eel AChE. Polyacrylamide gel stained for acetylcholinesterase activity. Lane a: AChE prepared from electric organ in this laboratory. Lane b: AChE purified in this laboratory starting from a commercial preparation (Sigma).




FIGURE 3



Figure 4: Light Micrograph of Acetylcholinesterase Crystals.

DISTRIBUTION LIST

12 copies Director
Walter Reed Army Institute of Research
Walter Reed Army Medical Center
ATTN: SGRD-UWZ-C
Washington, DC 20307-5100

1 copy Commander
US Army Medical Research and Development Command
ATTN: SGRD-RMI-S
Fort Detrick, Frederick, Maryland 21701-5012

2 copies Defense Technical Information Center (DTIC)
ATTN: DTIC-DDAC
Cameron Station
Alexandria, VA 22304-6145

1 copy Dean
School of Medicine
Uniformed Services University of the
Health Sciences
4301 Jones Bridge Road
Bethesda, MD 20814-4799

1 copy Commandant
Academy of Health Sciences, US Army
ATTN: AHS-CDM
Fort Sam Houston, TX 78234-6100

E M D

5-87

Dt i c