



N G LEV RESULTION ES HAR

ŗ

DTK EILE CULLY

SPECTROSCOPIC CHARACTERIZATION OF TITANIUM DOPED SAPPHIRE

AD-A180 345

Ð

\$

Final Report



submitted by

Professor Renata Reisfeld Department of Inorganic and Analytical Chemistry The Hebrew University of Jerusalem Jerusalem 91904 Israel

to

US Army Contract No. DAJA 45-85-M-0344

1 October 1985 - 30 September 1987

DISTRIBUTION STATEMENT A Approved for public release Distribution Unlimited

REPORT DOCUMENT						Form	Approved
			ΔD_Δ180 345				ate Jun 30, 19
UNCLASSIFIED	IFICATION		AD-A	100	UTU		
28 SECURITY CLASSIFICATION	N AUTHORITY		J DISTRIBUTION	AVAILABILIT	Y OF REP	DRT	
2b. DECLASSIFICATION / DOW	NGRADING SCHEDU	LE	APPROVED FOR PUBLIC RELEASE;				
NOT APPLICABLE	ION REPORT NUMBE	R(S)	5 MONITORING	ORGANIZATI	DN REPOR	NUMBER	5)
			R&D 4697-M	MS-01 FIN	AL REPO	RT	
6. NAME OF PERFORMING	ORGANIZATION	6b. OFFICE SYMBOL	7a. NAME OF MO	ONITORING C	RGANIZAT	ION	
THE HEBREW UNIVERS	SITY OF JERUSA	LEM (" Oppicedic)	USA	RDSG-UK			
6c. ADDRESS (City, State, an	d ZIP Code)		76. ADDRESS (Cit	ty, State, and	ZIP Code)		
DEPT. OF INORGANIC JERUSALEM 91904	C CEHMISTRY		FPC	O. BOX 65 O NEW YOR	K, 0951	0-1500	
ISRAEL							INADED
Ba. NAME OF FUNDING/SPC ORGANIZATION	JNSORING	(If applicable)	9. PROCUREMIEN				JIVIBER
EUROPEAN RESEARCH	OFFICE	USARDSG-UK	CON	NTRACT #	DAJA 45	5-85-M-0	344
8c. ADDRESS (City, State, and	d ZIP Code)		PROGRAM	PROJECT	TAS	ĸ	WORK UNI
223 OLD MARYLEBONI	E RD.		61102A 11	NO.	157-12	•	ACCESSION
11. TITLE (Include Security (Classification)						
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13a. TYPE OF REPORT FINAL	IC CHARACTERIZ	COVERED	UM DOPED SAP	PHIRE" ORT (Year, M	onth, Day)	15. PAGE	COUNT
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13a. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTA	IC CHARACTERIZ REISFELD 13b. TIME (FROM	COVERED	UM DOPED SAP	PHIRE"	onth, Day)	15. PAGE	COUNT
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13a. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTA NONE	IC CHARACTERIZ REISFELD 13b. TIME (FROM	COVERED	UM DOPED SAP	PHIRE" ORT (Year, M	onth, Day)	15. PAGE	COUNT
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTA NONE 17. COSATI	IC CHARACTERIZ	COVERED TO 181 SUBJECT TERMS	UM DOPED SAP	PHIRE" ORT (Year, M	onth, Day) ry and idea	15. PAGE	COUNT ck number)
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13a. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTA NONE 17. COSATI FIELD GROUP	IC CHARACTERIZ	COVERED TO 1815) OJECT TERMS	(Continue on rever	PHIRE" ORT (Year, M	onth, Day) ry and ide	15. PAGE	COUNT ck number)
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13a. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTA NONE 17. COSATI FIELD GROUP	IC CHARACTERIZ	COVERED	(Continue on rever	PHIRE" ORT (Year, M rise if necessai	onth, Day) ny and ide	15. PAGE	COUNT ck number)
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13a. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTA NONE 17. COSATI FIELD GROUP 19. ABSTRACT (Continue or	IC CHARACTERIZ	COVERED TO 1815) OJECT TERMS	(Continue on rever , c number)	PHIRE"	onth, Day) ry and ide	15. PAGE	COUNT ck number)
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13a. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTA NONE 17. COSATI FIELD GROUP 19. ABSTRACT (Continue of betract; Tit	IC CHARACTERIZ REISFELD 13b. TIME (FROM TION CODES SUB-GROUP CODES SUB-GROUP Treverse if necessary tanium (III)	ZATION OF TITANI	(Continue on rever , I) c number) shows a sho	PHIRE" ORT (Year, M rse if necessar N Ort-liv	onth, Day) ry and idea ed (3.	15. PAGE	COUNT ck number)
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTA NONE 17. COSATI FIELD GROUP 19. ABSTRACT (Continue or betract: Tit broad ² E - ²	IC CHARACTERIZ	TO	(Continue on rever , I / c number) shows a sho	PHIRE" ORT (Year, M rse if necessar r Ort-liv a 10 ti	ry and idea ed (3. mes 10	15 PAGE ntify by blo 1 (s) onger-1	COUNT ck number)
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13a. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTA NONE 17. COSATI FIELD GROUP 19. ABSTRACT (Continue of betract; Tit broad ² E - ² A, (*4s*) - ² T	IC CHARACTERIZ REISFELD 13b TIME (FROM	TOUVERED 1815/0/JECT TERMS 7, 4 cm y and identify by block in corundum ence around 7 around 410 nm	(Continue on rever ,, c number) shows a sho 40 nm, and a .Both the so	PHIRE" ORT (Year, M rse if necessar N ort-liv a 10 ti e broad	onth, Day) ny and idea ed (3. mes lo -band	15. PAGE ntify by blo 1 (a s) onger-1 emissi	COUNT ck number) ived ons
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13a. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTA NONE 17. COSATI FIELD GROUP 19. ABSTRACT (Continue or betract: Tit broad ² E - ² T A 1 ("4s") - ² T yith large St	IC CHARACTERIZ REISFELD 13b. TIME (FROM	TO TO TO TO TO TO TO TO TO TO	(Continue on rever ,,,,,,,,	PHIRE" ORT (Year, M rse if necessar N ort-liv a 10 ti e broad able la	ed (3. mes lo -band	15 PAGE ntify by blo 1 (is) onger-1 emissi The abs	COUNT ck number) ived ons orption
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13a. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTA NONE 17. COSATI FIELD GROUP 19. ABSTRACT (Continue of betract: Tit broad ² E - ² a (*4s*) - ² T with large St pectrum is s	IC CHARACTERIZ REISFELD 13b. TIME (FROM	TO TO	(Continue on rever , number) shows a sho 40 nm, and a .Both these ts for tunk aces of ti	PHIRE" ORT (Year, M rse if necessar N ort-liv a 10 ti: e broad able la tanium (ed (3. mes lo -band IV) ar	15 PAGE ntify by blo 1 (is) onger-1 emissi The abs re show	COUNT ck number) ck number) orption m not
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13a. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTA NONE 17. COSATI FIELD GROUP 19. ABSTRACT (Continue or betract: Tit broad ² E - ² T A ("4s") - ² T with large St pectrum is st to be involve	IC CHARACTERIZ REISFELD 13b. TIME (FROM	ZATION OF TITANI	(Continue on rever , I / c number) shows a sho 40 nm, and a .Both these ts for tunk aces of ti- ion, in con	PHIRE" ORT (Year, M rse if necessar n ort-liv a 10 ti e broad able la tanium (trast t	ed (3. mes lo -band sers.1 IV) ar o a pa	15 PAGE ntify by blo 1 (is s) onger-1 emissi the abs re show trasiti	COUNT ck number) ck number) ck number) ons orption m not c
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13a. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTA NONE 17. COSATI FIELD GROUP 19. ABSTRACT (Continue of betract; Tit broad ² E - ² 7 A ("4s") - ² 7 A (15) A (15	IC CHARACTERIZ REISFELD 13b TIME (FROM	TOUVERED 1815/0/JECT TERMS 1815/0/JECT TERMS 1815/0/JECT TERMS 1.100/JECT TERMS	(Continue on rever (Continue on rever ,,,,,,,, _	PHIRE" ORT (Year, M rse if necessar N ort-live a 10 til e broad able la tanium (trast t e sapph	ed (3. mes lo -band sers.1 IV) ar o a pa ire is	15 PAGE 15 PAGE 16	COUNT ck number) ck number)
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTA NONE 17. COSATI FIELD GROUP 19. ABSTRACT (Continue or betract: Tit broad $2E - \frac{2}{7}$ A 1 ("4s") - T ith large St pectrum is s to be involve bsorption an to Fe(II) Ti(1)	IC CHARACTERIZ REISFELD 13b TIME (FROM	ZATION OF TITANI	(Continue on rever ,, (Continue on rever ,, number) shows a sho 40 nm, and a .Both these its for tunk aces of ti ion, in con II, IV), like	PHIRE" ORT (Year, M rse if necessar ort-liv a 10 ti e broad able la tanium (trast t e sapph sfer ba	ed (3. mes lo -band sers.] IV) ar o a pa ire is nds.	15 PAGE nuify by blo 1 (ify b)	COUNT ck number) ck number) ons orption m not c due
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTA NONE 17. COSATI FIELD GROUP 19. ABSTRACT (Continue or betract: Tit broad $E - 27$ Cosati proad $E - 27$ a ("4s") - T with large St pectrum is s to be involve bsorption and to Fe(II) Ti(1) 20 DISTRIBUTION/AVAILA	IC CHARACTERIZ REISFELD 13b. TIME (FROM	ZATION OF TITANI	(Continue on rever ,	PHIRE" ORT (Year, M rse if necessar N ort-liv a 10 ti: e broad able la tanium (trast t e sapph sfer ba SECURITY CLA	ed (3. mes lo -band sers.1 IV) ar o a pa ire is nds.	15 PAGE ntify by blo 1 (s s) onger-1 emissi che abs ce show arasiti s blue k	COUNT ck number) ived ons orption m not c due
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13a. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTA NONE 17. COSATI FIELD GROUP 19. ABSTRACT (Continue of betract: Tit broad ² E - ² T A ("4s") - ² T A ("4	IC CHARACTERIZ REISFELD 13b TIME (FROM	TOUVERED TO TO TO TO TO TO TO TO TO TO	(Continue on rever , I / c number) shows a sho 40 nm, and a b.Both these ts for tune aces of ti- ion, in con II, IV), like ctron tran 21 ABSTRACT S UNCLASSIFI 225 TELEPHONE	PHIRE" ORT (Year, M rse if necessar N ort-liv a 10 ti e broad able la tanium (trast t e sapph sfer ba security CLA ED E (nelude Are	onth, Day) ed (3. mes lo -band sers.1 IV) ar o a pa ire is nds. (SSIFICATIO a (ode)12	15 PAGE ntify by blo 1 (is s) onger-1 emissi the abs re show irasiti blue k 20 OFFICE	COUNT ck number) ck number) ck number) ons orption m not c due
(U) "SPECTROSCOP 12. PERSONAL AUTHOR(S) PROFESSOR RENANTA 13. TYPE OF REPORT FINAL 16. SUPPLEMENTARY NOTA NONE 17. COSATI FIELD GROUP 19. ABSTRACT (Continue or betract; Tit broad ² E - ² 7 A ("4s") - ² 7 A (10) Construction (10) Constr	IC CHARACTERIZ REISFELD 13b. TIME (FROM	ZATION OF TITANI	(Continue on rever (Continue on rever ,,,,,,,, _	PHIRE" ORT (Year, M rse if necessar n ort-liv a 10 ti e broad able la tanium (trast t e sapph sfer ba SECURITY CLA (ED E (Include Are AMXS)	ed (3. mes lo -band sers.7 IV) ar o a pa ire is nds. .ssificatio a code) 2. N-IIK-EM	15 PAGE ntify by blo 1 (is s) onger-1 emissi The abs re show irasiti is blue k 20 OFFICE S	COUNT ck number) ived ons orption m not c due

The original proposal was scheduled for 1 Oct.1985 to 30 Sept.1986. However, since the crystals were obtained only in May 1986, an extension for the work was granted for an additional year.

The doped sapphire crystals were kindly supplied by Dr. R.N. Katz and Dr. J.L. Caslavsky of the Ceramics Research Division, Army Materials and Mechanics Research Center, Watertown, Massuchusetts 02172, USA.

The report consists of the paper which was sent to Chimia:

Unusual Luminescence of Titanium(III) in Aluminium Oxide, R.Reisfeld, M. Eyal and C.K. Jørgensen.

 (\cdot)



Acces	ion For	
NTIS DTIC Unanr Justifi	CRA&I TAB iounced cation	
By Dist.ib	ution/	
A	vailability	Codes
Dist	Avail and Specia	1/or il
A-1		

Unusual Luminescence of Titanium(III) in Aluminium Oxide

Renata Reisfeld, Marek Eyal and Christian K.Jørgensen *

Abstract: Titanium(III) in corundum shows a short-lived $(3.1 \mu s)$ broad ${}^{2}E - {}^{2}T_{2}$ luminescence around 740 nm,and a 10 times longer-lived ${}^{2}A_{1}("4s") - {}^{2}T_{2}$ centered around 410 nm.Both these broad-band emissions with large Stokes shift show prospects for tunable lasers.The absorption spectrum is studied in detail,and traces of titanium(IV) are shown not to be involved in the "violet" emission,in contrast to a parasitic absorption around 700 nm due to Ti(III,IV),like sapphire is blue due to Fe(II)Ti(IV) \rightarrow Fe(III)Ti(III) electron transfer bands.

* Correspondence: Prof.Dr.h.c. C.K.Jørgensen Département de Chimie minérale, analytique et appliquée Université de Genève Quai Ernest Ansermet 30, CH-1211 Geneva 4

Prof.Dr. R.Reisfeld,Dr.M.Eyal Department of Inorganic Chemistry Hebrew University 91904 Jerusalem (Israel)

COLUMN STORES

** Acknowledgements: We are very grateful to Dr.Jaroslav L.Caslavsky and Dr. Robert N.Katz,Ceramics Research Division,U.S.Army Materials Technology Laboratory,Watertown,Massachusetts 02172,for the highly valuable crystalline samples studied here.This work was supported in part by the U.S.Army European Research Office (London) under contract DAJA 45-85-M-0344.The grant no. 2.820-0.85 from the Swiss National Science Foundation made the collaboration possible between the universities in Jerusalem and Geneva.

Corundum, the α -modification of Al $_2O_3$, has almost exactly regular octahedral sites for the cation, though spectroscopic characteristics in polarized light clearly indicate the uniaxial hexagonal crystal type. Those trivalent ions which form hydrated alums can be incorporated on the octahedral site, and for instance, Al_2O_3 , Cr_2O_3 and Fe_2O_3 are fully miscible. The absorption spectra of trivalent 3d group ions in corundum were studied [1] when the first laser, the ruby $Cr_Al_{2-v}O_3$ with $x \sim 0.01$, was operated in 1960. Though the colourless corundum (nearly as hard as diamond, and used in optical components transparent in the far ultraviolet) frequently is called sapphire, this blue mineral contains simultaneously small amounts of iron(II) and titanium(IV) on the octahedral sites, and the broad absorption band [2,3] at 575 nm with a shoulder at 720 nm is due to electron transfer forming, to the first approximation, $3d^{5}$ iron(III) and $3d^{1}$ titanium(III) in the excited state, as frequently found [4-6] in compounds containing iron(II) at short distances from oxidizing ions, including iron (III). The black mineral ilmenite FeTiO, has an ordered super-structure of corundum. It may be noted that ilmenite-type Ni Mg_{1-x} TiO₃ and Ni Cd_{1-x} TiO₃ were studied by Reinen, allowing important conclusions [7] about the 3d⁸ states of nickel(II) corresponding to the absorption bands.

Some lasers, based on gaseous atoms or positive ions, emit spectral lines. This is almost true also for rare earths containing a partly filled 4f shell [8,9]. On the other hand, fluorescent organic colorants in slution, or excimer lasers based on ephemeric diatomic molecules in yases,emit broad luminescence bands,and under favourable conditions, can be used for tunable lasers, such as certain cases of 3d³ chromium(III) emitting a broad band in the near infrared from the first excited guartet state ${}^{4}T_{2}$ to the groundstate ${}^{4}\Lambda_{2}$. Since ${}^{4}T_{2}$ has one strongly anti-bonding d-like electron, the Føanck-Condon principle demands the co-excitation of a broad vibronic continuum belonging to the groundstate. The situation is quite different in the narrow-band emission of ruby, where the first doublet state ${}^{2}E$ involves the same three roughly nonbonding electrons as ${}^{4}A_{2}$. Tunable lasers emitting from ${}^{4}T_{2}$ in the isoelectronic vanadium(II) incorporated in rutile-type $V_{x}Mg_{1-x}F_{2}$ have been studied [10] but present difficulties of black-out due to colour centres formed during large-scale operation.

r 2

Titanium(III) has one 3d electron which can be excited (by light close to 500 nm) to the two anti-bonding orbitals with angular functions proportional to (x^2-y^2) and $(3z^2-r^2)$. If the octahedron TiX₆ was exactly regular, these two orbitals would have identical energies. Actually, the hexa-aqua ion in acidic, non-complexing solutions [11,12] shows a shoulder at 575 nm on the asymmetric peak at 492 nm. This can hardly be ascribed to a distorted environment, as would be conceivable in a doped crystal (especially in the case of quite differing ionic radii). It may be relevant that the isoelectronic vanadium(IV) forms the highly anisotropic [12,13] blue vanadyl aqua ion $OV(OH_2)_4^{+2}$ containing its unpaired electron in the orbital (xy) which is the only non-bonding 3d orbital in the case of strong π -anti-bonding effects of oxo ligands on the z Cartesian axis. The diamagnetic groundstate of $MO_{2}X_{4}$ complexes [14] of M=ruthenium(VI); #rhenium(V) and osmium(VI) corresponds to two electrons in (xy).It is generally argued that the splitting of the visible absorption band of titanium(III) is due to the Jahn-Teller effect. The slightly problematic side of this explanation is that Jahn-Teller-stable groundstates do not provide band splittings by transitions to even highly separated potential surfaces in the 16-dimensional space corresponding to 7 nuclei in MX, [13]. However,a quite weak deviation from O_b in the groundstate can produce strong energy separations by Franck-Condon projection on high-lying potential surfaces, much in the same way as the width of "ligand field" absorption bands, though originating in the thermal (and zero-point) vibration [15], frequently is 10 times kT, and largest for the lightest ligands. In the octahedral d-group complexes, the typical cases of strong \Im Jahn-Teller splitting occur for unbalanced occupation of (x^2-y^2) and $(3z^2-r^2)$ in the groundstate, actually by 0 and 1 electron in the quintet groundstates of 3d⁴ chromium(II) and manganese(III); and by 1 and 2 electrons in 3d⁹ copper(II).A few instances are known [16] of pronounced band splitting of iron(II) hexa-aqua ions and in the quintet groundstate hexafluoro complex of cobalt(III).Since the two σ -anti-bonding orbitals have each one electron, the phenomenon is here ascribed to Jahn-Teller effect in the (at most) π -anti-bonding (xy),(xz) and (yz) accommodating 4 electrons.It is known from chromium(III) complexes [17] that ar-antibonding effects are up to 30 percent of the σ -anti-bonding effect, the highest ratio is observed for fluoride ligands.Hence, it can be concluded that the weak Jahn-Teller effect on titanium(III) is induced by the π anti-bonding only.Because of the mechanism of the Stokes shift, the consequences are likely to be more pronounced for luminescence than for

р.З

absorption.Jahn-Teller effects for the excited ${}^{4}T_{2}$ of chromium(III) in cubic crystals have been detected [18] in the vibrational fine-structure (below 10 K) of the broad emission band.

Powell et al.[3] recently studied the absorption and emission spectra of $Ti_{x}Al_{2-x}O_{3}$ and found an asymmetric band (we find the maximum at 483 nm and a shoulder at 550 nm) surprisingly similar to the aqua ion in alums [11] and in acidic solutions. The near-infrared emission occurs in a broad band around 740 nm, satisfying one of the necessary conditions for tunable lasers [10]. This emission in corundum has been studied by two other groups [19-21] and one of the difficulties for obtaining a positive amplification along the laser path is a somewhat unpredictable absorption between 700 and 850 nm. Though it is difficult to study trace impurities in crystals almost as hard as diamond, it does not seem to be Fe(II), Ti(IV) providing the blue colour of sapphire [3] but rather adjacent Ti(III, IV) sites known [12] in 12 molar hydrochloric acid, where simultaneously present (very pale yellow) Ti(IV) and a sky-blue Ti(III) chloro complex (with maximum at 525 nm) form a dark brownishpurple 1:1 complex with a 8 times stronger, broad band at 482 nm, having a shoulder at 670 nm. The formation constant of this complex is 12 M^{-1} showing a pronounced affinity between the two (chloro-bridged) oxidation states. The diagnosis is that parasitic absorption can be avoided in titanium(III) doped corundum if all traces of Ti(IV) can be reduced to Ti(III), or at least kept at large average distance from the Ti(III) providing the main absorption.

As described in the experimental section, we studied three crystalline samples, among which one (R 10) does not present the flat background absorption slightly decreasing from 650 to 850 nm, as do the samples R 4 and R 8. In all three samples, excitation at 488 nm, monitoring emission at 732 nm, and also in R 10 excited at 488 nm and monitored at 650 and 850 nm, the decay curve (after an initial rise-time of $3 \cdot 10^{-7}$ s suggesting rapid energy transfer from states producing much stronger ultraviolet absorption than the "4s" state discussed below) is almost exactly exponential with the life-time (3.1 ± 0.1) µs. This is comparatively short, since the radiative life-time

$$T_{rad} = (e_2/e_1) 2.3 \cdot 10^{-8} s/[P(h\nu/eV)^2]$$
 (1)

is close to $3 \cdot 10^{-5}$ s for the Ti(III) aqua ion [15] having the oscillator strength P = $1.2 \cdot 10^{-4}$ if the debatable ratio between e_2 excited states and e_1 states in the ground level is put equal to 1.As seen in the

experimental section, the **E** values at 488 nm for Tr(111) in corundum are closely similar to 4.1 for the agua ion in solution, as well as the shape of the absorption band. Hence, it seems that the order of magnitude for the guantum yield of near-infrared luminescence is close to 0.1.

A much more spectacular luminescence of all three samples is a broad emission band (Figure 1) having the maximum at 410 nm and half the peak intensity at 365 and 460 nm. This "violet" emission was already reported by Powell et al.[3] at 420 nm with observed life-time \mathbf{T} =2.3 μ s. As seen in Table 1, we find life-times gliding from 15 toward 35 μ s for most of the emission wave-lengths studied, for excitation of all three samples at 337 nm. On the other hand, we did not find any dependence on the exciting wave-length of the position and shape of the emission band. The luminescence may very well be due to a superposition of two or several mechanisms, the slowest corresponding to the asymptotic value of \mathbf{T} close to 30 microseconds.

Searching the emitting state, we measured the excitation spectra for emission at 350,400 and 450 nm, and found in all three samples an (energy-wise) quite broad band at 265 nm. One may feel some doubts whether superposed competing absorption bands shift this excitation maximum at 265 nm from the average position of the emitting state. Going from the absorption maximum toward ultraviolet, the minimum $\mathfrak{E} \sim 2$ occurs at 405 nm, and \mathfrak{E} =5 is crossed close to 310 nm. The subsequent increase is very smooth, and reaches \mathfrak{E} =200 close to 245 nm. There are no evident shoulders between 300 and 250 nm, and $\mathfrak{E} \sim 25$ at 265 nm.

Since mixed oxides containing titanium(IV) absorb very strongly in the near ultraviolet, in contrast to the much less oxidizing zirconium(IV) and thorium(IV), one might expect an electron transfer state of Ti(IV) producing the "violet" emission in analogy to solid vanadates [22].This is rendered unlikely by the essentially similar absorption spectra in the ultraviolet, and the emission characteristics, of R 4 and R 8 compared with the sample R 10 lacking near-infrared absorption indicating absence of admixed Ti(IV).Hence, the "violet" emission seems to originate in Ti(III).

The most attractive hypothesis is a transition of the single 3d electron to a 4s-like orbital.Such a transition may be very weak, and if the asymptotic $\pi \sim 30 \ \mu$ s is not an artifact of energy trapping [23], equation (1) provides $\pi_{rad} \sim T$ for $P = (e_2/e_1) \cdot 4 \cdot 10^{-5}$.Since the quantum yield η cannot be above 1, this is a higher limit for P (which is to be multiplied by η if below 1).This means that the absorption corresponding

p. 5

to "violet" emission is a few times weaker than the band at 483 nm, and one can hardly hope a contribution to $\boldsymbol{\xi}$ below 2 to be detectable below 300 nm.

3d - "4s" transitions have only been detected in compounds of a few reducing d-group ions. The best characterized case [15] is the band of $Fe(OH_2)_{c}^{+2}$ at 248 nm (40300 cm⁻¹) having E=18 and $P=3.5\cdot10^{-4}$. The position may be compared with the two levels of $3d^54s$ of gaseous Fe^{+2} , 7s, situated 30089 cm⁻¹ and 5 s, at 40999 cm⁻¹ above the 3d⁶ groundstate. Two counteracting influences modify the 3d-4s separation: the effective positive charge is smaller [7-9] in the compound; but the 4s orbital is always anti-bonding with respect to any set of ligating atoms, as can be seen from gaseous Cu^+ of which the levels ${}^{3}D_{3}(21929), {}^{3}D_{2}(22847), {}^{3}D_{1}(23998)$ and ${}^{1}D_{2}(26264)$ of $3d^{9}4s$ are situated the number of cm^{-1} (given in parenthesis) above the closed-shell groundstate. These levels may be compared with a band [24] of the newly detected complex $Cu(NH_3)_3^+$ at 290 nm (34000 cm⁻¹) having $\mathbf{E} \sim 300$, whereas Cu(NH₃)⁺ absorbs less and begins at higher energy. The strong dependence of intensities of parity-forbidden transitions on the local symmetry is well-known from internal transitions in the partly filled 3d shell, e.g. of octahedral and tetrahedral cobalt(II) complexes. The 3d-"4s" transitions start at 31500 cm⁻¹ in Cu_vNa_{1-v}F and are quite weak [25].

Besides corundum, there are very few cases known of titanium(III) luminescence.Recently [26], a fluorophosphate glass was studied in Shanghai.The absorption increases strongly below 350 nm and shows a peak at 529 nm and a shoulder at 685 nm.Excitation at 308 nm produces a very broad emission centered around 530 nm with γ of order 10 μ s at 77 K, but depending to some extent on the emission wave-length, as one might expect from differing sites in the glass [9,23].It is likely that this green emission with an enormous Stokes shift (from a maximum in the excitation spectrum at 290 nm) is analogous to the "violet" emission studied here.

Experimental

The crystals obtained by the VSOM technique previously described [3] were polished with two parallel surfaces typically separated by 1 cm. The samples selected were:

R 4: density d=3.795 g/cm³, nominal titanium concentration $c_{Ti} = 0.01425$ mole/L. R 8: d=3.744 g/cm³ and $c_{Ti} = 0.01425$ mole/L.

P 10: d=3.928 g/cm³ and $c_{mi} = 0.01475$ mole/L.

p.6

Steady-state measurements:

Absorption spectra were measured of the crystal against air on a Perkin-Elmer double-beam spectrophotometer model Lambda 3.Assuming the c_{Ti} given above, the molar extinction coefficient \mathcal{E} at 488 nm was found to be 4.5 for R 4; 4.9 for R 8; and 3.9 for R 10.

Emission and excitation spectra were measured with the Perkin-Elmer L3 with a B&L home-made, and a Spex home-made, spectrofluorimeter.Excitation at 265 nm was performed with radiation from a high-pressure xenon lamp gone through a monochromator.

Life-time measurements:

A nitrogen laser (337 nm) was used directly for exciting "violet" emission, and combined with the dye no.8 emitting at 488 nm for exciting the red and near-infrared emission. Table 1 gives the folded life-time τ_1 (for e⁻¹ times the original intensity), τ_2 (half the time for e⁻²) and τ_3 (a-third the time for e⁻³), all in microseconds, of the emission studied at wave-lengths between 350 and 732 nm of the three samples.

Table 1.Life-times in microseconds of "violet" emission of titanium(III) in corundum excited at 337 nm, and emission measured at the nm value given. T_n are defined in the experimental section.

	1	1			
Sa	Imple	nm	τ_1	\mathbf{T}_2	T_3
R	4	350	1.8	6.3	10.3
R	4	400	3.1	12.6	19.4
R	4	450	11.7	18.9	24.2
R	4	500	24.1	28.4	29.2
R	4	542	18.8	31.3	30.3
R	4	600	23.7	32.5	29.4
R	4	650	15.2	29.8	30.8
R	4	732	4.2	17.1	28.2
R	8	542	23.1	32.6	29.4
R	10	542	15.7	30.5	34.8

Fig.1.Emission spectrum (full line; excited in an interval around 265 nm) and excitation spectrum (dashed line; emission measured at 450 nm) of sample R 4.

Fig.1.Emission spectrum (full line; excited in an interval around 265 nm) and excitation spectrum (dashed line; emission measured at 450 nm) of sample R 4.

Little - Kissis

「アンシント」 たんたんたいしん

\$5.0



- [1] D.S.McClure, J. Chem. Phys. 36 (1962) 2757.
- [2] K.Nassau, Am. Mineral. J. 63(1978)219.
- [3] R.C. Powell, J.L. Caslavsky, Z.AlShaieb, J.M. Bowen, J.Appl. Phys. 58 (1985) 2331.
- [4] M.B.Robin, P.Day, Adv. Inorg. Chem. Radiochem. 10(1967)247.
- [5] C.K.Jørgensen, Progress Inorg. Chem. 12(1970)101.
- [6] G.C.Allen, M.B. Wood, J.M. Dyke, J. Inorg. Nucl. Chem. 35(1973)2311.
- [7] C.K.Jørgensen:Oxidation Numbers and Oxidation States, Springer,Berlin(1969).
- [8] R.Reisfeld,C.K.Jørgensen:Lasers and Excited States of Rare Earths,Springer,Berlin(1977).
- [9] R.Reisfeld,C.K.Jørgensen, in K.Gschneidner, LeRoy Eyring (Ed.): Handbook on the Physics and Chemistry of Rare Earths, Vol.9, North-Holland, Amsterdam (1987).
- [10] P.Hammerling, A.B.Budgor, A.Pinto (Ed.): Tunable Solid State Lasers, Springer series in Optical Sciences 47, Springer, Berlin (1985).
 - [11] H.Hartmann, H.L.Schläfer, Z.Naturforsch. 6 a (1951) 760.
 - [12] C.K.Jørgensen, Acta Chem. Scand. 11(1957)73.
 - [13] C.K.Jørgensen, Topics Current Chem. 124(1984)1.
 - [14] C.K.Jørgensen, Rev. Chim. Minér. 14 (1977) 127.
 - [15] C.K.Jørgensen, Adv. Chem. Phys. 5(1963)33.
 - [16] F.A.Cotton, M.D.Meyers, J.Am. Chem. Soc. 82(1960) 5023.
 - [17] J.Glerup, O.Mønsted, C.E.Schäffer, Inorg. Chem. 15 (1976) 1399.
 - [18] R.Knochenmuss, C.Reber, M.V.Rajasekharan, H.U.Güdel, J.Chem.Phys. 85(1986)4280.
- [19] P.Lacovara, L.Esterowitz, R.Allen, Optics Lett. 10(1985)23.
- [20] P.Lacovara, L.Esterowitz, M.Kotka, IEEE J.Quantum Electron. <u>QE-21(1985)1614.</u>
 - [21] N.P.Barnes, D.K.Remelius, Digest of Technical Papers, CLEO Conference on Lasers and Electro-Optics, Baltimore, May 1985, p. 168. Optical Society of America, Washington D.C. (1985).
 - [22] G.Blasse, Structure and Bonding 42(1980)1.
 - [23] R.Reisfeld, M.Eyal, C.K.Jørgensen, A.H.Guenther, B.Bendow, Chimia <u>40</u>(1986)403.
 - [24] J.Bjerrum, Acta Chem. Scand. A 40(1986)233.
 - [25] D.S.McClure, C.Pedrini, J.Lumin. 31(1984)197.
 - [26] Liu Huimin, Gan Fuxi, J. Non-Cryst. Solids 80(1986)422.
 - [27] J.L.Caslavsky: The Growth of Large Diameter Single Crystals by Vertical Solidification of the Melt, in E.Kaldis(Ed.): Crystal Growth of Electronic Materials, p.113, Elsevier, Amsterdam and New York (1985).

