

AD-A101 356

ON THE CONVERGENCE RATES OF EMPIRICAL BAYES RULES FOR
TWO-ACTION PROBLEMS. (U) PURDUE UNIV LAFAYETTE IN DEPT
OF STATISTICS T C LIANG MAR 87 TR-87-8

1/1


UNCLASSIFIED

NO0014-84-C-1067

F/G 12/3

NL





DTIC FILE COPY



1

On The Convergence Rates Of Empirical Bayes Rules
For Two-Action Problems: Discrete Case *


by

TaChen Liang
Purdue University

Technical Report #87-8

AD-A181 356

PURDUE UNIVERSITY



DTIC
ELECTE
MAY 26 1987
S D

DEPARTMENT OF STATISTICS

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

87

5 21 423

On The Convergence Rates Of Empirical Bayes Rules
For Two-Action Problems: Discrete Case *

by

TaChen Liang
Purdue University

Technical Report #87-8

DTIC
ELECTE
MAY 26 1987
S D D

Department of Statistics
Purdue University

March 1987

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

* This research was partially supported by the Office of Naval Research Contract N00014-84-C-1067 and NSF Grant DMS-8606964 at Purdue University.

o

87-5-21-428

**On The Convergence Rates Of Empirical Bayes Rules
For Two-Action Problems: Discrete Case ***

by

TaChen Liang
Purdue University

Abstract

The purpose of this paper is to investigate the convergence rates of a sequence of empirical Bayes decision rules for the two-action decision problems where the distributions of the observations belong to a discrete exponential family. It is found that the sequence of the empirical Bayes decision rules under study is asymptotically optimal, and the order of associated convergence rates is $O(\exp(-cn))$, for some positive constant c , where n is the number of accumulated past experience (observations) at hand. Two examples are provided to illustrate the performance of the proposed empirical Bayes decision rules. A comparison is also made between the proposed empirical Bayes rules and some earlier existing empirical Bayes rules.

AMS Subject Classification: 62C12

Key words and phrases: Bayes rule; Empirical Bayes rule; Asymptotically optimal; Rates of convergence

* This research was partially supported by the Office of Naval Research Contract N00014-84-C-1067 and NSF Grant DMS-8606964 at Purdue University.



Codes

d/or
cial

A-1

1. Introduction

The empirical Bayes approach in statistical decision theory is appropriate when one is confronted repeatedly and independently with the same decision problem. In such instances, it is reasonable to formulate the component problem in the sequence as a Bayes decision problem with respect to an unknown prior distribution on the parameter space and then use the accumulated observations to improve the decision rule at each stage. This approach is due to Robbins (1956, 1964, 1983). Many such empirical Bayes rules have been shown to be asymptotically optimal in the sense that the risk for the n th decision problem converges to the optimal Bayes risk which would have been obtained if the prior distribution was fully known and the Bayes rule with respect to this prior distribution was used.

The usefulness of empirical Bayes rules in practical applications clearly depends on the convergence rates with which the risks for the successive decision problems approach the optimal Bayes risk. The purpose of this paper is to investigate the convergence rates of a sequence of empirical Bayes rules for two-action decision problems when the distributions of the observations belong to a discrete exponential family.

Let X be a random observation with probability function of the form

$$(1.1) \quad f(x|\theta) = h(x)\theta^x\beta(\theta), x = 0, 1, 2, \dots; 0 < \theta < Q,$$

where $h(x) > 0$ for all $x = 0, 1, 2, \dots$, and where Q may be finite or infinite. The observation X may be thought of as the value of a sufficient statistic based on several iid observations.

Consider the following testing: $H_0: \theta \geq \theta_0$ against $H_1: \theta < \theta_0$, where θ_0 is a known

positive constant. For each $i = 0, 1$, let i denote the action deciding in favor of H_i . For the parameter θ and action i , the loss function is defined as:

$$(1.2) \quad L(\theta, i) = (1 - i)(\theta_0 - \theta)I_{(0, \theta_0)}(\theta) + i(\theta - \theta_0)I_{[\theta_0, \infty)}(\theta),$$

where $I_A(\cdot)$ denotes the indicator function of the set A . In (1.2), the first item is the loss due to taking action 0 when $\theta < \theta_0$, and the second item is the loss of taking action 1 when $\theta \geq \theta_0$. It is assumed that θ is the value of a random variable Θ having an unknown prior distribution $G(\theta)$.

For a decision rule d , let $d(x) = P\{\text{accepting } H_0 | X = x\}$. That is, $d(x)$ is the probability of taking action 0 given $X = x$. Let D be the class of all decision rules. For each decision rule d , let $r(G, d)$ denote the associated Bayes risk. Then, $r(G) = \inf_{d \in D} r(G, d)$ is the minimum Bayes risk among the class D .

Based on the statistical model described above, the Bayes risk associated with the decision rule d is:

$$(1.3) \quad r(G, d) = \sum_{x=0}^{\infty} |\theta_0 - \varphi(x)| d(x) f(x) + C,$$

where

$$(1.4) \quad \varphi(x) = \frac{h(x)f(x+1)}{h(x+1)f(x)},$$

$$(1.5) \quad f(x) = \int_0^Q f(x|\theta) dG(\theta),$$

$$(1.6) \quad C = \sum_{x=0}^{\infty} \int_{\theta_0}^Q (\theta - \theta_0) f(x|\theta) dG(\theta).$$

We consider only priors G such that $\int_0^Q \theta dG(\theta) < \infty$ to insure that the risk is always finite.

Note that C is a constant which is independent of the decision rule d . Thus, from (1.3), a Bayes decision rule, say d_G , is clearly given by

$$(1.7) \quad d_G(x) = \begin{cases} 1 & \text{if } \varphi(x) \geq \theta_0, \\ 0 & \text{otherwise.} \end{cases}$$

Since the prior distribution G is unknown, it is not possible to apply the Bayes rule for the decision problem at hand. In this situation, we use the empirical Bayes approach. We note that Johns and Van Ryzin (1971) have studied the above decision problem via empirical Bayes approach. In this paper, a sequence of empirical Bayes decision rules $\{d_n^*\}$ is proposed for the above described decision problem. The associated asymptotic optimality property is investigated. It is found that the order of the rate of convergence of $\{d_n^*\}$ is $O(\exp(-cn))$ for some positive constant c , where n is the number of accumulated past experience (observations) at hand. Two examples are given to illustrate the performance of the proposed empirical Bayes decision rules. A comparison is also made between the proposed empirical Bayes rules and some earlier existing empirical Bayes rules.

2. The Proposed Empirical Bayes Rules and Its Asymptotic Optimality

For each $j = 1, \dots$, let (X_j, Θ_j) be a pair of random variables, where X_j is observable but Θ_j is not observable. Conditional on $\Theta_j = \theta$, X_j has probability function $f(x|\theta)$. It is assumed that $\Theta_j, j = 1, \dots$, are independently distributed with common unknown prior distribution G . Therefore, $(X_j, \Theta_j), j = 1, 2, \dots$, are iid. Let $X_n = (X_1, \dots, X_n)$ denote the n past observations and let $X_{n+1} \equiv X$ denote the current random observation.

According to (1.4) and (1.7), an empirical Bayes decision rule, say d_n^* , is proposed as follows. First, for each $x = 0, 1, 2, \dots$, let

$$(2.1) \quad f_n(x) = \frac{1}{n} \sum_{j=1}^n I_{\{x\}}(X_j) + \delta_n,$$

where δ_n is a positive value such that $\delta_n = o(1)$. Then, let

$$(2.2) \quad \varphi_n(x) = \frac{h(x)f_n(x+1)}{h(x+1)f_n(x)}.$$

We then define

$$(2.3) \quad \varphi_n^*(x) = [\max_{0 \leq y \leq x} \varphi_n(y)] \wedge Q,$$

where $a \wedge b = \min\{a, b\}$. Finally, the empirical Bayes decision rule d_n^* is defined as:

$$(2.4) \quad d_n^*(x) = \begin{cases} 1 & \text{if } \varphi_n^*(x) \geq \theta_0, \\ 0 & \text{otherwise.} \end{cases}$$

Note that the past data X_n is implicitly contained in the subscript n .

Definition 2.1. A decision rule d is said to be monotone if for $x, y \geq 0$ with $x \leq y$, $d(x) \leq d(y)$.

Note that from (2.3), $\varphi_n^*(x)$ is nondecreasing in x . Then, by (2.4), we see that $d_n^*(x)$ is a monotone decision rule.

In the following, the asymptotic optimality of the sequence of the proposed empirical Bayes decision rules $\{d_n^*\}$ will be investigated. The monotonicity of the decision rules $\{d_n^*\}$ will be used to obtain the related asymptotic optimality.

Consider an empirical Bayes decision rule $d_n(x)$. Let $r(G, d_n)$ be the Bayes risk associated with the rule d_n . Then,

$$(2.5) \quad r(G, d_n) = \sum_{x=0}^{\infty} [\theta_0 - \varphi(x)] E[d_n(x)] f(x) + C,$$

where the expectation E is taken with respect to X_n . Since $r(G)$ is the minimum Bayes risk, $r(G, d_n) - r(G) \geq 0$ for all n . Thus, the nonnegative difference $r(G, d_n) - r(G)$ is used as a measure of the optimality of the empirical Bayes decision rule d_n .

Definition 2.2. A sequence of empirical Bayes decision rules $\{d_n\}_{n=1}^{\infty}$ is said to be asymptotically optimal at least of order α_n relative to the (unknown) prior distribution G if $r(G, d_n) - r(G) \leq O(\alpha_n)$ as $n \rightarrow \infty$, where $\{\alpha_n\}$ is a sequence of positive numbers such that $\lim_{n \rightarrow \infty} \alpha_n = 0$.

Now, straightforward computation leads to that $\varphi(x)$ is increasing in x . Thus, we let $A(\theta_0) = \{x | \varphi(x) > \theta_0\}$ and $B(\theta_0) = \{x | \varphi(x) < \theta_0\}$. Define

$$(2.6) \quad M = \begin{cases} \min A(\theta_0) & \text{if } A(\theta_0) \neq \phi, \\ \infty & \text{if } A(\theta_0) = \phi, \end{cases}$$

$$(2.7) \quad m = \begin{cases} \max B(\theta_0) & \text{if } B(\theta_0) \neq \phi, \\ -1 & \text{if } B(\theta_0) = \phi, \end{cases}$$

where ϕ denotes the empty set.

By the increasing property of $\varphi(x)$ with respect to the variable x , $m \leq M$; also, $m < M$ if $A(\theta_0) \neq \phi$. Furthermore,

$$(2.8) \quad x \leq m \text{ iff } \varphi(x) < \theta_0 \text{ and } y \geq M \text{ iff } \varphi(y) > \theta_0.$$

The following theorem is our main result.

Theorem 2.1. Let $\{d_n^*\}$ be the sequence of empirical Bayes decision rules defined above.

Suppose that $\theta_0 < Q$. Also, assume that

$$(a) \int_0^Q \theta dG(\theta) < \infty \text{ and}$$

$$(b) m < \infty.$$

Then, $r(G, d_n^*) - r(G) \leq O(\exp(-cn))$ for some positive constant c .

Proof: Under Assumption (b) and by (2.8), direct computation leads to

(2.9)

$$r(G, d_n^*) - r(G) = \sum_{x=0}^m [\theta_0 - \varphi(x)] P\{\varphi_n^*(x) \geq \theta_0\} f(x) + \sum_{x=M}^{\infty} [\varphi(x) - \theta_0] P\{\varphi_n^*(x) < \theta_0\} f(x),$$

where $\sum_{x=0}^m \equiv 0$ if $m = -1$.

The nondecreasing property of $\varphi_n^*(x)$ implies

$$(2.10) \quad \begin{cases} P\{\varphi_n^*(x) \geq \theta_0\} \leq P\{\varphi_n^*(m) \geq \theta_0\} & \text{for all } x \leq m, \\ P\{\varphi_n^*(x) < \theta_0\} \leq P\{\varphi_n^*(M) < \theta_0\} & \text{for all } x \geq M. \end{cases}$$

Combining (2.9) and (2.10), we have

$$(2.11) \quad r(G, d_n^*) - r(G) \leq b_1 P\{\varphi_n^*(m) \geq \theta_0\} + b_2 P\{\varphi_n^*(M) < \theta_0\},$$

where $0 \leq b_1 = \sum_{x=0}^m [\theta_0 - \varphi(x)] f(x) < \infty$, $0 \leq b_2 = \sum_{x=M}^{\infty} [\varphi(x) - \theta_0] f(x) < \infty$, and the finiteness of both b_1 and b_2 is guaranteed since $\int \theta dG(\theta) < \infty$ by Assumption (a).

Therefore, it suffices to consider the asymptotic behaviors of both $P\{\varphi_n^*(m) \geq \theta_0\}$ and $P\{\varphi_n^*(M) < \theta_0\}$.

By the definition of $\varphi_n^*(x)$, when $\varphi_n^*(M) < Q$, then $\varphi_n^*(M) \geq \varphi_n(M)$, where $\varphi_n(\cdot)$ is the function defined in (2.2). In view of this fact and by (2.1) and (2.2),

$$(2.12) \quad \begin{aligned} P\{\varphi_n^*(M) < \theta_0\} &\leq P\{\varphi_n(M) < \theta_0\} \\ &= P\left\{\frac{1}{n} \sum_{j=1}^n A_j(M) < -t(M, \theta_0) + \Delta(M, \theta_0, n)\right\}, \end{aligned}$$

where

$$(2.13) \quad A_j(x) = h(x)[I_{\{x+1\}}(X_j) - f(x+1)] - \theta_0 h(x+1)[I_{\{x\}}(X_j) - f(x)],$$

$$(2.14) \quad t(x, \theta_0) = h(x)f(x+1) - \theta_0 h(x+1)f(x),$$

$$(2.15) \quad \Delta(x, \theta_0, n) = \delta_n[h(x+1)\theta_0 - h(x)].$$

Also, by the definition of $\varphi_n^*(x)$ and (2.1) and (2.2) again,

$$(2.16) \quad \begin{aligned} P\{\varphi_n^*(m) \geq \theta_0\} &= P\{\varphi_n(y) \geq \theta_0 \text{ for some } y = 0, 1, \dots, m\} \\ &\leq \sum_{y=0}^m P\{\varphi_n(y) \geq \theta_0\} \\ &= \sum_{y=0}^m P\left\{\frac{1}{n} \sum_{j=1}^n A_j(y) \geq -t(y, \theta_0) + \Delta(y, \theta_0, n)\right\}. \end{aligned}$$

Note that $A_j(x), j = 1 \dots n$, are iid; $E[A_j(x)] = 0$, and $a_1(x, \theta_0) \leq A_j(x) \leq a_2(x, \theta_0)$ where $a_1(x, \theta_0) = -h(x)f(x+1) - h(x+1)\theta_0 + h(x+1)\theta_0 f(x)$ and $a_2(x, \theta_0) = h(x) - h(x)f(x+1) + h(x+1)\theta_0 f(x)$. Also, since $\delta_n = o(1)$ and $m < \infty$, there exists some positive integer n_0 such that for all $n \geq n_0$, $|\Delta(y, \theta_0, n)| \leq \frac{1}{2}|t(y, \theta_0)|$ hold for all $0 \leq y \leq m$

and for $y = M$. Hence, for n being sufficiently large, $-t(M, \theta_0) + \Delta(M, \theta_0, n) < 0$ since $t(M, \theta_0) > 0$; and $-t(y, \theta_0) + \Delta(y, \theta_0, n) > 0$ for $0 \leq y \leq m$ since $t(y, \theta_0) < 0$ for $0 \leq y \leq m$.

In view of the above facts and by Theorem 2 of Hoeffding (1963),

$$\begin{aligned}
 & P\left\{\frac{1}{n} \sum_{j=1}^n A_j(M) < -t(M, \theta_0) + \Delta(M, \theta_0, n)\right\} \\
 & \leq \exp\{-2n[-t(M, \theta_0) + \Delta(M, \theta_0, n)]^2 a_3^{-1}(M, \theta_0)\} \\
 (2.17) \quad & \leq \exp\left\{-\frac{n}{2}[-t(M, \theta_0)]^2 a_3^{-1}(M, \theta_0)\right\}
 \end{aligned}$$

and for $0 \leq y \leq m$,

$$\begin{aligned}
 & P\left\{\frac{1}{n} \sum_{j=1}^n A_j(y) \geq -t(y, \theta_0) + \Delta(y, \theta_0, n)\right\} \\
 & \leq \exp\{-2n[-t(y, \theta_0) + \Delta(y, \theta_0, n)]^2 a_3^{-1}(y, \theta_0)\} \\
 (2.18) \quad & \leq \exp\left\{-\frac{n}{2}[-t(y, \theta_0)]^2 a_3^{-1}(y, \theta_0)\right\},
 \end{aligned}$$

where $a_3(x, \theta_0) = a_2(x, \theta_0) - a_1(x, \theta_0) = h(x) + h(x+1)\theta_0$.

Let

$$(2.19) \quad c = \frac{1}{2} \min \{t^2(y, \theta_0) a_3^{-1}(y, \theta_0) | 0 \leq y \leq m \text{ or } y = M\}.$$

It is clear that $c > 0$ since $m < \infty$ from Assumption (b) and $t^2(y, \theta_0) a_3^{-1}(y, \theta_0) > 0$, for all $0 \leq y \leq m$ and for $y = M$. Then from (2.11), (2.12) (2.16) to (2.19), we have

$$(2.20) \quad r(G, d_n^*) - r(G) \leq b_1 \sum_{y=0}^m \exp(-cn) + b_2 \exp(-cn) = O(\exp(-cn)).$$

Hence, the proof of this theorem is completed.

3. Examples and Remark

The following two examples have been considered by Johns and Van Ryzin (1971) and used to illustrate the performance of their proposed empirical Bayes decision rules for the two-action problem. We cite them and use the same to illustrate the performance of the proposed empirical Bayes decision rules $\{d_n^*\}$.

Example 1. (The Geometric Distribution). Suppose that

$$f(x|\theta) = \theta^x(1 - \theta), x = 0, 1, 2, \dots; 0 < \theta < 1;$$

and that the prior distribution has the probability density function $g(\theta)$ where

$$g(\theta) = (\alpha + 1)(1 - \theta)^\alpha, 0 < \theta < 1, \alpha > -1.$$

Then, $h(x) \equiv 1$ and $f(x) = \frac{(\alpha+1)\Gamma(x+1)\Gamma(\alpha+2)}{\Gamma(x+\alpha+3)}$. Thus, $\varphi(x) = \frac{h(x)f(x+1)}{h(x+1)f(x)} = \frac{x+1}{x+\alpha+3}$ which tends to 1 as $x \rightarrow \infty$. Taking $0 < \theta_0 < 1$, then, $A(\theta_0) = \{x|\varphi(x) \geq \theta_0\} \neq \emptyset$. Therefore, $m < M \equiv \min A(\theta_0) < \infty$. Hence, by Theorem 2.1, $r(G, d_n^*) - r(G) \leq O(\exp(-cn))$ for some positive constant c .

Example 2. (The Poisson Distribution). Let

$$f(x|\theta) = e^{-\theta}\theta^x/\Gamma(x+1), x = 0, 1, 2, \dots; \theta > 0.$$

Letting the prior density function be $g(\theta) = e^{-\theta}, \theta > 0$, we then have $f(x) = \frac{1}{\Gamma(x+1)}$ $\int_0^\infty \theta^x e^{-2\theta} d\theta = (\frac{1}{2})^{x+1}$, and $h(x) = \frac{1}{\Gamma(x+1)}$. Thus, $\varphi(x) = \frac{h(x)f(x+1)}{h(x+1)f(x)} = \frac{x+1}{2}$ which tends to infinity as x tends to infinity. Therefore, for any finite $\theta_0 > 0, m < \infty$. Then by Theorem 2.1, $r(G, d_n^*) - r(G) \leq O(\exp(-cn))$ for some positive constant c .

Johns and Van Ryzin (1971) considered several situations about the behavior of the tail probability of the prior probability density function, under which, their proposed empirical Bayes decision rules may achieve the best possible convergence rate $\alpha_n = n^{-1}$. We also apply those conditions to the sequence of the empirical Bayes decision rules $\{d_n^*\}$. We state the result as a corollary without citing the statement of those conditions. The reader is referred to Johns and Van Ryzin (1971) for detail.

Corollary 3.1. Let $\{d_n^*\}$ be the sequence of the empirical Bayes decision rules defined in Section 2. Suppose that $\int_0^Q \theta dG(\theta) < \infty$. Then, either under the assumptions in Theorem 3 or under the assumptions in Theorem 4 of Johns and Van Ryzin (1971), we have $r(G, d_n^*) - r(G) \leq O(\exp(-cn))$ for some positive constant c .

Proof: We need only to verify that $A(\theta_0) \neq \phi$ under each assumption. This can be done directly by noting the Lemmas 4, 5 and 6 of Johns and Van Ryzin (1971).

References

- Hoeffding, W. (1963). Probability inequalities for sums of bounded random variables *J. Amer. Statist. Assoc.* **58**, 13-30.
- Johns, M. V. Jr. and Van Ryzin, J. (1971). Convergence rates for empirical Bayes two-action problems I. discrete case. *Ann. Math. Statist.* **42**, 1521-1539.
- Robbins, H. (1956). An empirical Bayes approach to statistics. *Proc. Third Berkeley Symp. Math. Statist. Probab.* **1** 157-163, University of California Press.

Robbins, H. (1964). The empirical Bayes approach to statistical decision problems. *Ann. Math. Statist.* 35 1-20.

Robbins, H. (1983). Some thoughts on empirical Bayes estimation. *Ann. Statist.* 11 713-723.

UNCLASSIFIED

BEFORE COMPLETING FORM

1. REPORT NUMBER Technical Report #87-8	2. GOVT ACCESSION NO. AD-A121356	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ON THE CONVERGENCE RATES OF EMPIRICAL BAYES RULES FOR TWO-ACTION PROBLEMS: DISCRETE CASE	5. TYPE OF REPORT & PERIOD COVERED Technical	
7. AUTHOR(s) TaChen Liang	6. PERFORMING ORG. REPORT NUMBER Technical Report #87-8	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Purdue University Department of Statistics West Lafayette, IN 47907	8. CONTRACT OR GRANT NUMBER(s) N00014-84-C-0167	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Washington, DC	10. PROGRAM ELEMENT, PROJECT, TASK, AREA & WORK UNIT NUMBERS	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	12. REPORT DATE March 1987	
	13. NUMBER OF PAGES 13	
	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
	15a. DECLASSIFICATION DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release, distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Bayes rule; Empirical Bayes rule; Asymptotically optimal; Rates of convergence.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The purpose of this paper is to investigate the convergence rates of a sequence of empirical Bayes decision rules for the two-action decision problems where the distributions of the observations belong to a discrete exponential family. It is found that the sequence of the empirical Bayes decision rules under study is asymptotically optimal, and the order of associated convergence rates is $O(\exp(-cn))$, for some positive constant c , where n is the number of accumulated past experience (observations) at hand. Two examples are provided to illustrate the performance of the proposed empirical Bayes decision rules. A comparison is also made between the		

DD FORM 1473
1 JAN 73

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

20. proposed empirical Bayes rules and some earlier existing empirical Bayes rules.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

END

7-87

DTIC