

AD-A181 855

REPORT NO T15-87

# OPERATION EVEREST II: EFFECTS OF A SIMULATED ASCENT TO 29,000 FEET ON NUTRITION AND BODY COMPOSITION

U S ARMY RESEARCH INSTITUTE  
OF  
ENVIRONMENTAL MEDICINE  
Natick, Massachusetts

MAY 1987

DTIC  
SELECTED  
JUL 01 1987  
S D  
A R



Approved for public release; distribution unlimited.

UNITED STATES ARMY  
MEDICAL RESEARCH & DEVELOPMENT COMMAND

REPORT DOCUMENTATION PAGE

Form Approved  
OMB No 0704-0188  
Exp. Date Jun 30, 1986

1a. REPORT SECURITY CLASSIFICATION			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution is unlimited		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			4. PERFORMING ORGANIZATION REPORT NUMBER(S)		
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION U.S. Army Research Institute of Environmental Medicine		6b. OFFICE SYMBOL (if applicable) SGRD-UE-MN	7a. NAME OF MONITORING ORGANIZATION U.S. Army Medical Research & Development Command		
6c. ADDRESS (City, State, and ZIP Code) Natick, MA 01760-5007			7b. ADDRESS (City, State, and ZIP Code) Fort Detrick, MD 21701-5012		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS		
PROGRAM ELEMENT NO. 6.3A		PROJECT NO. 3M263763D 819	TASK NO. AI	WORK UNIT ACCESSION NO. DA305222	
11. TITLE (Include Security Classification) OPERATION EVEREST II: Effects of a simulated ascent to 29,000 feet on nutrition and body composition					
12. PERSONAL AUTHOR(S) Madeleine S. Rose, Charles S. Houston, Charles S. Fulco, Geoffrey Coates Dawn Carlson, John E. Sutton, Allen Cymerman					
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM Sep 85 TO Nov 85	14. DATE OF REPORT (Year, Month, Day) 1987 May 29		15. PAGE COUNT 128
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Body composition, percent body fat, hydrostatic weighing, acclimatization, high altitude, nutrition, food intake, weight loss		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Progressive body weight loss occurs during high mountain expeditions, but whether it is due to hypoxia, inadequate diet, malabsorption, or to the multiple stresses of the harsh environment is unknown. To determine whether hypoxia alone causes such body weight loss, six men were studied during progressive decompression to 240 torr for 40 days in a hypobaric chamber where hypoxia was the major variable. The subjects were provided a platable <u>ad libitum</u> diet that was modified for individual preferences. Complete food consumption data were available for only 38 of the 40 days. Mean caloric intake for 38 days at altitude was 2639±848 kcal/day (Mean±SD). Caloric intake decreased 42.3% from 3136 kcal during the first 7 days of exposure to altitude to 1789 kcal during the last 3 days at 282-240 Torr. During the same time periods the carbohydrate (CHO) intake decreased from 62.1% to 53.2% (p < 0.001). All subjects lost body weight with a mean loss of 7.44±2.24 kg (8.90% of the initial body weight) Loss of body weight was greater than could be accounted for by comparing calculated energy expenditures to actual caloric intake. Fluid balance data showed no evidence of hypohydration (key words)					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Madeleine S. Rose		22b. TELEPHONE (Include Area Code) (617) 651-4874		22c. OFFICE SYMBOL SGRD-UE-MN	

19 ABSTRACT (continued)

Pre- to post-exposure hydrostatic weighing indicated that 4 subjects lost 2.7% body fat whereas two subjects gained 0.65%. Computed Tomography scans indicated that most of the body weight loss was from muscle mass. The data indicate that prolonged exposure to increasing hypoxia does not increase CHO preference and that body weight could not be maintained despite provision of a highly palatable ad libitum diet.

### HUMAN RESEARCH and DISCLAIMER STATEMENTS

Human subjects participated in these studies after giving their free and informed voluntary consent. Investigators adhered to AR 70-25 and USAMRDC Regulation 70-25 on Use of Volunteers in Research.

The views, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other official documentation.

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



OPERATION EVEREST II:  
EFFECTS OF A SIMULATED ASCENT TO 29,000 FEET  
ON NUTRITION AND BODY COMPOSITION

MADELEINE S. ROSE, PhD<sup>1</sup>

CHARLES S. HOUSTON, MD<sup>2</sup>

CHARLES S. FULCO, MAT<sup>1</sup>

GEOFFREY COATES, MD<sup>3</sup>

DAWN CARLSON, MS<sup>1</sup>

JOHN R. SUTTON, MD<sup>3</sup>

ALLEN CYMERMAN, PhD<sup>1</sup>

<sup>1</sup>From USARIEM, Natick, MA 01760

<sup>2</sup>From Arctic Institute of North America, Washington, DC 20007

<sup>3</sup>From McMaster University, Hamilton, Ontario, Canada L8N 3Z5

Supported in part by the U.S. Army Research and Development Command  
under contract DAMD17-85-C-5306, the Arctic Institute of North America.

#### ACKNOWLEDGEMENTS

This paper is part of a series describing a study of acclimatization to hypobaric hypoxia conducted at and with help from the U.S. Army Research Institute of Environmental Medicine, Natick, Massachusetts. Principal Investigators were: Charles S. Houston, John R. Sutton, and Allen Cymerman. To the many who participated we express our appreciation, especially to the nine subjects who prefer to remain anonymous; their patience and sufferance made the project possible. This study would not have been possible without the help of William Sullivan, Peter Bangs, Sue Porter, and Esther Bartfeldt who helped feed the subjects. The data could not have been presented here without the computer expertise of Joseph Williams, Christopher Walsh, R. Brian Wenzinger, Edward Roche, Tom Dugan, and Robert White. The authors wish to thank Lisa Hodgess for her technical assistance. We are indebted to Dr. Philip Costello and Ms. Laura Wong of the Dept of Radiology New England Deaconess Hospital, Boston, MA for their invaluable assistance with the Computed Tomographic scans.

## TABLE OF CONTENTS

	Page
Acknowledgements	iii
Table of Contents	iv
List of Tables	v
List of Figures	vi
Abstract	viii
Introduction	1
Methods	3
Results	7
Discussion	11
Conclusions	21
References	22
Tables	26
Figures	46
Figure Legends	47
Appendices	83
Appendix 1 - Menus	84
Appendix 2 - Fluid Data	108

## LIST OF TABLES

NO.	TITLE	PAGE
1	Physical Characteristics of Subjects	27
2	Periods	28
3	Energy Intake (kcal/day)	29
4	Percent Decrease in Energy Intake	30
5	Daily Energy Expenditure from Exercise	31
6	Average Energy Requirements	32
7	Energy Balance and Weight Change in 38 Days	33
8	Weight Loss in 38 Days	34
9	Fluid Balance	35
10	Extremes of Energy Intake	36
11	Protein Intake (grams/day)	37
12	Percent Calories from Protein	38
13	Mean Nutrient Intake for 38 Days	39
14	Percent Calories From Carbohydrates	40
15	Percent Calories From Fat	41
16	Skinfold Measurements (mm) Pre- and Post-Exposure to 40 Days of Progressive Decompression to 240 torr	42
17	Average Fluid Balance (ml/day)	43
18	Trends in Food Ratings and Number of Food Groups Consumed	44
19	Trends in Ratings of Different Food Groups	45



## TABLE OF FIGURES

NO.	TITLE	PAGE
1	Energy Expenditure From Exercise: Subject 1	47
2	Energy Expenditure From Exercise: Subject 3	48
3	Energy Expenditure From Exercise: Subject 4	49
4	Energy Expenditure From Exercise: Subject 5	50
5	Energy Expenditure From Exercise: Subject 6	51
6	Energy Expenditure From Exercise: Subject 7	52
7	Energy Expenditure From Exercise: Subject 8	53
8	Energy Expenditure From Exercise: Subject 9	54
9	Duration of Exercise: Subject 1	55
10	Duration of Exercise: Subject 3	56
11	Duration of Exercise: Subject 4	57
12	Duration of Exercise: Subject 5	58
13	Duration of Exercise: Subject 6	59
14	Duration of Exercise: Subject 7	60
15	Duration of Exercise: Subject 8	61
16	Duration of Exercise: Subject 9	62
17	Energy and Weight Measurements: Subject 1	63
18	Energy and Weight Measurements: Subject 3	64
19	Energy and Weight Measurements: Subject 4	65
20	Energy and Weight Measurements: Subject 5	66
21	Energy and Weight Measurements: Subject 6	67
22	Energy and Weight Measurements: Subject 7	68
23	Energy and Weight Measurements: Subject 8	69

TABLE OF FIGURES (cont'd)

NO.	TITLE	PAGE
24	Energy and Weight Measurements: Subject 9	70
25	Composition of Diet	71
26	Body Composition Changes: Percent Body Fat by Hydrostatic Weighing	72
27	Body composition changes: Body Fat Weight as Determined by Hydrostatic Weighing	73
28	Body composition changes: Circumference Measurement of Arm	74
29	Body composition changes: Circumference Measurement of Thigh	75
30	Body composition changes: CT Scan of Arm	76
31	Body composition changes: CT Scan of Thigh	77
32	Body composition changes: CT Scan of Arm	78
33	Body composition changes: CT Scan of Thigh	79

## ABSTRACT

Progressive body weight loss occurs during high mountain expeditions, but whether it is due to hypoxia, inadequate diet, malabsorption, or to the multiple stresses of the harsh environment is unknown. To determine whether hypoxia alone causes such body weight loss, six men were studied during progressive decompression to 240 torr for 40 days in a hypobaric chamber where hypoxia was the major variable. The subjects were provided a palatable ad libitum diet that was modified for individual preferences. Complete food consumption data were available for only 38 of the 40 days. Mean caloric intake for 38 days at altitude was  $2639 \pm 848$  kcal/day (Mean  $\pm$  SD). Caloric intake decreased 42.3% from 3136 kcal during the first 7 days of exposure to altitude to 1789 kcal during the last 3 days at 282-240 Torr. During the same time periods the carbohydrate (CHO) intake decreased from 62.1% to 53.2% ( $p < 0.001$ ). All subjects lost body weight with a mean loss of  $7.44 \pm 2.24$  kg (8.90% of the initial body weight). Loss of body weight was greater than could be accounted for by comparing calculated energy expenditures to actual caloric intake. Fluid balance data showed no evidence of hypohydration. Pre- to post-exposure hydrostatic weighing indicated that 4 subjects lost 2.7% body fat whereas two subjects gained 0.65%. Computed Tomography scans indicated that most of the body weight loss was from muscle mass. The data indicate that prolonged exposure to increasing hypoxia does not increase CHO preference and that body weight could not be maintained despite provision of a highly palatable ad libitum diet.

## INTRODUCTION

### CALORIC INTAKE AND BODY WEIGHT

Loss of body weight is common during exposure to high altitude (1-13). Acute mountain sickness (AMS) frequently occurs after rapid ascent to moderate altitude and may cause body weight loss due to anorexia, nausea, vomiting, and dehydration, or it may cause weight gain due to fluid retention. Headache, weakness, and disordered sleep are other symptoms of AMS which interfere with normal activity. Altitude sickness is dependent on elevation, rate of ascent, and acclimation but usually decreases after the first three days (14-22). However appetite suppression and decreased caloric intake can persist for as long as several weeks (1,5,8,9,11). As exposure continues for weeks, normal values for food and water intake have been approached (1,9,18,23). Acclimatization can lessen the severity of AMS at altitude to lead to a lessening of anorexia (1,4,15,20-22) and of weight loss. However, even after the symptoms of AMS have lessened, body weight can remain reduced. Appetite suppression may decrease caloric and protein intake by as much as 30% and 40% respectively (1,5,8,9,11,24) and thus may explain the body weight loss which persists after AMS symptoms have subsided.

Studies of sustained body weight loss are numerous and contradictory and have attributed this loss to reduced dietary intake resulting from anorexia (1,5-13); loss of stored body fat (5,7,11,25,26); loss of body fluid (1-5,11,25,27-30); malabsorption of ingested fat and/or carbohydrate especially at altitudes in excess of 4300 m (4,31,32); and an elevated rate of energy expenditure under basal, resting, and mild to moderate exercise conditions (4,17,18,20,33-39). Body weight loss ranging from 3-5.4% (5,11,40,41) with rates of 145-330 g/day have been reported (11,23,42).

On an actual climb, it is difficult to determine whether body weight loss is due to increased energy expended in hard physical work, the cold environment, the limited availability or palatability of food, dehydration, malabsorption, or combinations of these stresses. Furthermore, lack of time, space, equipment, and interest make precise intake/output measurements very difficult. As a result, expeditions usually only estimate food intake (23,40).

#### CARBOHYDRATE PREFERENCE

There are a paucity of reports and a conflict of views regarding food preference at altitude (9,13,23,24,29,37,40). Anecdotal evidence from climbers suggests that they prefer sweets at altitude (9,13,23,24,29,40). However, researchers have suggested that the preference for carbohydrate was really due to a greater availability and ease of preparation (23,24). Few studies have determined dietary preferences for carbohydrate at high altitudes when subjects are given (9,37) palatable foods ad libitum.

#### FOOD RATING

Researchers have not studied food ratings to determine whether the problem is that food does not taste good or whether AMS causes the subject to reduce his food intake no matter how good the food tastes.

#### BODY COMPOSITION

Although loss in body weight at altitude is common, the composition of the weight loss is unclear. Some investigations show a reduction in body fat (5,11,17,23,25,43,44) while others show a loss in lean body weight (1,45).

The purpose of the present study was to determine if dietary preferences changed and if body weight could be maintained by offering palatable foods ad libitum during a simulated ascent of Mount Everest (240 torr).

## METHODS

Nine subjects in excellent physical condition were selected to participate in this study. One subject was excluded at the beginning of the study because of an upper respiratory infection and 2 subjects (#7 & #5) were withdrawn at 18,000 and 25,000 feet, respectively. All of the subjects are described by Houston (46). The subjects were confined in a hypobaric chamber for 40 days during progressive decompression to 240 Torr to simulate previously successful ascents of Mount Everest. Complete dietary data were available on only 6 subjects for 38 days due to incomplete data on the first and last days of confinement.

The subjects were allowed access to any quantity and variety of palatable food. The menus and food preparation were under the supervision of a registered dietitian. The subjects selected three meals per day from menus consisting of approximately 3000 kcal/day (60% CHO, 15% PRO, 25% FAT). The menus included a full breakfast selection, soup and/or sandwich lunches, commercially prepared dinners, simple home-cooked dinners, and food accompaniments to balance meals (Appendix 1). Foods were weighed before being served to the subjects. The returned foods were measured and subtracted from the amount served to accurately measure food intake. Any discrepancies between the menu and food items on the tray were resolved.

A variety of foods were available in the chamber for between meal snacking. Assorted fluids were available ad libitum and subjects were encouraged to drink to prevent dehydration. The subjects recorded any foods and fluids that were ingested between the three meals. The food and fluid intakes were analyzed using the University of Massachusetts Nutrient Data Bank program.

Food ratings were obtained as the subjects consumed the food. They were asked to rate each food item on a hedonic rating scale from 0 to 9 with 9 being "Like Extremely", 1 was "Dislike Extremely", and 0 was "Didn't Try".

The subjects were weighed daily on a physician's scale calibrated to the nearest 1/4 pound. Body heights (cm) were taken at the beginning of the study.

Energy expenditure could not be measured exactly; therefore, the Harris Benedict formula (47) was used to estimate basal energy expenditure (BEE). Because the subjects were confined in a small space, only 20% of BEE was added for daily living activities. The subjects kept a daily log of their voluntary exercises on the treadmill, stationary ladder ergometer, and cycle ergometer. The subjects were encouraged to exercise to a level comparable to that of an actual mountain climbing expedition. The calories expended from these exercises were added to the  $BEE + 20\%BEE$  value. Since so many of the tests required exercise, they also were included in the log book of exercise activity. Some of the tests that included extensive exercise regimens were:  $VO_2max$  (MAX), Inert Gas (GAS), Muscle Biopsy (MUSCLE), and Swan-Ganz Catheterizations (SG). The energy expenditure calculations were not adjusted for possible changes in rate of energy expenditure due to altitude.

Body composition information was collected at sea level pre- and post-altitude exposure. Body weight, whole body density (hydrostatic weighing), residual lung volume, and anthropometric measurements were obtained preprandial 3 to 6 days prior to and within 20 hours after the chamber confinement. During the hydrostatic weighings, a minimum of six practice trials were performed before three actual readings were recorded to assure a

true baseline. Residual volumes were obtained while the subjects were underwater during the each of the last three readings. Body density was converted to percent body fat (%BF) using the Siri formula (48).

Anthropometric measurements were obtained before noon on each of the measurement days by the same investigator throughout the entire study. Exercise was not permitted 1 hour prior to the anthropometric measurements. This assessment consisted of seven skinfold and 10 soft tissue circumference measurements obtained while subjects were standing. Two non-consecutive measurements were obtained at each site. A third measurement was taken if the difference between the first two was greater than 1%. The mean of the two or three measurements was used in the statistical analysis. Skinfolds were obtained at seven sites on the right side of the body (triceps, scapula, chest, midaxillary, suprailiac, abdomen, and thigh) using a Harpenden skinfold caliper with a caliper pressure of 10 g/mm<sup>2</sup>. Circumferences of the neck, shoulders, chest, hips, thigh, calf, upper arm, forearm, and two abdominal sites were measured using a Gulick spring-loaded tape. The measurement sites were located using standard landmarks.

To quantitatively assess the pattern of fat and muscle distribution (49), computed tomographic (CT) scans were taken of the upper arms and thighs at sea level, pre- and post-altitude exposure. Five 1 cm thick slices were scanned in each limb. In the thigh the center slice was positioned on a mark at mid-point between the tibial tubercle and the anterior superior iliac crest; i.e. mid thigh. In the arm the center slice was positioned on a mark 60% of the distance down from the acromium to the medial condyle of the ulnar. This sampled the thickest part of the upper arm. The area of the



bone, muscle, and fat on each slice was measured with a planimeter interfaced to an IBM personal computer. To allow for possible changes in the CT scanner magnification between the pre and post hypoxia images the fat and muscle content of each slice was expressed as a ratio of fat/bone and muscle/bone. It was assumed that there would be no significant change in the diameter of the ulna or femur in the 42 days between measurements. The data from the 10 slices in both limbs were pooled in each individual. The data presented represent the means of these pooled slices in all individuals.

#### STATISTICS

Descriptive statistics were run on the 6 subjects that completed the 38 days of the study. Mean  $\pm$  SD were calculated for each descriptive measurement. Data for the 2 subjects (#5 & #7) that were withdrawn from the study were included in the charts but were not included in the calculations of the means. Analysis of variance was used to test for significant differences between time periods. The Student Newman Keuls post hoc tests were used to determine where the significant differences occurred.

## RESULTS

### CALORIC INTAKE AND BODY WEIGHT

The physical characteristics of the subjects on day 2 of the study are described in Table 1. The 38 days for which complete dietary data were available were divided into 7-day periods except for the last period which only included 3 days (Table 2). Grouping the data into 6 periods made it easier to interpret overall trends.

Caloric intake decreased over the 38 days at altitude. Caloric intake during period 1 averaged  $3136 \pm 557$  kcal (Mean $\pm$ SD) and gradually decreased to  $1789 \pm 682$  kcal during period 6 (Table 3). Mean caloric intake averaged  $2639 \pm 848$  kcal/day. The percent decrease in energy intake between Periods 1 and 6 was 42.3% ranging from 23.3 to 62.7% (Table 4). Two subjects (#4 & #6) were able to limit the percent decrease in calories to 28.2 and 23.3%, respectively.

The number of calories expended, time, and number of times per day that the subjects engaged in exercise varied widely (Table 5). The subjects expended more calories exercising during the first half of the study (Figures 1-8). The amount of time spent exercising was higher at the lower altitudes or at the beginning of the study (Figures 9-16).

Average energy requirements ranged from 2372 to 4391 kcal/day (Table 6). Comparing the calculated energy requirements to the energy intake showed that the actual weight loss was greater than expected (Table 7) by as much as 8.2 kg. The mean caloric intake for 4 subjects was less than their calculated energy requirements but the other 2 subjects ingested more calories than were needed to meet their energy requirements (Table 7).

Mean body weight loss for the 38 days was  $7.44 \pm 2.24$  kg ( $p < 0.01$ ) ranging from 4.2 to 9.9 kg (Table 8). The subjects lost  $8.9 \pm 2.0\%$  of their sea level body weights. The weight loss ranged from 4.2% to 11.6% of original body weights in 38 days. Body weight was lost at a mean rate of 196 g/day (range 110.5 - 260.5 g/day). Body weight and caloric intake (Figures 17-24) followed the same general downward trend. Overall fluid balance was positive to discount fluid loss as a reason for weight loss (Table 9).

Comparing the value of the highest caloric intake to the lowest value for each subject (Table 10) showed a mean difference of 3268 kcal which could cause a 0.45 kg weight loss/day if continued. The difference between the highest and lowest caloric intakes was significant ( $p < 0.01$ ). The highest values occurred before the minimum values during the course of the study.

Five of the subjects had mean protein intakes that exceeded the RDA of 0.8 g/kg body weight (BW) (Table 11). The protein intake for these 5 subjects fell below the RDA on several days but there were surrounding days with more than adequate protein intakes. For these subjects the days of low protein intakes occurred more frequently in the last two periods of the study. The sixth subject (#1) had a mean protein intake of 83.7 g which was 83.0% of his sea level protein needs. The subject's protein intake was deficient 63.2% or 24 of the 38 days of the study and started occurring in the first period. The mean protein intakes for the six subjects showed a significant decrease ( $p < 0.001$ ) during the second half of the study (Table 11). As the caloric intake decreased, the protein intake also decreased ( $r = 0.77$ ,  $p < 0.001$ ). The percent of calories from protein (%PRO) did not differ significantly except for period 2 which was significantly greater

( $p < 0.01$ ) than the other periods (Table 12). The mean protein intake for all 38 days was  $15.4 \pm 3.8\%$  of the total calories.

The mean nutrient intake of the 6 subjects met the Recommended Dietary Allowances (RDA) (50) for all vitamins and minerals (Table 13). However two subjects (#4 & #6) were ingesting vitamin and mineral supplements which artificially increased the mean intake. Excluding the vitamin/mineral supplement from the data showed that the food intake provided enough nutrients to meet the RDA for energy and all vitamins and minerals except for vitamin B<sub>6</sub>, folacin, magnesium, and zinc.

#### CARBOHYDRATE PREFERENCE

The subjects started the study consuming a higher percentage of total calories as carbohydrate (%CHO) in their diets (62.1% CHO); but the %CHO dropped to 54.5% in the second period and remained in that general range for the remainder of the study (Table 14). There was a significant difference ( $p < 0.001$ ) between period 1 and the following periods. As the %CHO decreased, the percent of calories as fat (%FAT) increased ( $r = -0.90$ ,  $p < 0.001$ ) (Table 15, Figure 25).

#### FOOD RATING

The overall rating of the food showed that the subjects liked the foods very much. The mean rating was very high at 8.4 on the hedonic rating scale. The mean rating for the first half of the study was 8.4 (range 3-10) compared to 8.7 for the second half (range 4-10). The mean ratings showed a slight increase during the second half of the study; however, the amount of food that the subjects ate decreased.

## BODY COMPOSITION

According to the hydrostatic weighing measurements, there was an overall 1.6% decrease in %BF pre- to post-exposure to altitude but the difference was not significant (Figure 26). Four of the 6 subjects lost an average of 2.7%BF whereas the other 2 subjects (#8 & #9) gained an average of 0.65%BF. Mean energy expenditure was greater than the energy intake for the 4 subjects whose %BF decreased (Table 7). The 2 subjects that gained %BF were consuming sufficient calories so that they should have gained instead of losing body weight. Calculations of body fat weight (kg) showed a consistent decrease in body fat (Figure 27) for all subjects. Circumference measurements significantly decreased ( $p < 0.05$ ) in the arms and thighs of all subjects (Figures 28-29). Skinfold measurements, which are an indication of subcutaneous fat, decreased in the triceps and thighs except for the 2 subjects (#8 & #9) whose %BF increased according to hydrostatic weighing measurements (Table 16). The sum of all 7 skinfold measurements followed the same trend as hydrostatic weighing with skinfolds decreasing for 4 subjects and increasing for 2 subjects. The CT scan fat/bone data confirmed the trends of the hydrostatic weighing and skinfold data (Figures 30-31). Circumference, %BF, skinfold, and CT scan data showed that body fat decreased except for the 2 subjects (#8 & #9) who had increases in %BF.

Muscle mass decreased (Figures 32-33) in all subjects. CT scan muscle/bone measurements showed that the difference was not significant in the arm ( $p < 0.073$ ) but was significant ( $p < 0.002$ ) in the thigh measurement. Since bone measurements were not expected to change during the 40-day period of the study, change in the ratio was attributed to changes in muscle mass.

## DISCUSSION

### CALORIC INTAKE AND BODY WEIGHT

Averaging the data into 7-day periods showed that there was a definite downward trend in caloric intake over the six periods. The largest decreases in caloric intake occurred at the higher altitudes. The average decrease in caloric intake of 42.3% from period 1 to period 6 (Table 3) was comparable to the 20-40% range found in the literature for sojourns of up to 26-days duration (5,8,9,24). Two subjects (#4 & #6) were able to maintain their caloric intake at 71.8 and 76.7% of their beginning values. The small decrease in calories for the 2 subjects (#4 & #6) could be due to the fact that they ate less than their energy requirements in the beginning of the study. The body weight losses for the 2 subjects (#4 & #6) were among the higher values with percent body weight loss being equal to 11.6 and 9.7% (Table 8). Since their energy expenditures due to exercise were greater than those of the other subjects, the large body weight losses are to be expected (Figures 1-8). Energy requirements have been shown to be relatively higher for heavy exercise at altitude compared to that at sea level (36). Since these subjects were heavy exercisers, it is possible that the energy expenditure was increased above that predicted by the Harris Benedict equation. Increased energy expenditure coupled with their decreased energy intake could lead to large energy deficits which would lead to the catabolism of body protein and/or fat (1,5,9,11,24,25,38,43). The decrease in caloric intake was less than 50% for four of the subjects. The 4 subjects started the study eating more than their energy requirements. Boyer and Blume (23) had reported that some climbers ate excess amounts of food before starting to

climb a mountain in anticipation of anorexia and body weight loss. The 4 subjects may have been attempting to eat as much as possible at the beginning of the study to increase their body stores and to prevent the debilitating effects of a major weight loss. Their plans seemed to have worked since these subjects lost less weight in comparison to subjects #4 and #6 even though they suffered more from AMS.

Altitude sickness is affected by altitude, speed of ascent, and length of stay. Mild AMS subsides in a few days; high altitude pulmonary edema and high altitude cerebral edema either subside or grow worse and require descent. The longer the stay at altitudes up to 18-20,000 feet, the more complete the acclimatization. However, caloric intake and body weight tend to fall during long stays at altitudes higher than 20,000 feet (4).

The subjects were allowed to acclimate for 3-day periods at 15,000, 18,000, 20,000, and 25,000 feet. The subjects in the present study tended to feel better after each of the three day acclimation periods (46), but nevertheless all subjects lost body weight continuously throughout the study. This suggests that hypobaric hypoxia rather than the combined stresses of the mountain environment may be responsible for a major portion of the deterioration of food intakes during mountain expeditions. Due to the nature of this study, body weight loss was not due to increased climbing activity or to limited food availability. The amount of voluntary exercise decreased drastically for most of the subjects after reaching 23,000 feet on day 26. These results suggested that deterioration in physical condition could override the effects of acclimatization as Pugh (4) and Blume (24) had suggested as early as 1962.

The 8.9% body weight loss in the present study was greater than previously reported. Previous studies have shown weight loss of 3-3.49% in 8 days (5,41); 5.4% in 12 days (11); and 5% in 5 weeks (40) during exposure to altitude. The caloric intake and rate of weight loss for the present study were similar to those of previous studies (11,23,40,42). If the duration of those other studies were extended, the percent body weight loss would be similar to the present study.

All subjects continuously lost body weight during the 38 days at altitude. The mean energy intake for 4 subjects was less than their calculated energy requirements so weight loss was expected. However the other 2 subjects (#8 & #9) ingested more calories than were needed to meet their energy expenditure and should have gained body weight (Table 7). The weight loss for all subjects was greater than would be expected from the energy intake and expenditure figures. The reason for the disparity in body weight losses may simply be a question of the accuracy of the estimation of energy expenditure since energy intake was accurately determined. The subjects with caloric intakes greater than their energy requirements did tend to have smaller body weight losses. Other researchers have suggested that body weight loss may be due to water loss, increased metabolic rates, and fat and/or carbohydrate malabsorption (4,23). Hypohydration may be due to decreased fluid intake (1); increased water loss (2,3,4,11,27); or loss of fluid from the lungs due to hyperventilation (28,29). However, other studies have shown that normal body hydration can be maintained (5,25,30). The fluid data for the present study showed that the subjects were in a slightly positive balance (Table 9). Fluid intake exceeded the output by several



hundred milliliters. However losses of fluid from hyperventilation and sweating were not taken into account and would probably decrease the positive fluid balance to euhydration. Body fluid losses were probably minimized because the relative humidity in the hypobaric chamber ranged from 60 to 80% depending on the altitude. In general the fluid balance was negative during the last two periods of the study (Table 17, Appendix 2). The negative fluid balance could account for the continued weight loss of subjects #8 and #9 who should have gained weight according to energy balance calculations. Several researchers have suggested that energy expenditure increases at altitude (17,18,39). The effects of altitude on basal (4,17,33), resting (34,35,39), and mild to moderate exercise (35,36) conditions have been studied and show an increase of 7-11% of energy expenditure at altitude. The increase in energy expenditure has been attributed to the increased work of breathing or the decreased efficiency of work performance. However data from other studies show that the increase in basal energy expenditure is transient and returns to normal after about a week of acclimatization (4,20,37,38).

The possibility that malabsorption due to hypoxia was an important cause of body weight loss has been debated (4,23,31,32) but was not investigated in this study. On mountain expeditions food intake and energy output have only been estimated (23,24) and not at altitudes as high as attained in the present study. Rai et al. (32) and Sridharan (31) showed no disturbance in digestibility and utilization of dietary fat and/or carbohydrate up to 4700 m. Other studies have reported malabsorption of fat and/or xylose (4,23) at higher altitudes. Increased energy expenditure/metabolic rates and/or malabsorption could account for the body weight losses but neither were studied.

The confinement of the 8 subjects in a hypobaric chamber in the present study could have affected body weight loss. Loss of appetite, body weight, and strength could be attributed to confinement in the hypobaric chamber, isolation from the real world, limited physical activity, and boredom. However evidence was not available on the interaction of these factors with body weight and appetite loss. A study of 245 crew members confined to 4 submarines showed that 67% of the personnel lost weight, 25% gained weight, and 8% maintained their weight during long sea voyages even though the food was considered very palatable and exercise decreased (51). The authors attributed the weight loss of a third of the crew members to active dieting but the psychological and physical stresses of confinement may have contributed to the weight loss of the remainder of the crew members (51).

Negative nitrogen balances have been reported in humans exposed to high altitudes (1,41). Under normal conditions, the RDA for protein of 0.8 g/body weight (50) is more than adequate for maintaining nitrogen balance. When energy intakes were less than 2000 kcal/day, nitrogen balance studies on protein intakes of greater than 50-60 g/day were negative during altitude exposure (1,8,38,41). Negative nitrogen balances during caloric deficiency were probably due to protein being used to meet the energy needs of the body (45). Studies have shown that whenever men consumed adequate calories and protein during altitude exposure, positive nitrogen balances were possible (37,42). However decreased appetite usually leads to a decrease in dietary caloric and protein levels (5,8,9,24,41). Nitrogen balance studies were not conducted but five subjects received more than the RDA for protein while one subject (#1) ingested less than the RDA for protein. In the present study

the mean caloric intakes for 2 subjects (#8 & #9) were adequate to meet energy requirements but the levels were inadequate for the other 4 subjects (Table 7). The subjects were able to ingest an average of 2152-3022 kcal but their energy requirements ranged from 2314 to 2901 kcal. Four subjects in the present study were probably in negative nitrogen balance because of their caloric deficits. The 2 subjects receiving adequate calorie and protein intakes were probably in positive nitrogen balance (42).

Hannon (52) has concluded that extended periods of low protein intakes are unlikely to present serious nutritional problems since the protein intakes during the most pronounced stages of hypophagia usually exceeded the RDA of 0.8 g/kg BW. This conclusion was not supported by the results of the present study. Protein intake averaged 0.86 g/kg BW for the last period of the study but 2 subjects (#1 & #3) had protein intakes as low as 0.52-0.67 g/kg BW for the last 2 periods of the study. These 2 subjects were probably in negative nitrogen balance. Protein intake decreased on one day to 26.2 g for one of those subjects (#1) whose protein intake for the entire study was deficient 63.2% of the time.

Hypophagia can last from 8-28 days (1,5,8,23,38) at altitude although clinical evidence of nutritional deficiency other than a negative nitrogen balance has not been reported. It has been suggested that subjects starting in a good nutritional state would not develop vitamin and mineral deficiency problems for several months (52). However the duration of the hypophagia and anorexia of AMS may indicate a need for vitamin/mineral supplementation for subjects on lengthy expeditions who are exercising heavily and whose diets are unbalanced.

## CARBOHYDRATE PREFERENCE

Several studies have shown that subjects prefer carbohydrate at high altitude, usually at the expense of fat (4,18,37). Consolazio et al. (37) showed that this was true during the first week at altitude but a similar pattern was not apparent during prolonged exposure. The present study confirmed these findings. Though a high carbohydrate (60%) diet was offered throughout the study, the subjects started at 62.1% and then decreased their carbohydrate intake to 53.2%, although the menu choices were adequate to have maintained or even increased carbohydrate intakes. Others (14,23) have found intakes of carbohydrate to increase from 42-55% at sea level to 52-63% at altitude. In the present study, a wide variety of appetizing foods were readily available to the subjects and they consumed a more balanced diet than usually available to mountaineers. The composition of the diet for the entire study was 54.5% CHO, 15.2% PRO, and 31.4% FAT. In 1969, Kryzwicki et al. (11) observed similar body weight losses whether the subjects consumed a high carbohydrate (68%) or a low (42%) carbohydrate diet during 12 days at 4300 m. Increased dietary carbohydrate intake did not decrease body weight loss. Consolazio et al. (18) found a difference in work performance for the heaviest work levels with a high carbohydrate diet at high altitude. Carbohydrate may increase the blood oxygen transport capacity and improve metabolic efficiency (53,54). This may effectively decrease the physiological altitude by 1-2000 feet (55) and make physiological exertion relatively easier. The subjects in the present study did not increase or maintain their intake of carbohydrate.

The subjects did not eat pure fat such as butter but they did eat foods containing large amounts of fat such as ice cream, cream soup, whole milk, meat, and french fries. Although they asked for high carbohydrate snacks such as candy, granola bars, etc, these food also contain large amounts of fat. The preference may not be for pure carbohydrate but a desire for palatability resulting from the combination of carbohydrate and fat.

#### FOOD RATINGS

The high hedonic ratings indicted that the subjects liked the foods. The food was palatable but at higher altitudes the subjects' appetites were diminished compared to their appetites at lower altitudes (Table 18). Toward the end of the study the subjects ate less food but the foods selected were more calorically dense. The low calorie vegetables and salads were not requested as often. Caloric intake was greater than energy requirements on some days indicating that the subjects were attempting to increase their caloric intake (Figures 17-24). Some subjects were able to consume enough food to keep their caloric intake above their energy requirements right up to the 36th day of the study. It was possible to encourage the subjects to ingest more calories but they could not sustain the high intake. Sore throats, feeling full, not hungry, and the problem of eating while struggling to breathe were reasons given by the subjects for their reduced food consumption. Many foods which were favorites at sea level were not tolerated at higher altitudes.

The mean ratings for the different food groups generally increased (Table 19) as the study progressed. The fruit and juice, potato, bread, vegetable, and meat groups had more food items whose rating increased toward the end of

the study (Table 19). The ratings for potatoes increased for all subjects. However the potatoes were usually combined with cheese and other high protein and high fat additives and many variations were possible. The ratings decreased for the foods that were available throughout the study. The subjects were offered the foods at least five times during the course of the study. They were provided other foods as requested. The decrease was more likely due to boredom rather than an inability to tolerate the foods.

The food ratings of one subject (#9) increased for all foods during the study (Table 19). This subject also was able to minimize his weight loss (Figure 24). His mean caloric intake was greater than his energy requirements so his ability to keep his weight loss minimized was probably due to the high food intake rather than water retention especially since his fluid balance was negative during the last period of the study.

#### BODY COMPOSITION

There was a mean decrease in body weight, %BF, body fat weight, and LBM. Percent body fat measurements by hydrostatic weighing, skinfold measurements, and CT scanning increased in 2 subjects (#8 & #9). The fact that these 2 subjects (#8 & #9) ate more calories than they needed for energy expenditure and reduced their exercise provided support for the possibility that subcutaneous fat was maintained or increased slightly. In general the present study confirmed the results of previous studies (25,43) that showed losses in skinfold thickness and %BF (5,11,17,44). Although there was an mean decrease in skinfold thickness and %BF in the present study, there was a wide range from 2 subjects (#8 & #9) who showed increases in skinfold measurements and %BF to subjects who showed losses up to 33% in skinfold

thickness. Fulco (56) reported an increase in the %BF determined by hydrostatic weighing after an 18-day exposure to 14,100 ft in a hypobaric chamber. However, the increase in %BF may be misleading. The 1.1% increase in %BF in the Fulco (56) study may have occurred because the amount of fat did not change even though the subjects lost 1.5 kg body weight. The subjects in the present study lost body weight also. Analyzing the body fat weight of the subjects in the present study showed that all subjects lost body fat but the loss was very small for 2 subjects (#8 & #9). Skinfold measurements increased verifying the increase in subcutaneous fat (Table 16). Mountain climbing expeditions have never reported increases in %BF. This difference is probably due to the fact that the subjects in the hypobaric chamber were not exercising as much or in the same manner as would occur during an actual assault on a mountain.

About 25.8% of the 7.44 kg of weight lost by the subjects was from fat and 74.2% from the LBM. The most common percentage of fat loss was approximately 40% but adding the values of the 2 subjects (#8 & #9) who gained %BF lowered the percent attributed to fat considerably. Since the subjects were not dehydrated and bone was not expected to change, the 74.2% loss from LBM was from muscle mass. The CT scans confirmed the decreases in muscle mass.

The CT scans and the measurements of %BF, skinfold, and circumference presented an integrated picture of the changes in body composition. The overall trend was of decreasing muscle mass and body fat except for 2 subjects (#8 & #9) who actually gained fat.

## CONCLUSIONS

Conclusions from this study were: (1) energy output decreased as altitude increased; (2) food intake decreased as altitude increased; (3) fluid balance indicated little or no dehydration or edema in most subjects most of the time; (4) essential vitamins and minerals and protein were not deficient except for protein for one subject; (5) lack of oxygen, lack of adequate exercise, and confinement may have caused anorexia which contributed to body weight loss; (6) the amount of body weight loss can not be accounted for by balance of caloric intake vs energy expenditure indicating that there might have been malabsorption or increased energy expenditure that has not been taken into account; (7)  $\bar{X}$ BF decreased in 4 subjects but increased in 2 subjects whose caloric intake indicated that they should have gained weight; (8) body fat weight decreased in all subjects; and (9) LBM decreased.

The data indicate that prolonged exposure to increasing hypoxia does not increase carbohydrate preference and that body weight could not be maintained despite provision of a highly palatable ad libitum diet.



## REFERENCES

1. Consolazio CF, Matoush LO, Johnson HL, Daws TA. Protein and water balances of young adults during prolonged exposure to high altitude (4,300 meters). *Am J Clin Nutr* 1968;21:154-161.
2. Howell A, Cove DH. Birmingham Medical Research Expeditionary Society 1977 Expedition: the diuresis and related changes during a trek to high altitude. *Postgrad Med J* 1979;55:471-474.
3. Jain SC, Bardhan J, Swamy YV, Krishna B, Nayar HS. Body fluid compartments in humans during acute high-altitude exposure. *Aviat Space Environ Med* 1980;51:234-236.
4. Pugh LGCE. Physiological and medical aspects of the Himalayan Scientific and Mountaineering Expedition, 1960-61. *Br Med J* 1962;2:621-627.
5. Surks MI, Chinn KSK, Matoush LO. Alterations in body composition in man after acute exposure to high altitude. *J Appl Physiol* 1966;21:1741-1746.
6. Consolazio CF, Johnson HJ, Krzywicki JH. Body fluids, body composition, and metabolic aspects of high-altitude adaptation. In: Yousef MK, Howath SM, Bullard RW, eds. *Physiological Adaptation: desert and mountain*. New York: Academic Press, 1972:227-241.
7. Whitten BK, Janoski AH. Effects of high altitude and diet on lipid components of human serum. *Fed Proc* 1969;28:983-986.
8. Whitten BK, Hannon JP, Klain GJ, Chinn KSK. Effect of high altitude (14,100 ft.) on nitrogenous components of human serum. *Metabolism* 1968;17:360-365.
9. Hannon JP, Klain GJ, Sudman DM, Sullivan FJ. Nutritional aspects of high altitude exposure in women. *Am J Clin Nutr* 1976;29:604-613.
10. Hingston RWG. Physiological changes at high altitudes and their relation to mountain sickness. *Indian J Med Res* 1921-1922;9:173-190.
11. Krzywicki HJ, Consolazio CF, Matoush LL, Johnson HL, Barnhart RA. Body composition changes during exposure to altitude. *Fed Proc* 1969;28:1190-1194.
12. Janoski AH, Whitten BK, Shields JL, Hannon JP. Electrolyte patterns and regulation in man during acute exposure to high altitude. *Fed Proc* 1969;28:1185-1189.
13. Keys A. The physiology of life at high altitudes. The international high altitude expedition to Chile, 1935. *Sci Mon* 1936;43:289-312.
14. Evans WO. Measurement of subjective symptomatology of acute high altitude sickness. *Psychol Rep* 1966;19:815-820.

15. Hansen JE, Harris CW, Evans WO. Influence of elevation of origin, rate of ascent and a physical conditioning program on symptoms of acute mountain sickness. *Mil Med* 1967;132:585-592.
16. Carson RP, Evans WO, Shields JL, Hannon JP. Symptomatology, pathophysiology, and treatment of acute mountain sickness. *Fed Proc* 1969;28:1085-1091.
17. Gill MB, Pugh LGCE. Basal metabolism and respiration in men living at 5800 m (19,000 ft). *J Appl Physiol* 1964;19:949-954.
18. Consolazio CF, Matoush LO, Johnson HL, Krzywicki HJ, Daws TA, Isaac GJ. Effects of high-carbohydrate diets on performance and clinical symptomatology after rapid ascent to high altitude. *Fed Proc* 1969;28:937-943.
19. Harris CW, Shields JL, Hannon JP. Acute altitude sickness in females. *Aerospace Med* 1966;37:1163-1167.
20. Consolazio CF, Nelson RA, Matoush LO, Hansen JE. Energy metabolism at high altitude (3475 meters). U.S. Army Med Res and Nutr Lab Rep No. 289, 1966.
21. Evans WO, Robinson SM, Horstman DH, Jackson RE, Weiskopf RB. Amelioration of the symptoms of acute mountain sickness by staging and acetazolamide. *Aviat Space Environ Med* 1976;47:512-516.
22. Robinson SM, Evans WO, Sterner, RT. Effects of intermediate altitude staging upon acute mountain sickness (Abst). *Fed Proc* 1974;33:307.
23. Boyer SJ, Blume FD. Weight loss and changes in body composition at high altitude. *J Appl Physiol* 1984;57:1580-1585.
24. Blume FD. Metabolic and endocrine changes at altitude. In: West JB, Lahiri S, eds. *High altitude and man*. Bethesda, MD: Am. Phys. Soc., 1984:37-45.
25. Hannon JP, Shields JL, Harris CW. Anthropometric changes associated with high altitude acclimatization in females. *Am J Phys Anthropol* 1969;31:77-84.
26. Klain GJ, Hannon JP. Effects of high altitude on lipid components of human serum. *Proc Soc Exp Biol Med* 1968;129:646-649.
27. Consolazio CF, Johnson HL, Krzywicki HJ, Daws TA. Metabolic aspects of acute altitude exposure (4,300 meters) in adequately nourished humans. *Am J Clin Nutr* 1972;25:23-29.
28. Buskirk ER, Mendez J. Nutrition, environment and work performance with special reference to altitude. *Fed Proc* 1967;26:1760-1767.

29. Pugh LGCE. Animals in high altitudes: man above 5,000 meters--mountain exploration. In: Field J, ed in chief. Handbook of Physiology. Sec. 4. Adaptation to the environment. Washington, D. C.: American Physiological Society, 1964:861-868.
30. Hannon JP, Chinn KSK, Shields JL. Alterations in serum and extracellular electrolytes during high-altitude exposure. J Appl Physiol 1971;31:266-273.
31. Sridharan K, Malhotra MS, Upadhayay TN, Grover SK, Dua GL. Changes in gastro-intestinal function in humans at an altitude of 3,500 m. Eur J Appl Physiol 1982;50:145-154.
32. Rai RM, Malhotra MS, Dimri GP, Sampathkumar T. Utilization of different quantities of fat at high altitude. Am J Clin Nutr 1975;28:242-245.
33. Hannon JP, Sudman DM. Basal metabolic and cardiovascular function of women during altitude acclimatization. J Appl Physiol 1973;34:471-477.
34. Billings CE, Brashear RE, Bason R, Mathews DK. Medical observations during 20 days at 3,800 meters. Arch Environ Health 1969;18:987-995.
35. Klausen, K. Cardiac output in man in rest and work during and after acclimatization to 3800 m. J Appl Physiol 1966;21:609-616.
36. Johnson HL, Consolazio CF, Daws TA, Krzywicki HJ. Increased energy requirements of man after abrupt altitude exposure. Nutr Rep Intern 1971;4:77-82.
37. Consolazio CF, Matoush LO, Nelson RA. The caloric requirements of men performing moderate physical activities at 3475 meters. U.S. Army Med Res and Nut Lab Rep No. 295, 1966.
38. Johnson HL, Consolazio CF, Matoush LO, Krzywicki HJ. Nitrogen and mineral metabolism at altitude. Fed Proc 1969;28:1195-1198.
39. Grover RF. Basal oxygen uptake of man at high altitude. J Appl Physiol 1963;18:909-912.
40. Richalet JP, Rathat C, Keromes A, Herry JP, Larmignat P, Garnier M, Philardeau P. Plasma volume, body weight, and acute mountain sickness (letter) Lancet 1983;2:525.
41. Surks MI. Metabolism of human serum albumin in man during acute exposure to high altitude (14,100 feet). J Clin Invest 1966;45:1442-1451.
42. Consolazio CF, Johnson HL, Krzywicki HJ, Daws TA. Adaptation to high altitude (4300 m). J Physiologie 1971;63:232-235.
43. Bharadwaj H, Singh AP, Malhotra MS. Body composition of the high altitude natives of Ladakh. A comparison with sea level residents. Human Biol 1973;45:423-434.

44. Hannon JP, Chinn KSK. Effects of high altitude on body fluid volumes. Fed Proc 1966;26:719.
45. Munro HN. Energy intake and nitrogen metabolism. In: Kinney JM, ed. Assessment of energy metabolism in health and disease. Columbus, OH: Ross Lab, 1980:105-110.
46. Houston CF. Operation Everest: Overview Man at Extreme Altitude. J Appl Physiol (In Press).
47. Roza AM, Shizgal HM. The Harris Benedict equation reevaluated: resting energy requirements and the body cell mass. Am J Clin Nutr 1984;40:168-182.
48. Siri WE. Body composition from fluid spaces and density. In: Brozek J, Henschel A, eds. Techniques for measuring body composition. Washington, DC: National Academy of Science, 1961:223-244.
49. Grauer WO, Moss AA, Cann CE, Goldberg HI. Quantification of body fat distribution in the abdomen using computed tomography. Am J Clin Nutr 1984;39:631-637.
50. Recommended Dietary Allowances, Committee on Dietary Allowances, Food and Nutrition Board, Division of Biological Sciences, Assembly of Life Sciences, National Research Council. 9th ed. Washington, DC: National Academy Press, 1980.
51. Bondi KR. Present thoughts on exercise, weight, and performance aboard nuclear submarines. In: Knight DR, Bondi KR eds. Proceedings of the third tripartite conference on submarine medicine (France, The United Kingdom, The United States of America). Naval Submarine Medical Research Laboratory, Groton, CT, Special Report 83-1, 1983, pp177-179.
52. Hannon JP. Nutrition at high altitude. In: Horvath SM, Yousef MK, eds. Environmental Physiology: Aging, health and altitude. Elsevier North Holland Inc, 1980:309-327.
53. Hansen JE, Hartley LH, Hogan RP. Arterial oxygen increase by high-carbohydrate diet at altitude. J Appl Physiol 1972;33:441-445.
54. Dramise JG, Inouye CM, Christensen BM, Fults RD, Canham JE, Consolazio CF. Effects of a glucose meal on human pulmonary function at 1600-m and 4300-m altitudes. Aviat Space Environ Med 1975;46:365-368.
55. Van Liere EJ, Stickney JC. Hypoxia. Chicago: Univ of Chicago Press, 1963.
56. Fulco CS, Cymerman A, Pimental NA, Young AJ, Maher JT. Anthropometric changes at high altitude. Aviat Space Environ Med 1985;56:220-224.

**TABLES**

Table 1  
Physical Characteristics of Subjects \*

	Mean + SD	Range
Age, years	27.5 + 2.2	25.0 - 31.0
Height, cm	184.0 + 10.3	171.4 - 196.8
Weight, kg (7 Oct)	82.97 + 9.78	74.09 - 100.91

\*Excluding subjects 5 & 7

Table 2

# OPERATION EVEREST II PERIODS

---

PERIOD	DAY OF STUDY
1	2 - 8
2	9 - 15
3	16 - 22
4	23 - 29
5	30 - 36
6	37 - 39

Table 3

# Energy Intake (Kcal/Day)

Subject Number	Period (Study Days)						Mean
	1 (2-8)	2 (9-15)	3 (16-22)	4 (23-29)	5 (30-36)	6 (37-39)	
1	3570	2874	2658	2450	1643	1332	2536
3	2864	2662	2447	1934	1289	1137	2152
4	2650	2833	2712	2658	2634	1903	2635
5	3252	2867	2499	2516	1868		2695
6	3107	3108	3112	2136	2471	2383	2755
7	2672	2024					2373
8	3447	3533	3353	2615	2695	1773	3022
9	3178	2935	2861	2191	2742	2206	2736
Mean <sup>a</sup>	3136 <sup>b</sup>	2991 <sup>b</sup>	2857 <sup>b</sup>	2331	2246	1789 <sup>a</sup>	2639
Std Dev <sup>c</sup>	557	664	788	804	874	682	847.6

<sup>a</sup> Excluding Subjects 5 & 7

<sup>b</sup> Significantly different from other periods

<sup>c</sup> Significantly different from periods 4 & 5

p < 0.001



Table 4

# Percent Decrease in Energy Intake

Subject Number	% Decrease in Kcalories
1	62.7 <sup>a</sup>
3	60.3 <sup>a</sup>
4	28.2 <sup>a</sup>
5	42.6 <sup>b</sup>
6	23.3 <sup>a</sup>
7	24.3 <sup>a</sup>
8	48.6 <sup>a</sup>
9	30.6 <sup>a</sup>
<u>Mean<sup>c</sup></u>	<u>42.3<sup>a</sup></u>
SD	17.2

- <sup>a</sup> Excluding subjects 5 & 7
- <sup>b</sup> Difference between Period 1 & 6
- <sup>c</sup> Difference between Period 1 & 5
- <sup>d</sup> Difference between Period 1 & 2

Table 5

# Operation Everest II

Daily Energy Expenditure from Exercise  
(Study Day 4 to 39)

Subject Number	Calories		Exercise Time		Frequency of Exercise
	Mean	SD	Mean	SD	
1	315	325	32.0	31.2	0.92
3	206	218	23.6	27.1	0.72
4	1986	1045	211.5	96.5	2.53
5	466	323	47.6	25.7	1.55
6	1053	509	118.1	50.8	2.17
7	889	654	91.5	48.8	1.25
8	638	699	69.2	63.7	2.00
9	236	234	28.2	27.5	1.00
Mean*	715	807	76.5	81.2	2.11

\* excluding subjects 5 & 7

Table 6

# OPERATION EVEREST II AVERAGE ENERGY REQUIREMENTS

SUBJECT NUMBER	BEE	BEE + 20% ACTIVITY	EXTRA ACTIVITY	BEE + 20% ACTIVITY + EXTRA ACTIVITY
1	2232	2678	315	2993
3	1846	2215	206	2421
4	2004	2405	1986	4391
5	1600	1920	466	2386
6	1918	2302	1053	3355
7	1773	2128	889	3017
8	1781	2137	638	2775
9	1781	2137	235	2372

Table 7

## Energy Balance and Weight Change in 38 Days

Subject Number	Energy Requirement (Kcal)	Energy Intake (Kcal)	Change in Body Weight (kg)	
			Expected	Actual
1*	2993	2536	-2.3	-9.9
3*	2421	2152	-1.3	-6.1
4*	4391	2635	-8.7	-9.8
5**	2386	2695	+2.7	-1.7
6*	3355	2755	-3.0	-8.2
7***	3017	2373	-2.4	-2.9
8*	2775	3022	+1.2	-6.5
9*	2372	2736	+4.0	-4.2

\* October 7 to November 13, 1985

\*\* October 7 to November 7, 1985

\*\*\* October 7 to October 19, 1985

Table 8

# Operation Everest II

## Weight Loss in 38 Days

Subject Number	Initial Weight Oct 7, 1985 (kg)	Final Weight Nov 13, 1985 (kg)	Difference (kg)	Weight Loss (%)
1	100.91	91.02	9.89	9.8
3	76.70	70.57	6.14	8.0
4	84.43	74.66	9.77	11.6
5	59.43	57.73 <sup>a</sup>	1.70	2.4
6	84.55	76.36	8.18	9.7
7	72.73	69.77 <sup>b</sup>	2.96	4.1
8	77.16	70.68	6.48	8.4
9	74.09	69.89	4.20	5.7
Mean <sup>c</sup>	82.97	75.53	7.44 <sup>c</sup>	8.9
SD	9.78	8.02	2.24	2.0

<sup>a</sup> Nov 7, 1985

<sup>b</sup> Oct 19, 1985

<sup>c</sup> Significantly different,  $p < 0.01$   
 . Excluding Subjects 5 & 7

TABLE 9

## OPERATION EVEREST II

## FLUID BALANCE

Subject Number	Fluid Intake (ml/day)	Fluid Output (ml/day)	Balance
1	2464.4	1881.3	583.1
3	1906.8	1835.3	71.5
4	3854.1	3540.4	313.7
6	5448.5	5252.5	196.0
8	3928.3	3432.1	496.2
9	2199.2	2078.3	120.9
Mean	3278.5	3074.6	203.9
SD	1618.2	1527.3	1220.7

Table 10

# Operation Everest II

Subject	Extremes of Energy Intake Maximum (Kcal)	Minimum (Kcal)	Difference
1	4331	988	3343
3	3551	558	2993
4	5580	1611	3969
5	3916	1670	2246
6	4450	1269	3181
7	3600	1394	2206
8	4485	884	3601
9	4001	1483	2518
Mean*	4399.7	1132.2	3267.5
SD	+676.3	+395.8	+500.7

\* excluding subjects 5 & 7

Table 11

# Protein Intake (Grams/Day)

Subject Number	Period (Study Days)						Mean
	1 (2-8)	2 (9-15)	3 (16-22)	4 (23-29)	5 (30-36)	6 (37-39)	
1	115.0	108.6	90.0	66.6	54.0	47.5	83.7
3	111.2	114.0	91.7	80.1	47.4	47.4	85.6
4	127.5	150.8	127.5	126.4	106.1	66.6	122.9
5	116.6	102.7	75.5	73.9	65.4		89.6
6	93.4	126.8	121.3	90.4	92.7	71.6	102.3
7	101.6	87.4					95.0
8	108.5	135.8	106.9	98.5	103.2	66.1	107.1
9	102.7	121.9	102.9	83.2	95.3	84.6	99.9
Mean <sup>a</sup>	109.7 <sup>b</sup>	126.3 <sup>b</sup>	106.7 <sup>b</sup>	90.9	83.1	64.0 <sup>a</sup>	100.2
Std Dev	19.9	35.2	33.8	33.5	36.0	26.3	36.3

<sup>a</sup> Excluding subjects 5 & 7

<sup>b</sup> Significantly different from other periods

<sup>c</sup> Significantly different from periods 4 & 5

p < 0.001



Table 12

# Operation Everest II

## Percent Calories from Protein

Subject Number	Period (Study Days)						Mean
	1 (2-8)	2 (9-15)	3 (16-22)	4 (23-29)	5 (30-36)	6 (37-39)	
1	12.9	14.8	13.1	10.9	13.2	14.7	13.1
3	15.6	17.2	15.6	17.0	14.7	16.1	16.0
4	19.3	22.0	19.1	20.3	16.9	13.9	19.1
5	14.3	14.2	12.0	11.9	14.0		13.2
6	12.1	16.3	15.8	18.1	14.9	11.5	15.1
7	15.8	17.2					16.4
8	12.7	15.4	12.7	15.1	15.9	14.6	14.4
9	12.9	16.3	14.6	15.0	13.8	15.4	14.6
Mean*	14.3	17.0**	15.1	16.0	14.9	14.3	15.4
Std Dev	3.1	3.6	3.7	4.8	3.9	2.6	3.8

\* Excluding subjects 5 &amp; 7

\*\* Significantly different,  $p < 0.01$

TABLE 13

OPERATION EVEREST II  
MEAN NUTRIENT INTAKE FOR 38 DAYS

	RDA	MEAN INTAKE
Energy (kcal)		2639 <sub>±</sub> 848
Protein (gm)	56	100 <sub>±</sub> 36.3
(%)		15.4 <sub>±</sub> 3.9
Total Fat (gm)		92.3 <sub>±</sub> 44.7
(%)		31.1 <sub>±</sub> 9.5
Fat, saturated (gm)		36.2 <sub>±</sub> 20.7
Fat, monounsaturated (gm)		30.9 <sub>±</sub> 17.3
Fat, polyunsaturated (gm)		13.7 <sub>±</sub> 9.6
Cholesterol (mg)		380 <sub>±</sub> 272
Carbohydrate (gm)		359 <sub>±</sub> 127
(%)		54.5 <sub>±</sub> 9.6
Vitamin A (mcg RE)	1000	2877 <sub>±</sub> 1951
Thiamin (mg)	1.4	6.58 <sub>±</sub> 9.2
Riboflavin (mg)	1.6	7.20 <sub>±</sub> 9.29
Niacin (mg NE)	18	35.5 <sub>±</sub> 17.9
Vitamin B <sub>6</sub> (mg)	2.2	6.25 <sub>±</sub> 9.1
Vitamin B <sub>12</sub> (mcg)	3.0	10.2 <sub>±</sub> 10.1
Vitamin C (mg)	60	505 <sub>±</sub> 457
Folacin (mcg)	400	67109 <sub>±</sub> 149319
Sodium (mg)		3645 <sub>±</sub> 1387
Potassium (mg)		4001 <sub>±</sub> 1657
Iron (mg)	10	26.5 <sub>±</sub> 16.0
Calcium (mg)	800	1444 <sub>±</sub> 691
Phosphorus (mg)	800	1654 <sub>±</sub> 622
Magnesium (mg)	350	378 <sub>±</sub> 190
Zinc (mg)	15	20.3 <sub>±</sub> 14.9

Table 14

# Percent Calories from Carbohydrates

Subject Number	Period (Study Days)						Mean
	1 (2-8)	2 (9-15)	3 (16-22)	4 (23-29)	5 (30-36)	6 (37-39)	
1	61.8	50.8	54.5	55.7	51.6	54.3	54.9
3	62.6	54.4	54.5	52.8	56.4	54.5	56.1
4	67.5	64.6	55.4	57.2	59.0	50.6	59.9
5	63.3	58.0	59.1	60.4	61.1		60.3
6	68.0	54.3	55.4	52.3	54.4	51.2	57.0
7	59.7	51.2					55.8
8	54.0	47.9	47.4	43.5	37.3	55.0	46.8
9	58.5	49.9	56.2	53.6	45.0	53.4	52.5
Mean*	62.1**	54.5	53.2	52.9	50.6	53.2	54.5
SD	7.1	9.0	9.1	9.6	10.5	7.0	9.6

\* Excluding subjects 5 &amp; 7

\*\* Significantly different,  $p < 0.001$

Table 15

# Operation Everest II

Percent Calories from Fat

Subject Number	Period (Study Days)						Mean
	1 (2-8)	2 (9-15)	3 (16-22)	4 (23-29)	5 (30-36)	6 (37-39)	
1	24.4	31.3	36.7	32.0	34.9	30.4	31.7
3	23.9	27.8	31.4	32.7	29.3	30.3	29.1
4	15.9	17.9	29.0	23.1	26.4	36.7	23.6
5	23.4	28.3	31.5	28.5	25.8		27.7
6	23.1	28.8	31.8	30.6	30.7	39.4	30.0
7	25.2	30.9					27.9
8	34.1	36.3	42.1	42.5	46.9	30.7	39.6
9	29.2	32.1	31.3	31.2	40.8	30.3	32.7
Mean*	25.1**	29.2	33.7	32.0	34.8	33.0	31.1
Std Dev*	7.5	8.5	9.7	9.0	10.5	7.9	9.5

\* Excluding subjects 5 & 7

\*\* Significantly different,  $p < 0.001$

TABLE 16

SKINFOLD MEASUREMENTS (mm) Pre- and Post-exposure to 40 Days of Progressive Decompression to 240 Torr

Site	Time	#1	#3	#4	#5 <sup>a</sup>	#6	#7 <sup>b</sup>	#8	#9
Triceps	Pre	9.8	9.9	9.7	7.9	7.0	7.9	8.6	8.8
	Post	9.1	9.2	7.9	7.3	5.7	-	8.8	8.2
Subscapular	Pre	11.7	13.8	7.7	6.3	11.0	7.9	7.5	8.7
	Post	11.1	12.8	6.5	6.3	9.3	-	7.8	11.2
Chest	Pre	6.3	6.3	4.2	3.8	4.4	4.9	4.4	5.9
	Post	6.1	6.5	4.2	4.4	3.9	-	5.9	5.7
Midaxillary	Pre	9.3	12.8	5.1	4.7	6.1	5.0	8.2	10.1
	Post	7.5	9.5	3.8	4.6	4.6	-	7.4	10.2
Suprailiac	Pre	13.9	22.7	12.2	5.9	8.5	12.7	8.4	14.3
	Post	12.5	18.4	6.1	6.5	6.9	-	12.9	13.8
Abdominal	Pre	22.9	21.1	10.5	7.1	8.7	11.5	8.0	20.7
	Post	19.8	17.4	9.4	8.2	6.4	-	13.5	23.1
Thigh	Pre	12.5	18.0	15.5	7.3	9.6	8.0	12.4	12.4
	Post	12.3	16.1	11.1	9.0	7.3	-	14.1	12.7
TOTAL	Pre	86.4	104.6	64.9	43.0	55.3	57.9	57.5	80.9
	Post	78.4	89.9	49.0	46.3	44.1	-	70.4	84.9

<sup>a</sup>Withdrawn on 33rd day.<sup>b</sup>Withdrawn on 16th day.

Table 17

# Operation Everest II

Average Fluid Balance (ml/day)

Subject Number	Period (Study Days)						Mean
	1 (2-8)	2 (9-15)	3 (16-22)	4 (23-29)	5 (30-36)	6 (37-39)	
1	1090.8	820.0	825.0	1035.0	47.9	-903.3	583.1
3	713.3	22.1	67.9	-21.0	37.5	-866.7	71.5
4	2125.0	-24.2	-217.1	292.0	-5.7	-613.3	313.7
5	1092.5	727.9	77.9	-724.0	-516.7		250.9
6	814.0	876.4	762.1	-734.0	-357.9	-900.0	196.0
7	419.0	-26.7					251.9
8	2471.7	1149.3	-266.4	-129.0	-202.1	-528.3	496.1
9	984.2	-26.4	402.1	359.2	323.6	-2868.3	120.8
Mean*	1382.3**	481.6	262.3	111.2	-27.7	-1113.3**	296.1
SD	1188.6	1049.7	1031.9	870.0	993.4	1641.0	1259.5

\* excluding subjects 5 & 7

\*\* Significantly different from other groups  
p < 0.05

TABLE 18

## TRENDS IN FOOD RATINGS AND NUMBER OF FOOD GROUPS CONSUMED

SUBJECT CONSUMED	MEAN RATING	NUMBER OF FOODS
1	+	-
3	=	-
4	+	-
6	+	-
8	+	-
9	+	-

+Measurement increased from the first half of the study to the second half.

=Measurement remained the same.

-Measurement decreased from the first half of the study.

TABLE 19

## TRENDS IN RATINGS OF DIFFERENT FOOD GROUPS

SUBJECT	VEGETABLE	SALAD	MEAT	BREAD	SWEETS	DESSERTS	POTATOES	CHICKEN
1		-	+	-	+		+	
3	--	+	+	+	+	-	+	
4	++	+	-	+	=		+	
6		+	+	-	+		+	+
8	++	+	-	+	-	+	+	+
9		+	+	+		+	+	

<sup>+</sup>Measurement increased from the first half of the study to the second half.

<sup>=</sup>Measurement remained the same.

<sup>-</sup>Measurement decreased from the first half of the study.



**FIGURES**

## FIGURE LEGENDS

- Figure 1. Energy (kcal) expended in ladder ergometer, treadmill walking, and biking exercises from information in the daily log book of exercise activity. Value included in energy expenditure calculations for Subject 1.
- Figure 2. Energy (kcal) expended in ladder ergometer, treadmill walking, and biking exercises from information in the daily log book of exercise activity. Value included in energy expenditure calculations for Subject 3.
- Figure 3. Energy (kcal) expended in ladder ergometer, treadmill walking, and biking exercises from information in the daily log book of exercise activity. Value included in energy expenditure calculations for Subject 4.
- Figure 4. Energy (kcal) expended in ladder ergometer, treadmill walking, and biking exercises from information in the daily log book of exercise activity. Value included in energy expenditure calculations for Subject 5.
- Figure 5. Energy (kcal) expended in ladder ergometer, treadmill walking, and biking exercises from information in the daily log book of exercise activity. Value included in energy expenditure calculations for Subject 6.
- Figure 6. Energy (kcal) expended in ladder ergometer, treadmill walking, and biking exercises from information in the daily log book of exercise activity. Value included in energy expenditure calculations for Subject 7.
- Figure 7. Energy (kcal) expended in ladder ergometer, treadmill walking, and biking exercises from information in the daily log book of exercise activity. Value included in energy expenditure calculations for Subject 8.
- Figure 8. Energy (kcal) expended in ladder ergometer, treadmill walking, and biking exercises from information in the daily log book of exercise activity. Value included in energy expenditure calculations for Subject 9.
- Figure 9. Time spent exercising each day for Subject 1.
- Figure 10. Time spent exercising each day for Subject 3.
- Figure 11. Time spent exercising each day for Subject 4.
- Figure 12. Time spent exercising each day for Subject 5.
- Figure 13. Time spent exercising each day for Subject 6.

- Figure 14. Time spent exercising each day for Subject 7.
- Figure 15. Time spent exercising each day for Subject 8.
- Figure 16. Time spent exercising each day for Subject 9.
- Figure 17. Subject 1: Comparison of daily energy intake, energy expenditure, and body weight changes. Arrows indicate exercise bouts during testing that may have interfered with eating ie: Muscle Biopsy (MUSCLE),  $VO_2\text{max}$  (MAX), Inert Gas (GAS), and Swan-Ganz Catheterization (SG).
- Figure 18. Subject 3: Comparison of daily energy intake, energy expenditure, and body weight changes. Arrows indicate exercise bouts during testing that may have interfered with eating ie: Muscle Biopsy (MUSCLE),  $VO_2\text{max}$  (MAX), Inert Gas (GAS), and Swan-Ganz Catheterization (SG).
- Figure 19. Subject 4: Comparison of daily energy intake, energy expenditure, and body weight changes. Arrows indicate exercise bouts during testing that may have interfered with eating ie: Muscle Biopsy (MUSCLE),  $VO_2\text{max}$  (MAX), Inert Gas (GAS), and Swan-Ganz Catheterization (SG).
- Figure 20. Subject 5: Comparison of daily energy intake, energy expenditure, and body weight changes. Arrows indicate exercise bouts during testing that may have interfered with eating ie: Muscle Biopsy (MUSCLE),  $VO_2\text{max}$  (MAX), Inert Gas (GAS), and Swan-Ganz Catheterization (SG).
- Figure 21. Subject 6: Comparison of daily energy intake, energy expenditure, and body weight changes. Arrows indicate exercise bouts during testing that may have interfered with eating ie: Muscle Biopsy (MUSCLE),  $VO_2\text{max}$  (MAX), Inert Gas (GAS), and Swan-Ganz Catheterization (SG).
- Figure 22. Subject 7: Comparison of daily energy intake, energy expenditure, and body weight changes. Arrows indicate exercise bouts during testing that may have interfered with eating ie: Muscle Biopsy (MUSCLE),  $VO_2\text{max}$  (MAX), Inert Gas (GAS), and Swan-Ganz Catheterization (SG).
- Figure 23. Subject 8: Comparison of daily energy intake, energy expenditure, and body weight changes. Arrows indicate exercise bouts during testing that may have interfered with eating ie: Muscle Biopsy (MUSCLE),  $VO_2\text{max}$  (MAX), Inert Gas (GAS), and Swan-Ganz Catheterization (SG).

- Figure 24. Subject 9: Comparison of daily energy intake, energy expenditure, and body weight changes. Arrows indicate exercise bouts during testing that may have interfered with eating ie: Muscle Biopsy (MUSCLE),  $VO_2$ max (MAX), Inert Gas (GAS), and Swan-Ganz Catheterization (SG).
- Figure 25. Composition of diet.
- Figure 26. Percent body fat changes pre- to post-exposure to 29,000 feet as determined by hydrostatic weighing.
- Figure 27. Body fat weight (kg) changes pre- to post-exposure to 29,000 feet calculated from body weight and percent body fat.
- Figure 28. Circumference measurements of the arm to determine girth changes pre- to post-exposure to high altitude.
- Figure 29. Circumference measurements of the thigh to determine girth changes pre- to post-exposure to high altitude.
- Figure 30. Computed tomographic (CT) scans of the arm to determine changes in fat due to exposure to high altitude.
- Figure 31. Computed tomographic (CT) scans of the thigh to determine changes in fat due to exposure to high altitude.
- Figure 32. Computed tomographic (CT) scans of the arm to determine changes in muscle mass due to exposure to high altitude.
- Figure 33. Computed tomographic (CT) scans of the arm to determine changes in muscle mass due to exposure to high altitude.

Figure 1  
Operation Everest II  
Energy Expenditure from Exercise  
Subject 1





Figure 2  
Operation Everest II  
Energy Expenditure from Exercise  
Subject 3



**Figure 3**  
**Operation Everest II**  
**Energy Expenditure from Exercise**  
**Subject 4**




Figure 4  
Operation Everest II  
Energy Expenditure from Exercise  
Subject 5







Figure 5  
Operation Everest II  
Energy Expenditure from Exercise  
Subject 6



**Figure 6**  
**Operation Everest II**  
Energy Expenditure from Exercise  
Subject 7




Figure 7  
Operation Everest II  
Energy Expenditure from Exercise  
Subject 8




Figure 8  
Operation Everest II  
Energy Expenditure from Exercise  
Subject 9




Figure 9  
Operation Everest II  
Duration of Exercise  
Subject 1




Figure 10  
Operation Everest II  
Duration of Exercise  
Subject 3




Figure 11  
Operation Everest II  
Duration of Exercise  
Subject 4




Figure 12  
Operation Everest II  
Duration of Exercise  
Subject 5







Figure 13  
Operation Everest II  
Duration of Exercise  
Subject 6



**Figure 14**  
**Operation Everest II**  
Duration of Exercise  
Subject 7




Figure 15  
Operation Everest II  
Duration of Exercise  
Subject 8




Figure 16  
Operation Everest II  
Duration of Exercise  
Subject 9




Figure 17  
ENERGY and WEIGHT MEASUREMENTS

SUBJECT 1




Figure 1B  
**ENERGY and WEIGHT MEASUREMENTS**  
 SUBJECT 3




Figure 19  
**ENERGY and WEIGHT MEASUREMENTS**  
 SUBJECT 4




Figure 20  
**ENERGY and WEIGHT MEASUREMENTS**  
 SUBJECT 5






Figure 21  
**ENERGY and WEIGHT MEASUREMENTS**  
 SUBJECT 6




Figure 22  
**ENERGY and WEIGHT MEASUREMENTS**

SUBJECT 7




Figure 23  
**ENERGY and WEIGHT MEASUREMENTS**  
 SUBJECT 8




Figure 24  
**ENERGY and WEIGHT MEASUREMENTS**

SUBJECT 9







Figure 25  
**OPERATION EVEREST II**  
**COMPOSITION of DIET**




**Figure 26**  
**BODY COMPOSITION CHANGES**  
**PERCENT BODY FAT BY HYDROSTATIC WEIGHING**



**Figure 27**  
**BODY COMPOSITION CHANGES**  
**BODY FAT WEIGHT AS DETERMINED BY HYDROSTATIC WEIGHING**




**Figure 28**  
**BODY COMPOSITION CHANGES**  
**CIRCUMFERENCE MEASUREMENT OF ARM**







**Figure 29**  
**BODY COMPOSITION CHANGES**  
**CIRCUMFERENCE MEASUREMENT OF THIGH**




**Figure 30**  
**OPERATION EVEREST II**  
**BODY COMPOSITION CHANGES**  
**CT SCAN OF ARM**




**Figure 31**  
**OPERATION EVEREST II**  
**BODY COMPOSITION CHANGES**  
**CT SCAN OF THIGH**



**Figure 32**  
**OPERATION EVEREST II**  
**BODY COMPOSITION CHANGES**  
**CT SCAN OF ARM**



**Figure 33**  
**OPERATION EVEREST II**  
**BODY COMPOSITION CHANGES**



APPENDICES

APPENDIX 1

breakfast (1)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Lunch (1)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Amount	Unit	Food Item
8	oz.	Orange Juice
2	each	Hard boiled eggs
1	cup	raisin bran
1	cup	lowfat milk
2	slices	toast
1	T.	Margarine
1	T.	Jelly
2	pkg	sugar
8	oz	coifee
8	oz	hot tea

Amount	Unit	Food Item
4	slices	whole wheat bread
3	oz.	Roast Beef Lean
2	leaves	lettuce
2	T.	mayonnaise
1	cup	Potato Sld
1	each	Fresh banana
1	pkg	sugar
8	oz	coffee
8	oz	hot tea



Dinner (1)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Snack (1)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Amount	Unit	Food Item
10 1/2	oz	Spaghetti w/meatsauce
1/2	cup	corn
1	cup	tossed salad
1	T.	Italian Dressing
2	slices	french bread
1	T.	margarine
1	serving	Chocolate Cake
1	pkg	sugar
8	oz	coffee
8	oz	hot tea

Amount	Unit	Food Item
1	cup	lowfat yogurt w/fruit

Breakfast (2)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Lunch (2)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Amount	Unit	Food Item
8	oz	Orange Juice
1	each	Banana
1	cup	oatmeal
1	cup	milk, lowfat
1	each	English Muffin
2	T.	margarine
1	T.	jam
2	pkg	sugar
8	oz	coffee
8	oz	hot tea

Amount	Unit	Food Item
4	slices	Whole Wheat Bread
1/2	cup	Tuna, water packed
2	T.	Mayonnaise
2	leaves	lettuce
1/2	each	tomato slices
1/2	cup	celery sticks
1	large	carrot stick
1	each	pear
1	cup	milk, lowfat
1	cup	apple juice

Dinner (2)

Subject Name \_\_\_\_\_  
Date \_\_\_\_\_

Snack (2)

Subject Name \_\_\_\_\_  
Date \_\_\_\_\_

Amount	Unit	Food Item	Amount	Unit	Food Item
6 3/4	oz	Southern Fried Chicken Patty	3	cups	popcorn, plain
		w/Vegetable Medley			
1/2	cup	mashed potatoes			
1/4	cup	gravy			
1	cup	carrot raisin sld			
1	each	hard roll			
1	T.	margarine			
1	cup	ice cream			
1	pkg	sugar			
8	oz	coffee			
8	oz	hot tea			

Breakfast (3)

Lunch (3)

Subject Name \_\_\_\_\_

Subject name \_\_\_\_\_

Date \_\_\_\_\_

Date \_\_\_\_\_

Amount	Unit	Food Item	Amount	Unit	Food Item
8	oz	apple juice	4	slices	whole wheat bread
1/2	each	grapefruit	3	oz	turkey breast
1	cup	shredded wheat	2	leaves	lettuce
1	cup	milk, lowfat	2	T.	mayonnaise
1	each	bagel	5	each	radishes
1	oz	cream cheese	1	large	carrot stick
2	pkg	sugar	1	cup	fruit salad
8	oz	coffee	2	each	cookies
8	oz	hot tea	1	cup	milk, lowfat

**Dinner (3)**

Subject Name \_\_\_\_\_  
Date \_\_\_\_\_

**Snack (3)**

Subject Name \_\_\_\_\_  
Date \_\_\_\_\_

Amount	Unit	Food Item	Amount	Unit	Food Item
9 1/4	oz	beef steak w/carrots	3/4	cup	raisin, unsalted nut, coconut mix
1	each	baked potato			
1/4	cup	sour cream			
1	cup	tossed salad			
1	T.	1000 island dressing			
2	each	dinner roll			
2	T.	margarine			
8	oz	grape juice			
1	serving	surprise pie			
1	pkg	sugar			
8	oz	coffee			
8	oz	tea			

**Breakfast (4)**

Subject name \_\_\_\_\_

Date \_\_\_\_\_

**Lunch (4)**

Subject name \_\_\_\_\_

Date \_\_\_\_\_

Amount	Unit	Food Item	Amount	Unit	Food Item
8	oz	orange juice	10	each	crackers, unsalted
1	each	fresh banana	1	owl	Chef's salad
1/2	cup	grapenuts	2	T.	italian dressing
1	cup	milk, lowfat	1	each	fresh apple
2	slices	toast	1	cup	milk, lowfat
1	T.	margarine			
1	T.	jelly			
2	pkg.	sugar			
8	oz	coffee			
8	oz	hot tea			

Dinner (4)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Snack (4)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Amount	Unit	Food Item
10	oz	Chicken cacciatore w/spaghetti
1/2	cup	green beans
1	cup	tossed salad
1	T	blue cheese dressing
2	slices	french bread
1	T.	margarine
1	serving	Carrot Cake
8	oz	cranberry juice cocktail
1	pkg	sugar
8	oz	coffee
8	oz	hot tea

Amount	Unit	Food Item
3	cups	popcorn, plain

Breakfast (5)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Lunch (5)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Amount	Unit	Food Item
8	oz	grape juice
1	each	fresh orange
1	cup	wheat flakes
1	cup	milk, lowfat
2	each	doughnut, cake type
2	pkg	sugar
8	oz	coffee
8	oz	hot tea

Amount	Unit	Food Item
4	slices	whole wheat bread
3	oz	roast beef, lean
2	leaves	lettuce
2	T.	Mayonnaise
1	cup	tossed salad
1	T.	french dressing
1	each	fresh fruit
1	cup	milk, lowfat



Dinner (5)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Snack (5)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Amount	Unit	Food Item	Amount	Unit	Food Item
15 1/4	oz	sliced turkey w/gravy & stuffing	2	T.	peanut butter
		w/broccoli and carrots	4	each	graham crackers
1/8	cup	cranberry sauce			
2	each	dinner roll			
1	T.	margarine			
1/2	cup	coleslaw			
1	serving	Surprise Cake			
2	pkg	sugar			
8	oz	coffee			
8	oz	hot tea			

**Breakfast (6)**

Subject name \_\_\_\_\_

Date \_\_\_\_\_

**Lunch (6)**

Subject name \_\_\_\_\_

Date \_\_\_\_\_

---

Amount	Unit	Food Item
8	oz	orange juice
1/2	each	grapefruit
1/2	cup	grape nuts
1	cup	milk, lowfat
1	each	toasted bagel
1	oz	cream cheese
2	pkg	sugar
8	oz	coffee
8	oz	hot tea

---

Amount	Unit	Food Item
1	each	hamburger bun
3	oz	hamburger
1	T.	mayonnaise
1/2	each	tomato slices
2	leaves	lettuce
1	slice	onion
2	T.	catsup
1/2	cup	cottage fries
1	cup	milk, lowfat
1	each	apple

Dinner (6)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Snack (6)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Amount	Unit	Food Item
8	oz	New England Clam Chowder
5	each	crackers, unsalted
11	oz	Lasagna
1/2	cup	peas
1	cup	tossed salad
1	T.	italian dressing
2	slices	Boston Brown Bread
1	T.	margarine
1	serving	Cheesecake
1	pkg	sugar
8	oz	coffee
8	oz	hot tea

Amount	Unit	Food Item
3/4	cup	raisin, unsalted nut,
		chocolate chip mix

Breakfast (7)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Lunch (7)

Subject name \_\_\_\_\_

Date \_\_\_\_\_

Amount	Unit	Food Item
8	oz	apple juice
1	each	fresh banana
1	cup	40% Bran flakes
1	cup	milk, lowfat
1	each	English Muffin
1	T	Margarine
1	T.	Jelly
2	pkg	sugar
8	oz	coffee
8	oz	hot tea

Amount	Unit	Food Item
4	slices	whole wheat bread
2	egg	egg salad
2	leaves	lettuce
1/2	each	tomato wedges
1/2	cup	cauliflower, raw
1	large	carrot stick
1	cup	fruit salad
2	each	oatmeal cookies
1	cup	milk, lowfat
8	oz	orange juice

Dinner (7)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Snack (7)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Amount	Unit	Food Item	Amount	Unit	Food Item
9	oz	Sweet & Sour chicken	2	T.	peanut butter
		w/veg. Oriental	10	each	soda crackers
1/2	cup	rice			
1	cup	tossed salad			
1	T.	1000 dressing			
1	serving	apple pie			
1	each	hard roll			
1	T.	margarine			
2	pkg	sugar			
8	oz	coffee			
8	oz	hot tea			

Dinner (8)

Name \_\_\_\_\_

Date \_\_\_\_\_

Amount	Unit	Food Item
8	oz	grilled pork chops
1/4	cup	gravy
1	ea	baked potato
1/4	cup	sour cream
1/2	ea	baked acorn squash w/brown sugar and butter
1	cup	tossed salad
1	T	french dressing
2	ea	dinner rolls
1	T.	Margarine
1	cup	ice cream
8	oz	coffee
8	oz	hot tea

**SPECIAL MENUS**

Special Breakfast

Special Lunch

Subject Name \_\_\_\_\_

Subject Name \_\_\_\_\_

Date 19 Oct 85

Date 19 Oct 85

Amount	Unit	Food Item	Amount	Unit	Food Item
1	cup	orange juice	1	each	hard roll
1	each	banana	4	oz	roast chicken
1	cup	Fruit and Fiber Cereal	2	T.	mayonnaise
1	each	muffin	2	leaves	lettuce
1	T.	margarine	1	stalk	celery stick stuffed w/peanut butter
1	cup	milk, lowfat	5	each	radishes
2	pkgs	sugar	2	each	cookies
1	cup	coffee	2	cups	apple juice
1	cup	hot tea	1	cup	milk, lowfat



Special Dinner

Subject Name \_\_\_\_\_

Date 19 Oct 85

Amount      Unit      Food Item

slice      Pizza (Pans will be sent into chamber.

List amounts after you eat the pizza).

cheese

hamburger

green peppers, onions, mushrooms, and jalapenos

tossed salad

1      cup

1      T.      1,000 island dressing

1      cup      sherbert

1      pkg      sugar

1      cup      coffee

1      cup      hot tea

breakfast (l)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Lunch (l)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Amount	Unit	Food Item	Amount	Unit	Food Item
3	oz.	Orange Juice	1	cup	Cream of Tomato Soup
2	each	Scrambled eggs	2	cups	Beef & Green Bean Casserole
1	cup	Hash Browned Potatoes			or
1	cup	raisin bran	2	cups	Green Bean Casserole
1	cup	lowfat milk			
2	slices	toast			
1	T.	Margarine	1	each	Fresh banana
1	T.	Jelly	1	pkg	sugar
2	pkg	sugar	8	oz	coffee
8	oz	coffee	8	oz	hot tea
8	oz	hot tea	1	pkg	salt
1	pkg	salt			

Dinner (7)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Snack (7)

Subject Name \_\_\_\_\_

Date \_\_\_\_\_

Amount	Unit	Food Item	Amount	Unit	Food Item
1/4		Chicken w/Spicy Mushroom Gravy	2	T.	peanut butter
1	cup	rice	10	each	soda crackers
		or			
1/4		Roasted Chicken			
1	cup	Sweet Pctato Casserole			
1	cup	tossed salad			
1	T.	1000 dressing			
1	serving	apple pie			
1	each	hard roll			
1	T.	margarine			
2	pkg	sugar			
8	oz	coffee			
8	oz	hot tea			

Breakfast (8)

Name \_\_\_\_\_

Date \_\_\_\_\_

Lunch (8)

Name \_\_\_\_\_

Date \_\_\_\_\_

Amount	Unit	Food Item	Amount	Unit	Food Item
8	oz	orange juice	2	cups	Lipton's Chicken Noodle Soup
2	slices	french toast	2	cups	Tuna Noodle Casserole
2	pats	butter			
1/4	cup	syrup			
1	oz	ham, grilled	1	stalk	Celery Stuffed w/Peanut Butter
1/2	cup	100% Granola Cereal			
1	cup	milk, lowfat	8	oz	cranberry juice
8	oz	coffee	1	ea	fresh orange
8	oz	tea	1	cup	milk, lowfat
2	t.	sugar	1	pkg	salt
1	pkg	salt			

Dinner (3)

Name \_\_\_\_\_

Date \_\_\_\_\_

Amount	Unit	Food Item
1		Broiled Steak w/ Onions & Mushrooms
1	ea	baked potato
1/4	cup	sour cream
1/2	ea	baked acorn squash w/brown sugar and butter
1	cup	tossed salad
1	T	french dressing
2	ea	dinner rolls
1	T.	Margarine
1	cup	ice cream
8	oz	coffee
8	oz	hot tea
1	pkg	salt

Dinner (7)

Name \_\_\_\_\_

Date \_\_\_\_\_

Amount	Unit	Food Item	Amount	Unit	Food Item
1/4		chicken, roasted season w/spices or w/ homemade Shake N Bake	2	salad leaves or	lettuce
8	ounce or	sliced turkey	1	cup	spinach
1	cup or	tuna	1	leaf or	chinese cabbage
2	ounces or	cheese + 2 oz different cheese	1	wedge	red cabbage
1/4	pound	tofu	1	each	tomato
1	ea or	baked potato	1	stalk	celery
1/2	ea	baked squash + 1 T. brown sugar + 2 pats butter	1/4	ea	red onion or 2 stalks green onion
1	or cup	brown rice	1/4	ea	cucumber or green pepper
1	ea or	green apple or 1/4 melon	1/2	ea	zucchini
1	pint or	Ben & Jerry's Ice Cream	1	large	carrot
2	cups or	mixed fruit	1	cup	parsley or basil or cilantio
1/8	serving or	pie	1/4	cup	bean sprouts or alfalfa sprouts
4	ea	cookies	1/4	cup	garbanzo beans or kidney beans
3	slices or	Boston Brown Bread			
1	ea or	hard roll			
2	ea	rolls			

APPENDIX 2

## APPENDIX 2 LEGENDS

Daily fluid intake and output raw data.

N=Subject Number,

C\_Date=Calendar Date,

B\_Intake=Fluid intake (oz) at breakfast,

L\_Intake=Fluid intake (oz) at lunch,

D\_Intake=Fluid intake (oz) at dinner,

ADLIB\_6218=Intake (oz) of fluids available ad libitum 0600-1800 hours,

ADLIB\_1826=Intake (oz) of fluids available ad libitum 1800-0600 hours,

Urine\_6218=Urinary output 0600-1800 hours,

Urine\_1826=Urinary output 1800-0600 hours,

T\_Intake=Total fluid intake (cc)/24 hr period (B\_Intake+L\_Intake+D\_Intake+  
ADLIB\_6218+ADLIB\_1826),

T\_Output=Total fluid output (cc) in a 24 hr period (Urine\_6218+Urine\_1826),

Balance=Fluid balance derived from T\_Intake minus T\_Output.



N	C_DATE	B_INTAKE_L	ADLIB_0610	ADLIB_1000	URINE_0610	URINE_1000	INTAKE_T	OUTPUT	BALANCE
1	07-OCT-05	24	0	16	2376	859	2640	3225	-585
2	07-OCT-05	24	49	0	1899	859	1695	1765	-25
3	07-OCT-05	24	32	53	1699	1000	4000	3350	1310
4	07-OCT-05	24	00	00	2400	1000	5000	4000	1000
5	07-OCT-05	24	00	00	2400	1100	5000	3775	1300
6	07-OCT-05	16	00	00	2025	1100	3000	3000	1010
7	07-OCT-05	22	49	20	1860	1000	3000	3000	1010
8	07-OCT-05	24	0	104	000	500	1440	1100	300
9	07-OCT-05	24	34	0	800	800	2040	1600	2040
10	07-OCT-05	22	32	0	800	800	2040	1600	2040
11	08-OCT-05	24	16	0	800	400	1000	1000	0
12	08-OCT-05	24	20	53	1000	1000	1000	1000	0
13	08-OCT-05	24	44	48	400	1000	5000	2000	2040
14	08-OCT-05	22	44	0	1175	1175	5100	3000	2100
15	08-OCT-05	0	00	32	2200	4000	4000	4000	0
16	08-OCT-05	0	12	0	1300	900	2400	2200	200
17	08-OCT-05	24	10	00	450	1000	7000	2200	5070
18	08-OCT-05	24	10	00	1750	400	4000	3100	1000
19	08-OCT-05	22	44	34	400	1000	2400	2000	400
20	08-OCT-05	16	0	32	1000	1000	5200	2000	2000
21	08-OCT-05	24	20	16	2000	5000	5000	4700	900
22	08-OCT-05	16	24	70	2100	700	2520	2000	-400
23	08-OCT-05	16	20	12	2300	2100	2440	2000	1040
24	08-OCT-05	24	0	0	000	300	2040	2000	900
25	08-OCT-05	22	24	32	000	1000	3720	3100	600
26	08-OCT-05	20	20	0	1000	800	2400	2000	400
27	08-OCT-05	16	0	0	800	800	2100	2000	0
28	08-OCT-05	24	00	30	2000	2000	4000	4000	0
29	08-OCT-05	12	44	40	1075	1075	4500	4500	0
30	08-OCT-05	32	72	00	1475	1475	4500	4500	0
31	08-OCT-05	16	0	100	000	000	2400	2400	0
32	08-OCT-05	24	20	16	2125	550	2040	2000	40
33	08-OCT-05	24	20	0	725	725	4320	4320	0
34	08-OCT-05	20	00	0	2400	2400	2400	2400	0
35	08-OCT-05	16	0	16	1100	1100	2400	2400	0
36	08-OCT-05	24	00	42	1100	2200	6450	3000	1150
37	08-OCT-05	16	40	40	1500	1500	4320	4320	0
38	08-OCT-05	32	32	30	2400	1700	4200	4100	0
39	08-OCT-05	0	04	0	1000	1000	2520	2500	20
40	08-OCT-05	24	24	20	1000	1000	6000	3000	3000
41	08-OCT-05	32	00	00	050	050	975	975	0
42	08-OCT-05	24	0	12	1100	500	2700	1700	1000
43	08-OCT-05	24	0	104	575	575	4000	3000	1000
44	08-OCT-05	20	00	32	1925	1925	3000	3000	0
45	08-OCT-05	24	00	24	975	1000	4440	3575	865
46	08-OCT-05	16	104	40	2000	1700	4000	4000	0
47	08-OCT-05	24	40	20	1175	1300	5200	2475	1000
48	08-OCT-05	32	72	32	075	1325	2000	2000	0
49	08-OCT-05	24	40	20	000	000	1000	1000	0
50	08-OCT-05	24	0	30	1000	1000	2000	2000	0
51	08-OCT-05	16	0	0	1000	1000	2000	2000	0
52	08-OCT-05	24	16	32	2000	900	3040	3000	40
53	08-OCT-05	24	16	32	1400	2000	4000	3450	1000
54	08-OCT-05	24	16	44	2200	1275	3000	3000	0
55	08-OCT-05	24	16	20	2200	2075	6700	4275	1400
56	08-OCT-05	0	0	0	1000	1000	1200	1200	0
57	08-OCT-05	16	0	32	2400	1000	4000	4000	0
58	08-OCT-05	24	16	0	1000	1000	2700	2000	700
59	08-OCT-05	24	16	0	1125	1125	2340	2000	340
60	08-OCT-05	24	16	0	1200	1200	1750	1750	0
61	08-OCT-05	16	16	0	1700	1000	3000	2000	1000
62	08-OCT-05	24	32	22	1300	1075	3000	3175	100
63	08-OCT-05	16	44	16	2000	2000	5340	4000	1300
64	08-OCT-05	0	0	0	575	575	1075	1075	0
65	08-OCT-05	24	16	32	600	600	4000	3000	1000
66	08-OCT-05	24	16	0	1400	1400	2000	2000	0
67	08-OCT-05	24	16	0	1000	1000	2000	2000	0
68	08-OCT-05	24	16	0	1000	1000	2000	2000	0
69	08-OCT-05	24	16	0	1000	1000	2000	2000	0
70	08-OCT-05	24	16	0	1000	1000	2000	2000	0
71	08-OCT-05	24	16	0	1000	1000	2000	2000	0
72	08-OCT-05	24	16	0	1000	1000	2000	2000	0
73	08-OCT-05	24	16	0	1000	1000	2000	2000	0
74	08-OCT-05	24	16	0	1000	1000	2000	2000	0
75	08-OCT-05	24	16	0	1000	1000	2000	2000	0
76	08-OCT-05	24	16	0	1000	1000	2000	2000	0
77	08-OCT-05	24	16	0	1000	1000	2000	2000	0
78	08-OCT-05	24	16	0	1000	1000	2000	2000	0
79	08-OCT-05	24	16	0	1000	1000	2000	2000	0
80	08-OCT-05	24	16	0	1000	1000	2000	2000	0
81	08-OCT-05	24	16	0	1000	1000	2000	2000	0
82	08-OCT-05	24	16	0	1000	1000	2000	2000	0
83	08-OCT-05	24	16	0	1000	1000	2000	2000	0
84	08-OCT-05	24	16	0	1000	1000	2000	2000	0
85	08-OCT-05	24	16	0	1000	1000	2000	2000	0
86	08-OCT-05	24	16	0	1000	1000	2000	2000	0
87	08-OCT-05	24	16	0	1000	1000	2000	2000	0
88	08-OCT-05	24	16	0	1000	1000	2000	2000	0
89	08-OCT-05	24	16	0	1000	1000	2000	2000	0
90	08-OCT-05	24	16	0	1000	1000	2000	2000	0



C DATE	B INTAKE	L INTAKE	ADLIB	ADLIB	ADLIB	URINE	URINE	URINE	T INTAKE	T OUTPUT	T BALANCE
22-OCT-85	24	12	32	24	768	1888	3248	2688	748		
23-OCT-85	16	16	28	84	1728	1288	3488	2828	588		
24-OCT-85	44	16	84	22	1888	4828	4828	4828	-88		
25-OCT-85	8	12	28	32	1888	2488	3488	3488	-1288		
26-OCT-85	8	8	148	84	3688	1888	6288	5488	888		
27-OCT-85	24	8	88	88	1188	1888	6248	3888	3188		
28-OCT-85	16	16	28	16	1488	888	2848	2848	-18		
29-OCT-85	16	8	68	16	1888	888	2788	2788	848		
30-OCT-85	8	8	16	8	1288	1288	2488	2488	-1448		
31-OCT-85	24	8	44	8	3188	1888	4488	4488	-2888		
01-OCT-86	16	8	16	8	1288	1288	2488	2488	-1448		
02-OCT-86	16	8	68	16	1888	888	2788	2788	848		
03-OCT-86	8	8	16	8	1288	1288	2488	2488	-1448		
04-OCT-86	24	8	44	8	3188	1888	4488	4488	-2888		
05-OCT-86	16	8	68	16	1888	888	2788	2788	848		
06-OCT-86	16	8	68	16	1888	888	2788	2788	848		
07-OCT-86	8	8	72	24	1988	1888	5848	5488	-388		
08-OCT-86	24	8	16	84	688	1888	2188	2188	18		
09-OCT-86	16	24	8	56	768	488	2848	1188	1488		
10-OCT-86	16	24	8	32	488	1588	2648	1988	218		
11-OCT-86	24	8	81	12	1888	1288	4888	3888	1888		
12-OCT-86	8	8	8	12	1888	1288	4888	3888	1888		
13-OCT-86	16	16	68	84	1488	1888	6188	5828	-888		
14-OCT-86	8	8	96	88	3888	2828	4828	4828	-888		
15-OCT-86	8	8	8	88	1888	2888	4488	4488	-1788		
16-OCT-86	24	8	8	8	888	828	1288	1288	-128		
17-OCT-86	12	8	8	48	1828	788	2788	2788	1888		
18-OCT-86	16	8	8	24	288	788	1888	1888	738		
19-OCT-86	16	32	8	22	3488	2888	5828	6188	-288		
20-OCT-86	24	12	8	32	1828	2828	4828	4828	378		
21-OCT-86	8	8	48	88	888	3488	3488	3488	1848		
22-OCT-86	16	16	64	32	1888	2188	3788	3788	148		
23-OCT-86	16	22	24	24	388	878	3888	3888	2188		
24-OCT-86	8	8	8	8	2888	1288	3288	3288	828		
25-OCT-86	16	8	32	32	1888	1888	3888	3888	-178		
26-OCT-86	32	8	64	32	2288	2828	5848	4828	488		
27-OCT-86	16	8	78	8	2888	2728	4888	4728	78		
28-OCT-86	8	8	64	124	2888	3178	6388	6178	1188		
29-OCT-86	16	8	32	32	2918	2828	2888	2888	-1888		
30-OCT-86	8	8	28	8	888	1278	1888	2878	-278		
31-OCT-86	16	8	8	48	288	888	1888	1888	1678		
01-OCT-87	8	8	8	32	288	888	1288	1288	488		
02-OCT-87	8	8	8	67	1988	1778	3728	3728	1848		
03-OCT-87	8	8	64	58	2488	2178	4888	4828	188		
04-OCT-87	8	8	88	188	4888	3888	6388	7888	-1248		
05-OCT-87	32	8	48	84	2188	2478	4888	4828	-18		
06-OCT-87	24	8	8	8	688	828	1448	1178	288		
07-OCT-87	16	8	8	42	888	1388	2328	2188	128		
08-OCT-87	8	28	32	38	1888	888	3888	2888	718		
09-OCT-87	24	16	48	48	2888	2878	4888	4828	-788		
10-OCT-87	8	8	64	44	2488	1178	3488	3578	-88		
11-OCT-87	8	8	48	48	1788	2278	3888	4828	-428		
12-OCT-87	16	8	88	8	1188	1888	2488	2488	-88		
13-OCT-87	8	8	8	44	388	1388	1888	1888	1478		
14-OCT-87	16	16	8	32	388	1288	1928	1888	328		
15-OCT-87	16	8	32	32	2178	2888	2888	4178	-1288		
16-OCT-87	32	24	68	68	3288	2878	6888	6178	-178		
17-OCT-87	8	8	64	144	2328	3288	6188	5828	888		
18-OCT-87	8	8	48	44	1888	1788	3888	3888	-288		
19-OCT-87	8	24	32	32	888	978	2828	2828	848		
20-OCT-87	16	8	8	28	1878	1878	3248	3248	2188		
21-OCT-87	16	34	8	24	888	888	1488	1488	4448		
22-OCT-87	8	48	72	32	72	2478	4148	4148	8		
23-OCT-87	8	8	16	8	16	1488	2488	2488	8		
24-OCT-87	8	8	8	16	8	1488	2488	2488	8		



N	C_DATE	B_INTAKE	L_INTAKE	ADLIB_0210	ADLIB_1020	URINE_0210	URINE_1020	T_INTAKE	T_OUTPUT	BALANCE
3	10-NOV-85	14	16	0	0	200	1250	900	1400	-600
3	10-NOV-85	16	32	04	04	3350	1525	5000	4975	1000
4	10-NOV-85	0	10	120	0	2750	2675	7000	5325	2355
0	10-NOV-85	0	16	0	0	275	550	1000	1225	455
0	10-NOV-85	24	24	24	0	1100	1050	2400	2750	-350
1	11-NOV-85	0	0	0	0	1000	1150	720	2150	-1430
3	11-NOV-85	11	0	0	0	975	250	570	1225	-655
4	11-NOV-85	0	40	0	0	1350	1725	2400	2075	-375
0	11-NOV-85	16	0	0	0	3100	2575	2640	5675	-3035
4	11-NOV-85	16	0	0	0	1175	1600	1920	2075	-755
0	11-NOV-85	16	0	100	0	1850	3025	4200	5475	-1275
1	12-NOV-85	24	0	0	0	400	500	1000	1200	500
3	12-NOV-85	16	0	0	0	1300	720	720	1775	-1055
4	12-NOV-85	16	0	0	0	1075	475	3000	1675	1925
0	12-NOV-85	16	0	144	120	2100	3375	8500	6475	2105
0	12-NOV-85	16	0	40	32	2200	250	2400	2450	-50
0	12-NOV-85	16	0	0	0	2200	1925	720	4125	-3405
1	12-NOV-85	0	0	0	0	2000	1700	1920	3700	-1700
3	13-NOV-85	0	0	0	0	1000	250	900	1850	-900
4	13-NOV-85	0	0	0	32	2050	2000	900	4950	-3000
0	13-NOV-85	0	0	0	104	4500	3750	6400	9250	-1770
0	13-NOV-85	0	0	32	32	1100	1600	1920	2700	-700
0	13-NOV-85	0	0	0	0	1000	2425	300	4225	-925
1	14-NOV-85	16	0	0	0	375	400	400	400	0
3	14-NOV-85	0	16	16	0	1500	3100	2100	3100	0
4	14-NOV-85	0	0	04	0	1800	1920	1920	1920	0
0	14-NOV-85	0	0	04	0	400	400	400	400	0
0	14-NOV-85	0	0	0	0	1075	720	720	720	0
0	05-OCT-85	16	0	16	20	450	350	1800	800	1000
0	05-NOV-85	16	0	32	45	750	3250	2030	4000	-970
0	05-NOV-85	16	20	32	32	1675	1200	3000	2975	25

DISTRIBUTION LIST

NO. OF COPIES

Defense Technical Information Center ATTN: DTIC-DDA Alexandria, VA 22304-6145	12
Commander U.S. Army Medical Research and Development Command SGRD-RMS SGRD-PLC Fort Detrick Fredrick, MD 21701-5012	1 1
Commandant Academy of Health Sciences, U.S. Army ATTN: AHS-CDM ATTN: HSHA-CDM ATTN: HSHA-CDS Fort Sam Houston, TX 78234	1 1 1
Dir of Biol & Med Sciences Division Office of Naval Research 800 N. Quincy Street Arlington, VA 22217	1
CO, Naval Medical R&D Command National Naval Medical Center Bethesda, MD 20014	1
HQ AFMSC/SGPA Brooks AFB, TX 78235	1
Under Secretary of Defense Research and Engineering ATTN: OUSDRE(RAT)E&LS Washington, DC 20310	1
Dean School of Medicine Uniformed Services University of Health Sciences 4301 Jones Bridge Road Bethesda, MD 20014	1
Commander U.S. Army War College Carlisle Barracks, PA 17013	1
Commander U.S. Army Soldier Support Center Ft. Benjamin Harrison, IN 46216	1

Assistant Secretary of Defense (Health Affairs) 1  
ATTN: ASD(HA) PA&QA  
Washington, DC 20310

Assistant Secretary of Defense (Aquisition & Logistics) 1  
ATTN: OASD(A&L)SD  
Washington, DC 20310

Commander 1  
U.S. Army Troop Support Command  
ATTN: AMSTR-E  
4300 Goodfellow Boulevard  
St. Louis, MO 63120-1798

Commander 1  
U.S. Army Test and Evaluation Command  
ATTN: AMSTE-EV-S  
Aberdeen Proving Ground, MD 21005-5055

Commander 1  
U.S. Army Operational Test Evaluation Agency  
ATTN: CSTE-ZX  
5600 Columbia Pike  
Falls Church, VA 22041

Commander U.S. Army Training and Doctrine Command 1  
ATTN: ATCD-S  
Fort Monroe, VA 23651

Commander 1  
U.S. Army TRADOC Combined Arms Test Activity  
ATTN: ATCT-PO  
Ft. Hood, TX 76544

Commander 1  
U.S. Army Materiel Command  
ATTN: AMCDE-S  
Alexandria, VA 22333

Commander 1  
U.S. Army Combined Arms Center  
ATTN: ATZL-TIE  
Fort Leavenworth, KS 66027-5130

HQDA 1  
OTSG  
ATTN: DASG-DBD  
Rm 617, Bldg 5  
Skyline Place  
5111 Leesburg Pike, Falls Church VA 22041-3258

HQDA 1  
ATTN: DASG-RDZ  
Washington, DC 20310-2300

HQDA 1  
DCSLOG  
ATTN: DALO-TST  
Washington, DC 20310-2300

Commandant  
U.S. Army Quartermaster School  
ATTN: ATSM-CDT 1  
ATTN: ATSM-SFS-FM 1  
Fort Lee, VA 23807

Commandant  
U.S. Army Troop Support Agency  
ATTN: DALO-TAF 1  
ATTN: DALO-TAF-F 1  
FT. Lee, VA 23801

Commander  
U.S. Army Natick Research, Development and  
Engineering Center  
ATTN: STRNC-W 1  
ATTN: STRNC-Y 1  
ATTN: STRNC-T 1  
ATTN: STRNC-E 1  
ATTN: STRNC-TEAA 1  
Natick, MA 01760-5000

HQ U.S. Marine Corps 1  
Code LFS-4  
Washington, DC 20380-0001