

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS
2a. SECURITY CLASSIFICATION AUTHORITY DTIC SELECTED		3. DISTRIBUTION / AVAILABILITY OF REPORT This document has been approved for public release and sale; its distribution is unlimited.
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE 05 1987		5. MONITORING ORGANIZATION REPORT NUMBER(S)
4. PERFORMING ORGANIZATION REPORT NUMBER(S) ONR-TR- 27		

6a. NAME OF PERFORMING ORGANIZATION Howard University	6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION Office of Naval Research Chemistry Division, Code 1113
6c. ADDRESS (City, State, and ZIP Code) Department of Chemistry Washington, DC 20059		7b. ADDRESS (City, State, and ZIP Code)  Arlington, VA 22217

8a. NAME OF FUNDING / SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-80-C-805
8c. ADDRESS (City, State, and ZIP Code)		10. SOURCE OF FUNDING NUMBERS
	PROGRAM ELEMENT NO	PROJECT NO.
		TASK NO.
		WORK UNIT ACCESSION NO

11. TITLE (include Security Classification)  
The Photochemistry of Cyano and Dicyanoacetylene at 193 nm (Unclassified)

12. PERSONAL AUTHOR(S)  
J.B. Halpern, L. Petway, R. Lu, W.M. Jackson, V.R. McCrary and W. Nottingham

13a. TYPE OF REPORT Technical	13b. TIME COVERED FROM 9/86 to 9/87	14. DATE OF REPORT (Year, Month, Day) 1987, July 28	15. PAGE COUNT 30
----------------------------------	--	--	----------------------

16. SUPPLEMENTARY NOTATION  
Prepared for submission to the Journal of Chemical Physics

17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Photodissociation, cyanoacetylenes, Photochemistry
FIELD	GROUP	SUB-GROUP	

19. ABSTRACT (Continue on reverse if necessary and identify by block number)  
This paper reports on a study of the photochemistry of cyano and dicyanoacetylene at 193 nm. In particular, those photolysis channel have been characterized which lead to the production of excited state fragments and CN in the ground state. In addition to the direct production of CN and C<sub>2</sub>H, there is a second channel leading to C<sub>3</sub>N and H atoms. The results also show that at high laser intensity the HC<sub>3</sub>N dissociates by two photon photolysis and that the C<sub>3</sub>N undergoes secondary photolysis.

20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> OTHER		21. ABSTRACT SECURITY CLASSIFICATION Unclassified
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. George E. Walrafen		22b. TELEPHONE (include Area Code) 202 636-6900
		22c. OFFICE SYMBOL

87 8 4 071

OFFICE OF NAVAL RESEARCH

Contract N00014-80-C-805

R&T Code 4131 023

TECHNICAL REPORT NO. 27

The Photochemistry of Cyano- and Dicyanoacetylene at 193 nm.

by

J.B. Halpern, L. Petway, R. Lu, W.M. Jackson, and V.R. McCrary  
and W. Nottingham

Prepared for submission to the Journal of Chemical Physics

Department of Chemistry  
Laser Chemistry Division  
Howard University  
Washington, DC 20059

July 28, 1987

Reproduction in whole, or in part, is permitted for any purpose  
of the United States Government.

This document has been approved for public release and sale; its  
distribution is unlimited

THE PHOTOCHEMISTRY OF CYANO- AND DICYANOACETYLENE AT 193 NM

By

J. B. Halpern<sup>\*</sup>, L. Petway<sup>#</sup>, R. Lu<sup>@</sup>, W. M. Jackson<sup>§</sup>,  
and V. R. McCrary<sup>&</sup>

Department of Chemistry  
Howard University  
Washington, DC 20059

and

W. Nottingham

Department of Chemistry  
The University of the District of Columbia  
Washington, DC

- \* To whom correspondence should be sent
- # Present address: Lawrence Livermore National Laboratories  
Livermore, California
- @ Present address: Dalian Institute of Chemical Physics  
Dalian, People's Republic of China
- § Present address: Department of Chemistry  
University of California  
Davis, California
- & Present address: AT&T Bell Laboratories  
Murray Hill, New Jersey

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



## ABSTRACT

This paper reports on a study of the photochemistry of cyano- and dicyanoacetylene at 193 nm. In particular, those photolysis channels have been characterized which lead to the production of excited state fragments and CN in the ground electronic state. In addition to direct production of CN and  $C_2H$  there is a second dissociation channel leading to  $C_3N$  and H atoms. The results also show that at high laser intensity the  $HC_3N$  dissociates by two photon photolysis and that the  $C_3N$  undergoes secondary photolysis.

## INTRODUCTION

In 1920 Moreau and Bongrand synthesized the first members of the cyanoacetylene family, cyano- [1] and dicyanoacetylene [2]. Since that time several additional cyano polyenes of the series  $H-(C\ C)_n-CN$  and  $NC-(C\ C)_n-CN$  have been found. Chemists, and particularly spectroscopists have studied the cyanoacetylenes because of their linear structure. Interest in these molecules increased in 1971, when Turner detected  $HC_2CN$  in the interstellar medium using the rotational spectrum [3] first measured in the laboratory by Westenberg and Wilson [4].  $HC_2CN$  has also been observed in Titan's atmosphere by the Voyager missions to Saturn [5] and has been shown to play an important role in the odd nitrogen chemistry of that enormous moon [6].

Our laboratory has been studying the photochemistry of XCN species and  $C_2N_2$  for some time [7]. The work reported on here was started because of the astrochemical relevance of cyanoacetylene photochemistry, and also to see if there were any dynamic effects in photolysis that could be related to the increase in the molecule's length as compared to  $C_2N_2$ .

This article reports on the photolysis of cyano- and dicyano-acetylene at 193 nm. Measurements were made of emission spectra from excited fragments and laser induced fluorescence spectra of ground state CN fragments. The photolysis was accomplished by an unfocussed ArF laser whose intensity was varied over more than two orders of magnitude.

Previous work on cyanoacetylene photochemistry has used VUV light. Okabe and Diebler have studied the photodissociative excitation and photoionization of  $\text{HC}_2\text{CN}$  below 150 nm [8]. Sabety-Dzvonik, et al. characterized the quantum state distribution of CN produced in the 160 nm photolysis of  $\text{HC}_2\text{CN}$  [9] and  $\text{C}_4\text{N}_2$  [10]. Recently, the quantum state distribution from photolysis of  $\text{C}_4\text{N}_2$  between 165 and 145 nm has been measured by us [11].

The UV spectrum of  $\text{HC}_2\text{CN}$  from 290 to 200 nm has been measured and assigned by Job and King [12,13]. Conners, et al. have measured the VUV spectrum below 170 nm, but not assigned it [14]. Miller and Hannon have measured the UV spectrum of  $\text{C}_4\text{N}_2$  between 290 and 220 nm, and partially assigned the vibrational structure [15]. Conners, et al, also measured the VUV spectrum of  $\text{C}_4\text{N}_2$  below 200 nm [14].

## EXPERIMENTAL

The experimental apparatus has been described in detail elsewhere [16]. As shown in Figure 1, an ArF excimer laser (Lambda Physik EMG101) was used to photolyze the gas samples in a static gas reaction cell, and in effusive and pulsed molecular beams. The CN ( $X^2\Sigma^+$ ) fragments formed in the photolysis were detected by laser induced fluorescence (LIF) excited in the violet system of CN by a nitrogen pumped dye laser. The nitrogen laser could be pulsed at an arbitrary and adjustable delay after the excimer laser.

In some of the experiments a pulsed molecular beam source with a 0.5 mm nozzle (Laser Technics LPV) was set perpendicular to both of the counterpropagating laser beams. The cyanoacetylenes were diluted in helium or argon. The pressure behind the valve was of the order of an atmosphere. After expansion into the vacuum chamber, the rotational temperature of the polyatomic molecules is relaxed towards absolute zero. Although the vibrational degrees of freedom are also cooled, the vibrational energy transfer process is not especially efficient. Experiments were also done with an unexpanded beam giving a low pressure effusive flow of the cyanoacetylenes.

The fluorescence excited in the interaction region was detected by a photomultiplier tube (EMI 9789QB) placed at right angles to both the molecular beam and the lasers. The fluorescence was averaged by a boxcar analyzer (PAR 160) and displayed on a strip chart recorder. Direct fluorescence from

multiphoton excited fragments was measured through a 0.25 m monochromator (Jobin-Yvon H20-UV) with a slit width of 1 mm and a dispersion of 4 nm/mm. The LIF signal was measured with an appropriate band pass filter between the photomultiplier and the cell.

The excimer laser power was monitored by a Scientech power meter. Corrections were made for variation in the dye and excimer laser intensities.

The  $\text{HC}_2\text{CN}$  sample, synthesized by the method of Moreau and Bongrand [1] as modified by Miller and Lemmon [17] was furnished by Dr. Hideo Okabe.  $\text{C}_4\text{N}_2$  was synthesized by the method of Moreau and Bongrand [2]. GC/MS analysis of both samples showed no impurities.



## RESULTS

## A. Photolysis of Cyanoacetylene

Photodissociation of  $\text{HC}_2\text{CN}$  in a static gas cell at a pressure of 20 mtorr with  $50 \text{ mJ/cm}^2$  of ArF laser light produced a fairly strong fluorescence. The temporal profile of the total fluorescence, measured with a boxcar analyzer with a resolution of 30 ns is shown in Figure 2. In Figure 2 there is a strong emission that is over within 200 ns, as well as a weak emission with a longer lifetime. The actual decay of the sharp peak may be faster but is broadened by a jitter of about 50 ns in the firing of the excimer laser, due to an old thyratron. In any case, the bimodal characteristic of the fluorescence decay is evident. The decay rate of the weak emission is about 3 s.

Figure 3 shows the dispersed fluorescence about 100ns after the excimer laser fired. This spectrum is dominated by the CN violet bands. Their intensity varies as the second power of the laser flux. There is also some emission from the  $\text{C}_2^*$  Swan bands and a diffuse emission below 350 nm. The fluorescence at times greater than a microsecond after the photolysis was much weaker. In addition to peaks in the CN violet bands, there is some diffuse emission between 300 and 400 nm.

The LIF measurements of the rotational and vibrational distribution of CN ( $X^2\Sigma^+$ ) fragment have been done at a variety of dissociating laser intensities. The delay time between the excimer and the dye laser was fixed at 500 ns, which maintained a

collision free condition while eliminating interference from the immediate fluorescence. At the highest laser intensity ( $50 \text{ mJ/cm}^2$ ), hot vibrational and rotational excitation spectra were observed. One can see in Figure 4b that CN radicals with vibrational levels as high as  $v'' = 4$  were excited at the highest ArF laser intensity. The population of levels other than  $v'' = 0$  disappeared when the excimer laser intensity was decreased to about  $5 \text{ mJ/cm}^2$ , and dramatic changes were seen in the rotational envelope of the excitation spectrum. The LIF signal of vibrationally and rotationally hot CN ( $X^2\Sigma^+$ ) varies as the second power of the excimer laser intensity. Figure 4a shows a CN fragment excitation spectrum taken at lower excimer laser energy. Under these conditions both vibrational and rotational distributions were cold. The 1-1 band disappeared into the noise and lines originating from vibrational levels greater than two could not be found. The intensity of the low rotational lines increased linearly with the laser flux. The spectrum of Figure 4a can be converted into the quantum state distribution shown in Figure 5.

An attempt was made to dissociate  $\text{HC}_2\text{CN}$  in a seeded molecular beam to eliminate the contributions of hot bands to the dissociation. A mixture of 40 torr of  $\text{HC}_2\text{CN}$  with 720 torr He was used as the gas behind the pulsed beam valve. The ArF laser beam crossed the molecular beam 15 mm downstream from the exit of the pulsed nozzle. No quantitatively meaningful LIF spectrum of the CN ( $X^2\Sigma^+$ ) radical could be obtained from the pulsed beam studies at low laser intensity. A CN radical excitation spectrum could

only be observed in the pulsed molecular beam studies when the ArF laser power was very high. Though the spectra were noisy, the envelope is similar to those of Figure 4b.

#### B. Photolysis of Dicyanoacetylene

The direct fluorescence from the photolysis of  $C_4N_2$  at 193 nm was much weaker than that observed from  $HC_2CN$  photolysis under similar conditions. On the other hand, the LIF signals from CN ( $X^2\Sigma^+$ ) were much stronger than those seen in the  $HC_2CN$  measurements described above. As a result of this the fluorescence could be ignored when the ArF laser power was kept below  $20 \text{ mJ/cm}^2$  pulse. Better LIF spectra have been measured. The excitation spectrum is shown in Figure 6, along with the Boltzmann like fits to the observed quantum state distributions, shown in Figure 7. The latter are single exponential functions described by thermodynamic temperatures of 1100 and 730 K for CN fragments in the  $v'' = 0$  and 1 levels respectively.

We have also attempted to photolyze  $C_4N_2$  in a pulsed molecular beam. Unexpectedly, the direct emission relative to the LIF spectra became much stronger than it was in the static cell experiments. When  $C_4N_2$  was seeded in Ar, the emission was so strong, and had such a long decay time that one could not measure the nascent fragment LIF distribution immediately after photolysis. Figure 8 and 9 show the emission time profile and the spectrum respectively from the molecular beam with 30 torr of  $C_4N_2$  diluted in 730 torr of Ar.

When He was used as a carrier gas instead of Ar the fluorescence became weaker. With 30 torr of  $C_4N_2$  diluted in 250 torr of He, a usable LIF spectrum of CN ( $X^2+$ ) fragments was measured. Only the rotational distribution of the  $v'' = 0$  level could be analyzed from this spectrum, but within experimental error, the rotational temperature was the same as in the static photolysis.

## DISCUSSION

## A. Spectroscopy

The absorption spectrum of cyanoacetylene between 260 and 200 nm has been studied by Job and King [12,13], and the VUV absorption between 165 and 105 nm has been examined by Conners, et al [14]. Job and King found a structured, but very weak absorption system at 260 nm, and a stronger system, described as a structured continuum, starting at 230 nm and extending to lower wavelengths. It was noted that the 260 nm bands became diffuse below 240 nm. The 230 nm system of cyanoacetylene consists of a number of diffuse sets of bands with a spacing of  $2100 \text{ cm}^{-1}$  between the sets. The  $2100 \text{ cm}^{-1}$  interval corresponds to the excited state stretching frequency  $\nu_2$ . The bands have a halfwidth of  $5 \text{ cm}^{-1}$  at 230 nm, which increases to  $25 \text{ cm}^{-1}$  at 200 nm. The lack of a resolvable rotational absorption spectrum makes it impossible to assign the excited electronic state species. Argument by analogy is no help, as the equivalent state in  $\text{C}_2\text{N}_2$  is  $1^-$  [18], while in substituted acetylenes it is  $1\Delta_u$  [19].

Job and King's assignments can be extended to show that the state excited by the 1 nm bandwidth ArF laser at 193 nm is the  $2_0^3 4_0^1 5_0^1$  level at  $51,821 \text{ cm}^{-1}$ . The first state lies  $2120 \text{ cm}^{-1}$  above the assigned  $2_0^2 4_0^1 5_0^1$  level at  $49,701 \text{ cm}^{-1}$ . The excitation of one quanta in the  $4$  mode makes the transitions  $2_0^3 \rightarrow 2_0^2$  or  $2_0^3 \rightarrow 2_0^1$  allowed.

Because of the similarity of the low frequency bending modes in the ground and excited state, some hot bands would also be excited in the room temperature sample by the ArF laser. We have verified that there is a transition at  $51,821 \text{ cm}^{-1}$ , by measuring an absorption spectrum of cyanoacetylene in a spectrophotometer with  $0.1 \text{ nm}$  resolution. In addition to the vibronic structure there is a relatively strong continuum, rising smoothly from  $230 \text{ nm}$ . At  $193 \text{ nm}$  the absorption of cyanoacetylene is  $4 \times 10^{-19} \text{ cm}^2$  [11]. Most of this comes from the continuum.

Delon, et al., [20] have studied vibrational cooling in a nozzle beam of intermediate sized molecules, i.e. molecules having several vibrational degrees of freedom. For  $\text{HC}_2\text{CN}$  in a moderate expansion (a back pressure of 2 atmospheres, a nozzle diameter of  $25 \mu$  and a mixing ration of 10:1 with Argon as a buffer gas) there was little or no vibrational cooling. Starting from a source temperature of  $300 \text{ K}$  vibrational temperatures of  $100$  and  $235 \text{ K}$  were seen in the  $\nu_7$  and  $\nu_6$  modes after the expansion. Efficient rotational cooling was observed with a measured temperature of  $15 \text{ K}$ . Since our expansion conditions are similar, we can assume that the thermally excited bending modes of ground state  $\text{HC}_2\text{CN}$  and  $\text{C}_4\text{N}_2$  will at best be only partially relaxed by the expansion.

The UV absorption spectrum of  $\text{C}_4\text{N}_2$  has been studied by Miller and Hannon [15] and the VUV spectrum has been studied by Conners, Roebber and Weiss [14]. As in  $\text{HC}_2\text{CN}$ , Miller and Hannon found that there are two UV absorption systems in  $\text{C}_4\text{N}_2$ . A weak,

red degraded system has its origin at 280 nm, and there is a stronger system with an origin at 264 nm. They list four bands of this system at 264, 250, 237 and 226 nm. No other bands are listed, but there is a notation that the spectrophotometer that was used could not maintain its narrowest resolution (0.1 nm) for the last band. There is some possibility that their lamp was running out of intensity below 225 nm, and there may be one or more other bands. The second band, at 250 nm, is by far the strongest.



They were able to resolve sharp vibrational structures in the 280 nm system using a 2 m spectrograph with a 5.2 Å/mm dispersion, but in no case did they have enough resolution to measure rotational structure. Judging from the low resolution spectrum and assignments given by Miller and Hannon, the bands at 264 and 250 nm are equally sharp, while the 237 nm band is only a bit more diffuse, and the 226 nm band is obviously broadened. None of this is conclusive as to the dissociation limit, but it is consistent with the thermochemical limit given below.

Conners, et al., see an irregular absorption growing from 50,100 to 55,000  $\text{cm}^{-1}$  (198 to 180 nm) where a far stronger system begins. They offer no assignment, but using a CNDO/2 method SCF-MO calculation, they find that there are five possible states in the region which may be responsible for this absorption. Briefly they are  ${}^1\Delta_g^-$ ,  ${}^1\Delta_g$ ,  ${}^1\Delta_u^-$ ,  ${}^1\Delta_u$  (different from the pair of the same symmetries responsible for the 264 absorption) and  ${}^1\Delta_g$ . The 198 nm absorption is not broadened by a pressure of many

atmospheres of helium, which means that it is due to a valence transition. The absorption cross-section at 193 nm is about  $1 \times 10^{-19} \text{ cm}^2$ . From the shape of the spectrum this absorption is not a continuation of the 264 nm system, and indeed the peaks do not fall where one would expect the observed progression in the  $\nu_1$  mode of the 264 system to be. The electronic origin of this third electronic state must be near 198 nm. Thus, absorption of 193 nm ( $51,813 \text{ cm}^{-1}$ ) light would not excite any high frequency stretching modes ( $\nu_1 - \nu_5$ ) of the  $\text{C}_4\text{N}_2$ , but might involve excitation of a few low frequency stretches, and, of course, thermally excited hot bands of such modes. For most of the possible electronic species, at least one quanta of the  $\pi_g \nu_6$  or  $\nu_7$  modes would have to be excited.

## B. Thermochemistry

The thermochemistry of cyano and dicyanoacetylene is summarized in Table I. The one photon allowed processes for cyanoacetylene photolysis at 193 nm are




where the heat of reaction 2 is based on our measurement of the heat of formation of  $\text{C}_3\text{N}$  from the photodissociative excitation of  $\text{C}_4\text{N}_2$  [11].

Although the absorption of  $\text{C}_4\text{N}_2$  at 193 nm is four times smaller than that of  $\text{HC}_2\text{CN}$ , the CN LIF signal is much weaker. We have measured the branching ratio between processes (1) and (2).



It is at least 1:20 at 193 nm. We note that the absorption of a second photon by  $C_3N$  would lead to



The thresholds for processes 1, 2, and 3 would be 200, 244 and 193 nm respectively.


For 193 nm photolysis processes 1 and 3 can only produce CN in the ground state. With respect to 193 nm photolysis, process 1 is  $1,730 \text{ cm}^{-1}$  exothermic and process 3 is about thermoneutral, but with an uncertainty of about  $2,000 \text{ cm}^{-1}$  associated with the heat of formation of  $C_3N$ . Thus no vibrational excitation of the CN stretch can be produced directly from processes 1 and 3.

Process 2, is about  $10,800 \text{ cm}^{-1}$  exothermic with respect to 193 nm photolysis, but, again, there is a large associated uncertainty. There is evidence from this work and that of Sato that the  $C_3N$  is created with a significant amount of internal energy [21]. This point will be discussed further below. Some of this energy can remain in the CN fragment following photolysis of  $C_3N$  leading to the excitation spectrum observed at high laser intensities by a two step sequential process. The maximum energy available to the CN fragment would be  $10,656 \pm 1050 \text{ cm}^{-1}$ . It is energetically possible that some of the CN could be in the excited ( $A^2\Pi_i$ ) state.


For reasons which have been dealt with in detail in Xie, et al [21], it is extremely unlikely that a ground state CN radical

can absorb a second photon at 193 nm. Only a few levels of the CN ( $A^2\Pi_i$ ) state can do so, and the resulting fluorescence is well characterized, leading to emission from the F and E states of CN. Even at the highest intensities, no such emissions were seen in the photolysis of either  $HC_2CN$  or  $C_4N_2$ .

Production of  $CN(B^2\Sigma^+)$  can only proceed via the two photon process




or the overall three photon process of reaction 2 followed by



The amount of energy available to the  $CN(B^2\Sigma^+)$  fragment after two photon absorption at 193 nm is 332 kJ/mole or  $27,750 \text{ cm}^{-1}$  in process (4), , while that available after process (2) followed by (5) is 322 kJ/mol or  $26,915 \text{ cm}^{-1}$ .

In the photolysis of  $C_4N_2$  the first step will be



followed, possibly by reaction (3). At 193 nm, process 6 would leave  $8723 \text{ cm}^{-1}$  of excess energy in the modes of motion of the fragments. Reaction 6 has a threshold of  $232 \pm 1 \text{ nm}$ .

Excited CN could be formed by the two photon excitation of  $C_4N_2$  or reaction 5.

### C. Photochemistry

In the following discussion we will restrict ourselves to the photochemistry of the various species at 193 nm, except where specifically noted. Starting with cyanoacetylene, we first consider the LIF spectrum of CN fragments obtained at the lowest laser intensity. As can be seen in Figure 4a there is a vanishingly small population in  $v'' = 1$ , if any at all. Only the relatively poor signal to noise of the spectrum prevents us from asserting that there is no vibrational excitation of the product. This is in agreement with the proposition that all of the observed  $\text{CN}(X^2 \Sigma^+)$  population originates from process 1, where there is only about  $1700 \text{ cm}^{-1}$  of excess energy available to be distributed among the remaining modes of motion of the fragments.

The quantum state distribution of the  $\text{CN}(X^2 \Sigma^+, v'=0)$  fragments is seen in Figure 5. The last observable rotational state,  $N=24$ , corresponds to an energy of  $1134 \text{ cm}^{-1}$ . The line drawn through the points is the result of a least squares fit to the data of

$$(7) I/(N'+N''+1) = A_1 \exp[-B_0 N(N+1)/k_B T_1] + A_2 \exp[B_0 N(N+1)]$$

where

$$A_1/A_2 = 5.2$$

$$T_1 = 332 + 65 \text{ K}$$

$$T_2 = 2170 + 500 \text{ K}$$

$I$  is the LIF signal,  $B_{v''}$  is the rotational constant of the  $v''$  level and  $k_B$  is Boltzmann's constant.

Measurements have been made of the signal intensity of the  $R_{00}$  (4) line as a function of photolysis laser intensity. The signal varies linearly with the laser intensity, indicative of a single photon process. All of the above points to process (1) being the source of CN radicals in the low intensity photolysis.

Figure 4b shows the LIF spectrum obtained at high laser intensity. Contributions are measured from vibrational levels up to at least  $v'' = 3$ , with some indication of population in  $v'' = 4$ . The rotational distributions are hot, as can be seen from the absence of a true band gap between the  $P_{00}$  and  $R_{00}$  bands.

The process leading to the observed  $CN(X^2_{\Delta}^+)$  distribution at high laser intensities is most likely reaction 2 followed by 3. The total excess energy available to the  $CN(X^2_{\Delta}^+)$  after these two steps is about  $10656 \pm 1050 \text{ cm}^{-1}$ . This is consistent with the cutoff in the vibrational distribution at  $v'' = 3$  or 4 ( $\nu_e$  for CN is  $2064 \text{ cm}^{-1}$ ). The absence of a clear band gap between the  $P_{00}$  and  $R_{00}$  bands also shows that at least some rotational states are populated with energies up to about  $8000 \text{ cm}^{-1}$ . The second order dependence of the LIF signal for higher vibrational and rotational levels is consistent with this mechanism.

The only other channel which could lead to these vibrationally and rotationally excited CN radicals would be process 4 followed by radiative decay to the  $CN(X^2_{\Delta}^+)$  state.

This would be more than 3.5 eV exothermic, and should be able to form CN in much higher vibrational states.

We are then left to ask about the relative importance of the direct process 1 and the sequential process 2 and 3 for the production of CN. We know that process 2 dominates over process 1 at low intensities by a factor of about twenty. This comes from both a direct measurement of the branching ratio at 193 nm [11] and the observation that the CN LIF signal is much greater following the photolysis of  $C_4N_2$  than following the photolysis of  $HC_3N$ , even though the absorption cross-section of  $C_4N_2$  is four times lower. Thus the question reduces to what is the cross-section of process 3 at 193 nm. At all intensities, CN is produced in a single step in process 1, and it requires at least two additional photons to remove a CN radical from the states that are energetically accessible. Given an absorption cross-section of  $4 \times 10^{-19} \text{ cm}^2$ , even at the highest ArF laser intensities less than 1% of the  $HC_2CN$  molecules are photolyzed via reactions 1 and 2.

Simple kinetics shows that the ratio of CN produced by the one photon process to that produced by the two photon process will be

$$(8) \quad R = q_1 / (E_{\text{laser}} \sigma_1)$$

where  $q_i$  is the relative quantum yield of a reaction and  $\sigma_i$  the cross-section.

In Figure 4b, the envelope of the  $P_{00}$  band is similar to, but not exactly the same as that in Figure 4a, and the population found in  $v'' = 1$  is smaller, but of the same order of magnitude as that found in  $v'' = 0$ . At a laser flux of  $5 \times 10^{15}$  photons/cm<sup>2</sup>-pulse  $R$  is less than or equal to  $1/20$ . Increasing the flux by an order of magnitude increased the relative yield of radicals from the two photon mechanism by about the same factor. If we estimate the relative yields of the two pathways to produce CN to be of the order of unity at  $5 \times 10^{16}$  photons/cm<sup>2</sup>-pulse, and use our measure of the relative yields of processes 1 and 2, then we arrive at a crude estimate for the absorption of  $C_2CN$  at 193 nm of  $2 \times 10^{-18}$  cm<sup>2</sup>. The branching between reactions 1 and 2 accounts for the much smaller LIF signal when  $HC_2CN$  is photolyzed as against when  $C_4N_2$  is irradiated.

Excited fragments are also created in the photolysis. Consideration of the thermochemistry shows that absorption of at least two photons is needed to form excited radicals. From Figures 1 and 2, we can see that the emission in the first microsecond after the photolysis is dominated by the CN violet system. There is also some emission observed from the Swan bands of  $C_2$ . The intensity of the CN ( $B_{2+} \rightarrow X_{2+}$ ) fluorescence varies as the second power of the laser intensity. This favors the direct two photon absorption mechanism as the source of excited CN. Dissociation of  $C_3N$  to excited CN would require three photons overall. Moreover, one would expect to see the about the same pattern and strength of emission following the

photolysis of  $C_4N_2$  in the bulb experiments if this were the mechanism.

In Figure 2, the strong peak between 380 and 390 nm comes from the  $\Delta v = 0$  progression of the CN violet system. Weaker, associated  $\Delta v = +1$  and  $-1$  progressions of this system can be seen at 360 and 415 nm. The peaks between 400 to 410 nm and 465 to 475 nm are from excited  $C_2$ . Some other bands from  $C_2$  would be hidden below the CN violet bands.

The lifetime of the  $C_2$  Swan transitions is 170 ns [23], which would fall into the 200 ns region of the fast emission in Figure 1. Black carbon deposits on the window are also consistent with  $C_2$  production. We have not looked for ground state  $C_2$ , or  $C_2$  in the  $a^3\Pi$  state.

Sato studied the emission from cyanoacetylene [21]. In one experiment he varied the excitation frequency from 265 to 215 nm and recorded the dispersed emission. In the region from 265 to 250 nm, relatively strong direct fluorescence was seen centered at the irradiation wavelength. At 240 nm, where there is no absorption, there was no response. Below 230 nm, there was a sharp emission at the excitation frequency, which may have been scattered light and a broad emission between 300 and 400 nm. Quite a large range of vibronic states must be excited judging from the broad region of emission.

The photolysis of thermal  $C_4N_2$  at 193 yielded a fairly strong LIF signal from the  $CN(X^2\Sigma^+)$ . This can be seen in Figure

6. The signal intensity in the bandhead region was measured as a function of laser intensity and found to be linear. Thus, the primary process in the photolysis of  $C_4N_2$  is Reaction 6. The quantum state distribution shown in Figure 7. Thermochemically the CN fragment can have up to  $8700\text{ cm}^{-1}$  of excess energy. The LIF spectrum shows that CN ( $X^2_1$ ) is formed only in  $v'' = 0$  and  $v'' = 1$  levels. There is no evidence of  $v'' = 2$  or higher vibrational levels. Vibrational populations are probably controlled by Franck-Condon factors in the parent molecule. The highest amount of internal energy in any populated state is between  $3000$  and  $3500\text{ cm}^{-1}$ . About 17% of the CN is created in the  $v'' = 1$  level. This indicates that there is a dynamic restriction in the motion of the fragments as they separate.

The direct emission following the photolysis of  $C_4N_2$  at room temperature is much weaker than that from  $HC_2CN$ . This grows much stronger when  $C_4N_2$  is diluted in argon and expanded through a pulsed nozzle. Dilution in helium decreases the fluorescence. Most likely this emission is related to the formation of van der Waals complexes between the  $C_4N_2$  and the argon. There are two peaks in the spectrum. The one at 386 nm is from CN ( $B^2_1$ ). The band at 325 nm is very broad and very much like that from the photolysis of  $HC_2CN$  seen by Seki. We conclude that this emission is from excited  $C_3N$  formed in the multiphoton photolysis of Ar- $C_4N_2$  complexes.



## CONCLUSIONS

The single photon photolysis mechanism of cyano and dicyanoacetylene has been studied. Quantum state distributions of CN fragments have been measured. Multiphoton photolysis schemes of these two molecules have also been determined, and some characteristics of the fragments measured.

A broad emission from the photolysis of  $C_4N_2$ -Ar complexes has been tentatively identified as coming from excited  $C_3N$ .

It has been shown that secondary photolysis of the  $C_3N$  radical leads to the formation of CN radicals, from which we infer that the photolysis is also a source of  $C_2$  radicals. Some excited  $C_2$  has been seen from higher order processes. An estimate of the cross-section of the  $C_3N$  radical at 193 nm has been made.

Cyano-acetylene is commonly found in the interstellar medium, nitrogen-methane atmospheres, and probably in comets. On a simple calculation, it should be the second most common odd nitrogen species in these environments. It has been shown here that the primary photochemical process is a two step mechanism leading to the formation of  $C_2$  and CN radicals.

## ACKNOWLEDGEMENT

The authors wish to acknowledge the support of the Office of Naval Research for this work. J. B. Halpern was also supported by the National Science Foundation. We thank Hideo Okabe for many helpful discussions and the gift of the cyanoacetylene sample, and George E. Miller for sharing the results of his work before publication.

TABLE I

Heats of Formation and Heats of Reaction for Ground State  
Molecules and Radicals

$\Delta H_f$	kJ/mole		
$\text{HC}_2\text{CN}$	356	$\pm 4$	
$\text{C}_4\text{N}_2$	533	$\pm 4$	[23]
H	216.03	$\pm 0.004$	
CN	422	$\pm 4$	
$\text{C}_2\text{H}$	531	$\pm 1$	
$\text{C}_2$	829.3	$\pm 3$	
$\text{C}_3$	833	$\pm 1.5$	
$\text{C}_3\text{N}$	629	$\pm 17$	[10]
N	471	$\pm 4$	

Reaction	$\Delta H_R$	Threshold
$\text{HC}_2\text{CN} + h\nu \text{ ---> H} + \text{C}_3\text{N}$	490 kJ/mol	244 nm
$\text{---> CN(X)} + \text{C}_2\text{H}$	599 kJ/mol	200 nm
$\text{C}_4\text{N}_2 + h\nu \text{ ---> CN(X)} + \text{C}_3\text{N}$	515 kJ/mol	232 nm
$\text{C}_3\text{N} + h\nu \text{ ---> C}_2 + \text{CN(X)}$	621 kJ/mol	193 nm
$\text{---> C}_3 + \text{N}$	656 kJ/mol	182 nm

All heats of formation are from Reference 24 unless otherwise noted.

## REFERENCES

1. C. Moreau and J. C. Bongrand, *An. Chim. (Paris)* 14 (1920) 47.
2. C. Moreau and J. C. Bongrand, *An. Chim. (Paris)* 14 (1920) 5.
3. B. Turner, *Astrophys. J. Lett.*, 163 (1971) L35.
4. A. Westenberg and E. B. Wilson, Jr., *J. Am. Chem. Soc.*, 72 (1950).
5. See *Science* 212 (1981), *Nature* 292 (1981) and *J. Geophys. Res.* 87 (1982).
6. Y. L. Yung, M. Allen and J. Pinto, *Astrophys. J. Supp.*, 55 (1984) 465.
7. a) G. E. Miller, J. B. Halpern and W. M. Jackson, *J. Chem. Phys.*, 70, 2373 (1979).  
b) J. B. Halpern and W. M. Jackson, *J. Phys. Chem.*, 86, (1982) 973.  
c) J. B. Halpern and W. M. Jackson, *J. Phys. Chem.*, 86, (1982) 3528.
8. H. Okabe and V. H. Diebler, *J. Chem. Phys.*, 59 (1973) 2430.
9. J. B. Halpern and G. E. Miller, private communication.
10. M. J. Sabety-Dzvonik, R. J. Cody and S. Glicker, *J. Chem. Phys.*, (1986).
11. M. J. Sabety-Dzvonik, R. J. Cody and W. M. Jackson, *Chem. Phys. Lett.*, 44 (1976) 131.
12. V. A. Job and G. W. King, *J. Mol. Spectrosc.*, 19 (1966) 155.
13. V. A. Job and G. W. King, *J. Mol. Spectrosc.*, 19 (1966) 178.
14. R. E. Conners, J. L. Roebber and K. Weiss, *J. Chem. Phys.*, 60 (1974) 5011.
15. F. A. Miller and R. B. Hannon, Jr., *Spectrochem. Acta* 12 (1958) 321.
16. V. R. McCrary, R. Lu, D. Zakheim, J. A. Russell, J. B. Halpern, and W. M. Jackson, *J. Chem. Phys.*, 83, (1985) 3481.
17. F. A. Miller and D. H. Lemmon, *Spectrochem. Acta A*, 23 (1967) 1415.
18. G. B. Fish, G. J. Cartwright, A. P. Walsh and P. A. Warsop, *J. Mol. Spectrosc.*, 41, (1972) 20.

20. R. L. Delon and J. S. Muentzer, *Chem. Phys. Lett.*, 111 (1984) 147.
21. K. Sato, private communication
22. X. Xie, V. R. McCrary, J. B. Halpern, E. Pugh and W. M. Jackson, *J. Phys. Chem.*, 90 (1986) 2670.
23. A. Smith, *Astrophys. J.*, 156 (1969) 791.
24. G. T. Armstrong and S. Marantz, *J. Phys. Chem.*, 64 (1960) 1776.
25. H. Okabe, *Photochemistry of Small Molecules*, (Wiley, New York, 1978) pp 375

## FIGURE CAPTIONS

Figure 1 - A schematic diagram of the experimental apparatus. The apparatus consisted of a vacuum chamber with windows through which laser beams could enter and fluorescence emission of the fragments could be observed. For LIF experiments the ArF and dye laser beams entered the cell along the same axis but were counterpropagating. Fluorescence was observed through a filter by a photomultiplier. Emission was measured by replacing the filter with a small (0.25 m) monochromator which could be tuned over the region 200-800 nm. The photomultiplier signal was captured by a boxcar analyzer, and displayed on a strip chart recorder.

Figure 2 - Temporal profile of the total emission signal following the photolysis of  $\text{HC}_2\text{CN}$  by an unfocussed ArF laser if  $50 \text{ mJ/cm}^2$  fluence.

Figure 3 - Dispersed fluorescence from the high intensity photolysis of  $\text{HC}_2\text{CN}$ , from the initial part of the temporal profile shown in Figure 2.

Figure 4 - LIF spectra of the ground state CN fragments produced in the photolysis of  $\text{HC}_2\text{CN}$  at 193 nm. (a) shows the LIF spectrum when the laser fluence as  $50 \text{ mJ/cm}^2$  and (b) shows the spectrum at low laser energy.


Figure 5 - Quantum state distribution of CN from the photolysis of  $\text{HC}_2\text{CN}$  at low laser energy. The solid line is the result of a fit to a sum of exponentials.

Figure 6 - LIF spectra of CN fragments from the photolysis of  $\text{C}_4\text{N}_2$  at 193 nm under conditions of low laser intensity.

Figure 7 - Quantum state distribution of CN as measured from the spectrum shown in Figure 7.

Figure 8 - Emission time profile of CN from the 193 nm photolysis of expansion cooled  $\text{C}_4\text{N}_2$ .


Figure 9 - Dispersed emission from fragments following the photolysis of expansion cooled  $\text{C}_4\text{N}_2$  at 193 nm.




**SCHEMATIC OF LIF APPARATUS**




Time Dependence of direct emission from  $\text{HC}_2\text{CN}$




**Direct Fluorescence Observed from CN  
from Photolysis of Cyanoacetylene**



$\nu''$



(a)




384


386


388

Excitation wavelength, nm




$C_4N_2$






Time Dependence of direct fluorescence from  $C_4N_2$



Fluorescence Spectrum from  $C_2N_2$





ABSTRACTS DISTRIBUTION LIST, 051A

Dr. M. A. El-Sayed  
Department of Chemistry  
University of California  
Los Angeles, California 90024

Dr. E. R. Bernstein  
Department of Chemistry  
Colorado State University  
Fort Collins, Colorado 80521

Dr. J. R. MacDonald  
Chemistry Division  
Naval Research Laboratory  
Code 6110  
Washington, D.C. 20375-5000

Dr. G. B. Schuster  
Chemistry Department  
University of Illinois  
Urbana, Illinois 61801

Dr. J. B. Halpern  
Department of Chemistry  
Howard University  
Washington, D.C. 20059

Dr. M. S. Wrighton  
Department of Chemistry  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Dr. W. E. Moerner  
I.B.M. Corporation  
Almaden Research Center  
650 Harry Rd.  
San Jose, California 95120-6099

Dr. A. B. P. Lever  
Department of Chemistry  
York University  
Downsview, Ontario  
CANADA M3J1P3

Dr. George E. Walrafen  
Department of Chemistry  
Howard University  
Washington, D.C. 20059

Dr. Joe Brandelik  
AFWAL/AADO-1  
Wright Patterson AFB  
Fairborn, Ohio 45433

Dr. Carmen Ortiz  
Consejo Superior de  
Investigaciones Cientificas  
Serrano 121  
Madrid 6, SPAIN

Dr. Kent R. Wilson  
Chemistry Department  
University of California  
La Jolla, California 92093

Dr. G. A. Crosby  
Chemistry Department  
Washington State University  
Pullman, Washington 99164

Dr. Theodore Pavlopoulos  
NOSC  
Code 521  
San Diego, California 91232

Dr. John Cooper  
Code 6173  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Joseph H. Boyer  
Department of Chemistry  
University of New Orleans  
New Orleans, Louisiana 70148

Dr. Harry D. Gafney  
Department of Chemistry  
Queens College of CUNY  
Flushing, New York 11367-0904

D/1113/87/2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1