



2.2.2

MICROCCPY RESOLUTION TEST CHAR NATIONAL BUREAU OF STANDARDS - 1963 -



AD-A190 015





THE PHYSICAL OCEANOGRAPHY OF THE

NORTHERN BAFFIN BAY-NARES STRAIT REGION

by

Victor G. Addison, Jr.

December 1987

Thesis Advisor

R.H. Bourke

Approved for public release; distribution is unlimited.

Prepared for: Dierctor, Arctic Submarine Laboratory Naval Ocean Systems Center San Diego, CA 92152

88 3 10 022

NAVAL POSTGRADUATE SCHOOL Monterey, California

Rear Admiral R. C. Austin Superintendent

Kneale T. Marshall Acting Provost

This thesis was prepared in conjunction with research sponsored by the Arctic Submarine Laboratory, Naval Ocean Systems Center, San Diego, California under Work Order N66001-86-WR-00131. Reproduction of all or part of this report is authorized.

Released by:

achel

Gordon E. Schacher Dean of Science and Engineering

CLA	SSIF	TED			
TV	CLASS	F (A T (ON CF	THIS	PAC

	REPORT DOCUM	VENTATION	PAGE		
PORT SELURITY CLASSIFICATION		D RESTRICTIVE	MARKINGS		
CURITY CLASSIFICATION ALTHORITY		DISTRIBUTION	AVAILABILITY OF	REPORT	
		Approved	for public 1	release:	
ECLASSIFICATION: DOWNGRADING SCHEE	DULÉ	distribut	ion is unlin	mited.	
FORMING ORGANIZATION REPORT NUM	BER(S)	S MUNITORING	ORGANIZATION RE	EPORT NUMB	R(S)
PS 68-87-008					
				······	
AME OF PERFORMING ORGANIZATION	6D OFFICE SYMBOL (If applicable)	78 NAME OF MC	DNITORING ORGAN	NIZATION	
eval Postgraduate School	Code 68	Arctic Su	bmarine Labo	oratory	
DRESS (City, State, and ZIP Code)		70 ADDRESS (Cit	y, State, and ZIP ((ode)	
onterey, CA 93943-5000		Code 19, Naval Oce San Diego	Bldg. 371 an Systems (. CA 92152	Center	
AME OF FUNDING SPONSORING	BD OFFICE SYMBOL (It applicable)	9 PROCUREMEN	T INSTRUMENT IDE	ENTIFICATION	NUMBER
retic Submarine Jahoratory	Code 19	N6600186W	R00131		
DDRESS (City, State and ZIP Lode)		10 SOURCE OF F	UNDING NUMBER	、	
ldg. 371		PROGRAM	PROJECT	TASK	WOH- UNIT
aval Ocean Systems Center		LEWENT NO	NU	NU	ACCESSION NO
an Diego, CA 92152		l	I	<u>I</u>	
ice include security classification)					
				CTON	
HE PHYSICAL OCEANOGRAPHY O	F THE NORTHERN BA	FFIN BAI-NAR	ES SIKALL KE	LGIUN	
HE PHYSICAL OCEANOGRAPHY O	F THE NORTHERN BA	FFIN BAI-NAR		LGION	
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) ddision, Victor G., Jr.	F THE NORTHERN BA	FFIN BAI-NAR	LS SIKAII RE		
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) Idision, Victor G., Jr. YPE OF REPORT 13D TIME	F THE NORTHERN BA	14 DATE OF REPO	RT (Year, Month, I	Day) 15 PA	GE COUNT
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) ddision, Victor G., Jr. TYPE OF REPORT aster's Thesis FROM_	COVERED	14 DATE OF REPO 1987 Decem	RT (Year, Month, I	Day) 15 PA	ge count 110
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) ddision, Victor G., Jr. TYPE OF REPORT aster's Thesis IPPLEMENTARY NOTATION	F THE NORTHERN BA	14 DATE OF REPO 1987 Decem	ES SIRAII RE RT (Year, Month, I ber	Day) 15 PA	ge count 110
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) ddision, Victor G., Jr. VPE OF REPORT aster's Thesis VPPLEMENTARY NOTATION repared in conjunction with	F THE NORTHERN BA	14 DATE OF REPO 1987 Decem R.C. Paquet	ES SIRAII RE RT (Year, Month, I ber te	Day) 15 PA	ge count 110
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) ddision, Victor G., Jr. TYPE OF REPORT aster's Thesis IPPLEMENTARY NOTATION repared in conjunction with COSAT: CODES	F THE NORTHERN BA	14 DATE OF REPO 1987 Decem R.G. Paquet	RT (Year, Month, I ber te	Day) 15 PA	GE COUNT 110 Nock number)
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) ddision, Victor G., Jr. TYPE OF REPORT aster's Thesis JPPLEMENTARY NOTATION repared in conjunction with COSAT: CODES JELD GROUP SUB-GROUP	COVERED TO TO TO TO TO TO TO TO TO TO	14 DATE OF REPO 1987 Decem R.G. Paquet	RT (Year, Month, I ber te e 11 necessary and STR J	Day) 15 PA	GE COUNT 110 Nock number)
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) Idision, Victor G., Jr. YPE OF REPORT ISD TIME SET S Thesis IPPLEMENTARY NOTATION repared in conjunction with COSATI CODES IELD GROUP SUB-GROUP	COVERED TO TO TO A R.H. Bourke and 18 SUBJECT TERMS (C Baffin Bay Nar 25 Strai	14 DATE OF REPO 1987 Decem R.G. Paquet Continue on revers	ES SIRAII RE RT (Year, Month, I ber te e " necessary and SIR J North	Day) 15 PA	GE COUNT 110 Nock number) KLIN
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) Idision, Victor G., Jr. YPE OF REPORT ISTET'S Thesis IPPLEMENTARY NOTATION repared in conjunction with COSATI CODES ELD GROUP SUB-GROUP	F THE NORTHERN BA COVERED TO h R.H. Bourke and 18 SUBJECT TERMS (C Baffin Bay Nares Strain	14 DATE OF REPO 1987 Decem R.G. Paquet Continue on revers	ES SIRAIT RE RT (Year, Month, I ber te e " necessary and SIR J North	Day) 15 PA I identity by I JOHN FRAN h Water	GE COUNT 110 Work number)
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) Idision, Victor G., Jr. YPE OF REPORT ISTRACT S Thesis IPPLEMENTARY NOTATION COSATI CODES IELD GROUP SUB-GROUP ISTRACT (Continue on reverse if necessary A dense perverte of continue	COVERED TO TO TO TO TO TO TO TO TO TO	14 DATE OF REPO 1987 Decem R.G. Paquet Continue on revers t umber/ re-death (CT	RT (Year, Month, I ber te e " necessary and SIR J North	Day) 15 PA I mentiny by A JOHN FRANT h Water	GE COUNT 110 Nock number) (LIN
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) Idision, Victor G., Jr. YPE OF REPORT ISSET'S Thesis IPPLEMENTARY NOTATION repared in conjunction with COSATI CODES IELD GROUP SUB-GROUP ISTRACT (Continue on reverse if necessary A dense network of condu- in Bay northward to 82°00	COVERED TO TO TO TO TO TO TO TO TO TO	14 DATE OF REPO 1987 Decem R.G. Paquet Continue on revers t umberi re-depth (CT e to the Lin	ES SIRAIT RE RT (Year, Month, I ber te e " necessary and SIR J North D) measureme colo Sea in	Day) 15 PA Froentiny by A JOHN FRANT h Water	GE COUNT 110 Work number; KLIN conducted from
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) Idision, Victor G., Jr. YPE OF REPORT ISSET'S Thesis IPPLEMENTARY NOTATION repared in conjunction with COSATI CODES IELD GROUP SUB-GROUP ISTRACT (Continue on reverse if necessary A dense network of conduction in Bay northward to 82,09 lical oceanographic survey	COVERED TO TO TO TO TO TO TO TO TO TO	14 DATE OF REPO 1987 Decem R.G. Paquet Continue on revers t umber/ re-depth (CT e to the Lin n the northe	ES SIRAIT RE RT (Year, Month, A ber te e " necessary and SIR J North D) measureme ccln Sea, ir rn Baffin Ba	Day) 15 PA I mentity by 1 JOHN FRAN h Water ents was on a most con	GE COUNT 110 Work number) (LIN conducted from mprehensive Strait (NBB-NS
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) Idision, Victor G., Jr. VPE OF REPORT ISTRACT'S Thesis IPPLEMENTARY NOTATION repared in conjunction with COSAT: CODES IELD GROUP SUB-GROUP ISTRACT (Continue on reverse if necessar A dense network of conduction in Bay northward to 82,09 Sical oceanographic survey on. These data indicate	COVERED TO TO TO TO TO TO TO TO TO TO	TFIN BAI-NAK 14 DATE OF REPO 1987 Decem R.G. Paquet (ontinue on revers t t t t t t t t t t t t t	ES SIRAIT RE RT (Year, Month, I ber te e " necessary and SIR J North D) measureme ccln Sea, ir rn Baffin Ba diate Water	Day) 15 PA I identity by I JOHN FRAN h Water ents was o h most con av-Nares ((NSAIW)	GE COUNT 110 Work number) KLIN conducted from mprehensive Strait (NBB-NS and Arctic
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) Idision, Victor G., Jr. YPE OF REPORT ISSET'S Thesis IPPLEMENTARY NOTATION REDATED IN CODES IELD GROUP SUB-GROUP ISTRACT (Continue on reverse if necessar A dense network of conduct in Bay northward to 82,09 Hical oceanographic survey on. These data indicate I n Polar Water (ABPW) to be	COVERED TO TO TO TO TO TO TO TO TO TO	TFIN BAI-NAK 14 DATE OF REPO 1987 Decem R.G. Paquet Continue on revers t t t t t t t t t t t t t	RT (Year, Month, I ber te e " necessary and SIR J North D) measureme ccln Sea, ir rn Baffin Ba diate Water aters via th	Day) 15 PA Day) 15 PA I Gentury by D JOHN FRAN h Water ents was o n most con av-Nares S (NSAIW) a ne Canadia	GE COUNT 110 Work number) (LIN conducted from aprehensive Strait (NBB-NS and Arctic an Archipelago
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) Idision, Victor G., Jr. YPE OF REPORT ISSET'S Thesis IPPLEMENTARY NOTATION repared in conjunction with COSATI CODES IELD GROUP SUB-GROUP ISTRACT (Continue on reverse if necessary A dense network of conduction in Bay northward to 82, 09 Sical oceanographic survey on. These data indicate In Polar Water (ABPW) to be reas the West Greenland (We	COVERED TO TO TO TO TO TO TO TO TO TO	TFIN BAI-NAK 14 DATE OF REPO 1987 Decem R.G. Paquet Continue on revers t umber/ re-depth (CT e to the Lin n the northe ntic Interme ctic Basin w of the comp	RT (Year, Month, I ber te e " necessary and SIR J North D) measureme ccln Sea, ir rn Baffin Ba diate Water aters via th aratively di	Day) 15 PA Day) 15 PA I mentity by 1 JOHN FRANT h Water ents was of a most con sv-Nares (NSAIW) a he Canadia lute West	GE COUNT 110 Nock number) (LIN conducted from mprehensive Strait (NBB-NS and Arctic an Archipelago t Greenland
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) Idision, Victor G., Jr. YPE OF REPORT ISSER'S Thesis IPPLEMENTARY NOTATION repared in conjunction with COSATI CODES IELD GROUP SUB-GROUP ISTRACT (Continue on reverse if necessar A dense network of conduction ISTRACT (Continue on reverse if necessar A dense network of conduction ISTRACT (Continue on reverse if necessar A dense network of conduction ISTRACT (Continue on reverse if necessar A dense network of conduction ISTRACT (Continue on reverse if necessar A dense network of conduction ISTRACT (Continue on reverse if necessar A dense network of conduction In Bay northward to 82,09 Istical oceanographic survey Ion. These data indicate I In Polar Water (ABPW) to be reas the West Greenland (We rent Atlantic Intermediate	COVERED TO TO TO TO TO TO TO TO TO TO	rFIN BAI-NAK 14 DATE OF REPO 1987 Decem R.G. Paquet (ontinue on revers t umber) re-depth (CT e to the Lin n the northe ntic Interme ctic Basin W of the comp nd West Gree	ES SIRAIT RE RT (Year, Month, A ber te e " necessary and SIR J North D) measureme ccln Sea, ir rn Baffin Ba diate Water aters via th aratively di nland Curren	Day) 15 PA I menuny by 1 JOHN FRAN h Water ents was of av-Nares (NSAIW) ine Canadia llute Wes ht Polar 1	GE COUNT 110 Work number) KLIN conducted from mprehensive Strait (NBB-NS and Arctic an Archipelago t Greenland Water (WGCPW)
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) ddision, Victor G., Jr. VPE OF REPORT aster's Thesis UPPLEMENTARY NOTATION repared in conjunction with COSATI CODES IELD GROUP SUB-GROU	F THE NORTHERN BA COVERED TO TO TO TO TO TO TO TO TO TO	A DATE OF REPO 14 DATE OF REPO 1987 Decem R.G. Paquet (ontinue on revers t (ontinue on revers t (ontinue on revers t t on revers t t on the northe ntic Interme ctic Basin w of the comp nd West Gree found seaso	ES STRAIT RE RT (Year, Month, A ber te e " necessary and SIR J North D) measureme ccln Sea, in rn Baffin Ba diate Water aters via th aratively di nland Curren nally throug	Day) 15 PA I moentity by A JOHN FRAN h Water ents was on av-Nares S (NSAIW) a ne Canadia llute West the Polar I chout polar	GE COUNT 110 Work number) KLIN conducted from aprehensive Strait (NBB-NS and Arctic an Archipelago t Greenland Water (WGCPW) thern Baffin
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) Idision, Victor G., Jr. YPE OF REPORT ISSET'S Thesis IPPLEMENTARY NOTATION repared in conjunction with COSATI CODES ELD GROUP SUB-GROUP	COVERED TO TO TO TO TO TO TO TO TO TO	R.G. Paquet R.G. Paquet R.G. Paquet Continue on revers t t umber) re-depth (CT e to the Lin n the northe ntic Interme ctic Basin w of the comp nd West Gree found seaso WGC, which	ES STRAIT RE RT (Year, Month, I ber te e " necessary and SIR J North D) measureme ccln Sea, in rn Baffin Ba diate Water aters via th aratively di nland Curren nally throug attains a ma	Day) 15 PA Day) 15 PA 1 Gentury by 1 JOHN FRAN h Water ents was of h most con av-Nares 1 (NSAIW) a he Canadia lute West ht Polar H chout northan aximum bat	GE COUNT 110 Nock number) KLIN Conducted from aprehensive Strait (NBB-NS and Arctic an Archipelago t Greenland Nater (WGCPW) thern Baffin roclinic
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) Idision, Victor G., Jr. YPE OF REPORT ISSER'S Thesis IPPLEMENTARY NOTATION repared in conjunction with COSAT: CODES ELD GROUP SUB-GROUP	COVERED TO TO TO TO TO TO TO TO TO TO	R.G. Paquet R.G. Paquet R.G. Paquet Continue on revers t wmmoer/ re-depth (CT e to the Lin n the northe ntic Interme ctic Basin w of the comp nd West Gree found seaso WGC, which lle Bay (0.2	RT (Year, Month, A ber te e " necessary and SIR J North D) measureme ccln Sea, ir rn Baffin Ba diate Water aters via th aratively di nland Curren nally throug attains a ma Sv), south	Day) 15 PA Day) 15 PA Down 15 PA DOHN FRAN h Water ents was of the Canadia lute Wes thout north stimum bas of the Ca	GE COUNT 110 Nock number) KLIN Conducted from mprehensive Strait (NBB-NS and Arctic an Archipelago to Greenland Water (WGCPW) thern Baffin roclinic arey Islands
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) Idision, Victor G., Jr. YPE OF REPORT Ister's Thesis IPPLEMENTARY NOTATION repared in conjunction with COSATI CODES IELD GROUP SUBGROUP SUBSTRACT (Continue on reverse if necessar A dense network of conduct In Bay northward to 82,09 Sical oceanographic survey Ion. These data indicate I In Polar Water (ABPW) to be reas the West Greenland (We rent Atlantic Intermediate tions. Baffin Bay Surface Recurvature of component Sport of 0.7 Sv, occurs pro- Sv) and ultimately in Sm	COVERED TO TO TO TO TO TO TO TO TO TO	R.G. Paquet R.G. Paquet R.G. Paquet R.G. Paquet Continue on revers t wmber/ re-depth (CT e to the Lin n the northe ntic Interme ctic Basin w of the comp nd West Gree found seaso WGC, which lle Bay (0.2). The Baff	ES STRAIT RE RT (Year, Month, I ber te e " necessary and SIR J North D) measureme ccln Sea, it rn Baffin Ba diate Water aters via th aratively di nland Curren nally throug attains a ma Sv), south in Current o	Day) 15 PA Day) 15 PA 1 Gentiny by 1 JOHN FRAN h Water ents was of a most con av-Nares (NSAIW) a he Canadia lute West thout nort aximum ban of the Ca	GE COUNT 110 Work number) KLIN conducted from mprehensive Strait (NBB-NS and Arctic an Archipelago t Greenland Water (WGCPW) thern Baffin roclinic arey Islands a an A
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) Idision, Victor G., Jr. VPE OF REPORT IDD TIME STRACT'S Thesis INPLEMENTARY NOTATION REDARED IN CODES IELD GROUP SUB-GROUP	COVERED TO TO TO TO TO TO TO TO TO TO	R.G. Paquet R.G. Paquet R.G. Paquet (ontinue on revers t continue on revers t umber) re-depth (CT e to the Lin n the northe ntic Interme ctic Basin w of the comp nd West Gree found seaso WGC, which lle Bay (0.2). The Baff y net outflo	te r (Year, Month, A ber te e " necessary and SIR J North D) measureme ccln Sea, in rn Baffin Ba diate Water aters via th aratively di nland Curren nally throug attains a ma Sv), south in Current o w from Smith Circulati	Day) 15 PA Dow) 15 PA 1 JOHN FRAN h Water ents was of a most con av-Nares 3 (NSAIW) a he Canadia lute West thout norther chout norther at Polar 1 chout norther at polar 5 chout norther at polar	GE COUNT 110 Work number) KLIN conducted from aprehensive Strait (NBB-NS and Arctic an Archipelago t Greenland Water (WGCPW) thern Baffin roclinic arey Islands s as an and Lancaster th. Jones and
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) Idision, Victor G., Jr. VPE OF REPORT ISSTERIES INCE IPPLEMENTARY NOTATION Repared in conjunction with COSATI CODES IELD GROUP SUB-GROU	COVERED TO TO TO TO TO TO TO TO TO TO	R.G. Paquet R.G. Paquet R.G. Paquet Continue on revers t continue on revers t t continue on revers t t continue t continue t continue t continue t continue t continue t continue t comp nd West Gree f ound seaso WGC, which lle Bay (0.2). The Baff y net outflo respectively	ES STRAIT RE RT (Year, Month, I ber te e " necessary and SIR J North D) measureme ccln Sea, ir rn Baffin Ba diate Water aters via th aratively di nland Curren nally throug attains a ma Sv), south in Current o w from Smith . Circulati	Day) 15 PA Day) 15 PA Down 15 PA DOHN FRAN h Water ents was of h Water ents was of the Canadia (NSAIW) a me Canadia (NSAIW) a me Canadia (NSAIW) a the Can	GE COUNT 110 Work number Strait (NBB-NS and Arctic an Archipelago t Greenland Water (WGCPW) thern Baffin roclinic arey Islands a as an and Lancaster ith, Jones and
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) Idision, Victor G., Jr. VPE OF REPORT ID TIME Aster's Thesis IPPLEMENTARY NOTATION repared in conjunction with COSATI CODES IELD GROUP SUB-GROUP	COVERED TO TO TO TO TO TO TO TO TO TO	R.G. Paquet R.G. Paquet R.G. Paquet Continue on revers t continue on revers t t umber) re-depth (CT e to the Lin n the northe ntic Interme ctic Basin w of the comp nd West Gree found seaso WGC, which lle Bay (0.2). The Baff y net outflo respectively 21 ABSTRACT SE UNCLASSIN	ES STRAIT RE RT (Year, Month, A ber te e " necessary and SIR J North D) measureme ccln Sea, in rn Baffin Ba diate Water aters via th aratively di nland Curren nally throug attains a ma Sv), south in Current o w from Smith . Circulati CURITY CLASSIFICA	Day) 15 PA Day) 15 PA DOHN FRAN h Water ents was of h Water ents was of the Canadia (NSAIW) a me Canadia (NSAIW) a me Canadia (NSAIW) a he Canadia (NSAIW) a	GE COUNT 110 Work number) KLIN Conducted from aprehensive Strait (NBB-NS and Arctic an Archipelago t Greenland Water (WGCPW) thern Baffin roclinic arey Islands s as an and Lancaster ith, Jones and
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) ddision, Victor G., Jr. VPE OF REPORT aster's Thesis VPPLEMENTARY NOTATION repared in conjunction with COSATI CODES IELD GROUP SUB-GROUP BSTRACT (Continue on reverse if necessar A dense network of conduct tin Bay northward to 82,09 Sical oceanographic survey ion. These data indicate I In Polar Water (ABPW) to be reas the West Greenland (We rent Atlantic Intermediate tions. Baffin Bay Surface Recurvature of component asport of 0.7 SV, occurs put ISV) and ultimately in Smith redge jet in Smith Sound and ands at rates of 0.3 SV, 0.2 ISTRIBUTION: AVAILABILITY OF ABSTRACC UNCLASSIFIED/UNLIMITED SAME AND NAME OF RESPONSIBLE INDIVIDUAL H BOURKE	COVERED TO TO TO TO TO TO TO TO TO TO	R.G. Paquet R.G. Paquet R.G. Paquet (ontinue on revers t continue on revers t t umber) re-depth (CT e to the Lin n the northe ntic Interme ctic Basin w of the comp nd West Gree found seaso WGC, which lle Bay (0.2). The Baff y net outflo respectively 21 ABSTRACT SE UNCLASSIN 240 TELEPHONE (408) 644	ES STRAIT RE RT (Year, Month, A ber te e " necessary and SIR J North D) measureme ccln Sea, ir rn Baffin Ba diate Water aters via th aratively di nland Curren nally throug attains a ma Sv), south in Current o w from Smith . Circulati CURITY (LASSIFICA TIED Include Area Code b-3270	Day) 15 PA Day) 15 PA 1 Gentury by A JOHN FRAN h Water ents was of a most con av-Nares 3 (NSAIW) a ne Canadia lute West at Polar 1 chout nord at Polar 1 chout nord at Polar 5 chout nord at Polar 6 chout nord at Polar 6 at Polar 7 at Pola	GE COUNT 110 Nock number) KLIN conducted from aprehensive Strait (NBB-NS and Arctic an Archipelago t Greenland Water (WGCPW) thern Baffin roclinic arey Islands s as an and Lancaster ith, Jones and SYMBOL
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) ddision, Victor G., Jr. TYPE OF REPORT aster's Thesis JPPLEMENTARY NOTATION repared in conjunction with COSATI CODES IELD GROUP SUB-GROUP BSTRACT (Continue on reverse if necessar A dense network of conduing fin Bay northward to 82,09 Bical oceanographic survey ion. These data indicate I In Polar Water (ABPW) to but reas the West Greenland (Water Atlantic Intermediate tions. Baffin Bay Surface , Recurvature of component asport of 0.7 SV, occurs publics) and ultimately in Smith Sound and the survey of a sound and the survey in Smith Sound and asport of 0.3 SV, 0.1 ISTRIBUTION, AVAILABILITY OF ABSTRACCUNCLASSIFIED/UNLIMITED SAME AND SAME OF RESPONSIBLE INDIVIDUAL .H. Bourke	COVERED TO TO TO TO TO TO TO TO TO TO	R.G. Paquet R.G. Paquet R.G. Paquet R.G. Paquet Continue on revers t wmber/ re-depth (CT e to the Lin n the northe ntic Interme ctic Basin W of the comp nd West Gree found seaso WGC, which lle Bay (0.2). The Baff y net outflo respectively 21 ABSTRACT SE UNCLASSIN 220 TELEPHONE (408) 644 UNCLASSIN	ES STRAIT RE RT (Year, Month, I ber te e " necessary and SIR J North D) measureme ccln Sea, ir rn Baffin Ba diate Water aters via th aratively di nland Curren nally throug attains a ma Sv), south in Current o w from Smith . Circulati CURITY CLASSIFICA TIED Include Area Code b-3270	Day) 15 PA Day) 15 PA Down 15 PA 10 HN FRAN h Water ents was of the Canadia (NSAIW) a ne Canadia lute West the Canadia lute West the Canadia lute West the Canadia lute West the Canadia chout north aximum bas of the Ca originates h, Jones, lon in Small Alion	GE COUNT 110 Nock number) (LIN conducted from mprehensive Strait (NBB-NS and Arctic an Archipelago to Greenland Nater (WGCPW) thern Baffin roclinic arey Islands 5 as an and Lancaster ith, Jones and Symbol
HE PHYSICAL OCEANOGRAPHY O RSONAL AUTHOR(S) ddision, Victor G., Jr. VPE OF REPORT aster's Thesis IPPLEMENTARY NOTATION repared in conjunction with COSAT: CODES IELD GROUP SUB-GROUP BSTRACT (Continue on reverse if necessar A dense network of conduction fin Bay northward to 82,09 sical oceanographic survey ion. These data indicate In Polar Water (ABPW) to be reas the West Greenland (We rent Atlantic Intermediate tions. Baffin Bay Surface Recurvature of component isport of 0.7 SV, occurs pr SV) and ultimately in Smith redge jet in Smith Sound and ids at rates of 0.3 SV, 0. ISTRIBUTION: AVAILABILITY OF ABSTRACE UNCLASSIFIED/UNLIMITED SAME AU INCLASSIFIED/UNLIMITED SAME AU	COVERED TO TO TO TO TO TO TO TO TO TO	A DATE OF REPO 14 DATE OF REPO 1987 Decem R.G. Paquet Continue on revers t umber/ re-depth (CT e to the Lin n the northe ntic Interme ctic Basin w of the comp nd West Gree found seaso WGC, which lle Bay (0.2). The Baff y net outflo respectively 21 ABSTRACT SE UNCLASSIN 220 TELEPHONE ((408) 640 the emousted poorte	ES STRAIT RE RT (Year, Month, A ber te e " necessary and SIR J North D) measureme ccln Sea, it rn Baffin Ba diate Water aters via th aratively di nland Curren nally throug attains a ma Sv), south in Current o w from Smith . Circulati CURITY (LASSIFICA TIED INCLUGE AREA CODE D-3270 	Day) 15 PA Day) 15 PA Down 15 PA 1 Gentury by 1 JOHN FRAN h Water ents was of the Canadia lute West thout northold the Canadia lute West thout northold the Canadia chout northold the Canadia lute West thout northold the Canadia chout northold the Canadia the Can	GE COUNT 110 Nock number) (LIN conducted from mprehensive Strait (NBB-NS and Arctic an Archipelago t Greenland Nater (WGCPW) thern Baffin roclinic arey Islands s as an and Lancaster ith, Jones and SYMBOL N OF THIS PAGE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (Then Data Entered)

Block 18 (continued)

Geostrophic Estuarine Circulation Lancaster Sound Jones Sound Smith Sound Kane Basin Melville Bay

Block 19 (continued)

120000000 ... NYYYYYYYY

20101100

¥.

211**0**

Lançaster Sounds can be described in terms of the Geostrophic Estuarine Circulation Model (GEC). The North Water is caused by the combined influences of near-surface layer enthalpy and mechanical ice removal.



5 N 0102- LF- 014- 6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Date Entered)

Approved for public release; distribution is unlimited.

Ball Lin Lin Soll Li

The Physical Oceanography of the Northern Baffin Bay-Nares Strait Region

by

Victor G. Addison, Jr Lieutenant, United States Navy B.S., State University of New York, Stony Brook, 1979

Submitted in partial fulfillment of the requirements for the degree of

MASTER OF SCIENCE IN METEOROLOGY AND OCEANOGRAPHY

from the

NAVAL POSTGRADUATE SCHOOL December 1987

Author:

Victor (Addison, Jr.

Approved by:

🚬 R.H. Bourke, Thesis Advisor

61 he doutter R.G. Faquette,/Second Reader

kum ililine

C.A. Collins. Chairman Department of Oceanography

her 'n

Gordon E Schacher Dean of Science and Engineering

ABSTRACT

A dense network of conductivity-temperature-depth (CTD) measurements was conducted from Baffin Bay northward to $82^{\circ}_{1}09'_{1}N$ at the entrance to the Lincoln Sea, in the most comprehensive physical oceanographic survey ever performed in the northern Baffin Bay-Nares Strait (NBB-NS) region. These data indicate Nares Strait Atlantic Intermediate Water (NSAIW) and Arctic Basin Polar Water (ABPW) to be derived from Arctic Basin waters via the Canadian Archipelago, whereas the West Greenland Current (WGC) is the source of the comparatively dilute West Greenland Current Atlantic Intermediate Water (WGCAIW) and West Greenland Current Polar Water (WGCPW) fractions. Baffin Bay Surface Water (BBSW) is found seasonally throughout northern Baffin Bay. Recurvature of component branches of the WGC, which attains a maximum baroclinic transport of 0.7 Sv, occurs primarily in Melville Bay (0.2 Sv), south of the Carey Islands (0.1 Sv) and ultimately in Smith Sound (0.2 Sv). The Baffin Current originates as an ice-edge jet in Smith Sound and is augmented by net outflow from Smith, Jones, and Lancaster Sounds at rates of 0.3 Sv, 0.3 Sv and 1.1 Sv, respectively. Circulation in Smith, Jones and Lancaster Sounds can be described in terms of the Geostrophic Estuarine Circulation Model (GEC). The North Water is caused by the combined influences of near-surface layer enthalpy and mechanical ice removal.

4

nts. (112-2)

TABLE OF CONTENTS

Ι.	INT	RODUCTION
	Α.	PURPOSE
	B .	BACKGROUND13
		1. Bathymetry
		2. General Circulation16
		3. Water Masses
		4. The North Water
II.	MET	HODS AND MEASUREMENTS
	Α.	CRUISE SUMMARY
	Β.	INSTRUMENTATION
	С.	DATA REDUCTION AND ANALYSIS
III.	THE	RMOHALINE STRUCTURE
	Α.	INTRODUCTION
	B .	BAFFIN BAY SURFACE WATER
	С.	POLAR WATER
	D .	ATLANTIC INTERMEDIATE WATER
IV.	CIR	CULATION AND TRANSPORT
	Α.	INTRODUCTION
	B .	DYNAMIC TOPOGRAPHY
	С.	THE WEST GREENLAND CURRENT
	D .	CIRCULATION IN NARES STRAIT
	Ε.	THE BAFFIN CURRENT
	F .	CIRCULATION IN THE SOUNDS
		1. Introduction

3.0 10.

	2. Smith Sound
	3. Jones Sound
	4. Lancaster Sound
	5. Geostrophic Estuarine Circulation
	G. BAROCLINIC TRANSPORTS82
V .	THE NORTH WATER PROCESS
VI.	DISCUSSION
VII.	CONCLUSIONS
APPENDI	COMPUTATIONS ASSOCIATED WITH THE GEC MODEL
LIST OF	REFERENCES
INITIAL	DISTRIBUTION LIST

LIST OF TABLES

Ι.	NEAR-SURFACE AND INTERMEDIATE-DEPTH WATERS OF THE NORTHERN
	BAFFIN BAY-NARES STRAIT REGION
II.	GEOSTROPHIC ESTUARINE CIRCULATION MODEL CALCULATIONS

1. 2. 4. 4. A.

LIST OF FIGURES

1.1	A chart of the northern Baffin Bay-Nares Strait region14
1.2	The West Greenland Current and the Baffin Current (from Muench, 1971, p. 2)17
1.3	The approximate monthly mean extent of the North Water (from Dunbar, 1970, p. 279)21
2.1	The locations of CTD stations conducted during the September 1986 cruise of CCGS SIR JOHN FRANKLIN24
2.2	The locations of transects used in the analysis
3.1	A composite temperature profile illustrating the shallow thermocline characteristic of BBSW
3.2	The horizontal distribution of maximum temperatures in the BBSW layer
3.3	A T/S transect northeastward across Melville Bay33
3,4	A meridional T/S transect from Baffin Bay northward to Kane Basin
3.5	A composite salinity profile illustrating the dilution of WGCPW relative to ABPW
3.6	A composite density profile illustrating the dilution of WGCPW relative to ABPW
3.7	A composite T/S curve illustrating knee salinities characteristic of WGCFW and ABPW
3.8	A composite T/S curve illustrating the absence of a sharp knee in stations north of Smith Sound41
3.4	The horizontal distribution of ABPW and WGCPW in the NBB-NS region
3.10	A composite temperature profile illustrating the interleaving of ABPW and WGCPW in Smith Sound. Jones Sound and Melville Bay
3.11	A T/S transect northward across Melville Bay
3.12	A T/S transect across Kane Basin
3.13	A T/S transect across the southern entrance to Kane Basin 48

8

3.14	A T/S transect across the mouth of Jones Sound
3.15	A T/S transect across the mouth of Lancaster Sound
3.16	A composite density profile illustrating the magnitude of glucial meltwater-induced stair-stepped layering in Kane Basin
3.17	A T/S transect in Robeson Channel
3.18	The horizontal distribution of NSAIW and WGCAIW in the NBB-NS region
3.19	A meridional T/S transect from Kane Basin northward through Kennedy and Robeson Channels
3.20	A T/S transect Jones Sound eastward to Thule
4.1	Surface circulation in the NBB-NS region
4.2	The surface dynamic topography referenced to 200 decibars, in dynamic centimeters
4.3	A baroclinic velocity cross section through Melville Bay63
4.4	A baroclinic velocity cross section from Jones Sound eastward to Thule
4.5	A meridional baroclinic velocity cross section from Baffin Bay northward to Kane Basin
4.6	A baroclinic velocity cross section through a southern part of Kennedy Channel67
4.7	A baroclinic velocity cross section through the southern portion of the Kane Basin Gyre69
4.8	A baroclinic velocity cross section through Smith Sound72
4.9	A baroclinic velocity cross section through Jones Sound74
4.10	A baroclinic velocity cross section through Lancaster Sound. 76
4.11	A schematic cross section of coastal upper layer flow (from Leblond, 1980, p. 191)77
4.12	Surface dynamic height anomalies in Lancaster Sound (in dynamic cm) relative to 300 dbar level (from Fissel et al., 1982, p. 187)80
4.13	Dynamic topography of the surface relative to 500 dbars in Jones Sound for various years (from Muench, 1971, p. 92)83

- 5.1 Enthalpy of the near-surface (upper 75 m) layer in the NBB-NS region (referenced to -1.8°C)......90

ACKNOWLEDGEMENTS

Funding for the work described in this thesis was provided by the Arctic Submarine Laboratory, Naval Ocean Systems Center, San Diego, California under Work Order N-66001-86-WR00131.

I wish to thank Dr. R.H. Bourke for initially providing me with the opportunity to participate in this study, and, thereafter, for his invaluable assistance and encouragement. Dr. R.G. Paquette is also gratefully acknowledged for his advice and assistance and, in particular, for this continued support during periods of severe personal hardship.

The success of the cruise can be attributed to the enthusiasm and dedication of the Captain and crew of the CCGS SIR JOHN FRANKLIN. I also wish to thank A.M. Weigel, K.O. McCoy and Dr. P. Jones for their innumerable personal contributions to this effort. Miss J.E. Bennett is also gratefully acknowledged for her indispensable assistance in the preparation of this manuscript.

Finally. I wish to thank my lovely wife, Karen, for being there when I needed he:.

I. INTRODUCTION

A. PURPOSE

During September 1986, the CCGS SIR JOHN FRANKLIN conducted an oceanographic measurement program from Baffin : ay northward into Nares Strait, to the southern reaches of the Lincoln Sea. This expedition marked the first comprehensive use of continuous profiling conductivitytemperature-depth (CTD) measurements throughout the region, and the northernmost transit (82°09'N) of a surface ship through Nares Strait since the 1971 voyage of the CCGS LOUIS S. ST. LAURENT. Sponsorship of this endeavor was provided by the Arctic Submarine Laboratory, with diplomatic concurrence from the governments of Canada and Denmark.

The principal objective of the cruise was to examine the distribution of physical oceanographic variables from Baffin Bay northward, in order to determine the circulation and water mass characteristics of the region. The successful transit of the region by the SIR JOHN FRANKLIN provides a data set from which it is possible, essentially for the first time, to thoroughly define the baroclinic circulation and water mass structure of the northern Baffin Bay-Nares Strait region.

Specific objectives of the analysis will be to: (1) define the trajectory, transport and extent of the West Greenland and Baffin Currents; (2) describe the characteristics and extent of the water masses found throughout the region: (3) characterize the role of topographic steering and bathymetric boundaries on baroclinic

circulation and water mass penetration; (4) present a model which explains the observed circulation in Smith, Jones, and Lancaster Sounds; and (5) present a hypothesis for the formation and maintenance of the North Water, a large polynya located in Northern Baffin Bay.

B. BACKGROUND

The northern Baffin Bay-Nares Strait (hereinafter referred to as NBB-NS) region is historically significant for a number of seemingly unrelated reasons (Figure 1.1). Lancaster Sound, noted as the traditional Atlantic entrance to the Northwest Passage and as a net source of Arctic Ocean water inflow to the region, remains relatively ice free during winter. Indeed, northern Baffin Bay as a whole was thought to remain conspicuously unencumbered by ice during the winter due to the presence of the North Water. The Humboldt Glacier, the largest glacier in the northern hemisphere, is located on the eastern side of Nares Strait and is the source for many of the icebergs which are observed throughout the NBB-NS region. Once calved from the glacier, many of these icebergs travel southward past Cape York, in defiance of the northward flowing West Greenland Current. The most comprehensive oceanographic analysis of the northern Baffin Bav region is provided by Muench (1971). Muench's work serves as a summary of all significant oceanographic research conducted prior to 1971, and has been referenced in all subsequent analyses of the NBB-NS region.

Prior to 1972, all baroclinic oceanographic analyses in the NBB-NS region were derived from bottle sampling techniques. The CCGS LOUIS S. ST. LAURENT obtained discrete measurements of temperature and salinity in Nares Strait during August 1971, reaching a latitude of 82°56'N





D.

A chart of the northern Baffin Bay-Nares Strait region. Ice concentrations greater than 7/10 (during the FRANKLIN 86 cruise) are represented as shaded areas.

(Sadler, 1976). During 1972, scientific personnel aboard the LOUIS S. ST. LAURENT obtained the first winter oceanographic data from Davis Strait utilizing an in situ salinity/temperature/depth unit (STD) (Muench and Sadler, 1973). Transports through Nares Strait have been investigated by Sadler (1976), providing the most accurate current meter derived estimates of Lincoln Sea inflow to the region. In 1978 and 1979, the Eastern Arctic Marine Environment Studies (EAMES) program utilized current meters and CTD stations to investigate the southward flowing Baffin Current (Fissel et al., 1982). Research concerning the processes leading to the formation of the North Water has most recently been conducted by Steffen and Ohmura (1985).

1. <u>Bathymetry</u>

The bottom topography of the NBB-NS region is complex and, as such, has the potential to exert significant control over physical oceanographic processes. The influence of glaciers on all aspects of the morphology of this region is quite obvious.

Northern Baffin Bay is fed by Lancaster and Jones Sounds to the west, and Smith Sound to the north. Subaling to depths of less than 200 m occurs at some point in both Lancaster and Jones Sounds, effectively restricting deeper Arctic Ocean inflow. A sill which ranges in depth from 160 m to 200 m is present at the southern end of Kennedy Channel, providing similar constraints on Lincoln Sea inflow to the region. Similarly, shoaling to a depth of 250 m occurs at the Smith Sound entrance to Kane Basin.

The deep, relatively flat central portion of northern Baffin Bay is punctuated in its eastern particulate a hollow a 200 mm bank

located in Melville Bay. A deep (500 m to 900 m) channel passes offshore from Cape York, terminating northward in Smith Sound. To the west, the mouths of Jones and Lancaster Sounds reach depths of 700 m, but the bottom northeast of Devon Island shoals to less than 200 m approaching Smith Sound.

2. General Circulation

The baroclinic circulation pattern in the NBB-NS region is dominated by the northward flowing West Greenland Current (WGC) and the southward flowing Baffin Current (Figure 1.2). Augmenting the transport of the Baffin Current are net outflows from Jones and Lancaster Sounds, coupled with southward flow through Nares Strait. The net baroclinic transport through northern Baffin Bay has been computed to be approximately 2.0 Sv $(10^6 m^3 s^{-1})$ southward (Muench, 1971). Muench postulated that this transport is driven by a higher surface elevation in the Arctic Ocean relative to that in Baffin Bay, presumed to be a consequence of differing water structures in their respective upper 250 m layers.

Bottom topography, predominantly northerly winds, and meltwater admixture are thought to play variable roles in the maintenance of a cyclonic circulation pattern in northern Baffin Bay. The WGC, a significantly barotropic current, may be enhanced baroclinically by near-shore low salinity wedges of meltwater. Alternately, the Baffin Current exhibits the increased baroclinic intensity characteristically found in western boundary currents. The westward turning of the WGC off Cape York, and the southward turning of Lancaster Sound outflow are both influenced by topographic steering (Snewshold 1).







and the house

Circulation features in the NBB-NS region may be subject to seasonal variations. Lemon and Fissel (1982) noted a general winter weakening of near-surface baroclinic currents by a factor of two or more. The WGC, which transports northward only 10% of the baroclinic volume transported southward by the Baffin Current, is not always present during late summer (Muench, 1971).

The directional components of some features of the surface and baroclinic flow fields are temporally variable. Muench stated that reversals of baroclinic flow have been observed in Smith and Jones Sounds, while reversals of surface flow have occurred in Smith Sound. Lancaster Sound, however, consistently provides net inflow to Baffin Bay on the order of twice that of Smith or Jones Sounds (Muench, 1971). Anticyclonic eddies (sometimes referred to as counter currents) have been reported near Bylot Island (Fissel et al., 1982), while cyclonic eddies have been observed off Cape York (Muench, 1971).

Current speeds in the NBB-NS region, although variable, are consistently greatest in magnitude near eastern Lancaster Sound. Fissel et al. (1982), using current meters, found near-surface velocities of 0.75 m/sec in the portion of the Baffin Current located east of Lancaster Sound. Directly measured current speeds in Nares Strait are extremely variable in magnitude, with maximum near-surface values ranging from 0 1 to 0.60 m/sec (Sadler, 1976). Maximum baroclinic velocities in the WGC are on the order of 0 1 m/sec (Muench, 1971).

The nature and vertical extent of the flow regimes in the NBB-NS region are spatially complex. Althou hothe vertical extent of volume flow in the WGC probably reaches a letth of 200 moment.

baroclinic flow is limited to the upper 100 m (Muench, 1971). The Baffin Current nominally extends to a depth of 500 m, with flow reaching the bottom (700 m to 840 m deep) in the region of its intrusion into Lancaster Sound (Fissel et al., 1982). Significant baroclinic flow in this current is associated with the upper few hundred meters (Muench, 1971). The flow in Nares Strait has a substantial baroclinic component which reaches to depths of 300 m (Sadler, 1976).

3. <u>Water Masses</u>

Adopting the usual convention for Arctic regions, Muench (1971) divided the waters of northern Baffin Bay into three layers: a cold $(<0^{\circ}C)$ upper Arctic Water layer; a warmer $(>0^{\circ}C)$ intermediate-depth Atlantic Water layer; and a cold $(<0^{\circ}C)$ Deep Water Layer. The Arctic Water, confined to the upper 200 m to 300 m, has an upper limit in salinity of approximately 34.0. The Atlantic Water extends from the bottom of the Arctic Water layer to depths of 700 m in Lancaster Sound and 1300 m in northern Baffin Bay, exhibiting salinities of 34.2 to 34.5. Baffin Bay Deep Water extends from the bottom of the Atlantic Layer to the seabed, and is characteristically isohaline at 34.48.

The density of water masses in the NBB-NS region, as in other Arctic regions, is primarily determined by salinity. Surface and nearsurface layers, therefore, are usually characterized by strong haloclines and associated pycnoclines. The delineation of a surface (upper 75 m) layer of Arctic Water, which is modified by boundary layer processes, has been suggested by Fissel et al. (1982). Since diverse processes such as solar heating, meltwater admixture, and wind mixing

play roles in the formation of this laver, its temperature and salinity characteristics vary seasonally.

The thermohaline characteristics of the water masses of the region can generally be traced to their sources, while their horizontal extent is often determined by bathymetric effects. Arctic Water is water of Arctic Ocean origin which either entered Baffin Bay via Davis Strait and was modified by cooling and admixture of runoff within northern Baffin Bay, or was modified by cooling and freshening within the Arctic Ocean prior to entering Baffin Bay from the north. Mixing of inflowing Arctic Ocean Water and resident Baffin Bay Arctic Water of the same density occurs in southern Smith Sound, Jones Sound, and Lancaster Sound. Shallow sills present in Lancaster Sound, Jones Sound and Kennedy Channel prevent southward flow of Arctic Ocean Atlantic Water into Baffin Bay. Atlantic Water in northern Baffin Bay, therefore, originates from the Atlantic Ocean via Davis Strait (Muench, 1971).

4. The North Water

Extensive interest has been focused on a persistent polynya, termed the North Water, which is located in northern Baffin Bay (Figure 1.3). Dunbar (1970) concluded that the polynya is defined by a stable northern boundary at the Smith Sound entrance to Kane Basin, and a consistent western ice edge which forms along Ellesmere Island. The locations of the eastern and southern boundaries are seasonally dependent. Significant penetration of the North Water occurs westward into Lancaster Sound in June – The phenomenon is less well defined during the summer season after ice block up commences – fontrary to earlier reports, the North Water occurs to completely open during.



Figure 1.3

The approximate mean monthly extent of the North Water in: March (...), April (_____), Mav (---), June (____); (from Dunbar, 1970, p. 279).

winter, and is characterized by extensive areas of new ice formation (Dunbar, 1973). Steffen and Ohmura (1985) used thermal infrared measurements from an airplane in winter to determine that the North Water was mostly covered with new and young ice, and probably devoid of first year ice until March. The ice surface, they surmised, was approximately 20°C warmer than that of the surrounding fast ice.

The North Water essentially remains an unexplained phenomenon. Muench (1971) computed a heat budget for the region and concluded that insufficient heat is present in the Atlantic Water layer in northern Baffin Bay to prevent ice formation. He similarly concluded that there was not enough heat present in the surface Arctic Water layer either, and postulated mechanical ice removal by northerly winds and currents as the effects responsible for the formation of the North Water. Dunbar (1973) reinforced this hypothesis by noting that the age and thickness of ice increases from the Smith Sound ice arch southward. Steffen and Ohmura (1985), however, computed more accurate heat budgets for the region and concluded that oceanic heat of unknown origin is responsible for the formation and maintenance of this polynya.

22

II. METHODS AND MEASUREMENTS

A. CRUISE SUMMARY

Between 7 and 27 September 1986, the CCGS SIR JOHN FRANKLIN conducted the most extensive physical oceanographic survey ever performed in the NBB-NS region. The SIR JOHN FRANKLIN covered over 4000 km in the course of conducting 145 CTD stations (Figure 2.1; refer to Figure 2.2 for station number identification). Numerous transects were performed while transiting from Lancaster Sound northward to Robeson Channel, and subsequently southward to Melville Bay. Ice reconnaissance was provided by the SIR JOHN FRANKLIN's embarked helicopter, which was instrumental in enabling the ship to penetrate Robeson Channel to 82°09'N; the furthest northward of any ice breaker since 1972.

B. INSTRUMENTATION

The instruments used were the Applied Micro Systems Limited Conductivity-Temperature-Depth recorder, model STD-12. This instrument has a stated accuracy of 0.02 mS/cm in electrical conductivity, 0.01°C in temperature and no stated accuracy in depth. The resolutions were stated to be 0.003 mS/cm, 0.001°C, and 0.05 dbar, respectively.

Three instruments, numbered 422, 433 and 467, respectively, were utilized during the cruise. CTD #433 is the property of the Naval Postgraduate School (NPS), and was calibrated both before and after the cruise with resultant errors of 0.003°C in temperature and 0.007 mS/cm in conductivity. Calibrations for the other two instruments were not initially available, and were obtained empirically in the form of first



AN USE OF A CONTRACT OF A C

Figure 2.1 The locations of CTD stations conducted during the September 1986 cruise of CCGS SIR JOHN FRANKLIN.

degree polynomials by intercomparison with CTD #433 during simultaneous lowerings. It is concluded that the absolute error in the variables issmaller than 0.02°C, 0.02 mS/cm, and 2.0 dbars. The root mean square error in salinity, therefore, is approximately 0.03.

The data were recorded internally in coded binary form, and were downloaded into a Compaq computer for conversion into engineering units and storage on 5.25 in. floppy diskettes. Each CTD was programmed to sample the data stream every 0.8 dbar of pressure change. Significantly finer resolution in pressure was not convenient because of computer memory limitations and downloading time constraints.

The primary navigation aid was the ship's Magnavox MX 1107 Satellite Navigation System. An average of two fixes per hour provided a mean navigational accuracy of 0.5 km.

C. DATA REDUCTION AND ANALYSIS

Upon completion of the cruise, the data were transferred to mass storage cartridges for further processing with the NPS IBM 3033 computer. Editing of spurious and mis-sequenced data points, the cause of which is discussed by Tunnicliffe (1985), was subsequently performed. Removal of dynamic response errors in temperature and salinity was accomplished as prescribed by Bourke et al. (1986). The edited profile data were then sub-sampled every 5 m for use in the production of baroclinic velocity profiles and related volume transport estimates.

Transects used in the data analysis for the construction of vertical sections are depicted in Figure 2.2. Temperature-salinity (T/S) transects are oriented as noted, with temperature values $(^{\circ}C)$ represented as solid lines and salinity values (Practical Salinity



Figure 2.2

The locations of transects used in the analysis. The dotted line denotes stations used for construction of meridional cross sections.

Scale) represented as dashed lines. Baroclinic velocity cross sections are oriented with velocities (cm/sec) "into" and "out of" the page represented as solid (positive values) and dashed (negative values), respectively.

Surface dynamic heights and baroclinic velocity profiles were derived from the geostrophic approximation, assuming a level of no motion of 200 dbars. The technique of Helland-Hansen (1934) was deemed inappropriate for use in the NBB-NS region because of the inherently shellow and irregular bathymetry, thereby precluding the selection of a deeper reference level for regional application. Although a 200 dbar reference level is suitable for estimating direction in near-surface baroclinic currents, magnitudes were found to be underestimated by as much as 25% when compared to those derived with a level of no motion of 400 dbars.

Baroclinic volume transports were calculated using variable reference levels, determined by the maximum common sample depth in a station pair. As the sample depths were occasionally limited by mechanical rather than bathymetric considerations, the associated transports may be subject to underestimation accordingly. The transports were colculated by vertical trapezoidal integration of baroclinic velocities over 5 m intervals.

27

III. THERMOHALINE STRUCTURE

A. INTRODUCTION

In order to more accurately characterize the near-sulface and intermediate depth waters of the NBB-NS region, a deline dign of five discrete water masses by vertical distribution of temperature and salinity is required (Table I). Baffin Bay Surface Water (BBSW) is derived locally and found seasonally in northern Baffin Bay at depths of up to 75 m. It is present in both the Baffin Current and the WGC In general, the water masses which uniquely comprise the WGC are found to be less saline than their counterparts which enter the NBB-NS region through the Canadian Archipelago. Although the salinity difference between the Polar Water fractions is small. it is useful as a classification tool. Mixing of Arctic Basin Polar Water (ABPW) and West Greenland Current Polar Water (WGCPW) should be expected to occur along isopycnal surfaces in regions where their domains overlap. Although the salinity distributions of Nares Strait Atlantic Intermediate Water (NSAIW) and West Greenland Current Atlantic Intermediate Water (WCCAIW) do not intersect, it is bathymetry which prevents mixing of these two source fractions.

B. BAFFIN BAY SURFACE WATER

Baffin Bay Surface Water (BBSW) is locally derived from the combined effects of meltwater admixture and solar heating. The large, ubiquitous glaciers in the NBE-NS region serve as a continuous source of meltwater during periods of insolation. Admixture of glacial runoif with Baffin Bay Polar Water (BBPW) creates a significant bala line and resultant pycnocline in the upper 50 m to 10 m of the water column. Concurrently, TABLE I NEAR-SURFACE AND INTERMEDIATE DEPTH WATERS OF THE NORTHERN BAFFIN BAY-NARES STRAIT REGION

WATER MASS	CHARACTERISTICS	COMMENTS
•Baffin Bay Surface Water (BE	BSW) T: > 0°C S: < 32.8	•Surface water originally defined by Coachman and Aagaard (1974)
		-BBSW suggested by Fissel et al. (1982)
-Arctic Basin Polar Water (ABPW)	T: < 0°C S: < 34.8 T C 0.00	•Polar water originally defined by Aagaard and Coachman (1968)
	I-3 KHEE: 33.3 - 33.7 D: < 250 m	 I-S knee corresponds to bottom of halocline
·West Greenland Current Polar Water (WGCPW)	T: < 0° S: < 34.2	
	T-S knee: 33.4 - 33.5	
.Nares Strait Atlantic	T: > 0°C	•AIW originally defined by Aagaard
Intermediate water (NSAIM)	0: 250 m - 800 m D: 250 m - 800 m	and Coachman (1908)
.West Greenland Current	T: > 0°C	Generally warmer than NSAIW
Atlantic Intermediate Water (WGCAIW)	S: 33.8 - 34.5 D: 250 m - 800 m	

the effects of solar heating are essentially confined to this surface layer due to its inherent buoyancy. The admittedly arbitrary delineation of BBSW by the 0°C isotherm serves to distinguish it from BBPW, rather than imply that the effects of insolation are restricted to a discrete temperature range. BBSW is best identified, however, by the presence of a strong, shallow thermocline (Figure 3.1). A seasonal reversal of the processes which create BBSW, altimately results in its elimination.

The distribution of BBSW is a regional phonomenon. The horizontal distribution of maximum temperatures in the BSSW laver shows distinct gradients in the meridional and offshore directions (Figure 3.2). This result implies that BBSW is formed primarily by coastal processes, and is subsequently advected throughout most of northern Baffin Bay. The complete absence of BBSW north of Smith Sound is due to the widespread occurrence of pack ice, which serves to raise the albedo of the region and reduce insolation.

The considerable concentration of BBSW located in the shelf waters of Melville Bay contributes significantly to the baroclinicity of the WGC. A qualitative examination of a T/S transect through Melville Bay not only indicates that a major portion of the baroclinicity occurs in the upper 100 m, but that the isohalines (isopycnals) related to the presence of BBSW assume a negative slope in the core of the WGC between Stations 122 and 123 (Figure 3.3). The seasonal presence of BBSW. therefore, would be expected to enhance the baroclinic transport of the WGC.



A composite temperature profile illustrating the shallow thermocline characteristic of BBSW. Refer to Figure 2.2 for station location.



らいシン

ίΛ.

Figure 3.2 The horizontal distribution of maximum temperatures in the BBSW layer.


Temperature contours are indicated by solid lines, and salinity contours by dashed lines.

The presence of BBSW at an ice edge can be considered to represent an equilibrium state. The northernmost extent of BBSW is precisely correlated with the location of the pack ice edge. A meridional T/S transect from Baffin Bay northward to Kane Basin graphically illustrates this interaction between relatively warm surface water and an ice edge (Figure 3.4). A filament of BBSW dives subsurface between Stations 44 and 47 beneath water of lower density at the ice edge. An equilibrium state can exist if sufficient thermal advection is present to prevent ice growth. Mitigation of thermal advection or atmospheric cooling will cause the ice edge to advance or retreat, respectively.

C. POLAR WATER

Polar Water in the NBB-NS region is derived from two distinct sources. Direct inflow from the Arctic Ocean through the Canadian Archipelage is generically classified as Arctic Basin Polar Water (ABPW), whereas northward flow via the WGC through Davis Strait is termed West Greenland Current Polar Water (WGCPW). Subtle contrasts in the thermohaline signatures of these two water masses are indicative of their different origins, and serve as qualitative tracers of their respective flow fields.

Although both forms of Polar Water have origins in the Arctic Ocean, the flow of WGCPW follows a circuitous route around the southern tip of Greenland before entering the NBB-NS region. The cumulative influence of shelf and surface-driven processes on the thermohaline structure of WGCPW is, therefore, of greater magnitude than the effect of similar processes on ABPW. The primary modification which occurs is due to meltwater admixture, resulting in a net dilution of the water column.





Viewed simplistically, WGCPW can be considered a diluted form of ABPW The comparative decrease in salinity and, hence, density of WGCPW below the surface layer is most apparent in vertical property profiles (Figures 3.5 and 3.6). The maximum salinity of WGCPW is approximately 34.2, while ABPW reaches 34.8.

The presence of Polar Water with an underlying warm layer produces a characteristic inflection, or knee, when temperature is plotted against salinity. This knee can be viewed as a boundary between the bottom of the arctic halocline. formed as a consequence of convective overturn in winter, and a deeper thermocline, which marks the transition to an underlying warm layer. The salinity and depth at which this knee occurs are functions of the inherent thermohaline characteristics of the particular Polar Water mass. The combination of relatively dilute WGCPW and underlying Atlantic Intermediate Water results in a knee with a salinity of 33.4 to 33.5, at depths ranging typically from 50 m to 120 m. As ABPW enters the NBB-HS region through Lancaster Sound, for instance it is subject to interleaving and mixing along isopvcnal surfaces. This comparatively undiluted inflow is characteristically more dense than the resident Polar Water of northern Baffin Bay, and consequently sinks. The resultant combination of comparatively saline ABPW and underlying Atlantic Intermediate Water produces a knee with salinities between 33.5 and 33.7, at depths of up to 250 m. A composite T/S curve illustrates the contrasting knew salinity ranges of WGCPW and ABPW (Figure 3.7)

A reduction in significant insolation in the year conface layer, or the absence an underlying warm layer of the old to be called thermal







A composite density profile illustrating the dilution of WGCPW (Stations 122 and 140) relative to ARPW (Stations 6 and 87).



Ī

A composite T/S curve illustrating knee salinities characteristic of WGCPW (Stations 138 and 142), and ABPW (Stations 4 and 6).

gradient and a bend, rather than a sharply defined knee, in the curve. This condition is apparent north of Smith Sound due to the widespread occurrence of pack ice, with an associated increase in albedo (Figure 3.8).

The thermohaline characteristics of the two Polar Water masses can be used to roughly estimate the horizontal extent of their flow regimes. A plan view of Polar Water distributions indicates that inflow of WGCPW represents the minority fraction of Polar Water in the NBB-NS region (Figure 3.9). WGCPW is prevalent along the West Coast of Greenland northward to Cape York, where a bifurcation in flow creates one filament which extends to Smith Sound and another which projects westward into Jones Sound. Northward passage of WGCPW into Kame Basin was not observed.

ABPW is present from the Lincoln Sea to the southern end of Kane Basin, and throughout the central and western portions of Baffin Bay. The presence of ABPW at stations north of 77°N not only suggests that Smith Sound is a mixing site for the two Polar Water masses, but is indicative of southward passage of ABPW from Nares Strait into northern Baffin Bay. Inflow of ABPW is restricted to the southernmost portion of Jones Sound, implying significant mixing there as well — Inflow of ABPW to the region is greatest in Lancaster Sound, as indicated by the consistently high knee salinities encountered there. A filament of ABPW can be seen in the southwestern corner of Melville Bay at Stations 132 and 133, which marks the location of a frontal boundary between an extreme northeastern branch of the Baffin furtent, possibly derived from Lancaster Sound outflow, and a contrared thering in and of the Wet. The







Figure 3.9 The horizontal distribution of ABPW and WGCPW in the NBB-NS region.

extensive interleaving which would be expected in regions such as these, where ABPW and WGCPW coexist, is evident in composite temperature profiles (Figure 3.10).

The presence of subsurface temperature extremes in the Polar Water layer is associated with the contrasting effects of convective overturn in winter and insolation in summer. Convective overturn during ice formation creates a cold, isohaline layer which can project to a depth of greater than 50 m locally. Lateral infusion of brine produced during the freezing of shelf waters can increase the vertical extent of this isohaline layer to over 100 m. Subsequent surface heating and meltwater mixture produces significant enthalpy increases in the shallow pvcnocline layer only, isolating an underlying cold lens at depths typically between 50 m and 150 m.

South of Smith Sound cold lenses are sharply defined by strong temperature gradients. This is typically the case in Melville Bay, where cold, isohaline water is sandwiched between an overlying layer of BBSW and an underlying layer of Atlantic Intermediate Water (Figure 3.11).

The absence of cold lenses north of Smith Sound is indicative of the inherently persistent ice cover. The associated high albedo results in steadily decreasing temperatures towards the surface and, consequently, no subsurface cold lenses are apparent north of Station 72, located at approximately 79.2°N (Figure 3.12).

Southward decreases in ice concentration result in the formation of diffuse cold lenses in northern Smith Sound. A T/S transect taken at the southern entrance to Kane Basin remeals the remnants of a subsurface





4.,









cold lens on the western side, and alternating warm and cold layers to the east (Figure 3.13).

Pronounced warm lenses are located on the southern side of Lancaster Sound, and along both boundaries of Jones Sound. Conditions in Jones Sound, with ambient air temperatures averaging -4°C and the presence of new ice observed along both shores, indicate that the local occurrence of subsurface lenses of BBSW is due to the initiation of convective overturn (Figure 3.14). The warm lens in southern Lancaster Sound is a sharply defined feature located at a depth of approximately 10 m (Figure 3.15). The depth of this warm lens indicates that it is probably an outflow jet of ABPW which dives subsurface upon encountering a cooler, but comparatively less saline layer of resident Polar Water. This effect further illustrates the point that ABPW is more saline than the residual Polar Water mixture of northern Baffin Bay, and is subject to mixing along isopycnals at deeper depths.

The major source of meltwater in the NBB-NS region is the Humboldt Glacier, located on the eastern side of Kane Basin. All stations in Kane Basin south of approximately $79^{\circ}56'N$ (Station 84) have low surface salinities, typically approaching 31.0, and exhibit significant stairstepped lavering in vertical density structure (Figure 3.16). This not only illustrates the magnitude of glacial influence in the region, but suggests that extensive recirculation of near-surface waters may occur in Kane Basin. A significant local source of meltwater is apparent in the northwest corner of Robeson Channel, where surface salinities were generally below 30.0, the lowest encountered in the NBB-NS region (Figure 3.17).



Figure 3.13 A T/S transect across the southern entrance to Kane Basin. Note the remnant of a cold lens at Station 55, and alternating warm and cold layers to the east.



DAM SCOULDE

Figure 3.14 A T/S transect across the mouth of Jones Sound.



Figure 3.15 A T/S transect across the mouth of Lancaster Sound.



X





Figure 3 1 A T/S transect in Robeson Channel.

D. ATLANTIC INTERMEDIATE WATER

The two Atlantic Intermediate Water fractions found in the NBB SS region exhibit inherently different salinity characteristics and exclusive horizontal distributions. Nares Strait Atlantic Intermediate Water (NSAIW) is derived from the Atlantic Layer of the Arctic Basin and flows southward from the Lincoln Sea into Nares Strait. West treenland Current Atlantic Intermediate Water (WGCAIW) flows northward through eastern Davis Strait, advected by the WGC into northern Baffin Bay.

WCCAIW is warmer but less saline than NSAIW. Following the same line of reasoning which requires subdivision of the Polar Water tractions. WCCAIW is subject to more significant dilution effects by shelf and surface-driven processes compared to direct inflow from the Arctic Basin. Since CTD cast depths were not deep enough to reach the bottom of the WCCAIW layer, the upper limit in salinity presented in Table I for this water mass should be considered an estimate. The maximum salinity encountered in WCCAIW was 34.44, at a depth of 562 m -Station 124), whereas a maximum salinity of 35.24 was recorded for NSAIW at a depth of 481 m (station 97). The comparatively lower temperatures in NSAIW are a reflection of the general north-south temperature gradient in the region. The highest Atlantic Laver temperature recorded in NSAIW was 0.20°C at a depth of 715 m Station 75), in contrast with a maximum temperature of 1.81°C, recorded at a depth of 445 m (Station 133) for NGCAIW

The horizontal distributions of MSAIN and MODAIN in the NBB-NS region are primarily determined by the region data better a configure . For







Southward passage of NSAIW into Kane Basin is prevented by shoaling of the sea floor in southern Kennedy Channel (Figure 3.19). Although the major portion of WGCAIW flow is turned evclonically to the southwest by the coastlines of Melville Bay and Cape York, northward transport of the remainder is abruptly terminated by a shallow bank (200 m) located in the vicinity of the Carev Islands. Northward flow of WGCAIW could continue to follow a deep narrow channel around this bank (Figure 3.20) but was not observed to do so. Once turned to the southwest, the flow of WGCAIW throughout northern Baffin Bay is not restricted by bathymetry. It should be noted than even with the hypothetical occurrence of westward flow from Cape York, penetration into Jones Sound by WGCAIW would be impossible because of shallow banks located east of the mouth. The flow into Lancaster Sound, however, is not similarly restricted. It is apparent that southeastern Lancaster Sound is a site of significant interleaving of dissimilar water masses, as the associated T/S cross section reveals the presence of BBSW, ABPW, and WGCAIW in profile (refer to Figure 3.15). It is presumed that WGCAIW is present at stations 7, 8, 9, 10 and 11, although the shallow CTD casts at these stations precludes absolute confirmation.



19 A meridional IS transect from Each northward through Fennedy and Robeson Channels. The meridional distribution of NSALW is indicated by the extent of the 0°C isotherm.



G



IV <u>CIRCULATION AND TRANSFORT</u>

A. INTRODUCTION

Macroscale circulation in the NPR-NS region is dominated by the northward flowing WOC and the southward flowing Baffin Current. Although net southward flow in the region is maintained primarily by baroclinic teroing (Muench, 1971), bathymetry exerts considerable influence on the formation of mesoscale circulation patterns. An examination of the surface circulation of the region, as deduced from baroclinic velocity and dynamic height calculations, reveals the rature of this influence (Figure 4.1). Reference to this surface circulation diagram is implied in the ensuing discussions

B. DYNAMIC TOPOGRAPHY

An examination of the surface dynamic topography of the NBB-NS region referenced to 200 dbars indicates a general westward increase in dynamic heights (Figure + 2). This trend is associated with the influence of the Coriolis force on the net southward flow.

The shoreward gradient of the dynamic height contours, in northern Baffin Bay lends a exclonic appearance to the circulation pattern Meltwater runoff may be responsible for the increase in dynamic heights along the coastline of Melville Bay. The complex circulation patterns associated with the Wellare inferred from the reversal of gradients and bending of contours in the vicibily of the teversal of gradients and bending of contours in the vicibily of the level both. Northbestwird flow around tape fork is directed codes will to the level in the distribuof the case. Islands, we better the level for the level in the distribueacthward flow to 2000 to 200



Figure 4.1 Surface circulation in the NBB-NS region as deduced from baroclinic velocity and dynamic height calculations.



Figure 4.2

The surface dynamic popography referenced to 200 decibars, in dynamic centimeters.

the perimeter of a shallow bank. suggesting a possible topographic influence. Westward intrusions of surface flow are apparent at the northern sides of the entrances to both Lancaster and Jones Sounds. Although net flow is eastward in both cases, the closer spacing of contours in Lancaster Sound is associated with much greater baroclinic velocities. A significant portion of the ABPW outflow from Jones Sound into northern Baffin Bay is turned anticyclonically around the southeast shore of Devon Island, augmenting the westward intrusion of flow into Lancaster Sound.

The sharp bending of contours in Smith Sound marks the boundary between two intersecting circulation regimes. A northward flowing branch of the WGC turns cyclonically, coincident with the formation of an ice-edge jet at the southern limit of the ice pack. The 0.36 dynamic meter contour is representative of the impingement of the relatively warm surface current on the ice pack, with a resultant density gradient normal to the ice edge. The deflection of southward flow from Nares Strait at the front is also apparent, but determination of associated circulation patterns within Kane Basin is not possible at this resolution. The meandering 0.28 dynamic meter contour in Kane Basin, however, is suggestive of the potential for gyre formation in this region.

C. THE WEST GREENLAND CURRENT

Circulation of the WGC in northern Baffin Bay appears to be governed by conservation of potential monticity. As a typical eastern boundary current, the WGC may be a sumelify because the basis of the barotropic component of flow. Conservation of potential volticity.

requires barotropic flow to follow lines of constant f.H. where f is the Coriolis parameter, and H is the water depth. Since mesoscale variations in the Coriolis parameter can be considered negligible, flow will tend to parallel isobaths

Justification of the preceeding assumptions in the case of the WC is provided by a number of observations. Although the direction of mean flow in the WGC is closely correlated with the 500 m isobath. recurvature to the west and south is noted at three primary locations. In Melville Bay the flow is observed to recurve southeastward around a shallow bank. A cross section of baroclinic velocities in this area not only indicates significant deep flow around the periphery of the bank. but confirms the filamental nature of the return current (Figure 4.3). The two distinct flow regimes on the southwest side of the bank are, as stated in Chapter 3, of different origins. Flow between Stations 128 and 129 is a recurved branch of the WCC which, assuming conservation of relative vorticity, probably forms a cyclonic gyre around the perimeter of the bank. Southwest of an apparent frontal boundary, however, a second axis of flow exists which has been shown to principally ontain ABPW

Shoaling of the bottom in the vicinity of the Carev Island, causes complete recurvature of WGCAIW. The barcelinic velocity cross section from Jones Sound westward to Thule reveals that the core of the WGC is found east of Station 22 (Figure + + - Since water of Atlantic Laver origin has been confirmed between statisms 21 and 24 secured flow of WGCAIW must pass between Stations 1 and 1 and 24 secured flow of velocities between Stations 1 and 1 and

6.



from the geostrophic approximation, assuming a level of no A bareelinic velocity cross section through Melville Bav. Velocities between successive station pairs were derived motion at 200 dbars. Northward flow is fudleated by positive isotachs, and southward flow by negative tsotachs. Figure 4.3



Coburg Island (not visible) and the 150 m bank are located topographically-influenced cvclonic gyres associated with eastward to Thule. The main cores of the Baffin Current and the WCC are located between Stations 18 and 10, and Stations 22 and 23, respectively. The axes of flow for A baroclinic velocity cross section from Jones Sound at Stations 17 and 19, respectively. Figure 1 4

at 375 m, indicates that southward flow for this station pair is limited to the upper 55 m of the water column. Northward flow between Stations 22 and 23, calculated with a level of no motion at 400 m. decreased from 0.12 m/s at the surface to less than 0.03 m s at a depth of 115 m. The implication of this result is that most baroclinicity in the WGC is confined to the surface and near-surface layers. Barotropic currents are presumed to dominate below this level, hence, the postulated influence of topographic steering on the WGC.

The remaining branch of the WGC is recurved at the northern extent of the 500 m isobath in Smith Sound. A meridional velocity cross section through the region further substantiates the classification of the baroclinic component of this flow as an ice edge jet (Figure 4.5) Baroclinic velocities between Stations 41 and 44, calculated with a level of no motion at 400 m. decreased from 0.44 m/s at the surface to approximately 0.04 m/s at 200 m. Below 200 m. predominantly barotropic flow is topographically steered to the west and south by rapid shoaling of the bottom.

). CIRCULATION IN NARES STRAIT

The macroscale circulation regime in Nares Strait generates net southward flow of ABPW into Smith Sound. Mesoscale circulation patterns in the region, however, can not be characterized so succinctly.

Inflow from the Lincoln Sea proceed: southward through Robeson and Kennedy Channels. A baroclinic velocity cross section through a southern part of Kennedy Channel indicates the relatively uniform nature of this flow (Figure + 6). Recalling that while a title of NSALV is terminated by rapid shoaling just north of this transact creter to



B





Figure 4.6 A barocli

A baroclinic velocity cross section through a southern part of Kennedy Channel.

Figure 3.19), it is probable that the relatively high current speeds in this area are associated with constriction of flow. Furthermore, shoaling is more extreme on the eastern side of the channel resulting in correspondingly higher speeds. In contrast, the high baroclinic current speed in Robeson Channel is probably due to the influence of the previously described local concentration of meltwater.

Circulation in Kane Basin is characterized by southward flow in two peripheral branches surrounding a central cyclonic gyre. Circulation along the eastern side of Kane Basin is topographically steered southward along the inherently shallow seabed. Augmented by runoff from the Humboldt Glacier, this eastern branch continues southwestward and merges with recurved WGC water in Smith Sound. The flow of the western branch is also bathymetrically influenced. A portion of this branch turns cyclonically along the 200 m isobath forming a gyre, while the remainder continues southward, merging with both the western branch and the recurved WGC. The southern portion of the gyre is quite apparent in a baroclinic velocity cross section through Kane Basin (Figure 4.7).

In contrast to its effect on the WGC, topographic steering does not appear to play as strong a role in Kane Basin. This implies that circulation within Kane Basin has a significant baroclinic component of flow. This presumption is most apparent between Stations 74 and 76, at the northern edge of the gyre. Baroclinic velocities between these stations reverse from southward at the surface, to northward below 65 m depth. This vertical shear in velocities is indicative of baroclinic instability, which might serve as a potential contract sense for either the Kane Basin Gyre or smaller understable to the
Elgure 4.7 A baroclinic velocity cross section through the southern portion of the Kane Basin Gyre.



E. THE BAFFIN CURRENT

The Baffin Current originates in Smith Sound, where southwestward flow from Nares Strait merges with recurved WGC water — During it: progression southward, the Baftin Current is augmented by outflow from Jones and Lancaster Sound , and by WGC water which has recurved south of Smith Sound.

As a typical western boundary current, the Baifin Current can be considered highly baroclinic. A boroclinic velocity profile between Stations 28 and 29, assuming a level of no motion at 400 m. indicates surface flow of 0.19 m/sec steadily decreasing to 0.5 m/sec at 200 m. Unlike the WGC, significant baroclinicity in the Baffin Current is not confined to the upper 100 m.

Interaction between the Coriolis force and bathymetry provides the primary steering influence for the Baffin Current. Throughout the course of the current, the tendency for westward turning by the Coriolis force is offset by the presence of a lateral boundary. At the northern entrance to Jones and Lancaster Sounds, however, the removal of this restriction permits the observed westward intrusion of flow. It should be emphasized, however, that the barotropic component of flow associated with the Paffin current is not negligible. An examination of the baroclinic velocity cross section from Jones Sound eastward to Thule provides three clear cases of topographic steering rin conjunction with Figure 4.4, refer to Figure 4^{-1} . Although chronic et the Baffin Current turns westward into fores bound, the matrix is et the Baffin proceeds southward between Stations 1, a^{-1} is a figure 4 to the source for bathymetry. Also apparent are two cyclonic gyres; one around Coburg Island, and the other in the vicinity of a 150 m bank.

Eastward outflow from the south side of Jones Sound subsequently merges with the main core of the Baffin Current along the coast of Deron Island. In contrast to the narrow filament entering Jones Sound, the major branch of the Baffin Current forms the westward intrusion of flow into Lancaster Sound.

F. CIRCULATION IN THE SOUNDS

1. Introduction

Smith. Jones and Lancaster Sounds are all considered regions of net volume inflow to northern Baffin Bay. An examination of the circulation characteristics associated with each sound will provide a basis for their comparison.

2. Smith Sound

Within Smith Sound exists a frontal boundary separating southward flowing ABPW and recurved WGC water. Circulation in this region, therefore, is determined primarily by baroclinic influences. A baroclinic velocity cross section through Smith Sound reveals a weak core between Stations 41 and 4? which is actually the remaining branch of the northward flowing WGC (Figure 4.8). Complete recurvature of this branch occurs just north of Station 41. Southward flow between Stations 42 and 43 is a filament of ABPW from the east branch of circulation in Kane Basin. The interaction between recurved BBSW and southwestward flowing ABPW is marked by the presence of a distinct ice edge. Indeed, the impingement of warm BBSW on the ice of a distinct ice edge. Indeed, the impingement of warm BBSW on the ice of a circulation providing a strong



-

ŏ.



baroclinic influence on flow direction. The resultant formation of an ice edge jet between Stations 40 and 41 marks the origin of the Baffin Current. Augmentation of the Baffin Current by ABPW occurs beneath the shallow pychocline, which is associated with relatively warm, buoyant BBSW from the WGC (refer to Figure 3.4).

Significant bathymetric influence in Smith Sound is limited to positioning of the frontal boundary between the two circulation regimes. Although topographic steering along the 500 m isobath clearly influences re urvature of the WGC, baroclinic effects are responsible for the formation of the Baffin Current. Smith Sound, therefore, is the site where a significantly barotropic circulation regime is transformed into a predominantly baroclinic one.

3. Jones Sound

The combined influence of bathymetry and the Coriolis force on the circulation structure of Jones Sound is most apparent in a cross section of baroclinic velocities (Figure 4.9). A filament of the Baffin Current is deflected westward by the Coriolis force around the southeast corner of Ellesmere Island and into Jones Sound north of Station 15. Eastward outflow of ABPW is restricted to the south of Station 15, continuing along the east coast of Devon Island. The presence of a cyclonic gyre around Coburg Island (refer to Figure 4.1) links the flow paths accordingly.

While eastward outflow from Jones Sound tends to follow the 500 m isobath, the westward intrusion is driven by the Coriolis force over a 200 m bank, against prevailing westerly winds. The baroclinic



アンドンドレ



velocity cross section illustrates the relatively shallow nature of this westward intrusion of flow.

4. Lancaster Sound

Circulation in Lancaster Sound is characterized by a westware intrusion of flow between Stations 6 and 8, and an eastward return current of greater magnitude between Stat ones 1 and 6 (Figure 4.10) The increase in velocities towards the sides of the sound indicates the effect of the Coriolis force on the inflow and outflow branches. Baroclinic velocities in the outflow branch are the highest in the entire NBB-NS region, reaching a maximum of 0.78 m/sec at the surface between Stations 2 and 3 (assuming a level of no motion at 340 m).

5. <u>Geostrophic Estuarine Circulation</u>

Although their thermohaline and bathymetric features are inherently different, the general characteristics of circulation in Smith. Jones and Lancaster Sounds are remarkably similar. In each case, geostrophically balanced inflow is entrained, in estuarine fashion, by geostrophically balanced outflow. A quasi-vertical frontal feature initially separates the two current regimes, the slope of which is determined by their respective velocity structures (refer, for example, to Figure 4.10). The dynamic aspects of the ensuing model, hereinafter referred to as Geostrophic Estuarine Circulation (GEC), are derived from the work of Leblond (1980). Variations in the observed character of the circulation in each sound can be explained in terms of how closely local conditions approximate GEC.

In proposing a two layer "Soastal Current Model" to explain the observed circulation in some channels of the smaller archipelago.



WWWWWWWWWWWWWWW

ALLARIA SUSSIES UNDER



Leblond showed that two geostrophically balanced, upper-layer flows would coexist without interference in a channel whose width is sufficiently large compared to the local internal Rossby radius of deformation. Restriction of flow to the upper layer creates a sloped interfacial surface, resulting in a characteristically wedge-shaped velocity structure (Figure 4.11)



Figure 4.11 A schematic cross section of coastal upper layer flow (out of the page) of speed u, driven by a sea surface slope $\eta(x)$. The lower layer of density ρ_2 is at rest because the interface h(v) slopes in a direction opposite to that of the free surface. Y_0 is the distance from the coast at which the thickness of the upper layer vanishes (from Leblond, 1980, p. 191).

Additionally, Leblond found that purely geostrophic flow can turn corners without separating from a coast if the radius of curvature of the coast is large enough that the Rossov number remains well below unity. As a conservative estimate of the width of a coastal geostrophic current, Y_0 is defined as $Y_0 = R/F$: where R is the internal Rossov radius of deformation and F is the internal Froude number — If the width of a channel is shown to exceed 2 (Y_0) , two geostrophically balanced upper-layer flows can theoretically exist without interference The results of applying I blond's equations to Smith. Jones and Lancaster Sounds are listed in Table II. with the requisite calculations presented in the Appendix – Since the computed Rossby number (R_0) at the mouths of all three sounds is less than 0.1. nonlinear effects may be locally neglected – It should be noted that values of Y_0 are only calculated for the outflow regimes which, in each sound, were the widest of the two currents

Conditions in Lancaster Sound, with a minimum channel width (L) of 80 km, and a geostrophic current width (Y_0) of 20.35 km, most closely approximate GEC. Referring to the baroclinic velocity cross section through Lancaster Sound (Figure 4.10), it clear that although the calculated value of Y_0 is considered an underestimation of the width of the outflow current, interference between the two circulation regimes appears negligible. The obvious wedge-shaped current structure implies that flow is primarily restricted to the upper 150 m. The broadening of the outflow wedge beyond the calculated value of γ_0 , however, indicates that this restriction is not complete. Indeed, the likelihood of significant flow occurring below 250 m depth is confirmed by the presence of We AIW in western Lancaster Sound.

Entrainment of inflow results in cyclonic cross-channel flow The connection of inflow and outflow currents in Lancaster Sound by cross-channel flow was most re-ently investigated by Fissel et al (1982), who found the westward extent of the inflow intrusion strongly marked by dynamic height anomalies. For use a loc of this result illustrates that, in GFC evolves of the test of the second with entrainment of infloc of each of the inflow intrusion. The

<u>`</u>~

TABLE II GEOSTROPHIC ESTUARINE CIRCULATION MODEL CALCULATIONS

.

: Minimum Channel ch Width (L)	80 km	60 km	35 km
Geostrophic Current Widt (Y ₀)	20. 4 km	30.3 кт	25.3 кш
Internal Rossby Radius	8.5 km	7.3 km	5.4 km
Densimetric Frcude Number (F)	0.42	0.24	0.21
Rossby Number (R _O)	0.06	0.07	0.02
	Lancaster Sound	Smith Sound	Jones Sound

four periods of CTD observations in 1979: (A) buly 18-19, Sound (in dynamic cm) relative to the 300 dbar level for Surface dynamic height anomalies in eastern Lancaster (B) July 29 - August 3; (C) September 2 4; and (D) September 14-23 (from Fissel et al., 1982, p. 1875) Figure 12



convergence of inflow and outflow regimes creates a local elevation in the sea surface, generating geostrophic flow normal to the resultant pressure gradient. With sufficient eastward outflow and the presence of lateral boundaries, characteristic cyclonic flow will result. Since convergence implies downwelling, a reduction in vertical density gradients may occur along this frontal feature, thereby enhancing the tendency for mixing.

In Lancaster Sound the eastward outflow branch of the circulation illustrates the increase in volume and velocity typically associated with estuarine entrainment. In GEC, however, the entrainment is geostrophically, not turbulently, driven.

The application of GEC to Smith Sound is influenced by both geometric constraints and significant horizontal density gradients. Referring to Table II, it is apparent that L is approximately equal to 2 (Y_0) . Since this is considered a minimum requirement, the potential for interference between the inflow and outflow regimes may be significant. This interference may manifest itself in the form of baroclinic instability and associated eddy generation.

Within Smith Sound the inflow and outflow regimes are not appreciably influenced by coastal boundaries. The absence of a characteristically wedge-shaped velocity structure in this region is probably a consequence of this fact. Geostrophy does influence the direction of the inflow and outflow branches, however, resulting in cyclonic cross-channel flow. As stated previously, recurvature of the remaining branch of the WGC occurs as the result of an extreme horizontal density gradient at the include theory. Entrainment of WGC

inflow in Smith Sound, while not directly attributed to convergence, is nonetheless geostrophically balanced.

The limitations in applying GFC to Jones Sound are considerable. Since $L < 2(Y_0)$. Jones Sound is wide enough to accommodate only one geostrophically-balanced current – Referring to the 33.0 isohaline in Figure 3.14. the formation of densily and, hence, baroclinic velocity wedges is evident at the sides of the channel; however, a complete blurring of the structure occurs between Stations 14 and 15. Since these current branches can not coexist in Jones Sound, it may be possible for complete intrusion of the westward inflow branch to occur at times, resulting in reversal of the net eastward flow in Jones Sound. Historical evidence of this occurrence has been cited by Muench (1971) (Figure 4.13). In 1962, for example, the flow is entirely westward, while in 1963 both inflow and outflow are present simultaneously but with the outflow dominant.

The macroscale circulation pattern in the NEB-NS region can also be envisioned as conforming to the GEC model Viewed simplistically, the inflowing waters of the WGC are geostrophically entrained at various latitudes, subsequently resulting in the augmented outflow which characterizes the Baffin Current. The predictably cyclonic circulation pattern associated with GEC is identifiable as a major macroscale feature in the region.

G. BAROCLINIC TRANSPORTS

Baroclinic transports derived is a various arbitrary reference levels should be used, only with the table of a sector forter continuity of volume. This is especially true in eastern boundary surrent regions.





like the WGC, where the baroclinicity is known to be weak. The main purpose of this section, therefore, will be to qualitatively compare the baroclinic components of flow for major currents in the NBB-NS region Unless otherwise specified, subsequent references to transports are summarized in plan view and assumed to be baroclinic in nature (Figure 4.14).

The WGC achieves maximum transport in Melville Bac, prior to its previously described partern of recurvature in norther: Baffin Bav and Smith Sound. Transport of the WGC consequently decreases from a maximum of 0.7 Sv in Melville Bay, to 0.4 Sv in the vicinity of the Carey Islands, and 0.2 Sv in Smith Sound. The major portion of baroclinic transport is concentrated at the extreme inshore stations in Melville Bay, further substantiating the assumption that glacial runoff is primarily responsible for this component of WGC flow.

The Baffin Current is partially derived from the net southward flow of Nares Strait. Southward transport increases from 0.2 SV in Robeson Channel to 0.7 SV at the mouth of Kennedy Channel. This increase may be attributed to admixture of meltwater runoff, in particular from Petermann's Glacier, located on the eastern shore of Robeson Channel. Southward transport through Kane Basin is primarily associated with the western circulation branch (0.3 SV), while southward transport in the eastern circulation branch is limited (0.1 SV). The swirl transport within the Kane Basin Gyre is approximately 0.4 St, indicative of its effect in recirculating meltwater from the Humboldt Clacier throughout the basin. The cyclonic ice-edge jet in Smith Sound, with associated transport of 0.5 SV, can be considered a schemet from Sares Strait



Figure 4.14 Major baroclinic transports in the NBB-NS region.

which is augmented by enryphics that the rescander of the West OPUS System **Net southward** outliew is an Ease. Basin through Smith Sound, therefore, is interied to be approximately 0.3 System.

The Baftip current, which becomes an identifiable feature in Smith Sound, continues southwart as one the cost of Ellesment Island where transport is measured as 1.9 sv. The committee of a superaphically induced evolution are is contrident with branching of the Battin Currentmortheast of Toburg Island - A westward to wing tilament (0.1 Sv) enters Jones Sound, where it is entrained and augmented by eastward outflow (0.4 Sv), resulting in net eastward transport of 0.3 Sv A southwestward flowing branch (2.5x) probably follows the 500 m isobath, subsequently merging with the main core of the Baffin Current. The relatively large transport (0.7 Sv) indicated southwest of Melville Bay is actually an algebraic summation of transports obtained between Stations 127 and 132 (refer to Figure +3) Southeastward transport of Baffin Current water (0.4 Sy), probably devived from a brath of Lancaster Sound outflow which is topographically steered eastward along the 1000 m isobath (Muenels 1971), is apparent southwest of Station I30. Flow of WGC water, which has recurved either around the shallow bank in Melville Bay (0.2 Sv) or south of the Carer Islands (0.1 Sv), is indicated between Stations 127 and 129

The main core of the Battin Current, sugmented by eastward outflow from Jones Sound, follows the southwest court of Devon Island westward into Lancaster Sound (0.6 System GEC entranement and subsequent augmentation of outflow vields groups and outflow of 1.1 System

Sh

transport in Lancaster Sound, therefore, is inferred to be 1.1 Sv eastward.

Ĭ,

• •

È

V. THE NORTH WATER PROCESS

The North Water Process is characterized by complex interaction between thermohaline and dynamis effects. The net result of this process is the demarcation of an 80,000 km² arctic region which is conspicuously devoid of first year ice from the onset of winter until March (Steffen and Ohmura 1985). Following a brief examination of some relevant factors, a qualitative description of the process will be presented.

In southern Baffin Bay freezing is generally accompanied by the formation of an isothermal isohaline near-surface layer approximately 75 m to 100 m in depth (Garrison et al., 1976). Sea-ice production in the NBB-NS region is similarly associated with the formation of such deep mixed layers (Muench, 1971). In the weakly stratified waters of Nares Strait, convective processes are sufficient to form this homogenous layer. The region south of Smith Sound, however, is characterized by the ubiquitous occurrence of BBSW. The resultant presence of such a strong pycnocline would ordinarily limit the depth of convective mixing associated with surface freezing to much less than 75 m. A combination of turbulent and convective processes must be required, therefore, to overcome the inherent stability of this buoyant layer and produce the observed deep mixed layers.

Given a particular set of climatic conditions, the enthalpy present in the near-surface layer determines the length of time required to initiate surface freezing. Assuming heat flux only through the surface and neglecting thermal advection, it is possible to evaluate this time delay as a function of heat budget parameter. Staffen and Ohmura

(1985) computed monthly and annual heat budgets for the North Water region and concluded, assuming a reference temperature of -1.8°C, that the amount of oceanic heat flux required to prevent ice formation is 337.4 MJM⁻² in October and 445.1 MJM⁻² in November. The enthalpy contained in the 75-m deep, near-surface layer was calculated. referenced to -1.8°C, for all FRANKLIN 86 stations (Figure 5.1). For 22 stations located southward of Smith Sound, between 76° N and 78° N, the enthalpy in September averaged 475 MJ. Applying the aforementioned heat budget at the onset of winter (October), implies that surface freezing would be delayed by approximately five weeks in this region. In contrast, the average enthalpy contained in the ice-covered, nearsurface layer north of 78°N (Nares Strait) was 120 MJ.

The presence of mechanical ice removal effects such as divergent currents and winds will cause the rate of sea-ice production to greatly exceed the rate of sea ice accumulation for a given area. This phenomena is graphically illustrated in the vicinity of the Kane Basin Gyre where cyclonic and, hence, divergent circulation results in an area of reduced ice concentration. Ice concentrations within the gyre were typically 3/10 or less, compared with surrounding concentrations of greater than 5/10. The cyclonic macroscale circulation pattern characteristic of northern Baffin Bay can be expected to provide a similar mitigating influence on the accumulation of sea-ice. The potential influence of persistent northerly winds on the reduction of sea-ice concentration in northern Baffin Bay has been discussed by Muench (1971) and Dunbar (1973).



Figure 5.1 Enthalpy of the near surface (upper 75 m) layer in the NBB-NS region (referenced to - 1.8°C).

والمراجع والمراجع والمراجع والمراجع والمحاج والمحاج والمحاج والمحاج والمحاج والمحاج والمحاج والمحاج والمحاج وال

The North Water Process results from the combined influences of near-surface layer enthalpy and mechanical ice removal. Since the North Water is characterized by reduced ice coverage, the associated decrease in albedo results in the largest annual net radiation for any region within the Arctic Circle (Steffern and Ohmura, 1985). Admixture of glacial runoff creates a strong, shallow pychocline which effectively confines insolation effects to the near-surface layer. Such inherent stratification would normally restrict convective overturn associated with surface freezing to shallower depths. Mechanical ice removal. however, results in a proportionally greater rate of brine formation which increases the vertical extent of mixing. Complete removal of the enthalpy in this near surface layer delays significant sea-ice formation accordingly. Divergent cyclonic currents and northerly winds then serve to reduce sea-ice accumulation for the remainder of winter. It is the self-perpetuating nature of the North Water Process which accounts for its persistence.

VI. LISCUSSION

The comprehensive nature of this analysis mandates further examination of certain relevant topics. A combination of elaboration and postulation will be employed in this regard.

The dilution of WGC water masses, relative to those which enter the NBB-NS region through the Canadian Archipelago, is a consequence of the more extensive shelf-driven modification effects inherent in the former. The assertion by Muench (1971), that the major proportion of Arctic Ocean Water is less dense than Baffin Bay Water, was derived from point data sources, under the assumption that there are significant differences in water structures between the upper 250 m layers of the Arctic Ocean and Baffin Bay. Muench further concludes that these waterstructure differences create a proportionally higher surface elevation in the Arctic Ocean, resulting in net southward transport.

An examination of the dynamic topography of the NBB-NS region reveals an obvious lack of north-south dynamic height gradients (refer to Figure 4.2). This is due to the fact that ABPW and WGCPW have a common origin in the Arctic Basin, and are quite similar in thermohaline character. What is apparent, however, is a distinct dynamic height gradient normal to the coastline. Although the increase in dynamic heights in the onshore direction can be viewed as a consequence of the influence of the Coriolis force on the prevailing baroclinic flow, it may alternately be considered as a driving force for the observed circulation in the NBB-NS region.

The ubiquitous glaciers in the restor is a subject solution of the definition of the second states of meltwaters an influence of the second states of the drive a

geostrophically-balanced current with the shoreline on the right. in the direction of flow. The implication of this hypothesis is that a progressive winter weakening in near-surface currents would be expected throughout the NBB-NS region. Lemon and Fissel (1982) provided evidence of just such an occurrence in northwestern Baffin Bay.

On a larger scale, the concentration of glacial meltwater within the NBB-NS region results in near-surface layer dilution relative to the saltier waters of the North Atlantic. It is the difference in nearsurface water structure between these regions that may drive the net southward baroclinic transport. A progressive winter decrease in this net southward transport would be expected accordingly.

VII. CONCLUSIONS

The physical oceanography of the northern Baffin Bay - Nares Strat region has been investigated using a dense network of CTD stations occupied by the CCGS SIR JOHN FRANKLIN in September 1986. The following major conclusions can be drawn:

- * Five water masses can be delineated in the region. The WGC is the source of WGCPW and WGCAIW, while ABPW and NSAIW are derived from the Arctic Basin via the Canadian Archipelago. WGCAIW and NSAIW are prevented by shoaling of the sea floor from entering Smith Sound and Kane Basin, respectively. BBSW is found seasonally throughout northern Baffin Bay.
- * The waters unique to the WGC are subject to more extensive dilution effects by shelf-driven processes and are comparatively less saline than their Canadian Archipelago derived counterparts.
- * Bathvmetry provides a significant influence on the flow of the WGC, which attains a maximum baroclinic transport of 0.7 SV in Melville Bay. Recurvature of component branches of the WGC occurs primarily in Melville Bay (0.2 Sv), south of the Carey Islands (0 1 Sv), and ultimately in Smith Sound (0.2 Sv).
- * The Baffin Current originates as an ine edge jet in Smith Sound subsequently proceeding southward along the coast of Elleumen Island and throughout worthsee are lated as a log part of





5.25

Baffin Current is augmented by net outflow from Kane Basin, Jones Sound and Lancaster Sound at rates of 0.3 Sv, 0.3 Sv and 1.1 Sv, respectively.

* Circulation in Smith, Jones and Lancaster Sounds can be described in terms of the GEC model, in which estuarine inflow, entrainment and outflow are geostrophically balanced processes.

- * Although a significant portion of the meltwater derived from the Humboldt Glacier is recirculated by the Kane Basin Gyre (0.4 Sv), the influence of glacial admixture on thermohaline structure and near-surface circulation is apparent throughout the NBB-NS region.
- * The North Water Process results from the combined influences of near-surface layer enthalpy and mechanical ice removal.

APPENDIX

COMPUTATIONS ASSOCIATED WITH THE GEC MODEL

The computation of geostrophic current widths in Smith, Jones and Lancaster Sounds is accomplished as prescribed by Leblond (1980). The requisite equations and calculations are presented here.

The "Coastal Current Model" of Leblond (1980) assumes geostrophic flow confined to the upper layer of a two-layer stratified fluid, in a channel of width L and depth H (refer to Figure 4.11). Assuming the upper layer is relatively thin with respect to the lower layer, the speed of interfacial waves in the fluid may be approximated as:

$$C^{2} = \frac{g(\rho^{2} - \rho_{1}) t(o)}{\rho_{1}}$$
(1)

where: g = the gravitational acceleration;

ETLACK

 $\label{eq:relation} \begin{array}{l} \rho_1 = \mbox{the density of the upper layer;}\\ \rho_2 = \mbox{the density of the lower layer; and}\\ \mbox{t(o) = the thickness of the upper layer at the channel wall.}\\ \mbox{In Smith Sound:} \quad \rho_1 = 1025.5 \ \mbox{kg/m}^3; \ \rho_2 = 1027.0 \ \mbox{kg/m}^3; \ \mbox{and t(o)}\\ = \ \mbox{75 m.}\\ \mbox{In Jones Sound:} \quad \rho_1 = 1026.2 \ \mbox{kg/m}^3; \ \ho_2 = 1027.0 \ \mbox{kg/m}^3; \ \mbox{and t(o)}\\ = \ \mbox{75 m.}\\ \mbox{In Lancaster Sound:} \quad \rho_1 = 1026.0 \ \mbox{kg/m}^3; \ \ho_2 = 1027.0 \ \mbox{kg/m}^3; \ \mbox{and t(o)}\\ = \ \mbox{150 m.} \end{array}$

The internal Rossby radius of deformation is defined as:

$$\mathbf{R} = \mathbf{C}/\mathbf{f} \tag{2}$$

where: f = the Coriolis parameter.

The internal Froude number is defined as:

$$\mathbf{F} = \mathbf{u}/\mathbf{C} \tag{3}$$

where: u = the speed of the upper layer flow.

In Smith Sound: u = 0.25 m/s.

In Jones Sound: u = 0.16 m/s.

In Lancaster Sound: u = 0.50 m/s.

The Rossby number is defined as:

$$R_{o} = \frac{R}{fr}$$
(4)

where: r = the radius of curvature of the flow.

In Smith Sound: r = 25 km. In Jones Sound: r = 75 km. In Lancaster Sound: r = 60 km. The width of the geostrophic current is then defined as:

$$Y_o = \frac{R}{F}$$
 (5)

LIST OF REFERENCES

Aagaard, K., and L.K. Coachman, The East Greenland Current north of the Denmark Strait, I, <u>Arctic</u>, 21, 181-200, 1968.

- Bourke, R.H., R.G. Paquette, and A.M. Weigel, MIZLANT 85 data report: results of an oceanographic cruise to the Greenland Sea, September, 1985, Tech. Rep. NPS 68-86-007, Dept. of Oceanography, Naval Postgraduate School, Monterey, California, 1986.
- Coachman, L.K. and K. Aagaard, Physical oceanography of Arctic and subarctic seas, in <u>Arctic Geology and Oceanography</u>, edited by Y Herman, pp. 1-72, Springer-Verlag, New York, 1974.

reserves, courses through

- Dunbar, I.M., Winter regime of the North Water. <u>Trans. R. Soc. Can.</u>, 4(9), 275-281, 1973.
- Fissel, D.B., D.D. Lemon, and J.R. Birch, Major features of the summer near-surface circulation of western Baffin Bay. 1978 and 1979. <u>Arctic</u>, 35, 180-200, 1982.
- Garrison. G.R., H.R. Feldman and P. Becker. Oceanographic measurements in Baffin Bay, Unpublished manuscript. Applied Physics Laboratory University of Washington. February. 1976.
- Leblond, P.H., On the surface circulation in some channels of the Canadian Arctic Archipelago, <u>Arctic</u>, 35, 189-197, 1980.
- Lemon, D.D. and D.B. Fissel, Seasonal variations in currents and water properties in northwestern Baffin Bay, 1978 and 1979, <u>Arctic</u>. 35, 211-218, 1982.
- Melling, H., R.A. Lake, D.R. Topham. and D.B. Fissel. Oceanic thermal structure in the western Canadian Arctic, <u>Continental Shelf</u> <u>Research</u>, 3, 233-258, 1984.
- Muench, R.D., The physical oceanography of the northern Baffin Bay region. Baffin Bay-North Water Proj. Arct. Inst. North Am. Sci. Rep., 1-72, 1971.
- Muench, R.D., and H.E. Sadler, Physical oceanographic observations in Baffin Bay and Davis Strait. <u>Arctic</u>, 26, 73-76, 1973.
- Sadler, H.E., Water, heat, and salt transports through Nares Strait, Ellesmere Island, <u>J. Fish, Res. Board Can.</u>, 33, 2286-2295, 1976.
- Steffen, K. and A. Ohmura, Heat exchange and surface conditions in the North Water, northern Baffin Bay, <u>Ann. Glaciology</u>, 6, 178-181, 1985.

Tunnicliffe, M.D., An investigation of the waters of the East Greenland Current, Master's Thesis, Naval Postgraduate School, Monterey, California, September 1985.

INITIAL DISTRIBUTION LIST

10000

 No. Copies

1.	Director Applied Physics Laboratory Attn: Mr. Robert E. Francois Mr. E.A. Pence Mr. G.R. Garrison Library University of Washington 1013 Northeast 40th Street Seattle, Washington 98105	1 1 1 1
2.	Director Arctic Submarine Laboratory Code 19, Building 371 Naval Ocean Systems Center San Diego, California 92152	5
3.	Superintendent Naval Postgraduate School Attn: Dr. R.H. Bourke, Code 68Bf Dr. R.G. Paquette, Code 68Pa Dr. D.C. Smith IV, Code 68Si Monterey, California 93943	7 1 1
4.	Polar Research Laboratory, Inc. 6309 Carpinteria Ave. Carpinteria, California 93103	1
5.	Chief of Naval Operations Department of the Navy Attn: NOP-02 NOP-22 NOP-964D2 NOP-095 NOP-098 Washington, District of Columbia 20350	1 1 1 1
6.	Commander Submarine Squadron THREE Fleet Station Post Office San Diego, California 92132	1
7.	Commander Submarine Group FIVE Fleet Station Post Office San Diego, California 92132	1
8.	Dr. John L. Newton 10211 Rookwood Drive San Diego, California 92131	1

9.	Director Marine Physical Laboratory Scripps Institution of Oceanography San Diego, California 92132	1
10.	Commanding Officer Naval Intelligence Support Center 4301 Suitland Road Washington, District of Columbia 20390	1
11.	Commander Space and N∈val Warfare Systems Command Department of the Navy Washington, District of Columbia 20360	1
12.	Director Woods Hole Oceanographic Institution Woods Hole, Massachusetts 02543	1
13.	Commanding Officer Naval Coastal Systems Laboratory Panama City, Florida 32401	1
14.	Commanding Officer Naval Submarine School Naval Submarine Base, New London Groton, Connecticut 06349-5700	1
15.	Assistant Secretary of the Navy (Research and Development) Department of the Navy Washington, District of Columbia 20350	1
16.	Director of Defense Research and Engineering Office of Assistant Director (Ocean Control) The Pentagon Washington, District of Columbia 20301	1
17.	Commander, Naval Sea Systems Command Department of the Navy Washington, District of Columbia 20362	1
18.	Chief of Naval Research Department of the Navy Attn: Code 102-OS Code 220 Code 1125 Arctic 800 N. Ouincy Street Arlington, Virginia 22217	1 1 1

19.	Project Manager Anti-Submarine Warfare Systems Project Office (PM4) Department of the Navy Washington, District of Columbia 20360	1
20.	Commanding Officer Naval Underwater Systems Center Newport, Rhode Island 02840	1
21.	Commander Naval Air Systems Command Headquarters Department of the Navy Washington, District of Columbia 20361	1
22.	Commander Naval Oceanographic Office Attn: Library Code 3330 Washington, District of Columbia 20373	1
23.	Director Advanced Research Project Agency 1400 Wilson Boulevard Arlington, Virginia 22209	1
24.	Commander SECOND Fleet Fleet Post Office New York, New York 09501	1
25.	Commander THIRD Fleet Fleet Post Office San Francisco, California 96601	1
26.	Commander Naval Surface Weapons Center White Oak Attn: Mr. M.M. Kleinerman	1
	Library Silver Springs, Maryland 20910	1
27.	Officer-in-Charge New London Laboratory Naval Underwater Systems Center New London, Connecticut 06320	1
28.	Commander Submarine Development Squadron TWELVE Naval Submarine Base New London Groton, Connecticut 06349	1
29.	Commander Naval Weapons Center Attn: Library China Lake, California 93555	1
-----	--	---
30.	Commander Naval Electronics Laboratory Center Attn: Library 27: Catalina Boulevard San Diego, California 92152	1
31.	Director Naval Research Laboratory Attn: Technical Information Division Washington, District of Columbia 20375	1
32.	Director Ordnance Research Laboratory Pennsylvania State University State College, Pennsylvania 16801	1
33.	Commander Submarine Force U.S. Atlantic Fleet Norfolk, Virginia 23511	1
34.	Commander Submarine Force U.S. Pacific Fleet Attn: N-21 Pearl Harbor, Hawaii 96860-6550	1
35.	Commander Naval Air Development Center Warminster, Pennsylvania 18974	1
36.	Commander Naval Ship Research and Development Center Bethesda, Maryland 20084	1
37.	Commandant U.S. Coast Guard Headquarters 400 Seventh Street, S.W. Washington, Nistrict of Columbia 20590	1
38.	Commander Pacific Area, U.S. Coast Guard 630 Sansome Street San Francisco, California 94126	1
39.	Commander Atlantic Area, U.S. Coast Guard 159E, Navy Yard Annex Washington, District of Columbia 20590	1

ALARTAR STATE AND THE ALARTAR STATES STATES AND AND A CONTRACT AND A

40.	Commanding Officer U.S. Coast Guard Oceanographic Unit Building 159E, Navy Yard Annex Washington, District of Columbia 20590	1
41.	Scientific Liaison Office Office of Naval Research Scripps Institute of Oceanography La Jolla, California 92037	1
42.	Scripps Institution of Oceanography Attn: Library P.O. Box 2367 La Jolla, California 92037	1
43.	School of Oceanography University of Washington Attn: Dr. L.K. Coachman Dr. S. Martin Mr. D. Tripp Library Seattie, Washington 98195	1 1 1 1
44.	School of Oceanography Oregon State University Attn: Library Corvallis, Oregon 97331	1
45.	CRREL U.S. Army Corps of Engineers Attn: Library Hanover, New Hampshire 03755-1290	1
46.	Commanding Officer Fleet Numerical Oceanography Center Monterey, California 93943	1
47.	Commanding Officer Naval Environmental Prediction Research Facility Monterey, California 93943	1
48.	Defense Technical Information Center Cameron Station Alexandria, Virginia 22304~6145	2
49.	Commander Naval Oceanography Command NSTL Station Bay St. Louis, Mississippi 39529	1

50.	Commanding Officer Naval Ocean Research and Development Activity Attn: Technical Director NSTL Station Bay St. Louis, Mississippi 39529	1
51.	Commanding Officer Naval Polar Oceanography Center, Suitland Washington, District of Columbia 20373	1
52.	Director Naval Oceanography Division Naval Observatory 34th and Massachussetts Ave. NW Washington, District of Columbia 20390	1
53.	Commanding Officer Naval Oceanographic Command NSTL Station Bay St. Louis, Mississippi 39522	1
54.	Scott Polar Research Institute University of Cambridge Attn: Library Sea Ice Group Cambridge, ENGLAND CB2 1ER	1
55.	Chairman Department of Oceanography U.S. Naval Academy Annapolis, Maryland 21402	2
56.	Dr. James Morison Polar Science Center 4057 Roosevelt Way, NE Seattle, Washington 98105	1
57.	Dr. Kenneth Hunkins Lamont-Doherty Geological Observatory Palisades, New York 10964	1
58.	Dr. David Paskowsky, Chief Oceanography Branch U.S. Department of the Coast Guard Research and Development Center Avery Point, Connecticut 06340	1
59.	Science Applications, Inc. Attn: Dr. Robin Muench 13400B Northrup Way Suite 36 Bellevue, Washington 98005	1

ADDRESS ADDRES

20

200

.,

105

60.	Institute of Polar Studies Attn: Library 103 Mendenhall 125 South Oval Mall Columbus, Ohio 43210	1
61.	Institute of Marine Science University of Alaska Attn: Library Fairbanks, Alaska 99701	1
62.	Department of Oceanography University of British Columbia Attn: Library Vancouver, British Columbia CANADA V6T 1W5	1
63.	Institute of Marine Science University of Alaska Attn: Dr. H.J. Niebauer Fairbanks, Alaska 99701	1
64.	Bedford Institute of Oceanography Attn: Dr. P. Jones Library P.D. Box 1006 Dartmouth, Nova Scotia CANADA B27 4A2	1
65.	Canol Pease Pacific Marine Environmental Lab/NOAA 7600 Sand Point Way N.E. Seattle, Washington 98115	1
66.	Department of Oceanography Dalhousie University Halifax, Nova Scotia CANADA B3H 4J1	1
67.	Dr. Richard Armstrong MIZEX Data Manager National Snow and Ice Data Center Cooperative Institute for Research in Environmental Sciences Boulder, Colorado 80309	2

Statistics -

- 68. Institute of Ocean Sciences Attn: Dr. Eddy Carmack Dr. E. L. Lewis Dr. G. Holloway
 P.O. Box 6000 Sidney, British Columbia CANADA V8L 4B2
- 7C. Dr. Knut Aagaard NOAA/PMEL NOAA Bldg. #3 7600 Sand Point Way, N.E. Seattle, Washington 98115
- Department of Ocean Engineering Attn: Library Mass. Institute of Technology Cambridge, Massachusetts 02139
- 72. Dr. Theodore D. Foster Center for Coastal Marine Studies University of California Santa Cruz, California 95064
- 73. Research Administration
 Code 012
 Naval Postgraduate School
 Monterey, California 93943-5000
- 74. Dr. Hugh D. Livingston
 Woods Hole Oceanographic Institution
 Woods Hole, Massachusetts 02543
- 75. Dr. T.O. Manley Lamont-Doherty Geological Observatory Palisades, New York 10964
- 76. Dr. A. Foldvik Geophysical Institute University of Bergen Bergen, NORWAY
- 77. Dr. Preben Gudmandsen Electromagnetics Institute Technical University of Denmark Building 349 DK-2800 Lyngby, DENMARK
- 78. Dr. D. Hanzlick Flow Industries, Inc. Kent, Washington 98064

1

1

1

1

1

1

1

1

1

1

- 79. Dr. W.D. Hibler Thayer School of Engineering Dartmouth College Hanover, New Hamsphire 03755
- 80. Director General Fleet Systems Attn: Ivan Cote' Canadian Coast Guard Tower A, Place de Ville Ottawa, Ontario Canada K1A ON7
- 81. Mr. Tom Cocke Office of Marine Science U.S. Department of State Washington, D.C. 20525
- 82. Commanding Officer CCGS SIR JOHN FRANKLIN Canadian Coast Guard Base P.O. Box 1300 St. John's Newfoundland Canada A1C 6H8

Pression Contraction

- 83. M. Kim O. McCoy SeaMetrics
 9320 Chesapeake Dr., #212 San Diego, CA 92123
- 84. CDR Erik Thomsen Danish Liaison Office Thule AFB Thule, Greenland
- LT V. G. Addison, Jr.
 46 Division Avenue Massapequa, N.Y. 11758
- 86. LT A.M. Weigel 4332 Great Oak Drive Charleston, SC 29418
- 87. Research Administration Code 012 Naval Postgraduate School Monterey, CA 93943-5000

88. Superintendent Naval Postgraduate School Attn: Library (Code 0142) Monterey, California 93943-5002 1

1

1

1

1

1

5

1

1

END DATE FILMED DTIC 4/88