

AD-A199 418

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

TECHNICAL REPORT No. 83

Propagators for Driven Coupled Harmonic Oscillators

by

Kyu-Hwang Yeon, Chung-In Um, Woo-Hyung Kahng and Thomas F. George

Prepared for Publication

in

Physical Review A

Departments of Chemistry and Physics  
State University of New York at Buffalo  
Buffalo, New York 14260

September 1988

Reproduction in whole or in part is permitted for any purpose of the  
United States Government.

This document has been approved for public release and sale;  
its distribution is unlimited.

DTIC  
ELECTE  
SEP 29 1988  
S H D

88 9 28 00 3

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS			
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited			
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/88/TR-83		5. MONITORING ORGANIZATION REPORT NUMBER(S)			
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260		7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-86-K-0043		
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Propagators for Driven Coupled Harmonic Oscillators					
12. PERSONAL AUTHOR(S) Kyu-Hwang Yeon, Chung-In Um, Woo-Hyung Kahng and Thomas F. George					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) September 1988	15. PAGE COUNT 23
16. SUPPLEMENTARY NOTATION Prepared for publication in Physical Review A					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	PROPAGATORS	LASER DRIVEN	
			HARMONIC OSCILLATORS	ENERGY EXPECTATION	
			COUPLED	PATH INTEGRALS	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)  Propagators for coupled and driven coupled harmonic oscillators are evaluated exactly by the path-integral method. The propagators for coupled harmonic oscillators are used to obtain explicitly the energy expectation values.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410	22c. OFFICE SYMBOL	

**Propagators for driven coupled harmonic oscillators**

Kyu-Hwang Yeon  
Department of Physics  
Chungbuk National University  
Cheong Ju, Chung Buk 360-763, Korea

Chung-In Um and Woo-Hyung Kahng  
Department of Physics  
College of Science  
Korea University  
Seoul 136-701 Korea

Thomas F. George \*  
Departments of Chemistry and Physics & Astronomy  
239 Fronczak Hall  
State University of New York at Buffalo  
Buffalo, New York 14260

Propagators for coupled and driven coupled harmonic oscillators are evaluated exactly by the path-integral method. The propagators for coupled harmonic oscillators are used to obtain explicitly the energy expectation values.

PACS Nos. 02.90+p, 03.65.W, 03.65Db

\* To whom correspondence should be addressed.

1. Introduction

Although the Feynman path-integral formulation<sup>1</sup> offers a general approach for treating quantum-mechanical systems, only several problems can be solved exactly. Two of these are the driven harmonic oscillator with a quadratic Hamiltonian<sup>2</sup> and the time-dependent damped driven harmonic oscillator.<sup>3</sup> A number of situations such as superconducting quantum interference devices,<sup>4</sup> quantum nondemolition measurements,<sup>5</sup> magnetohydrodynamics,<sup>6</sup> etc., can be described by driven copuled harmonic oscillators. Introducing the Caldirola-Kanai Hamiltonian,<sup>7</sup> one can obtain the time-dependent Schroedinger equation for the damped harmonic oscillator. However, it has been a matter of debate as to whether or not this Schroedinger equation represents the quantum mechanical dissipative system.<sup>8</sup> Some workers<sup>9</sup> claim affirmation while others<sup>10</sup> object to it. This problem has been critically reviewed by Greenberger<sup>11</sup> and Cervero and Villaroel.<sup>12</sup>

The purpose of this paper is to derive the propagator for a driven coupled harmonic oscillators (DCHO) system from our previous work<sup>13</sup> for both coupled and coupled driven harmonic oscillators by means of the path-integral method. We introduce two harmonic oscillators that are coupled together with another spring. We review the classical case and construct the form of the propagator for DCHO, respectively, in Secs. 2 and 3. Section 4 gives the exact derivation of the propagator for the coupled harmonic oscillators (CHO), and in Sec. 5 we evaluate the exact propagator for DCHO by using the results obtained in Sec. 4. The energy expectation values of CHO are evaluated in Sec. 6, and finally we give results and discussion in Sec. 7.

<input checked="" type="checkbox"/>
<input type="checkbox"/>
<input type="checkbox"/>



By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

## 2. Classical case

In this section we consider a system of two harmonic oscillators which are coupled together by means of another spring. We assume that the masses of the oscillators and three spring constants are all the same. Let the forces  $f_1(t)$  and  $f_2(t)$  exerted on the two oscillators and their displacements be  $x_1$  and  $x_2$ . Then the Hamiltonian for DCHO can be written as

$$H = \frac{1}{2m} (p_1^2 + p_2^2) + m\omega^2 (x_1^2 - x_1 x_2 + x_2^2) - f_1(t)x_1 - f_2(t)x_2 \quad (2.1)$$

where  $\omega^2 = k/m$ . Hamilton's equations of motion for Eq. (2.1) are

$$\dot{x}_1 = p_1/m \quad (2.2)$$

$$\dot{x}_2 = p_2/m \quad (2.3)$$

$$\dot{p}_1 = m\omega^2 (x_2 - 2x_1) + f_1(t) \quad (2.4)$$

$$\dot{p}_2 = m\omega^2 (x_1 - 2x_2) + f_2(t) \quad (2.5)$$

Equations (2.1)-(2.5) yield the Lagrangian,

$$\begin{aligned} L &= (p_1 \dot{x}_1 + p_2 \dot{x}_2) - H \\ &= \frac{m}{2} (\dot{x}_1^2 + \dot{x}_2^2) - m\omega^2 (x_1^2 - x_1 x_2 + x_2^2) + f_1(t)x_1 + f_2(t)x_2 \quad (2.6) \end{aligned}$$

with the corresponding equations of motion

$$x_1 + \omega^2(2x_1 - x_2) = f_1(t)/m \quad (2.7)$$

$$x_2 + \omega^2(2x_2 - x_1) = f_2(t)/m \quad (2.8)$$

The classical solutions of Eqs. (2.7) and (2.8) are given by

$$x_1(t) = A \sin(\omega t) + B \cos(\omega t) + C \sin(\sqrt{3}\omega t) + D \cos(\sqrt{3}\omega t) \\ + \int^t dr \int^r d\nu e^{i\omega(2r-\nu-t)} [f_1(\nu) + f_2(\nu)] \quad (2.9)$$

and

$$x_2(t) = A \sin(\omega t) + B \cos(\omega t) - C \sin(\sqrt{3}\omega t) - D \cos(\sqrt{3}\omega t) \\ + \int^t dr \int^r d\nu e^{i\omega(2r-\nu-t)} [f_1(\nu) - f_2(\nu)] \quad (2.10)$$

### 3. Path integral of driven coupled harmonic oscillators

In the path-integral formulation, the solution of the Schroedinger equation is given as the path-dependent integral equations with propagator  $K$ ,

$$\psi(x_1, x_2, t) = \int dx'_1 dx'_2 K(x_1, x_2, t; x'_1, x'_2, 0) \psi(x'_1, x'_2, 0) \quad (3.1)$$

which gives the wavefunction  $\psi(x_1, x_2, t)$  at time  $t$  in terms of the wave function  $\psi(x'_1, x'_2)$  at time  $t = 0$ . The propagator in Eq. (3.1) can be written by means of the Feynman path integral

$$K(x_1, x_2, t; x'_1, x'_2, 0) = \int_{(x'_1, x'_2, 0)}^{(x_1, x_2, t)} Dx(t) \exp[(i/\hbar)S(x_1, x_2, x'_1, x'_2; t)] \quad (3.2)$$

where

$$Dx(t) = \lim_{N \rightarrow \infty} \frac{1}{A} \prod_{j=1}^{N-1} [dx_{1j} dx_{2j} / A^2] \quad , \quad (3.3)$$

and  $S(x_1, x_2, x'_1, x'_2; t)$  is the action defined as the time integral over the Lagrangian  $L(\dot{x}_1, \dot{x}_2, x_1, x_2; t)$  between  $t = t$  and  $t = 0$ :<sup>1</sup>

$$S(x_1, x_2, x'_1, x'_2; t) = \int_0^t dt L(\dot{x}_1, \dot{x}_2, x_1, x_2; t) \quad . \quad (3.4)$$

In Eq. (3.3)  $A$  is the normalization factor given by

$$A = [2\pi i \hbar \epsilon / m]^{1/2} \quad , \quad \epsilon = \lim_{N \rightarrow \infty} (t/N) \quad . \quad (3.5)$$

Substituting Eq. (2.6) into Eq. (3.4), the action becomes

$$S(x_1, x_2, x'_1, x'_2; t) = S_c(x_1, x_2, x'_1, x'_2; t) + \int_0^t d\tau \frac{m}{2} (\dot{y}_1^2(\tau) + \dot{y}_2^2(\tau) - 2\omega^2 [y_1^2(\tau) - y_1(\tau)y_2(\tau) + y_2^2(\tau)]) \quad , \quad (3.6)$$

where  $S_c$  is the classical action and  $y_i$  is the deviation of  $x_i(t)$  from its classical path  $x_{ci}$  given as

$$y_i = x_i - x_{ci} \quad (i = 1, 2) \quad (3.7)$$

Then the propagator [Eq. (3.2)] can be expressed as

$$K(x_1, x_2, t; x'_1, x'_2, 0) = F(t) e^{iS_c/\hbar} \quad (3.8)$$

Here,  $F(t)$  is the multiplicative function given in the form

$$F(t) = \int_0^0 Dx(t) \left( \exp \left[ (im/2\hbar) \int_0^t dt (\dot{y}_1^2 + \dot{y}_2^2 - 2\omega^2(y_1^2 - y_1 y_2 + y_2^2)) \right] \right) \quad (3.9)$$

It is easy to show that  $F(t)$  has the same form for CHO and DCHO. Therefore, the propagator depends only on the classical action in both cases. In Eq. (3.9), change the variables  $x_1 \pm x_2$  into

$$z_1 = \frac{1}{\sqrt{2}} (x_1 - x_2) \quad (3.10)$$

$$z_2 = \frac{1}{\sqrt{2}} (x_1 + x_2) \quad (3.11)$$

we can reduce the condition  $(y_1, y_2) = (0, 0)$  to  $(z_1, z_2) = (0, 0)$ . Applying Eqs. (3.10) and (3.11) to Eq. (3.9), the multiplicative function becomes

$$F(t) = \int_0^0 Dz(t) \left( \exp \left[ (im/2\hbar) \int_0^t Dz(t) [(\dot{z}_1^2 - \omega^2 z_1^2) + (\dot{z}_2^2 - 3\omega^2 z_2^2)] \right] \right) \quad (3.12)$$



$$\begin{bmatrix} y_{11} \\ y_{12} \\ y_{21} \\ \cdot \\ \cdot \\ y_{N-1,2} \\ y_{N1} \\ y_{N2} \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 & \cdot & \cdot & \cdot & \cdot & 0 & 0 & 0 \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 & \cdot & \cdot & \cdot & \cdot & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{\sqrt{2}} & \cdot & \cdot & \cdot & \cdot & 0 & 0 & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & 0 & \cdot & \cdot & \cdot & \cdot & \frac{1}{\sqrt{2}} & 0 & 0 \\ 0 & 0 & 0 & \cdot & \cdot & \cdot & \cdot & 0 & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ 0 & 0 & 0 & \cdot & \cdot & \cdot & \cdot & 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} z_{11} \\ z_{12} \\ z_{21} \\ \cdot \\ \cdot \\ z_{N-1,2} \\ z_{N1} \\ z_{N2} \end{bmatrix} \quad (3.13)$$

In Eq. (3.12),  $J$  becomes unity.

If the action is separated into the functionals with only same variables in the path integral, then this integral can be represented by the multiplication of path integrals with each variable. Therefore, Eq. (3.12) becomes

$$F(t) = F_1(t) F_2(t)$$

$$\begin{aligned}
& - \left( \int_0^t Dz_1(\tau) \exp\left[\frac{im}{2\hbar} \int_0^\tau dt (\dot{z}_1^2 - \omega^2 z_1^2)\right] \right) \\
& \times \left( \int_0^t Dz_2(\tau) \exp\left[\frac{im}{2\hbar} \int_0^\tau dt (\dot{z}_2^2 - 3\omega^2 z_2^2)\right] \right) . \quad (3.14)
\end{aligned}$$

Since  $F_1(t)$  and  $F_2(t)$  are the path integrals of the harmonic oscillator, the evaluation of Eq. (3.14) gives

$$F(t) = \frac{m\omega}{2\pi i\hbar} \left[ \frac{\sqrt{3}}{\sin(\omega t) \sin(\sqrt{3}\omega t)} \right]^{1/2} . \quad (3.15)$$

Hence, the propagator of DCHO can be written as

$$K(x_1, x_2, t; x'_1, x'_2, 0) = \frac{m\omega}{2\pi i\hbar} \left[ \frac{\sqrt{3}}{\sin(\omega t) \sin(\sqrt{3}\omega t)} \right]^{1/2} e^{iS_c/\hbar} . \quad (3.16)$$

#### 4. Propagator for the coupled harmonic oscillators

To evaluate the exact propagator expressed by Eq. (3.16), we should first obtain the propagator for CHO. The classical action of CHO is

$$S_c = \int_0^t dt \left( \frac{m}{2} (\dot{x}_{c1}^2 + \dot{x}_{c2}^2) - m\omega^2 (x_{c1}^2 - x_{c1}x_{c2} + x_{c2}^2) \right) , \quad (4.1)$$

where  $x_{c1}$  and  $\dot{x}_{c1}$  are the classical path and velocity, respectively.

Integrating Eq. (4.1) over the time, we get

$$S_c = \frac{m}{2} (x_{c1}\dot{x}_{c1} + x_{c2}\dot{x}_{c2}) \Big|_0^t - \int_0^t dt \frac{m}{2} x_{c1} (\ddot{x}_{c1} + \omega^2 (2x_{c1} - x_{c2}))$$

$$\begin{aligned}
& - \int_0^t dt \frac{m}{2} x_{c2} (\ddot{x}_{c2} + \omega^2 (2x_{c2} - x_{c1})) \\
& - \frac{m}{2} [x_{c1}(t)\dot{x}_{c1}(t) + x_{c2}(t)\dot{x}_{c2}(t) - x_{c1}(0)\dot{x}_{c1}(0) - x_{c2}(0)\dot{x}_{c2}(0)] \quad (4.2)
\end{aligned}$$

Here the second and third terms become zero because of the equations of motion [see Eqs. (2.7) and (2.8)], given as

$$\ddot{x}_1 + \omega^2 (2x_1 - x_2) = 0 \quad (4.3)$$

$$\ddot{x}_2 + \omega^2 (2x_2 - x_1) = 0 \quad (4.4)$$

To obtain the exact expression of Eq. (4.2), we solve Eqs. (4.3) and (4.4) to obtain

$$x_1 = x_1(t) = A \sin(\omega t) + B \cos(\omega t) + C \sin(\sqrt{3}\omega t) + D \cos(\sqrt{3}\omega t) \quad (4.5)$$

$$x_2 = x_2(t) = A \sin(\omega t) + B \cos(\omega t) - C \sin(\sqrt{3}\omega t) - D \cos(\sqrt{3}\omega t) \quad (4.6)$$

and  $\dot{x}_1$  and  $\dot{x}_2$  are given, respectively, by

$$\begin{aligned}
\dot{x}_1 = \dot{x}_1(t) &= \omega(A \cos(\omega t) - B \sin(\omega t) + \sqrt{3}C \sin(\sqrt{3}\omega t) \\
& - \sqrt{3}D \sin(\sqrt{3}\omega t)) \quad (4.7)
\end{aligned}$$

$$\begin{aligned}
\dot{x}_2 = \dot{x}_2(t) &= \omega(A \cos(\omega t) - B \sin(\omega t) - \sqrt{3}C \cos(\sqrt{3}\omega t) \\
& + \sqrt{3}D \sin(\sqrt{3}\omega t)) \quad (4.8)
\end{aligned}$$

Equations (4.5)-(4.8) give

$$x'_1 - x_1(0) = B + D \quad (4.9)$$

$$x'_2 - x_2(0) = B - D \quad (4.10)$$

$$\dot{x}'_1 - \dot{x}_1(0) = \omega(A + \sqrt{3}C) \quad (4.11)$$

$$\dot{x}'_2 - \dot{x}_2(0) = \omega(A - \sqrt{3}C) \quad (4.12)$$

The time-dependent constants A, B, D and D obtained from Eqs. (4.5) and (4.6), and Eqs. (4.9) and (4.10) can be expressed as

$$A = \left[ \frac{1}{2} \sin(\omega t) \right] (x_1 + x_2 - (x'_1 + x'_2) \cos(\omega t)) \quad (4.13)$$

$$B = \frac{1}{2} (x'_1 + x'_2) \quad (4.14)$$

$$C = \left[ \frac{1}{2} \sin(\sqrt{3}\omega t) \right] (x_1 - x_2 + (x'_1 - x'_2) \cos(\sqrt{3}\omega t)) \quad (4.15)$$

$$D = \frac{1}{2} (x'_1 - x'_2) \quad (4.16)$$

Substitution of Eqs. (4.5) -(4.16) into (4.2) gives the classical action:

$$S_c = \frac{m\omega}{4} ((x_1^2 + x_2^2 + x_1'^2 + x_2'^2) [\cot(\omega t) + \sqrt{3} \cot(\sqrt{3}\omega t)]) \\ + 2(x_1 x_2 + x_1' x_2') [\cot(\omega t) - \sqrt{3} \cot(\sqrt{3}\omega t)]$$

$$\begin{aligned}
& - 2(x_1 x'_1 + x_2 x'_2) \left( \frac{1}{\sin(\omega t)} + \frac{\sqrt{3}}{\sin(\sqrt{3}\omega t)} \right) \\
& + 2(x_1 x'_2 + x_2 x'_1) \left[ -\frac{1}{\sin(\omega t)} + \frac{\sqrt{3}}{\sin(\sqrt{3}\omega t)} \right] .
\end{aligned} \tag{4.17}$$

Combining Eqs. (4.17) and (3.16), we obtain the propagator for CHO:

$$\begin{aligned}
K(x_1, x_2, t; x'_1, x'_2, 0) &= \frac{im\omega}{2\pi i\hbar} \left[ \frac{\sqrt{3}}{\sin(\omega t)} \sin(\sqrt{3}\omega t) \right]^{\frac{1}{2}} \\
&\times \exp\left( \frac{im\omega}{4\hbar} \left[ (x_1^2 + x_2^2 + x_1'^2 + x_2'^2) [\cot(\omega t) + \sqrt{3} \cot(\sqrt{3}\omega t)] \right. \right. \\
&+ 2(x_1 x_2 + x_1' x_2') [\cot(\omega t) - \sqrt{3} \cot(\sqrt{3}\omega t)] - 2(x_1 x'_1 + x_2 x'_2) \left. \left. \left[ \frac{1}{\sin(\omega t)} \right. \right. \right. \\
&\left. \left. \left. + \frac{\sqrt{3}}{\sin(\sqrt{3}\omega t)} \right] + 2(x_1 x'_2 + x_2 x'_1) \left[ -\frac{1}{\sin(\omega t)} + \frac{\sqrt{3}}{\sin(\sqrt{3}\omega t)} \right] \right) \right] .
\end{aligned} \tag{4.18}$$

##### 5. Propagator for driven coupled harmonic oscillators

When we set  $f_1(t) = f_2(t) = 0$ , DCHO reduces to CHO, whereby we can write the propagator for DCHO as

$$\begin{aligned}
K(x_1, x_2, t; x'_1, x'_2, 0) &= \exp[a(t)x_1^2 + b(t)x_1 x_2 + c(t)x_2^2 + d(t)x_1 \\
&+ g(t)x_2 + h(t)] .
\end{aligned} \tag{5.1}$$

Here  $a(t)$ ,  $b(t)$ ,  $c(t)$ ,  $d(t)$ ,  $f(t)$  and  $h(t)$  are time-dependent functions including  $x'_1$  and  $x'_2$ , which need to be determined. Equation (5.1) must satisfy the Schroedinger equation

$$i\hbar(\partial K/\partial t) = H K \quad (5.2)$$

Substitution of Eq. (5.1) into Eq. (5.2) gives the time-dependent coefficients

$$a(t) = \frac{i\hbar}{2m} [4a^2(t) + c^2(t)] + m\omega^2/i\hbar \quad (5.3)$$

$$b(t) = \frac{i\hbar}{2m} [4b^2(t) + c^2(t)] + m\omega^2/i\hbar \quad (5.4)$$

$$c(t) = \frac{2i\hbar}{m} [a(t)c(t) + b(t)c(t)] - m\omega^2/i\hbar \quad (5.5)$$

$$d(t) = \frac{i\hbar}{m} [2a(t)d(t) + c(t)g(t)] + \left(\frac{i}{\hbar}\right)f_1(t) \quad (5.6)$$

$$g(t) = \frac{i\hbar}{m} [2b(t)g(t) + c(t)d(t)] + \left(\frac{i}{\hbar}\right)f_2(t) \quad (5.7)$$

$$h(t) = \frac{i\hbar}{2m} [d^2(t) + g^2(t) + 2a(t) + 2b(t)] \quad (5.8)$$

Since Eqs. (5.3) and (5.4) have the same form, we get

$$a(t) = b(t) \quad (5.9)$$

Substituting Eq. (5.9) into Eq. (5.5) and changing the variables  $a$  and  $c$  into

$$\eta = a + c/2 \quad , \quad (5.10)$$

$$\zeta = a - c/2 \quad , \quad (5.11)$$

we obtain two ordinary differential equations:

$$\dot{\eta} - \frac{2i\mathcal{H}}{m} \eta^2 + \frac{m\omega^2}{2i\mathcal{H}} \quad (5.12)$$

$$\dot{\zeta} - \frac{2i\mathcal{H}}{m} \zeta^2 + \frac{3m\omega^2}{2i\mathcal{H}} \quad (5.13)$$

The solutions of Eqs. (5.12) and (5.13) are given by

$$\eta = \frac{i\omega m}{2\mathcal{H}} \cot(\omega t + \theta_1) \quad (5.14)$$

$$\zeta = \frac{\sqrt{3}i\omega m}{2\mathcal{H}} \cot(\sqrt{3}\omega t + \theta_2) \quad (5.15)$$

where  $\theta_1$  and  $\theta_2$  are the constants to be determined. The time-dependent coefficients  $a(t)$ ,  $b(t)$  and  $c(t)$  obtained in comparison with Eqs. (5.10), (5.11), (5.14) and (5.15) are given as

$$a(t) - b(t) = \frac{i\omega m}{4\mathcal{H}} [\cot(\omega t + \theta_1) + \sqrt{3} \cot(\sqrt{3}\omega t + \theta_2)] \quad (5.16)$$

$$c(t) = \frac{i\omega m}{2\mathcal{H}} [\cot(\omega t + \theta_1) - \sqrt{3} \cot(\sqrt{3}\omega t + \theta_2)] \quad (5.17)$$

Equations (5.16) and (5.17) do not include the driven forces  $f_1(t)$  and  $f_2(t)$ . Therefore, through setting  $f_1(t) = f_2(t) = 0$ , Eqs. (5.16) and (5.17) do not change at all and should be equal to the coefficients of  $x_1^2$  and  $x_2^2$  in Eq. (4.18). Comparison of these two equations shows  $\theta_1$  and  $\theta_2$  to be zero. Substituting Eq. (5.9) into Eqs. (5.6) and (5.7) and changing variables  $d$  and  $g$  into

$$\rho = d + g \quad (5.18)$$

$$\sigma = d - g \quad (5.19)$$

we obtain the two differential equations

$$\dot{\rho} = \frac{iM}{m} [2a(t) + c(t)] \rho + \frac{i}{M} [f_1(t) + f_2(t)] \quad (5.20)$$

$$\dot{\sigma} = \frac{iM}{m} [2a(t) + c(t)] \sigma + \frac{i}{M} [f_1(t) - f_2(t)] \quad (5.21)$$

Combining Eqs. (5.20) and (5.21) with Eqs. (5.16) and (5.17), we obtain the solutions

$$\rho = [1/\sin(\omega t)] \left( \int_0^t dr \frac{i}{M} [f_1(r) + f_2(r)] \sin(\omega r) + \alpha \right) \quad (5.22)$$

$$\sigma = [1/\sin(\sqrt{3}\omega t)] \left( \int_0^t dr \frac{i}{M} [f_1(r) - f_2(r)] \sin(\sqrt{3}\omega r) + \beta \right) \quad (5.23)$$

where  $\alpha$  and  $\beta$  are constants to be determined. We can obtain the time-dependent coefficients  $d(t)$  and  $g(t)$  by substituting Eqs. (5.22) and (5.23) into Eqs. (5.18) and (5.19):

$$\begin{aligned} d(t) = & [1/2M\sin(\omega t)] \int_0^t dr [f_1(r) + f_2(r)] \sin(\omega r) \\ & + [1/2M\sin(\sqrt{3}\omega t)] \int_0^t dr [f_1(r) - f_2(r)] \sin(\sqrt{3}\omega r) \\ & + [\alpha/2\sin(\omega t)] + [\beta/2\sin(\sqrt{3}\omega t)] \end{aligned} \quad (5.24)$$



$$\begin{aligned}
g(t) &= \left[ \frac{iK}{2} \sin(\omega t) \right] \int_0^t dr [f_1(r) + f_2(r)] \sin(\omega r) \\
&- \left[ \frac{iK}{2} \sin(\sqrt{3}\omega t) \right] \int_0^t dr [f_1(r) - f_2(r)] \sin(\sqrt{3}\omega r) \\
&+ \left[ \frac{\alpha}{2} \sin(\omega t) \right] - \left[ \frac{\beta}{2} \sin(\sqrt{3}\omega t) \right] \quad . \quad (5.25)
\end{aligned}$$

Substitution of Eqs. (5.16), (5.17), (5.24) and (5.25) into Eq. (5.8) yields

$$\begin{aligned}
h(t) &= - \frac{iK}{4m\omega} \left[ \alpha^2 \cot(\omega t) + (\beta^2/\sqrt{3}) \cot(\sqrt{3}\omega t) \right] \\
&- \left[ \frac{\alpha}{m\omega} \sin(\omega t) \right] \int_0^t dr [f_1(r) + f_2(r)] \sin[\omega(t-r)] \\
&- \left[ \frac{\beta/\sqrt{3}}{m\omega} \sin(\sqrt{3}\omega t) \right] \int_0^t dr [f_1(r) - f_2(r)] \sin[\sqrt{3}\omega(t-r)] \\
&+ \left[ \frac{1}{4iK} \frac{1}{m\omega} \sin(\omega t) \right] \int_0^t dr \int_0^t d\nu [f_1(r) + f_2(r)] \\
&\times [f_1(\nu) + f_2(\nu)] \sin[\omega(t-r)] \sin(\omega\nu) \\
&+ \left[ \frac{1}{4\sqrt{3}iK} \frac{1}{m\omega} \sin(\sqrt{3}\omega t) \right] \int_0^t dr \int_0^t d\nu [f_1(r) - f_2(r)] \\
&\times [f_1(\nu) - f_2(\nu)] \sin[\sqrt{3}\omega(t-r)] \sin(\sqrt{3}\omega\nu) - \ln[\sin(\omega t) \\
&\times \sin(\sqrt{3}\omega t) + \delta \quad . \quad (5.26)
\end{aligned}$$

Here,  $\delta$  is also a constant to be determined. When setting  $f_1(t) = f_2(t) = 0$ , Eqs. (5.24) and (5.25) should be reduced to the coefficients of  $x_1$  and

$x_2$ , and Eq. (5.26) should also be reduced to the terms in the exponent in Eq. (4.18). Comparison between them gives the constants  $\alpha$ ,  $\beta$  and  $\delta$ :

$$\alpha = \frac{m\omega}{i\hbar} (x'_1 + x'_2) \quad , \quad (5.27)$$

$$\beta = \frac{m\omega}{i\hbar} (x'_1 - x'_2) \quad , \quad (5.28)$$

$$\delta = \ln\left(\frac{3^{\frac{1}{2}} m\omega}{2\pi i\hbar}\right) \quad . \quad (5.29)$$

Substitution of the above results into Eq. (5.1) gives the propagator for DCHO:

$$\begin{aligned} K(x_1, x_2, t; x'_1, x'_2, 0) &= \frac{m\omega}{2\pi i\hbar} \left\{ \sqrt{3} / [\sin(\omega t) \sin(\sqrt{3}\omega t)] \right\}^{\frac{1}{2}} \\ &\times \exp \left\{ \frac{i m \omega}{4\hbar} \left[ (x_1^2 + x_2^2 + x_1'^2 + x_2'^2) [\cot(\omega t) + \sqrt{3} \cot(\sqrt{3}\omega t)] \right. \right. \\ &+ 2(x_1 x_2 + x_1' x_2') [\cot(\omega t) - \sqrt{3} \cot(\sqrt{3}\omega t)] - 2(x_1 x_1' + x_2 x_2') \{ 1/\sin(\omega t) \\ &+ \sqrt{3}/\sin(\sqrt{3}\omega t) \} \\ &+ 2(x_1 x_2' + x_1' x_2) \{ -1/\sin(\omega t) + \sqrt{3}/\sin(\sqrt{3}\omega t) \} \\ &+ \frac{2x_1}{m\omega} \left( [1/\sin(\omega t)] \int_0^t dr [f_1(r) + f_2(r)] \sin(\omega r) \right. \\ &+ \left. [1/\sin(\sqrt{3}\omega t)] \int_0^t dr [f_1(r) - f_2(r)] \sin(\sqrt{3}\omega r) \right) \end{aligned}$$

$$\begin{aligned}
& + \frac{2x_2}{m\omega} \left( [1/\sin(\omega t)] \int_0^t dr [f_1(r) + f_2(r)] \sin(\omega r) \right. \\
& - [1/\sin(\sqrt{3}\omega t)] \int_0^t dr [f_1(r) - f_2(r)] \sin(\sqrt{3}\omega r) \left. \right) \\
& + \frac{4x'_1}{m\omega} \left( [1/\sin(\omega t)] \int_0^t dr [f_1(r) + f_2(r)] \sin[\omega(t-r)] \right. \\
& + [1/\sin(\sqrt{3}\omega t)] \int_0^t dr [f_1(r) - f_2(r)] \sin(\sqrt{3}\omega r) \left. \right) \\
& + \frac{4x'_2}{m\omega} \left( [1/\sin(\omega t)] \int_0^t dr [f_1(r) + f_2(r)] \sin[\omega(t-r)] \right. \\
& - [1/\sin(\sqrt{3}\omega t)] \int_0^t dr (f_1(r) - f_2(r)) \sin(\sqrt{3}\omega r) \left. \right) \\
& - [1/m^2\omega^2\sin(\omega t)] \int_0^t dr \int_0^t d\nu [f_1(r) + f_2(r)][f_1(\nu) + f_2(\nu)] \\
& \times \sin[\omega(t-r)] \sin(\omega\nu) \\
& - [1/\sqrt{3}m^2\omega^2\sin(\sqrt{3}\omega t)] \int_0^t dr \int_0^t d\nu [f_1(r) - f_2(r)][f_1(\nu) - f_2(\nu)] \\
& \times \sin[\sqrt{3}\omega(t-r)] \sin(\sqrt{3}\omega\nu) \left. \right\} . \tag{5.30}
\end{aligned}$$

## 6. Energy expectation values of coupled harmonic oscillators

The Hamiltonian of CHO is

$$H = \frac{1}{2m} (p_1^2 + p_2^2) + m\omega^2 (x_1^2 - x_1 x_2 + x_2^2) \quad (6.1)$$

Using Eqs. (3.1) and (3.2) with Eq. (6.1), we obtain the Schroedinger equation,

$$i\hbar(\partial/\partial t) \psi(x_1, x_2, t) = H_{op} \psi(x_1, x_2, t) \quad (6.2)$$

where  $H_{op}$  is the Hamiltonian operator in which the momentum  $p_i$  is changed into  $p_i = (\hbar/i)(\partial/\partial x_i)$ . Since Eq. (6.2) can be separated into time and coordinate parts, we may write

$$K(t) = e^{-iH_{op}t/\hbar} \quad (6.3)$$

$$H_{op} |\ell, n\rangle = E_{\ell n} |\ell, n\rangle \quad (\ell, n = 1, 2, 3, \dots) \quad (6.4)$$

Here the states  $|\ell, n\rangle$  are the complete set with energy eigenvalues of  $H_{op}$ . Since the function with states  $|\ell, n\rangle$  can be expressed by

$$\phi_{\ell n}(x_1, x_2) = \langle x_1, x_2 | \ell, n \rangle \quad (6.5)$$

the propagator at  $t > 0$  becomes

$$K(x_1, x_2, t; x'_1, x'_2, 0) = \langle x_1, x_2 | e^{-iH_{op}t/\hbar} | x'_1, x'_2 \rangle$$

$$\begin{aligned}
& - \sum_{\ell} \sum_n \sum_{\ell'} \sum_{n'} \langle x_1, x_2 | \ell n \rangle \langle \ell n | e^{-iH_{op}t/\hbar} | \ell', n' \rangle \langle \ell', n' | x'_1, x'_2 \rangle \\
& - \sum_{\ell} \sum_n \phi_{\ell n}(x_1, x_2) e^{-iE_{\ell n}t/\hbar} \phi_{\ell n}^*(x'_1, x'_2) \quad . \quad (6.6)
\end{aligned}$$

Equation (6.6) should be the same as Eq. (4.18). Setting  $x'_1 = x_1$  and  $x'_2 = x_2$  in Eq. (4.18) and integrating over  $x_1$  and  $x_2$ , we get

$$\sum_{\ell} \sum_n \iint dx_1 dx_2 \phi_{\ell n}^*(x_1, x_2) e^{-iE_{\ell n}t/\hbar} \phi_{\ell n}(x_1, x_2) = e^{-iE_{\ell n}t/\hbar} \quad (6.7)$$

and

$$\begin{aligned}
& \iint dx_1 dx_2 \frac{m\omega}{2\pi i\hbar} \left\{ \sqrt{3} / [\sin(\omega t) \sin(\sqrt{3}\omega t)] \right\}^4 \\
& \times \exp\left\{ \frac{i m \omega}{2\hbar} [(x_1 + x_2)^2 - \sqrt{3}(x_1 - x_2)^2] [\cot(\omega t) - 1/\sin(\omega t)] \right\} \\
& = - \frac{1}{2} [\sin(\omega t/2) \sin(\sqrt{3}/2\omega t)]^{-1} \quad . \quad (6.8)
\end{aligned}$$

Hence, we have

$$\begin{aligned}
\sum_{\ell} \sum_n e^{-iE_{\ell n}t/\hbar} & = - \frac{1}{2} [\sin(\omega t/2) \sin(\sqrt{3}/2\omega t)]^{-1} \\
& = [e^{-i\omega t/2} / (1 - e^{-i\omega t})] [e^{-i\sqrt{3}\omega t/2} / (1 - e^{-i\sqrt{3}\omega t})]
\end{aligned}$$

$$- \sum_{\ell=0}^{\infty} \sum_{n=0}^{\infty} \exp(-i\omega t[(\ell + \frac{1}{2}) + \sqrt{3}(n + \frac{1}{2})]) \quad (6.9)$$

Therefore the expectation values of CHO becomes

$$E_{\ell n} = [(\ell + \frac{1}{2}) + \sqrt{3}(n + \frac{1}{2})] \hbar\omega \quad (6.10)$$

## 7. Results and discussion

In the previous sections we have obtained the exact propagators [Eqs. (4.18) and (5.30)] for CHO and DCHO by the path-integral method. The forms of the propagators are new. Setting  $f(t) = 0$ , Eq. (5.30) is reduced to Eq. (4.18). Although DCHO is a nonconservative system, the quantum-mechanical problem for the momentum operator does not appear because the canonical momentum is equal to the kinetic momentum in our derivation.<sup>13</sup>

Making use of Eq. (4.18), we have obtained the energy expectation values [Eq. (6.10)] for CHO, given by the sum of two energy expectation values corresponding to the quantum states of two oscillators. Even though we have not evaluated the wavefunction of CHO, we may easily surmise that the wavefunction will be given by the multiplication of two wavefunctions for two oscillators. In the case of DCHO, one cannot easily apply Eq. (5.20) to obtain the energy expectation values, since this equation cannot be expressed in the form of Eq. (6.6), and one should recognize that the energy operator is not equal to the Hamiltonian operator in a nonconservative system.<sup>9</sup>

The evaluations for the wavefunctions, energy expectation values for CHO and DCHO, and propagator and other physical quantities for  $n$  coupled and

n driven coupled harmonic oscillators (arbitrary n) are in progress and will be reported in the near future.

#### **Acknowledgments**

This research was supported in part by a grant to Korea University from the BSRI Program, Ministry of Education 1987, Republic of Korea, and in part by the Office of Naval Research, the Air Force Office of Scientific Research (AFSC), under Contract F49620-86-C-0009, and the National Science Foundation under Grant CHE-8620274. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

**References**

1. R. P. Feynman and A. R. Hibbs, "Quantum Mechanics and Path Integrals" (McGraw-Hill, New York, 1965).
2. J. T. Marshall and J. L. Pell, J. Math. Phys. 20, 1297 (1979); G. E. Prince and C. J. Eliezer, J. Phys. A 13, 815 (1980).
3. L. F. Landovitz, A. M. Levine and W. M. Schreiber, Phys. Rev. A 20, 1162 (1979); J. Math. Phys. 21, 2159 (1980); A. M. Levine, E. Ozizmir and W. M. Schreiber, J. Chem. Phys. 78, 291 (1983).
4. A. C. Rose-Innes and E. H. Rhoderick, "Introduction to Superconductivity" (Pergamon, Oxford, 1978).
5. V. B. Braginsky and Yu. I. Vorontsov, Usp. Fiz. Nauk 114, 41 (1974); W. G. Unruh, Phys. Rev. D 18, 1764 (1978); 19, 2888 (1979).
6. R. King, B. A. Huberman and J. Barchas, in "Synergetics of the Brain," ed. by H. Haken (Springer, Berlin, 1983).
7. P. Caldirola, Nuovo Cimento 18, 393 (1941); B 77, 241 (1983); E. Kanai, Prog. Theor. Phys. 3, 440 (1948).
8. H. Dekker, Phys. Rep. 80, 1 (1981); B. K. Cheng, J. Phys. A: Math. Gen. 17, 2475 (1984); K. H. Yeon, C. I. Um and T. F. George, Phys. Rev. A 36, 5287 (1987); E. Eckhardt, Phys. Rev. A 35, 5191 (1987).
9. V. V. Dodonov and V. I. Manko, Phys. Rev. A 20, 550 (1979).
10. I. R. Senitzky, Phys. Rev. 119, 670 (1960).
11. D. M. Greenberger, J. Math. Phys. 20, 762 (1979).
12. J. J. Cervero and J. Villaroel, J. Phys. A: Math. Gen. 17, 2963 (1984).
13. C. I. Um, K. H. Yeon and W. H. Kahng, J. Phys. A: Math. Gen. 20, 611 (1987); J. Korean Phys. Soc. 19, 1, 8 (1986); New Phys. (Korean Phys. Soc.) 26, 100, 117 (1985).



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 M. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Duda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. J. H. Weaver  
Department of Chemical Engineering  
and Materials Science  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. A. Reisman  
Microelectronics Center of North Carolina  
Research Triangle Park, North Carolina  
27709

Dr. M. Grunze  
Laboratory for Surface Science and  
Technology  
University of Maine  
Orono, Maine 04469

Dr. J. Butler  
Naval Research Laboratory  
Code 6115  
Washington D.C. 20375-5000

Dr. L. Interante  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Irvin Heard  
Chemistry and Physics Department  
Lincoln University  
Lincoln University, Pennsylvania 19352

Dr. K.J. Klaubunde  
Department of Chemistry  
Kansas State University  
Manhattan, Kansas 66506

Dr. C. B. Harris  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. F. Kutzler  
Department of Chemistry  
Box 5055  
Tennessee Technological University  
Cookeville, Tennessee 38501

Dr. D. DiLella  
Chemistry Department  
George Washington University  
Washington D.C. 20052

Dr. R. Reeves  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Steven M. George  
Stanford University  
Department of Chemistry  
Stanford, CA 94305

Dr. Mark Johnson  
Yale University  
Department of Chemistry  
New Haven, CT 06511-8118

Dr. W. Knauer  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. J. Murday  
Naval Research Laboratory  
Code 6170  
Washington, D.C. 20375-5000

Dr. J. B. Hudson  
Materials Division  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Theodore E. Madey  
Surface Chemistry Section  
Department of Commerce  
National Bureau of Standards  
Washington, D.C. 20234

Dr. J. E. Demuth  
IBM Corporation  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. M. G. Lagally  
Department of Metallurgical  
and Mining Engineering  
University of Wisconsin  
Madison, Wisconsin 53706

Dr. R. P. Van Duyne  
Chemistry Department  
Northwestern University  
Evanston, Illinois 60637

Dr. J. M. White  
Department of Chemistry  
University of Texas  
Austin, Texas 78712

Dr. D. E. Harrison  
Department of Physics  
Naval Postgraduate School  
Monterey, California 93940

Dr. R. L. Park  
Director, Center of Materials  
Research  
University of Maryland  
College Park, Maryland 20742

Dr. W. T. Peria  
Electrical Engineering Department  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson  
Department of Metallurgy and  
Materials Science  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Dr. S. Sibener  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Arnold Green  
Quantum Surface Dynamics Branch  
Code 3817  
Naval Weapons Center  
China Lake, California 93555

Dr. A. Wold  
Department of Chemistry  
Brown University  
Providence, Rhode Island 02912

Dr. S. L. Bernasek  
Department of Chemistry  
Princeton University  
Princeton, New Jersey 08544

Dr. W. Kohn  
Department of Physics  
University of California, San Diego  
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Richard Colton  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Dan Pierce  
National Bureau of Standards  
Optical Physics Division  
Washington, D.C. 20234

Dr. R. Stanley Williams  
Department of Chemistry  
University of California  
Los Angeles, California 90024

Dr. R. P. Messmer  
Materials Characterization Lab.  
General Electric Company  
Schenectady, New York 22217

Dr. Robert Gomer  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Ronald Lee  
R301  
Naval Surface Weapons Center  
White Oak  
Silver Spring, Maryland 20910

Dr. Paul Schoen  
Code 6190  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. John T. Yates  
Department of Chemistry  
University of Pittsburgh  
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene  
Code 5230  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. L. Kesmodel  
Department of Physics  
Indiana University  
Bloomington, Indiana 47403

Dr. K. C. Janda  
University of Pittsburgh  
Chemistry Building  
Pittsburg, PA 15260

Dr. E. A. Irene  
Department of Chemistry  
University of North Carolina  
Chapel Hill, North Carolina 27514

Dr. Adam Heller  
Bell Laboratories  
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann  
Department of Chemistry  
University of Southampton  
Southampton SO9 5NH  
UNITED KINGDOM

Dr. H. Tachikawa  
Chemistry Department  
Jackson State University  
Jackson, Mississippi 39217

Dr. John W. Wilkins  
Cornell University  
Laboratory of Atomic and  
Solid State Physics  
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis  
Department of Physics  
University of California  
Irvine, California 92664

Dr. D. Ramaker  
Chemistry Department  
George Washington University  
Washington, D.C. 20052

Dr. J. C. Hemminger  
Chemistry Department  
University of California  
Irvine, California 92717

Dr. T. F. George  
Chemistry Department  
University of Rochester  
Rochester, New York 14627

Dr. G. Rubloff  
IBM  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. Horia Metiu  
Chemistry Department  
University of California  
Santa Barbara, California 93106

Dr. W. Goddard  
Department of Chemistry and Chemical  
Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. P. Hansma  
Department of Physics  
University of California  
Santa Barbara, California 93106

Dr. J. Baldeschwieler  
Department of Chemistry and  
Chemical Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. J. T. Keiser  
Department of Chemistry  
University of Richmond  
Richmond, Virginia 23173

Dr. R. W. Plummer  
Department of Physics  
University of Pennsylvania  
Philadelphia, Pennsylvania 19104

Dr. E. Yeager  
Department of Chemistry  
Case Western Reserve University  
Cleveland, Ohio 41106

Dr. N. Winograd  
Department of Chemistry  
Pennsylvania State University  
University Park, Pennsylvania 16802

Dr. Roald Hoffmann  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

Dr. A. Stecki  
Department of Electrical and  
Systems Engineering  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. G.H. Morrison  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853