

4

Technical Report 1277  
March 1989

## Ionospheric Variability

A.K. Paul

AD-A209 251

DTIC  
ELECTE  
JUN 21 1989  
S D & D

Approved for public release; distribution is unlimited.

89 6 20 245

# **NAVAL OCEAN SYSTEMS CENTER**

**San Diego, California 92152-5000**

---

**E. G. SCHWEIZER, CAPT, USN**  
Commander

**R. M. HILLYER**  
Technical Director

## **ADMINISTRATIVE INFORMATION**

This document was prepared by Code 542 of the Naval Ocean Systems Center. The funding for this document was provided by the Independent Research Program (IR).

Released by  
J.A. Ferguson, Head  
Ionospheric Branch

Under authority of  
J.H. Richter, Head  
Ocean and Atmospheric  
Sciences Division

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT  Approved for public release; distribution is unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) <b>NOSC Technical Report 1277</b>		7a. NAME OF MONITORING ORGANIZATION	
6a. NAME OF PERFORMING ORGANIZATION <b>Naval Ocean Systems Center</b>	6b. OFFICE SYMBOL (if applicable) <b>NOSC</b>	7b. ADDRESS (City, State and ZIP Code)	
6c. ADDRESS (City, State and ZIP Code)  <b>San Diego, CA 92152-5000</b>		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION <b>Office of Chief of Naval Research</b>	8b. OFFICE SYMBOL (if applicable)	10. SOURCE OF FUNDING NUMBERS	
8c. ADDRESS (City, State and ZIP Code)  <b>Independent Research Program (IR) OCNR-10P Arlington, VA 22217-5000</b>		PROGRAM ELEMENT NO.  <b>0601152N</b>	TASK NO.  <b>ZT43</b>
		AGENCY ACCESSION NO.  <b>RR00N00</b>	<b>DN306 212</b>
11. TITLE (include Security Classification)  <b>IONOSPHERIC VARIABILITY</b>			
12. PERSONAL AUTHOR(S) <b>A.K. Paul</b>			
13a. TYPE OF REPORT <b>Final</b>	13b. TIME COVERED <b>FROM Oct 1985 TO Oct 1988</b>	14. DATE OF REPORT (Year, Month, Day) <b>March 1989</b>	15. PAGE COUNT <b>15</b>
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
		F-region variability	
		F-region tilts	
		sporadic E-layer tilts	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) <p>This report summarizes the results of a study of short term variations of the ionosphere. The digital ionograms used in this analysis were recorded at Brighton, Colorado, many of them at a high rate of 12 ionograms per hour and faster. The results show that oscillations of the electron density distribution take place in the F-region with varying amplitudes and with periods of a fraction of an hour. The oscillations are coupled with varying tilts. Rapid changes, mainly of the electron density, are also observed at E-layer heights and many records indicate that the echoes come from tilted sporadic-E patches with relatively small dimensions.</p> <p><i>Key words: F Region, Sporadic E Layer, (1d) =</i></p>			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>	
22a. NAME OF RESPONSIBLE PERSON <b>A.K. Paul</b>		22b. TELEPHONE (include Area Code) <b>(619) 553-3074</b>	22c. OFFICE SYMBOL <b>Code 542</b>

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

DD FORM 1473, 84 JAN

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

# CONTENTS

<b>SUMMARY</b> .....	1
<b>INTRODUCTION</b> .....	1
<b>F-region Variations</b> .....	2
<b>Sporadic-E Variability</b> .....	6
<b>CONCLUSIONS</b> .....	8
<b>REFERENCES</b> .....	9

## FIGURES

- |    |   |   |
|----|---|---|
| 1. | Variations of the propagation parameter MUF(3000) for an active day (top) and a quiet day (bottom). . . . .   | 3 |
| 2. | Deviations from vertical propagation as indicated by the apparent positions of the reflection points. . . . .   | 4 |
| 3  | Temporal derivative of the MUF(3000). . . . .   | 5 |
| 4. | Top frequencies of sporadic E-layers embedded in the regular E-layer. . . . .   | 7 |
| 5. | Apparent average positions of sporadic E-layers over a 24-hour period indicating strong tilts. (x observed in the extraordinary propagation mode, o observed in the ordinary mode). . . . . | 8 |



Accession For

NTIS CRA&I ☒

DDC TAB ☐

Unannounced ☐

Justification

By

Date Recd

Contract, Order

Contract No.

Order No.

A-1

## SUMMARY

This report summarizes the results of a study of short-term variations of the ionosphere. The digital ionograms used in this analysis were recorded at Brighton, Colorado, many at a high rate of 12 ionograms per hour and faster. The results show that oscillations of the electron density distribution take place in the F-region with varying amplitudes and with periods of a fraction of an hour. The oscillations are coupled with varying tilts. Rapid changes, mainly of the electron density, are also observed at E-layer heights and many records indicate the echoes come from tilted sporadic-E patches with relatively small dimensions.

## INTRODUCTION

During the last sunspot maximum, 1980 to 81, a digital ionosonde developed by the National Oceanic and Atmospheric Administration (NOAA) (Grubb, 1979) and cosponsored by the National Science Foundation (NSF) was operated at Brighton, Colorado. The data recorded are of importance mainly for two reasons:

- (a) The design and the concept of the system provide data of high accuracy
- (b) Many of the ionograms were taken in rapid sequences of 12 ionograms per hour and faster.

This combination of high-accuracy and high-temporal resolution offers the opportunity to study details of ionospheric variations. At the time the ionospheric observation program in NOAA was terminated in 1981, very few data had been analyzed. When funds became available through the Naval Ocean Systems Center (NOSC) Independent Research (IR) program to conduct a systematic analysis of the data, the original records were borrowed from the World Data Center in Boulder, Colorado.



The ionosonde had the capability to record in digital form the travel time, amplitude, and phase of each echo reflected from the ionosphere at four receiving antennas. The receiving antennas were short dipoles located at the corners of a square with a diagonal length of 100 meters. In a typical mode of operation the two-channel receiver recorded the signals at two diagonally opposite antennas simultaneously and was switched to the other diagonal antenna pair in time to receive the echoes of the next pulse transmitted. The radio frequency was increased by 8 kHz and two more pulses were transmitted with the antenna switching sequence reversed. This way the change of the radio phase with frequency, which permits more accurate estimates of the echo travel time (virtual height), could be separated from the phase change with time, which gives the Doppler frequency or Doppler velocity. A phase comparison at the four antennas can be used to determine the angle of arrival of an echo, or in combination with the virtual height, to determine the apparent position of its reflection point.

## F-REGION VARIATIONS

The important and surprising findings of this data analysis are the ever present oscillations of the F-layer coupled with strong tilts of the reflecting surface. The oscillations are clearly visible in the temporal changes of a variety of standard parameters, such as the critical frequency, the virtual height for a constant frequency, the height of maximum of the layer, etc. We found the most accurate and probably most representative indicator for F-region variations is the maximum usable frequency for a 3000 km propagation path, the MUF(3000), a parameter which can easily be derived from the vertical ionograms (Paul, 1988a). Figure 1 shows two examples of the daytime variations of this parameter. While a typical period on February 16 (bottom) was approximately 20 minutes, some longer periods with larger amplitudes were present on January 15 (top) in addition to the shorter oscillations. No explanation for the different magnitude of the variations could be found, but there is definitely no correlation with magnetic activity. There is convincing evidence (Paul, 1988b) that the variations observed are caused by acoustic gravity waves traveling through the F-region which means the temporal variations are coupled with spatial variations. The only information about spatial variations in the horizontal dimensions which can be obtained from the data of this monostatic observation technique is the deviation from vertical propagation. In figure 2 we compare the apparent positions of the reflection point for the two data sets which were used in figure 1. The apparent positions shown here are for the point where the MUF(3000) transmission curve is tangent to the ordinary echo trace which corresponds to a height approximately halfway between the lower edge of the F-layer and the height of the electron density maximum. We see the points are spread over a larger area for the data of January 15 (top) than for those of February 16 (bottom). The median zenith distance for the data in the top of figure 2 is 2.2 degrees and 1.5 degrees for the data in the bottom half of this figure. We can compute an estimate of the temporal derivative of the MUF(3000) simply by dividing the differences of consecutive data by the 3-minute time interval and obtain curves as shown in figure 3. The median magnitude of this quantity is approximately 6.9 MHz/hour for January 15 and 4.2 MHz/hour for February 16. The ratio of those two numbers is very close to the ratio of the median zenith distances for the two days and is quite different from the ratio of the median deviations from the mean MUF(3000). This comparison indicates that it is the temporal derivative of the MUF(3000) and not the quantity itself which permits us to obtain an estimate of the deviation from vertical propagation and therefore of the magnitude of F-region tilts. This relation was found to be true consistently with all the data sets available. Application of this relation for tilt estimates based on temporal variations requires that the sampling rate be sufficiently high (10 ionograms per hour or more) to avoid aliasing.



Usually the zenith and azimuth angles change slowly with frequency (height) within an ionogram. Occasionally such changes indicate a periodic behavior over a fraction of a period within the frequency (height) range of observation and sometimes an estimate of a vertical wavelength can be obtained.

In many cases the zenith distance is only a lower limit for the tilt angles in the vicinity of the reflection area. For example, the expansion phase of an acoustic-gravity wave at the height of the maximum causes a local reduction of the maximum electron density as can be seen in the variation of the critical frequency foF2. This



**Figure 1.** Variations of the propagation parameter MUF(3000) for an active day (top) and a quiet day (bottom).





**Figure 2.** Deviations from vertical propagation as indicated by the apparent positions of the reflection points.





Figure 3. Temporal derivative of the MUF(3000).


means for a short period of time a horizontal gradient exists at the height of the maximum and the ray direction of signals reflected may become almost horizontal for frequencies close to the penetration frequency. The integrated effect of this kind of structure results in an increase of the zenith angle with increasing frequency which indicates a strong tilt, but still underestimates the actual tilt angle in the vicinity of the reflection point. While the variations of the critical frequency and of the MUF (3000) have some direct effect in the conventional sense on HF-propagation, local tilts, depending on their orientation relative to the propagation direction, can have a much stronger influence (e.g., overlong propagation distances).

The temporal variations of the MUF(3000) are a combination of a vertical motion of the F-layer and a change of the electron density. Both variations take place in some periodic fashion but are not in phase (Paul, 1988b). This phase lag indicates that a periodic deformation of the layer takes place and standard parameters like critical frequency, height of maximum, and half-thickness of the layer which are obtained by extrapolation are sometimes ill defined which also means isolated observations of such parameters may not be representative for a longer period of time.

## **SPORADIC-E VARIABILITY**

The significant results here are the short life time (observation time) and the large tilt angles of this layer (Paul, 1988c). Figure 4 shows ftEs, the highest frequency of echoes reflected from E-region heights for a 2-hour period during daytime. Ionograms were recorded continuously at a rate of 77 ionograms per hour. Most of the data simply show the regular E-layer critical frequency, foE. There are however several instances where ftEs suddenly increases to twice the value of foE or higher for the short duration of 1- or 2-sampling intervals. Only a few hours of data were recorded at such a high sampling rate, but ionograms recorded at a somewhat slower rate show a similar behavior. Of 4019 ionograms recorded in 3-minute intervals, 1013 showed sporadic-E traces and 15 percent of those ionograms were isolated events where no sporadic-E was recorded in the preceding and following ionogram. Approximately 32 percent of the sporadic-E observations formed short sequences of 5 or less consecutive ionograms. No explanation for those rapid changes could be found and it is not clear whether the formation and decay of a layer can take place in such a short period or small patches of ionization move rapidly through the observation field. On the other hand a few cases were found in this data set of 20 ionogram per hour recording rate where sporadic-E was observed in consecutive ionograms for several hours. While these long sequences showed continuity in time in terms of existence of sporadic-E, there were often significant changes of the apparent position from one ionogram to the next one. Those observations are likely to indicate that the proper condition for the formation of sporadic-E existed for a longer period of time and are less likely to prove that an individual layer lasted that long.

The existence of small patches of ionization is supported by the frequent observation of layers with large tilt angles, which for physical reasons can only exist over a limited height range. A total of 1911 sporadic-E layers were found in the 1013 ionograms mentioned above. In 999 cases the average zenith angle of the echoes was 10 degrees or less. In 409 cases the deviation from vertical propagation was between 10 and 20 degrees and more than 20 degrees for the remaining 503 cases. We point out that contrary to F-layer reflections, there is little bending of a ray reflected from



**Figure 4.** Top frequencies of sporadic E-layers embedded in the regular E-layer.

a sporadic E-layer, since the layer is so thin. For this reason the angle of arrival is a direct measure of the tilt angle. The presence of two or more Es-layers in one ionogram also implies a relative small size of each layer or a corrugated structure with a scale size not to exceed a few tens of kilometers.

No preferred orientation of the tilts could be found. The apparent reflection points showed a rather uniform azimuthal distribution. This can be seen in figure 5 where the average position of all Es-layers over a 24-hour period are plotted.

When large zenith distances are observed the so-called virtual heights have to be interpreted as ranges and have to be corrected correspondingly. This correction applied to the data sample described above lowered the median virtual height by 4 to 107 kilometers and also narrowed the height distribution of sporadic-E.




Figure 5. Apparent average positions of sporadic E-layers over a 24-hour period indicating strong tilts (x observed in the extraordinary propagation mode, o observed in the ordinary mode).

## CONCLUSIONS

Typical time scales for the F-layer are on the order of a fraction of an hour. For this reason a minimum sampling rate of 12 ionograms per hour is required in order to obtain a correct description of the temporal variation of the F-layer. Based on the rate of change of well defined standard parameters in combination with reasonable assumption about the velocity of acoustic gravity waves, we conclude that local tilts can exist at any time of the day (or night) which are of the same magnitude as those observed during sunrise or sunset (Paul, 1985). Deviations from vertical propagation of 10 degrees or more are frequently found, at least for frequencies close to the penetration frequency. The area over which such local tilts may exist is likely much smaller than the sunrise or sunset zone. While the orientation of the tilts during sunrise and sunset are predictable, this orientation is not true for tilts coupled with gravity waves since their velocities (magnitude and direction) cannot be determined from monostatic observations.

Typical time scales for sporadic-E can be much shorter than those for the F-region, but are of a different nature. While most of the F-region variations are periodic, no periodic phenomena were found in sporadic E-layers. Life times or observation times of sporadic-E cover a larger range than the F-layer periodicities, from less than 1 minute to several hours. Tilts of sporadic E-layers can be larger than F-layer tilts and have similar effects on HF propagation.

## REFERENCES

- Paul, Adolf K., 1985. "F-region tilts and ionogram analysis." *Radio Science* 20, 959-971.
- Paul, Adolf K., 1986. "Limitations and possible improvements of ionospheric models for radio propagation. Effects of the sporadic E-layer." *Radio Science* 21, 304, 308.
- Paul, Adolf K., 1988a. *The MUF(3000) as an indicator for F-region variations*. NOSC Technical Report.
- Paul, Adolf K., 1988b. "Medium scale structure of the F-region." Accepted for publication in *Radio Science*.
- Paul, Adolf K. 1988c. "On the variability of sporadic-E." submitted to *Radio Science*.